ON THE FUNCTION
$$\sum_{k=-\infty}^{\infty} \exp(-\frac{k^2}{s^2})$$

Ioan STURZU^{*}

Abstract: One is studying numerically and analitically some properties of the function $\mathbf{y}_0(s) = \sum_{k=-\infty}^{\infty} \exp(-\frac{k^2}{s^2})$

Key words: Series of functions; Poisson summation formula

The function $\mathbf{y}_0(s) = \sum_{k=-\infty}^{\infty} \exp(-\frac{k^2}{s^2})$ is related to the Poisson integral:

$$I(s) = \int_{-\infty}^{\infty} \exp(-\frac{x^2}{s^2}) dx = \sqrt{p} \cdot s$$

by the bonding relation: $I(s) < \mathbf{y}_0(s) < I(s) + 2$ which can be easy proved using the monotony of the function $\exp(-\frac{x^2}{s^2})$.

However, there are situations [1] when a deeper study of the properties of this function is needed. We shall start by studying the behaviour of y(s) in the limiting cases $s \to +0$ and $s \to +\infty$. If $s \to +0$ all the terms in the series go to θ excepting the term which corresponds to k=0. So $y(s) \to 1$ when $s \to +0$. When $s \to +\infty$ one can write y(s) as a Riemann series for I(s):

$$\frac{1}{s} \cdot \mathbf{y} \ (s) = \sum_{k=-\infty}^{\infty} \exp(-\frac{k^2}{s^2}) \cdot \frac{\Delta k}{s} \to \int_{-\infty}^{\infty} \exp(-x^2) dx = \sqrt{\mathbf{p}}$$

so $\mathbf{y}_0(s)$ behaves like I(s) when $s \to +\infty$. Let us write $\mathbf{y}_0(s)$ in the form: $\mathbf{y}_0(s) = \sqrt{\mathbf{p}} \cdot s + 1 - e(s)$; than, e(s) has the following asymptotic behaviour:

$$\frac{e(s)}{\sqrt{\mathbf{p}} \cdot s} \to 1, \text{ when } s \to +0 \text{ ; } e(s) \to 1, \text{ when } s \to +\infty.$$

^{*} Physics Department., *Transilvania* University of Bra°ov.

A reasonable numerical calculus of $y_0(s)$ gives for e(s) the plot from Fig. 1.



Fig.1. Numerical calculus for e(s)

One can easy see that e(s) behaves like $\sqrt{\mathbf{p}} \cdot s$ for $s \in [0;0.4]$ and is equal to 1 for $s \in [0.8; +\infty]$; so $\mathbf{y}_0(s) \approx 1$ for $s \in [0;0.4]$ and $\mathbf{y}_0(s) \approx \sqrt{\mathbf{p}} \cdot s$ for $s \in [0.8; +\infty]$. For the domain [0.4;0.8] one can obtain a Boltzmann-type fitting:

$$efit(s) = 1.00582 - \frac{0.71664}{1 + \exp(\frac{s - 0.36712}{0.10290})}$$

which is represented in Fig.2.

Let the displaced function of $y_a(s)$, $a \in [0;0.5]$:

$$\mathbf{y}_{a}(s) = \sum_{k=-\infty}^{\infty} \exp(-\frac{(k+a)^{2}}{s^{2}})$$

The result of a numerical calculus for $y_{1/2}(s)$ is represented in Fig.3, while the difference $y_0(s) - y_{1/2}(s)$ was numerically interpolated, and yield the formula:

$$diffit(s) = \frac{1}{1 + \exp[\text{sgn}(s - 0.45) \cdot \left| \frac{s - 0.45}{0.088} \right|^{1.29}]}$$

Results are consistent with those obtained from Poisson summation formula [2],[3]:

$$\sum_{k=-\infty}^{\infty} F(\frac{k}{s}) = s \cdot \sum_{k=-\infty}^{\infty} G(s \cdot k)$$

where G is the Fourier Transform of F.

For $F(x) = \exp[-(x+a)^2]$, $G(y) = \sqrt{p} \cdot \exp(-p^2 \cdot y^2 - i \cdot a \cdot y)$, so one has the general relation:

$$\mathbf{y}_{a}(s) = \sqrt{\mathbf{p}} \cdot s \sum_{k=-\infty}^{\infty} \exp(-\mathbf{p}^{2} \cdot s^{2} \cdot k^{2} - i \cdot a \cdot s \cdot k)$$

which goes to:



Fig.2. Numerical interpolation for e(s)

On the function ...



Fig.3. Numerical calculus for $\mathbf{y}_{1/2}(s)$

References

- 1. Sturzu, I.: Operational Quantum Physics without Projection Postulate (to appear)
- 2. Lighthill, M.J.: Introduction to Fourier Analysis and Generalised Functions, Cambridge University Press (1958)
- 3. Rosenfelder, R.: private communication

Asupra functiei
$$\sum_{k=-\infty}^{\infty} \exp(-\frac{k^2}{s^2})$$

Rezumat: Sunt studiate numeric si analitic unele proprietăti ale functiei

$$\mathbf{y}(s) = \sum_{k=-\infty}^{\infty} \exp(-\frac{k^2}{s^2})$$

Cuvinte cheie: serii de functii, formula de sumare Poisson

Recenzent: Prof. Univ. Dr. Spiridon DUMITRU