Gauge ^P representations for quantum-dynami
al problems: Removal of boundary terms

P. Deuar[∗] and P. D. Drummond[†]

Department of Physi
s, University of Queensland, QLD 4072, Brisbane, Australia

(Dated: 5th November 2018)

P representation techniques, which have been very successful in quantum optics and in other fields, are also useful for general bosonic quantum dynamical many-body calculations such as Bose-Einstein condensation. We introduce a representation called the gauge P representation which greatly widens the range of tractable problems. Our treatment results in an infinite set of possible time-evolution equations, depending on arbitrary gauge fun
tions that an be optimized for a given quantum system. In some ases, previous methods an give erroneous results, due to the usual assumption of vanishing boundary conditions being invalid for those particular systems. Solutions are given to this boundary-term problem for all the cases where it is known to occur: two-photon absorption and the single-mode laser. We also provide some brief guidelines on how to apply the stochastic gauge method to other systems in general, quantify the freedom of choice in the resulting equations, and make a omparison to related re
ent developments.

PACS numbers: 02.70.Rr, 05.10.Gg, 42.50.-p, 03.75.Fi

I. INTRODUCTION

One of the most difficult problems in theoretical physi
s is also on
eptually the simplest. How does one al
ulate the dynami
al time evolution or even the ground state of an intera
ting many-body quantum system? In essence, this is a natural part of practically any omparison of quantum theory with experiment. The difficulty is that the Hilbert space of all but the most trivial control come to the enormously a nite that the second the enormous omputer is needed to to solve problems that an easily become nearly infinite in dimensionality, if treated using an orthogonal basis expansion.

In this paper, we formally introdu
e and give examples of te
hniques for treating general bosoni many-body quantum systems, whi
h we all gauge P representations. These are an extension of the phase-spa
e method alled the positive-P representation $[1]$, and have been recently used in the context of interacting Bose gases $[2, 3]$. The advantages of the new technique are the following.

(1) The elimination of ertain types of mathemati
al terms known as boundary-term orre
tions, whi
h have aused problems in the positive-P representation for over a decade $[4-6]$. This is the main focus of the present paper.

(2) Greatly redu
ed sampling error in omputations. Gauge P representations have been used re
ently to reduce the sampling error in Kerr oscillator simulations [2].

(3) The extension of allowable problems to 'imaginarytime' canonical ensemble calculations. These problems will be treated elsewhere.

Related extensions to the positive-P representation

although restri
ted to the s
alar intera
ting Bose gas problem introduced also been introduced also been introduced also been introduced also been introduced as a se pro
edures have been introdu
ed by Carusotto, Castin, and Dalibard $[7, 8]$, and by Plimak, Olsen, and Collett [9]. These methods implicitly assume the absence of boundary term corrections. This paper unifies and substantially generalizes all these re
ent advan
es. It also shows how the gauge method an be used to solve the long-standing problem of boundary-term corrections in the positive P representation. Comparisons to the other methods are given in an Appendix.

Owing to the work of Wilson $[10]$, and many others $[11]$, we know that large Hilbert space problems can often be treated using sto
hasti or Monte Carlo te
hniques for the ground-state, particle masses, and finite-temperature orrelations. This is the basis for mu
h work in omputational quantum statisti
al me
hani
s, and in QCD as well. However, Wilson's and other related methods are restricted to static or 'imaginary-time' calculations, rather than quantum-dynami
al problems.

Methods like these that use orthogonal basis sets have not proven useful for quantum dynami
s; owing to the notorious phase problem that occurs when trying to sum over families of paths in real-time Feynman path integrals. For this reason, the many-body quantum timeevolution problem is often regarded as inherently insoluble due to its exponential omplexity. In fa
t, it was this very problem that motivated the original proposal of Feynman $[12]$ to develop quantum computers. In these (usually on
eptual) devi
es, the mathemati
al problem is solved by a physi
al system onsisting of evolving `qubits' or two-state physi
al devi
es. Fortunately, this method of doing calculations is not the only one, since no large enough quantum computer exists at present [13].

Histori
ally, an alternative route is the use of quasiprobability representations of the quantum state, whi
h either impli
itly or expli
itly make use of a non orthogonal basis. The term quasi-probability is used be
ause

^{*}Electronic address: deuar@physics.uq.edu.au

[†]Electronic address: drummond@physics.uq.edu.au; URL:www. physi
s.uq.edu.au/BEC

there can be no *exact* mapping of all quantum states to a classical phase space with a positive distribution [14] that also preserves all the marginal probabilities. These methods include the Wigner [15] (W), Glauber-Sudarshan (P) [16, 17], and Husimi (Q) [18, 19] representations. The classical phase-space representations can be classified acording to the operator ordering that sto
hasti moments orrespond to: the W is symmetri
ally ordered, the Q is anti-normally ordered, while the P representation is normally ordered. Apart from numerous laser physi
s and quantum optics calculations, these methods have also been used to some extent in quantum statisti
al me
hanics: for example, the theory of BEC phase fluctuations $[20]$.

None of these methods result in a stochastic time evolution with a positive propagator when there are nonlinearities. To achieve this, a better approach is to use a nonlassi
al phase spa
e of higher dimension. A omplex higher-dimensional `R representation' was proposed in Glauber's seminal paper on oherent state expansions [16]. The first probabilistic method of this type was the positive-P representation $[1]$ (+P), which has proved apable of performing sto
hasti time-domain quantum calculations in some many-body quantum systems [21]. This uses a basis of oherent states that are not orthogonal, thus allowing freedom of choice in the construction of the representation. The positive-P representation of a quantum state is therefore the most versatile out of a large group of quasi-probability distributions developed to aid quantum mechanical calculations. It has been successfully applied to mesoscopic systems such as quantum solitons $[21-23]$ and the theory of evaporative cooling [24], which correctly reproduces the formation of a BEC — as observed in experiment $[25-27]$.

Quasi-probability distributions of this type are omputationally superior to dire
t density matrix methods, which are susceptible to computational complexity blowup for large Hilbert spa
es. Provided ertain boundary terms vanish, the usual pro
edure is to generate a Fokker-Plan
k equation (whi
h will vary depending on the distribution hosen) from the master equation, and then to onvert this to a set of sto
hasti Langevin equations. For some simple cases, it may even be possible to arrive at appealing results directly from the Fokker-Planck equation (FPE). The resulting stochastic equations can be thought of just as quantum mechanics written in different variables. They have two main advantages over orthogonal basis-state methods, as follows.

First, the whole quantum dynamics can be written exactly in terms of a small number of stochastic equations. In a one-mode case, there is just one complex variable for P and Q and W, and two complex variables for $+P$. Although a simulation requires us to average over many realizations of the stochastic process, this is often more practical than solving the infinite set of deterministic equations required to solve directly for all the elements of a density matrix. Such an infinite set may be truncated, but this is only a good approximation for a system with

few parti
les, and no more than a few modes.

Second, for a many-mode problem the Hilbert space dimension is $N = n^M$ for the case of *n* particles distributed over M modes. This gives exponential growth as a fun
tion of the number of modes. However, the number of quasi-probability dynami
al equations grows only linearly with the number of modes, rather than exponentially in the case of direct methods. Other stochastic methods, known as quantum-trajectory methods, can be used to reduce the N^2 dimensionality of an $N \times N$ density matrix problem to that of the N -dimensional underlying Hilbert space — but this is clearly insufficient to solve the omplexity problem inherent in the exponential growth of the Hilbert spa
e dimension.

There are, however, some caveats when using these distributions. In particular, the vanishing of boundary terms is an important fundamental issue with quasiprobability distributions, and it is this issue that we fo
us on mostly in this paper. To get an overall picture, consider that once we have a time-evolution problem there are five typical requirements that are encountered in deriving sto
hasti equations for quasi-probability representations of many-body systems. These requirements occur in closed (unitary evolution) systems, in open systems (in general des
ribed by a master equation), or even using a distribution to solve for the canonical ensemble in imaginary time. As su
h, these requirements are generi to the use of sto
hasti equations with operator representations:

(1) Positive distribution. A well-behaved positive distributions for all quantum states, including especially the hosen initial ondition, is essential for a general algorithm. For example, a number state has a highly singular P distribution, and a W distribution that is negative in some regions of phase space [28], making either distribution impossible to interpret probabilisti
ally for these states. The R distribution is inherently omplex. Su
h problems do not occur for the Q or $+P$ representation these are positive, and well-behaved for all quantum states [1].

(2) Ultraviolet onvergen
e. While normally-ordered representations are well behaved at large momentum, non-normally-ordered representations of quantum fields — such as the Q or W representations — typically face the problem of ultraviolet divergen
e in the limit of large momentum cutoff [24]. This means that almost any observable quantity will involve the simulation of a (nearly) infinitely noisy classical field, leading to diverging standard deviations in two or more spa
e dimensions, even for linear systems. This rules out the Q and W distributions for quantum field simulations in higher than onedimensional environments.

(3) Second-order derivatives. Only FPEs with second or infinite-order derivatives can be translated into stochastic equations [29]. Normally-ordered methods such as the P and $+P$ representations can handle most commonly occurring nonlinearities and two-body interactions, with only se
ond-order derivatives. Non-normally ordered representations of quantum fields often lead to third- or higher-order partial derivatives in the Fokker-Planck equation with no stochastic equivalent. For example, the Wigner representation gives su
h problems for almost any nonlinear term in the master equation.

 (4) *Positive-definite diffusion*. A Fokker-Planck equation must have positive-definite diffusion, to allow simulation with stochastic processes [29]. When the master equation has nonlinear terms, this does not occur with any of the classical representations. However, the $+P$ representation is guaranteed to always produ
e positivedefinite diffusion [1], provided no higher derivative terms occur.

(5) Vanishing boundary terms. In the derivation of the Fokker-Plan
k equations, it is assumed that ertain boundary terms arising in partial integration an be negle
ted. This is not always the ase. Boundary terms due to power-law tails can occur when there are moving singularities that can escape to infinity in finite time. In the $+P$ method, such trajectories may cause systematic errors in stochastic averages [6], especially for nonintegrable dynami
al systems. These problems are exponentially suppressed when linear damping rates are in reased, but an be large at low damping.

The $+P$ method is often the representation of choice, because it satisfies conditions one to four. Gauge representations (G) combined with stochastic methods to be treated in this paper, share these advantages with the $+P$ representation. However, they can also satisfy the fifth requirement — for an appropriate gauge choice — hence allowing all of the mathemati
al problems in simulating time evolution to be treated. For this reason, the present paper will focus on solving boundary-term issues encountered with the $+P$ representation for certain nonlinear master equations. The overall picture is summarized in Table I, as applied to the two-boson anonlinear absorber cases treated here in Sec. IV:

We emphasize that the particular examples treated here have a small parti
le number and extremely low (or zero) linear damping. As su
h, they are soluble using other techniques, which allows us to test the accuracy of gauge te
hniques. Our purpose is to demonstrate the success of the stochastic gauge method in simple cases where boundary terms arise within the $+P$ representation. In this way, we can understand more complex situations where no exa
t result is known.

We will first derive and describe the stochastic gauge method in Secs. II and III, and subsequently work through two examples: First, solving the boundary-value problem for the driven one- and two-photon absorber in Sec. IV. Second, in Sec. V we will consider the one-mode laser at extremely low power, whi
h exhibits boundary term errors when very non-optimal starting conditions are used. This example will show that gauge methods an also be used to remove errors from this system, but some judgment must be employed to avoid hoosing a pathologi
al initial distribution. In the Appendix, we compare the methods derived here with recent related extensions of the positive-P representation by Carusotto

and co-workers $[7, 8]$, Plimak *et al.* [9], and Deuar and Drummond [2].

Finally, we point out a sixth requirement of containing the growth of sampling error: the averages calculated from the sto
hasti Langevin equations orrespond to quantum me
hani
al expe
tation values only in the limit of infinitely many trajectories. Provided boundary terms do not occur, the averages will approach the correct values — within an acceptable sampling error — for sufficiently many trajectories. If this number should inrease rapidly with time, the simulation will only be of use for a limited period $|2|$.

The problem of growing sampling error can occur even when there are no boundary terms, and may be regarded as the ultimate frontier in representation theory, just as similar issues dominate the theory of classical chaos. This is less of a fundamental issue, sin
e the sampling error an always be estimated and ontrolled by in
reasing the number of trajectories. This is simply a matter of moving to a lustered, parallel omputational model, or repeating the al
ulation many times. Nevertheless, it is of great practical significance. The sampling error problem requires areful gauge optimization, and remains an open area for investigation. An intelligent hoi
e of gauge an often vastly outweigh a brute for
e omputational approa
h, in terms of sampling error.

II. **GAUGE OPERATOR REPRESENTATIONS**

In gauge representations, the density matrix to be computed is expanded in terms of a oherent state basis. For definiteness, we shall focus on the coherent states of the harmoni os
illator, whi
h are useful in expanding Bose fields; but other choices are clearly possible. The expansion kernel is more general than that used in the positive-P representation. In order to define the notation, we start by introdu
ing a set of boson annihilation and creation operators \hat{a}_i , \hat{a}_i^{\dagger} . The operator $\hat{n}_i = \hat{a}_i^{\dagger} \hat{a}_i$ is therefore the boson number operator for the ith mode or site. Boson commutation relations of $[\hat{a}_i, \hat{a}_j^{\dagger}] = \delta_{ij}$ hold for the annihilation and creation operators.

A. Coherent states

If $\alpha = (\alpha_1, \ldots, \alpha_M)$ is a complex M-dimensional vector with $\alpha_i = x_i + iy_i$, and $\hat{\mathbf{a}} = (\hat{a}_1, \dots, \hat{a}_M)$ is an Mdimensional ve
tor of annihilation operators, then the Bargmann coherent state $\|\alpha\rangle$ is defined by

$$
\|\boldsymbol{\alpha}\rangle = \exp\left[\boldsymbol{\alpha}\cdot\widehat{\mathbf{a}}^{\dagger}\right]\left|0\right\rangle = \exp\left[|\boldsymbol{\alpha}|^{2}/2\right]|\boldsymbol{\alpha}\rangle , \qquad (1)
$$

where $|\alpha\rangle$ is the usual normalized coherent state which is a simultaneous eigenstate of all the annihilation operators. The inner product of two Bargmann coherent states is

$$
\langle \beta^* \parallel \alpha \rangle = \exp\left[\alpha \cdot \beta\right] \ . \tag{2}
$$

Table I: Comparison of phase-space representations as applied to stochastic treatments of a one- and two-boson nonlinear absorber.

Method	Form of Distribution	UV converges	Order of derivatives	Non-negative diffusion	Stochastic simulations	Boundary term removal	Simulated correctly
W	Real	No	4	Sometimes	No		
Q	Positive	No	4	$_{\rm Yes}$	No		
$_{\rm R}$	Complex	$_{\rm Yes}$	$\overline{2}$		No		
Ρ	Singular	$_{\rm Yes}$	$\overline{2}$	No	No		
$+P$	Positive	$_{\rm Yes}$	$\overline{2}$	$_{\rm Yes}$	Yes	No	Sometimes
G	Positive	$_{\rm Yes}$	$\overline{2}$	$\rm Yes$	$_{\rm Yes}$	Yes	Yes

It is important to notice here that $\|\alpha\rangle$ is an analytic function of the complex vector α . The following identities therefore follow immediately:

$$
\widehat{a}_i \|\alpha\rangle = \alpha_i \|\alpha\rangle \n\widehat{a}_i^{\dagger} \|\alpha\rangle = \frac{\partial}{\partial \alpha_i} \|\alpha\rangle .
$$
\n(3)

Since $\|\alpha\rangle$ is an analytic function, the notation $\partial/\partial\alpha_i$ is interpreted here as an analytic derivative, which can be evaluated in either the real or imaginary directions,

$$
\frac{\partial}{\partial \alpha_i} \|\alpha\rangle = \frac{\partial}{\partial x_i} \|\alpha\rangle = -i \frac{\partial}{\partial y_i} \|\alpha\rangle \quad . \tag{4}
$$

Sin
e the oherent states are an overomplete basis set, any operator can be expanded in more than one way using oherent states. For example, the simplest resolution of the identity operator is

$$
\widehat{I} = \frac{1}{\pi^M} \int |\alpha\rangle \langle \alpha| d^{2M} \alpha.
$$
 (5)

Thus, introducing a second M-dimensional vector β , we can expand any operator \widehat{O} directly as

$$
\hat{O} = \frac{1}{\pi^{2M}} \int \int |\alpha\rangle \langle \alpha| \hat{O} |\beta^* \rangle \langle \beta^* | d^{2M} \alpha d^{2M} \beta
$$

$$
= \int \int O(\alpha, \beta) |\alpha\rangle \langle \beta^* | d^{2M} \alpha d^{2M} \beta . \qquad (6)
$$

Here, we have introdu
ed

$$
O(\boldsymbol{\alpha}, \boldsymbol{\beta}) = \frac{1}{\pi^{2M}} \langle \boldsymbol{\alpha} | \widehat{O} | \boldsymbol{\beta}^* \rangle . \tag{7}
$$

B. P representations

The possibility of expanding any operator in terms of oherent states leads to the idea that su
h an expansion can be used to calculate observable properties of a quantum density matrix $\hat{\rho}$. Historically, this was first proposed by Glauber and Sudarshan $[16, 17]$, who suggested a diagonal expansion of the form

$$
\widehat{\rho} = \int P(\boldsymbol{\alpha}) \, |\boldsymbol{\alpha}\rangle \, \langle \boldsymbol{\alpha} | \, d^{2M} \boldsymbol{\alpha} \; . \tag{8}
$$

Unlike the direct expansion given above, this has no offdiagonal elements. Surprisingly, expansions of this type always exist, as long as the function $P(\alpha)$ is defined to allow highly singular generalized fun
tions and nonpositive distributions [28].

As these do not have a sto
hasti interpretation, the positive-P representation was introduced [1], which is defined as

$$
\widehat{\rho} = \int P^{(+)}(\alpha, \beta) \frac{|\alpha\rangle \langle \beta^*|}{\langle \beta^*| \alpha \rangle} d^{2M} \alpha d^{2M} \beta \tag{9}
$$

for an M -mode system.

It is always possible to obtain an explicitly positivedefinite distribution of this type $[1]$, with the definition

$$
P^{(+)}(\alpha, \beta) = \frac{1}{(4\pi^2)^M} \exp\left[-\left|\frac{\alpha - \beta^*}{2}\right|^2\right] \times \left(\frac{\alpha + \beta^*}{2}\right|\hat{\rho}\left|\frac{\alpha + \beta^*}{2}\right) . \quad (10)
$$

This form always exists, as do an infinite class of equivalent positive distributions. Even simpler ways to onstru
t the positive-P representation are available in some ases. For example, if the Glauber-Sudarshan representation exists and is positive, then one can simply construct

$$
P^{(+)}(\alpha,\beta) = P(\alpha)\delta^{2M}(\alpha - \beta^*)
$$
 (11)

The stochastic time evolution of the positive-P distribution does not generally preserve the above ompa
t forms, and may allow less ompa
t positive solutions instead. However, to obtain a time-evolution equation, it is ne
essary to use partial integration, with the assumption that boundary terms at infinity can be neglected. It is these less compact solutions, occurring during time evolution with a nonlinear Fokker-Plan
k equation, that lead to power-law tails in the distribution $-$ and hence boundary-term problems aused by the violation of the assumption that these terms vanish.

C. Gauge representations

A te
hnique for onstru
ting an even more general positive distribution is to introdu
e a quantum omplex amplitude Ω , which can be used to absorb the quantum phase fa
tor. This leads to the result that any Hermitian density matrix $\hat{\rho}$ can be expanded in an over-complete basis $\widehat{\Lambda}(\overrightarrow{\alpha})$, where $\overrightarrow{\alpha} = (\Omega, \alpha, \beta)$, and

$$
\widehat{\Lambda}(\overrightarrow{\alpha}) = \Omega \frac{\|\alpha\rangle \langle \beta^*\|}{\langle \beta^*\| \alpha \rangle} \n= \Omega \|\alpha\rangle \langle \beta^*\| \exp[-\alpha \cdot \beta] .
$$
\n(12)

We define the gauge representation $G(\overrightarrow{\alpha})$ as a real, positive function that satisfies the following equation:

$$
\widehat{\rho} = \int G(\vec{\alpha}) \left[\widehat{\Lambda}(\vec{\alpha}) \right] d^{4M+2} \vec{\alpha}
$$

$$
= \frac{1}{2} \int G(\vec{\alpha}) \left[\widehat{\Lambda}(\vec{\alpha}) + \text{H.c.} \right] d^{4M+2} \vec{\alpha}. \quad (13)
$$

The last line above follows from the fact that $\hat{\rho}$ is a Hermitian density matrix and $G(\overrightarrow{\alpha})$ is real. Here, H.c. is used as an abbreviation for Hermitian onjugate. The use of a omplex weight in the above gauge representation is similar to related methods introdu
ed re
ently for interacting Bose gases $[7, 8]$, except that we multiply the weight by a normalized (positive-P) projector, in order to simplify the resulting algebra.

As an existen
e theorem that shows that this representation always exists, onsider the omplex solution

$$
P_0(\boldsymbol{\alpha}, \boldsymbol{\beta}) = \frac{1}{\pi^{2M}} \langle \boldsymbol{\alpha} | \hat{\rho} | \boldsymbol{\beta}^* \rangle \langle \boldsymbol{\beta}^* | \boldsymbol{\alpha} \rangle \tag{14}
$$

obtained from Eq. (7), with a phase $\theta = \arg(P_0)$, and simply define

$$
G(\vec{\alpha}) = |P_0(\boldsymbol{\alpha}, \boldsymbol{\beta})| \delta^2 (\Omega - \exp[i\theta(\boldsymbol{\alpha}, \boldsymbol{\beta})]) . \qquad (15)
$$

In this type of gauge representation, $G(\vec{\alpha})$ is a positive distribution over a set of Hermitian density-matrix elements $\widehat{\Lambda} + \widehat{\Lambda}^{\dagger}$. It is simple to verify that, by construction

$$
\operatorname{Tr}\left(\widehat{\Lambda}\right) = \Omega\ .\tag{16}
$$

For the case of $\Omega = 1$, this representation reduces to the positive-P representation, and the kernel $\widehat{\Lambda}(\vec{\alpha})$ is a projection operator. Since the positive-P representation is a omplete representation, it follows that another way to construct the gauge P representation is always available, if one simply defines

$$
G(\vec{\alpha}) = P^{(+)}(\alpha, \beta)\delta^2(\Omega - 1) . \qquad (17)
$$

As a simple example, a thermal ensemble with n_0 bosons per mode gives a diagonal P distribution that is Gaussian, so that

$$
G_{th}(\vec{\alpha}) \propto \exp\left[-\left|\alpha\right|^2/n_0\right] \delta^{2M}(\alpha - \beta^*)\delta^2(\Omega - 1) . (18)
$$

One advantage of the proposed representation is that it allows more general expansions than the positive-P distribution, and also in
ludes the ase of the omplex P representation — which has proved useful in solving for non-equilibrium steady-states in quantum systems.

D. Operator identities

The utility of these methods arises when they are used to calculate time (or imaginary time $-$ for which the positive-P distribution annot be used) evolution of the density matrix. This occurs via a Liouville equation of generi form

$$
\frac{\partial}{\partial t}\hat{\rho} = \hat{L}(\hat{\rho}) , \qquad (19)
$$

where the Liouville superoperator typically involves preand post-multiplication of $\hat{\rho}$ by annihilation and creation operators. As an example, the equation for purely unitary time evolution under a Hamiltonian \hat{H} is

$$
i\hbar \frac{\partial}{\partial t} \hat{\rho} = \left[\hat{H}, \hat{\rho} \right] . \tag{20}
$$

Effects of the annihilation and creation operators on the projectors are obtained using the results for the actions of operators on the Bargmann states,

$$
\hat{\mathbf{a}}\hat{\Lambda}(\vec{\alpha}) = \alpha \hat{\Lambda}(\vec{\alpha}) \n\hat{\mathbf{a}}^{\dagger}\hat{\Lambda}(\vec{\alpha}) = [\partial_{\alpha} + \beta] \hat{\Lambda}(\vec{\alpha}) \n\hat{\Lambda}(\vec{\alpha}) = \Omega \partial_{\Omega} \hat{\Lambda}(\vec{\alpha}) .
$$
\n(21)

For brevity, we use $\overrightarrow{\partial} = (\partial_{\Omega}, \partial_{\alpha}, \partial_{\beta})$ to symbolize either $(\partial_i^x \equiv \partial/\partial x_i)$ or $-i(\partial_i^y \equiv \partial/\partial y_i)$ for each of the $i = 0, \ldots, 2M$ complex variables $\vec{\alpha}$. This is possible since $\widehat{\Lambda}(\overrightarrow{\alpha})$ is an analytic function of $\overrightarrow{\alpha}$, and an explicit hoi
e of the derivative will be made later.

Using the operator identities given above, the operator equations can be transformed to an integro-differential equation,

$$
\frac{\partial \widehat{\rho}}{\partial t} = \int G(\overrightarrow{\alpha}) \left[\mathcal{L}_A \widehat{\Lambda}(\overrightarrow{\alpha}) \right] d^{4M+2} \overrightarrow{\alpha} . \qquad (22)
$$

Here the anti-normal ordered notation \mathcal{L}_A indicates an ordering of all the derivative operators to the right. As an example, in the Hamiltonian case, if the original Hamilto- $\hat{H}(\widehat{\mathbf{a}},\widehat{\mathbf{a}}^{\dagger})$ is normally-ordered (annihilation operators) to the right), then

$$
\mathcal{L}_A = \frac{1}{i\hbar} \left[H_A(\boldsymbol{\alpha}, \boldsymbol{\partial}_{\alpha} + \boldsymbol{\beta}) - H_A(\boldsymbol{\beta}, \boldsymbol{\partial}_{\beta} + \boldsymbol{\alpha}) \right] . \tag{23}
$$

If no terms higher than second order occur, this procedure gives a differential operator with the following general expansion:

$$
\mathcal{L}_A^{(+)} = V + A_j^{(+)} \partial_j + \frac{1}{2} D_{ij} \partial_i \partial_j . \tag{24}
$$

where, to simplify notation, the Latin indices i, j, k will from now on be summed over $i = 1, \ldots, 2M$, since no derivatives with respect to Ω are used as yet. V is a term not involving derivative operators with respe
t to any of the variables in $\overrightarrow{\alpha}$. The drift term $A_j^{(+)}$ that is normally found using the positive-P representation is labeled with the superscript $(+)$ to identify it.

At this stage, the usual pro
edure in representation theory is to integrate by parts, provided boundary terms vanish. This gives a normally-ordered differential operator a
ting on the distribution itself, of form

$$
\frac{\partial}{\partial t}G(\vec{\alpha}) = \left[V - \partial_j A_j^{(+)} + \frac{1}{2}\partial_i \partial_j D_{ij}\right]G(\vec{\alpha}).
$$
 (25)

This type of generalized Fokker-Plan
k equation an be treated formally using te
hniques developed by Graham, involving time-symmetric curved-space path integrals [30]. For computational purposes, we require special hoi
es of the analyti derivatives to obtain a positivedefinite diffusion, so that the path integrals have equivalent stochastic equations [29]. We emphasize here that the equations resulting are quite different to those obtained from the direct insertion of a coherent state iden $tity$ into a Feynman path integral $-$ which results in severe convergence problems [31]. The usual positive-P representation equations are obtained at this stage provided there is no potential term $-$ and can be transformed to stochastic equations using the techniques described in the following section.

III. GAUGE FUNCTIONS

In gauge representations, the time evolution of the representation is modified from the usual positive-P representation equations, by the introdu
tion of a number of arbitrary and freely defined functions on the phase space. This freedom of choice is, of course, not present with an orthogonal basis, and is due to the non-orthogonal nature of a oherent basis set. Although we do not investigate other ases, it is worth noting that a similar gauge freedom is implicitly present whenever a non-orthogonal $expansion$ is used $-$ even if it involves different states from the choice of coherent states made here (e.g., the Fock state wave functions in Refs. [7, 8]).

A. Diffusion gauges

We first introduce the diffusion gauges, which were impli
itly present in the original positive-P representation, but were only re
ognized re
ently as allowing improvements in the sampling error. These gauges occur via

the non-unique decomposition of the complex diffusion matrix D , which determines the stochastic correlations in the final equations. Arbitrary functional parameters can therefore be inserted into the final stochastic equations in the noise coefficients, which may lead to further optimization of the simulation. This is be
ause the de composition of the complex diffusion matrix $D = BB^T$, which is needed to define a stochastic process, does not specify the resulting noise matrix B completely.

It has been re
ently shown by Plimak, Olsen and Collett [9] that for the Kerr oscillator using a decomposition different from the obvious diagonal one leads to impressive improvements in the signal-to-noise ratio of the simulation (briefly described in Appendix A 2). This somewhat surprising result leads us to try to quantify the amount of freedom of choice available from this source.

Since $D = D^T$, it can always be diagonalized by a omplex orthogonal transformation

$$
D = O\lambda^2 O^T = B^{(+)}B^{(+)T} , \qquad (26)
$$

where λ is the diagonal matrix whose square gives the eigenvalues of D. Thus $B^{(+)} = O\lambda$ can be considered the canonical, or "obvious" choice of decomposition, unique apart from the 2M signs of the diagonal terms. However, for any orthogonal \tilde{U} , if $B^{(+)}$ is a valid decomposition of D, then so is the matrix $B = B^{(+)}U$. Hence, any matrix in the whole orthogonal family $B = O\lambda U$ is a valid decomposition. This can be easily quantified using a basis

$$
\sigma_{kl}^{(ij)} = \delta_{ik}\delta_{jl} - \delta_{il}\delta_{jk} ,
$$

of the $M(2M-1)$ independent antisymmetric $2M \times 2M$ matrices $\underline{\sigma}^{(ij)}$. One simply introduces

$$
U = \exp\left(\sum_{i < j} g_{ij}(\vec{\alpha}, t) \underline{\underline{\sigma}}^{(ij)}\right) \tag{27}
$$

As an example, for a one-mode case there is one complex gauge function introduced this way, which is $g^d = g_{12}$. The resulting transformation is

$$
U = \exp\left(g^d \underline{\underline{\sigma}}^{(12)}\right)
$$

= $\cos(g^d) + \underline{\underline{\sigma}}^{(12)} \sin(g^d)$, (28)

where the antisymmetric matrix $\underline{\sigma}^{(12)}$ is proportional to a Pauli matrix,

$$
\underline{\underline{\sigma}}^{(12)} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} . \tag{29}
$$

Hence, if the noise was diagonal in the canonical form, the transformed (but equivalent) noise matrix be
omes

$$
B = \begin{bmatrix} \lambda_{11} \cos(g^d) & \lambda_{11} \sin(g^d) \\ -\lambda_{22} \sin(g^d) & \lambda_{22} \cos(g^d) \end{bmatrix} .
$$
 (30)

Now, the $2M$ -dimensional (complex) orthogonal matrix family contains $M(2M - 1)$ free complex parameters, so there are $M(2M - 1)$ diffusion gauge functions $g_{ij}(\overrightarrow{\alpha},t)$ that one can choose arbitrarily. This represents a large class of specific gauges that can be used directly in simulations, as opposed to the conditions on noise correlations usually given elsewhere [9].

As pointed out by Graham $[30]$, there is a close similarity between the theory of curved-space metrics, and path integrals with a space-varying diffusion matrix. In the present ontext, the spa
e is omplex, and we have a family of gauges that are generated on taking the matrix square root of the diffusion matrix. We have not yet used this matrix square root, but this de
omposition will be applied to obtain positive-definite equations via the hoi
e of analyti derivatives made in the following sections.

The above holds for square noise matrices B_s , but one is also free to add more noise coefficients in the manner $B_Q = [B_s, Q]$. Then

$$
B_s B_s^T = \tilde{D} = D - QQ^T , \qquad (31)
$$

and all the 2MW coefficients in the $2M \times W$ matrix Q are additional arbitrary omplex fun
tions. The freedom in B_s is the same as before [i.e. $M(2M-1)$ independent complex gauge functions, with the proviso that B_s is now given by $\widetilde{O} \widetilde{\lambda} U$ where the square of $\widetilde{\lambda}$ gives the eigenvalues of the modified matrix \tilde{D} . The matrix B_s would be unchanged if QQ^T were set to zero, although this hoi
e of Q does not appear to be useful; it just adds extra noise. In general it is not clear whether or not any advantage an be gained by introdu
ing the additional off-square gauge functions contained in Q .

If B is given a functional form dependent on the phasespa
e variables, it may lead to additional terms in the Stratonovi
h form of the equations, whi
h are onsidered later in this se
tion. In this situation one must be areful not to introdu
e additional boundary-term errors arising from an ex
essively rapid growth of the noise gauges.

There is a subtlety here whi
h one must take some are with. The complex noise matrix B is not the matrix that usually appears in the theory of sto
hasti equations. Instead, this matrix is subsequently transformed into an 'equivalent' stochastic form, by taking advantage of the analyti
ity of the Bargmann states. This means that the effect of the diffusion gauges on the final equations also makes use of the non-uniqueness of the coherent basis set itself.

B. Drift gauges

While the diffusion gauges can control sampling error due to the orrelations of noise terms, they annot eliminate boundary terms due to singular trajectories in the drift equations. The extra variable Ω allows the ∂_{Ω} identity to be used to convert any potential term V to a derivative term, and also to introduce a stochastic gauge to stabilize the resulting drift equations. This defines an infinite class of formally equivalent Fokker-Planck equations, in a similar way to related pro
edures in QED and QCD. To demonstrate this, we introduce $2M$ arbitrary complex drift gauge functions $\mathbf{g} = [\, g_i(\overrightarrow{\alpha},t) \,],$ to give a new differential operator \mathcal{L}_{GA} whose form differs from the original $\mathcal{L}_A^{(+)}$ by terms that vanish identically when applied to the kernel $\widehat{\Lambda}(\overrightarrow{\alpha})$,

$$
\mathcal{L}_{GA} = \mathcal{L}_A^{(+)} + \left[V + \frac{1}{2} \mathbf{g} \cdot \mathbf{g} \, \Omega \, \partial_{\Omega} + g_k B_{jk} \partial_j \right] \left[\Omega \partial_{\Omega} - 1 \right]. \tag{32}
$$

The total differential operator \mathcal{L}_{GA} has an anti-normal Fokker-Planck form. Extending the drift and diffusion matrices to include the extra variable Ω , we can write this — summing repeated a, b, c indices over $a = 0, \ldots, 2M$ as

$$
\mathcal{L}_{GA} = \left[A_a \partial_a + \frac{1}{2} D_{ab} \partial_a \partial_b \right] . \tag{33}
$$

The total complex drift vector is $\overrightarrow{A}=(A_0,A_1,\ldots,A_{2M});$ where

$$
A_0 = \Omega V
$$

\n
$$
A_j = A_j^{(+)} - g_k B_{jk}
$$
 (34)

The new diffusion matrix \underline{D} with elements D_{ab} is not diagonal, but it can be factorized. Explicitly, it is now a square $(2M + 1) \times (2M + 1)$ complex matrix, given by

$$
\underline{D} = \begin{bmatrix} \Omega^2 \mathbf{g} \mathbf{g}^T & \Omega \mathbf{g} B^T \\ B \mathbf{g}^T \Omega & B B^T \end{bmatrix}
$$

$$
= \begin{bmatrix} 0 & \Omega \mathbf{g} \\ 0 & B \end{bmatrix} \begin{bmatrix} 0 & 0 \\ \Omega \mathbf{g}^T & B^T \end{bmatrix} = \underline{B} \underline{B}^T . \qquad (35)
$$

Thus, we now have a new stochastic noise matrix with one added dimension,

$$
\underline{\underline{B}} = \begin{bmatrix} 0 & \Omega \mathbf{g} \\ 0 & B \end{bmatrix} . \tag{36}
$$

The operator (32) was chosen to give this form for \underline{B} , so that the only change in noise is for the Ω variable.

Positive-definite diffusion \mathbf{C} .

It is always possible to transform these secondderivative terms into a positive semi-definite diffusion ment for a stochastic equation. When $\underline{D} = \underline{B} \underline{B}^T$, divide
 $\underline{B} = \underline{B}^x + i\underline{B}^y$ into its real and imaginary parts. A similar procedure is followed for \overline{A} .

Recalling that the original kernel was analytic, thus allowing for more than one choice of derivatives, the choice for ∂_a can now be made definite by choosing it so that the resulting drift and diffusion terms are always real,

$$
A_a \partial_a \rightarrow A_a^x \partial_a^x + A_a^y \partial_a^y, \qquad (37)
$$

\n
$$
D_{ab} \partial_a \partial_b \rightarrow B_{ac}^x B_{bc}^x \partial_a^x \partial_b^x + B_{ac}^y B_{bc}^x \partial_a^y \partial_b^x + (x \leftrightarrow y).
$$

Hence, the gauge differential operator can now be written explicitly as

$$
\mathcal{L}_{GA} = \left[\widetilde{A}_{\mu} \partial_{\mu} + \frac{1}{2} \widetilde{D}_{\mu\nu} \partial_{\mu} \partial_{\nu} \right],
$$
 (38)

where the indices μ, ν cover the $(4M + 2)$ -dimensional phase-space of the real and imaginary parts of $\vec{\alpha}$, so that $\tilde{\alpha} = (\vec{x}, \vec{y})$, and $\partial_{\mu} = \partial/\partial \tilde{\alpha}_{\mu}$. The diffusion matrix $\underline{\tilde{D}} =$ $\underline{\widetilde{B}}\underline{\widetilde{B}}^T$ is now positive semi-definite, since, by construction

$$
\underline{\underline{\widetilde{B}}} = \begin{bmatrix} 0 & \underline{\underline{B}}^x \\ 0 & \underline{\underline{B}}^y \end{bmatrix} . \tag{39}
$$

so that the diffusion matrix is the square of a real ma $trix$ — explicitly,

$$
\underline{\underline{\widetilde{D}}} = \begin{bmatrix} 0 & \underline{\underline{B}}^x \\ 0 & \underline{\underline{B}}^y \end{bmatrix} \begin{bmatrix} 0 & 0 \\ (\underline{\underline{B}}^x)^T & (\underline{\underline{B}}^y)^T \end{bmatrix} . \tag{40}
$$

As \mathcal{L}_{GA} is now explicitly real as well as positive-definite by onstru
tion, it an be applied to the Hermitian onjugate kernel as well, resulting in the final time-evolution equation,

$$
\frac{\partial \widehat{\rho}}{\partial t} = \int G(\widetilde{\alpha}) \left[\mathcal{L}_{GA} \widehat{\Lambda}(\widetilde{\alpha}) \right] d^{4M+2} \widetilde{\alpha} . \tag{41}
$$

On integrating by parts, provided boundary terms vanish, at least one solution will satisfy the following (normally-ordered) positive-definite Fokker-Planck equa- tion — with the differential operators on the left, each a
ting on all terms to the right,

$$
\frac{\partial G}{\partial t} = \mathcal{L}_{GN} G \equiv \left[-\partial_{\mu} \widetilde{A}_{\mu} + \frac{1}{2} \partial_{\mu} \partial_{\nu} \widetilde{D}_{\mu \nu} \right] G. \quad (42)
$$

This implies that we have an equivalent set of Ito stochastic differential equations available, with 2M real Gaussian noises dW_i , which are

$$
d\Omega = \Omega(Vdt + g_k dW_k)
$$

\n
$$
d\alpha_j = (A_j^{(+)} - g_k B_{jk})dt + B_{jk} dW_k
$$
 (43)

The noises obey $\langle dW_i dW_j \rangle = \delta_{ij} dt$, and are uncorrelated between time steps.

Numeri
al simulations are usually done in the Stratonovich calculus, due to superior convergence properties [32], so the equivalent *complex* Stratonovich equation allows us to write efficient algorithms,

$$
d\alpha_a = dx_a + idy_a
$$

= $\left[A_a - \frac{1}{2} (B_{bk} \partial_b) B_{ak}\right] dt + B_{ak} dW_k$, (44)

where $(B_{bk}\partial_b) \equiv (B_{bk}^x \partial_b^x + B_{bk}^y \partial_b^y)$. The derivative terms above are the Stratonovich correction in the drift, corresponding to related terms obtained in curved-space path integrals.

These gauge terms are now utilized to stabilize oherent-state paths entering into highly nonlassi
al regions of phase space. This allows one to benefit from the overompleteness of oherent states, in redu
ing the sampling error and eliminating boundary terms.

D. Moments

The procedure for calculating observable moments is slightly different for the gauge representation than for the positive P. Any moment an be written in terms of the normally ordered operator products $\hat{a}^{\dagger n}\hat{a}^m,$ and their expe
tation values are given by

$$
\left\langle \hat{a}^{\dagger n} \hat{a}^{m} \right\rangle_{\text{quant}} = \frac{\left\langle \beta^{n} \alpha^{m} \Omega + (\alpha^{n} \beta^{m} \Omega)^{*} \right\rangle_{\text{stoch}}}{\left\langle \Omega + \Omega^{*} \right\rangle_{\text{stoch}}} \ . \tag{45}
$$

which differs from the positive-P situation whenever Ω differs from unity.

The average norm $\langle \Omega \rangle$ is always preserved if there is no potential term $(V = 0)$, since the resulting equation for the weight variable is

$$
d\Omega = \Omega g_k dW_k \tag{46}
$$

The decorrelation property of Ito equations [29] then implies that

$$
\langle d\Omega \rangle = \langle \Omega g_k \rangle \langle dW_k \rangle = 0 \ . \tag{47}
$$

E. Gauge properties

We turn briefly here to the question of gauge classification and properties. Just as in QED, the overomplete nature of the oherent-state expansion means that many equivalent, stable gauges exist. However, they may not be equivalent in terms of boundary terms. These are determined by the tails of the distribution function, which depends intimately on the gauge hosen for the time evolution. It is essential that the distribution tails are suf ficiently bounded to eliminate boundary terms arising in partial integration. It is sufficient to bound tails better than any inverse power law, for which it is conjectured to require (as a necessary condition) that all deterministic

trajectories are bounded over any finite time interval [6]. This issue is discussed in greater detail below, and in Ref. $[33]$.

The main criteria for a useful gauge are the elimination of boundary terms and the redu
tion of sampling error. However, there is an enlarged spa
e of variables for the Fokker-Plan
k equation here. For this reason, it is possible to stabilize trajectories in the usual positive-P phase space, while introducing new gauge-induced boundary terms in the Ω space. When it comes to the formation of boundary terms, the phase of Ω is generally innocuous provided the gauge is periodic in this variable, but the gauge distribution must be strongly bounded as $|\Omega| \to \infty$ to prevent new boundary terms from arising.

We can classify gauges according to their real or imaginary nature, and their functional dependence; which can be on just the phase-spa
e variables, just the quantum phase, or on both. This gives rise to nine gauge types, depending on the following criteria.

1. Gauge omplexity

Gauges are in general omplex fun
tions, whi
h leads to the following classification of gauge complexity:

- 1. Real gauge
- 2. Imaginary gauge
- 3. Complex gauge

In general, we find that trajectories can be stabilized by real, imaginary or omplex gauges, provided they have some (α, β) phase-space dependence.

It is worthwhile to note that the imaginary and real parts of the gauges affect the behavior of sampling error differently. In the Ito calculus, the evolution of the weight Ω due to the gauges is simply $d\Omega = \Omega q_k dW_k$. Typically, i.e., when there are no significant correlations between the phase of α (or β) and Ω , the weight factor appearing in moment calculations is just approximately $\text{Re}[\Omega]$. As a general rule, sampling errors are partially due to stochastic fluctuations in the phase-space trajectories, and partially due to stochastic fluctuations in the weight function. Thus there is a trade-off; a gauge that is strongly stabilizing may reduce phase-space fluctuations at the expense of in
reased weight varian
e, and vice versa.

To understand the different types of gauges in somewhat greater detail, we onsider the evolution of the weight variance for real and imaginary gauges, in a simple ase where gauge and weight are de
orrelated, with $\Omega = 1$ initially. Let $\Omega = \Omega' + i\Omega''$ and $g_k = g'_k + ig''_k$, then

$$
d\Omega' = (\Omega' g'_k - \Omega'' g''_k) dW_k ,
$$

\n
$$
d\Omega'' = (\Omega' g''_k + \Omega'' g'_k) dW_k .
$$
\n(48)

If we onsider the evolution of the squares of these terms, the Ito rules of stochastic calculus give

$$
d\langle [\Omega']^2 \rangle = \langle (\Omega' g'_k - \Omega'' g''_k)^2 \rangle dt ,
$$

$$
d\langle [\Omega'']^2 \rangle = \langle (\Omega' g''_k + \Omega'' g'_k)^2 \rangle dt .
$$
 (49)

Suppose for simplicity that the g_k and Ω are approximately uncorrelated, then we have two cases to consider.

1. Real gauge:

$$
d\langle[\Omega']^2\rangle = \langle[\Omega']^2\rangle d\tau , \qquad (50)
$$

where $d\tau = \langle q_k q_k \rangle dt$. This initially leads to linear growth in the varian
e, and hen
e in the sampling error. The real part of the gauge will ause noise directly in Ω' , producing asymetric spreading in Ω' , which can lead to a few rare very highly weighted trajectories for times $\tau \gtrsim 1$. The effect of the real gauge may be
ome misleading on
e the distribution be
omes highly skewed, as the rare tra je
tories that are important for moment calculations may be missed if the sample is too small. At long times, if $\langle g_k g_k \rangle$ is constant and uncorrelated with Ω , then the growth becomes exponential, with $\langle [\Omega']^2 \rangle = e^{\tau}$.

2. Imaginary gauge:

$$
d\langle [\Omega']^2 \rangle = \langle [\Omega'']^2 \rangle d\tau ,
$$

\n
$$
d\langle [\Omega'']^2 \rangle = \langle [\Omega']^2 \rangle d\tau .
$$
\n(51)

where $d\tau = \langle g''_k g''_k \rangle dt$. This leads initially to quadratic growth in the variance of Ω' , and hence a slower growth in the sampling error. If $\langle q_k q_k \rangle$ is constant and remains uncorrelated with Ω , then the growth is given by $\langle [\Omega']^2 \rangle = \cosh(\tau)$, $\langle [\Omega'']^2 \rangle =$ $\sinh(\tau)$. An imaginary gauge will cause mutual canceling of trajectories that have weights of randomly positive and negative sign once $\tau \gtrsim \pi$. This can also have deleterious effects for small samples, if the average sample weight becomes negative \sim of ourse, this annot be true over the entire sto
hasti population.

The generic behavior is more complex than in the examples given above, due to orrelations between the gauge

Clearly any type of gauge tends to ause growth in the norm varian
e. However, there is an ex
eption to this rule: the norm-preserving gauges. This lass of gauges is of spe
ial interest as they generate tra je
tories having an invariant normalization, so that $\text{Re}[d\Omega] \equiv 0$. From the equation for the norm varian
e, Eq. (49), it follows that a necessary and sufficient condition for a norm-preserving gauge is that $\Omega' g'_k = \Omega'' g''_k$. If $\Omega' = 1$ initially, this implies that $g_k = i\Omega^* f_k = i(1 - i\Omega'')f_k$, where f_k is a real function. Unless $g_k = 0$, norm-preserving gauges are generally fun
tions of both the phase-spa
e variables and the weight Ω . A preliminary study of these gauges has shown that these gauges can greatly reduce sampling error, although gauge-indu
ed boundary terms are also possible [2], depending on the choice of f_k .

2. Fun
tional dependen
e

From the above analysis, we see that gauges can functionally depend on any phase-spa
e variable, as well as the generalized quantum phase variable or weight Ω . This leads to three functional types:

- 1. Autonomous (depends on Ω only)
- 2. Spa
e dependent (depends on phase-spa
e only)
- 3. Mixed (depends on all components of $\vec{\alpha}$ including Ω)

Autonomous gauges appear to be the least useful sin
e they do not affect α or β behavior, but gauges of either purely spa
e-dependent or mixed type an be used.

A possible aveat with mixed gauges is that they may be mu
h harder to analyze, as two-way ouplings will occur between the normal phase-space variables α , β and the weight.

IV. NONLINEAR ABSORBER CASE

The nonlinear absorber is an example of a nonlinear master equation that can give either correct or incorrect results when treated with the usual positive-P representation methods, if the boundary terms are ignored. Generally, problems only arise when the linear damping has ex
eptionally small values or the number of bosons per mode is small (see Fig. 2), so this is not a practical problem in optics. However, for other physical systems such as a BEC this may be significant. It is a well-studied case, and a detailed treatment can be found in Ref. $[6]$. It also has the merit that exact solutions can be readily found using other means. By analyzing this example we can ensure that the modifications to the drift equations obtained from gauge terms, do eliminate boundary terms and give orre
t results.

Consider a cavity mode driven by coherent radiation, and damped by a zero temperature bath that auses both one and two photon losses. We have s
aled time so that the rate of two-photon loss is unity. Without this nonlinear pro
ess, nothing unusual happens. The s
aled onephoton loss rate is γ , and ε is the scaled (complex) driving field amplitude. The master equation is

$$
\frac{\partial \hat{\rho}}{\partial t} = \left[\varepsilon \hat{a}^{\dagger} - \varepsilon^* \hat{a}, \hat{\rho} \right] + \frac{\gamma}{2} (2 \hat{a} \hat{\rho} \hat{a}^{\dagger} - \hat{a}^{\dagger} \hat{a} \hat{\rho} - \hat{\rho} \hat{a}^{\dagger} \hat{a}) \n+ \frac{1}{2} (2 \hat{a}^2 \hat{\rho} \hat{a}^{\dagger 2} - \hat{a}^{\dagger 2} \hat{a}^2 \hat{\rho} - \hat{\rho} \hat{a}^{\dagger 2} \hat{a}^2).
$$
\n(52)

Following the treatment of Se
. II, we arrive at the gauge representation Stratonovi
h sto
hasti equations

$$
d\alpha = [\varepsilon - \alpha(\alpha\beta + ig + (\gamma - 1)/2)]dt + i\alpha dW,
$$

\n
$$
d\beta = [\varepsilon^* - \beta(\alpha\beta + i\overline{g} + (\gamma - 1)/2)]dt + i\beta d\overline{W},
$$

\n
$$
d\Omega = S_{\Omega}dt + \Omega \left[g dW + \overline{g} d\overline{W} \right].
$$
\n(53)

Here $S_{\Omega}dt$ is the appropriate Stratonovich correction term [given by the derivative terms in Eq. (44)], which depends on the particular gauges chosen.

With no gauge $(g = \overline{g} = 0)$, the positive-P Stratonovich equations are recovered,

$$
d\alpha = [\varepsilon - \alpha(\alpha\beta + \{\gamma - 1\}/2)]dt + i\alpha dW,
$$

\n
$$
d\beta = [\varepsilon^* - \beta(\alpha\beta + \{\gamma - 1\}/2)]dt + i\beta d\overline{W}.
$$
 (54)

We will concentrate on the various simplifications of this model, whi
h orrespond to existing literature, and simpler analysis.

A. Relevan
e to many-body problems

The nonlinearity seen here can occur directly in the form of a nonlinear ollisional damping term in a manybody system, so that it an be referred to generi
ally as `two-boson absorption'. This type of damping is ommon both to nonlinear photonic and atomic interactions.

It is of nearly the same form as for an `imaginary-time' thermal equilibrium state calculation for the usual model of an alkali-metal Bose gas or BEC $[34]$. There, for example, the intera
tion energy between identi
al bosons of mass m and s -wave scattering length a_s in D -dimensional spa
e is given by

$$
\hat{H} = \frac{2\pi\hbar^2 a_s}{m} \int d^D \mathbf{x} \hat{\psi}^{\dagger 2}(\mathbf{x}) \hat{\psi}^2(\mathbf{x}) , \qquad (55)
$$

provided that a_s is much smaller than other characteristic lengths of the system (which is usually the case). The master equation for an imaginary-time calculation is

$$
\frac{\partial \hat{\rho}_e}{\partial \tau} = -\frac{1}{2} \left\{ \hat{H} - \mu \hat{N}, \hat{\rho}_e \right\}_+ , \qquad (56)
$$

where $\hat{\rho}_e$ is the thermal canonical ensemble density matrix, μ is the chemical potential, N is the number operator for the entire system, and $\tau = 1/k_BT$ is an inverse temperature. Apart from the fact that it is not tracepreserving, this is a nonlinearity very similar to that ocurring in the nonlinear absorber master equation.

While boundary-term discrepancies only occur with this nonlinearity for low occupations per mode (see also Fig. 2), for a many-mode system at finite temperature one expe
ts a large number of modes to have just su
h a low occupation. Thus, it is important to check that boundary terms are indeed eliminated. Note that the gauge representation simulation is efficient over a wide range of occupation numbers. See, for example, Fig. 3. More details of appli
ations to both real and imaginary time many-body systems with many modes will be given elsewhere.

Two-boson absorber **B.**

In its simplest form, corresponding to $\gamma = \varepsilon = 0$, only two-boson absorption takes pla
e. We expe
t that for a

state $|\psi\rangle = \sum_n c_n |n\rangle$ all even boson number components will decay to vacuum, and all odd-numbered components will decay to $|1\rangle$, leaving a mixture of vacuum and oneboson states at long times.

The positive-P representation has been found to give erroneous results $[4, 35-37]$ due to the existence of moving singularities $[6]$, which cause power-law tails in the distribution leading to boundary terms. The moment usually on
entrated on in this system is the number of bosons $\hat{n} = \hat{a}^\dagger \hat{a}$, which corresponds to the statistical average of $n = \alpha \beta$ in the positive-P representation. This has a convenient closed equation (Stratonovich),

$$
dn = -n(n + i\tilde{g} - 1/2)d\tau + indW^+ \tag{57}
$$

with $dW^+ = (dW + d\overline{W})$, $\tau = 2t$, and $\tilde{g} = (g + \overline{g})/2$.

Let us examine the behavior of the above equation, when $\tilde{q} = 0$, i.e., in the standard, un-gauged formulation. The deterministic part of the evolution has a repellor at $n = 0$, and an attractor at $n = \frac{1}{2}$. The noise is finite, and of standard deviation $\sqrt{dt/2}$ at the attractor. We can see that the deterministic part of the evolution has a single trajectory of measure zero which can escape to infinity along the negative real axis,

$$
\alpha = -\beta = \frac{1}{\sqrt{\tau_0 - \tau}},\tag{58}
$$

where $\tau_0 = 1/\alpha(0)^2 = -1/n(0)$. This moving singularity is known to cause the power law behavior of the Fokker-Planck solution at large $|n|$, which means that integration by parts is not in fact valid $-$ which leads to incorrect results.

Indeed, it can be easily seen that in the steady-state limit, all trajectories in a simulation will head toward $n = \frac{1}{2}$, making $\lim_{t \to \infty} \langle \hat{n} \rangle = \frac{1}{2}$. Quantum mechanics, however, predicts that if we start from a state $\hat{\rho}_0$, the steady state will be

$$
\lim_{t \to \infty} \langle \hat{n} \rangle = \sum_{j=0}^{\infty} \langle 1 + 2j | \hat{\rho}_0 | 1 + 2j \rangle . \tag{59}
$$

For a coherent state $|\alpha_0\rangle$ input, say, this will be

$$
\lim_{t \to \infty} \langle \hat{n} \rangle = \frac{1}{2} \left(1 - e^{-2|\alpha_0|^2} \right) . \tag{60}
$$

Thus we can expect that the positive P simulation will give correct results only when $e^{|\alpha_0|^2} \gg 1$.

To orre
t the problem we have to hange the phasespace topology in some way to prevent the occurrence of moving singularities. We have found that a good gauge for a two-boson absorber nonlinearity in general is

$$
g = \overline{g} = \tilde{g} = i(n - |n|) . \tag{61}
$$

This replaces the $-n^2$ term in Eq. (57), which may become repulsive from zero, with $-n|n|$ which is always a restoring for
e, and so never leads to super-exponential es
ape.

Figure 1: Comparison of two-boson damping simulations. $Circles: positive P simulation; solid line: circular gauge simu$ lation; *dashed line*: exact calculation (truncated number-state basis). Simulation parameters: 40 000 tra je
tories; step size $= 0.005$; initial coherent state. Stratonovich semi-implicit method $[32]$.

With the gauge (61) , the Stratonovich equations become

$$
dn = -n(|n| - 1/2)d\tau + indW^{+},
$$

\n
$$
d\Omega = \Omega \{ [n + (n - |n|)^{2}]d\tau/2 + i(n - |n|)dW^{+} \}.
$$
\n(62)

Phase-space trajectories have changed now, but since it has all come from the same master equation, it still des
ribes the same system. Consider the equations for the polar decomposition of $n = re^{i\phi}$,

$$
dr = -r(r - 1/2)d\tau ,
$$

\n
$$
d\phi = dW^{+}.
$$
\n(63)

This is exa
t, and shows that now we have an attra
tor on the circle $|n| = \frac{1}{2}$, and a repellor at $n = 0$, with free phase diffusion in the tangential direction. Once trajectories reach the attractor, only phase diffusion occurs. Some more complicated evolution is occurring in the Ω variable. In any case, there are now no moving singularities anywhere in the phase spa
e, and simulations orrespond exactly to quantum mechanics.

Figure 1 ompares results for a trun
ated number-state basis calculation, a positive-P calculation, and a "circular" gauge (61) calculation for an initial coherent state of $\alpha_0 = 1/\sqrt{2}$. Figure 2 compares steady-state values for exact, positive-P, and gauge calculations for various initial oherent states in a wide range. It is seen that the gauge al
ulation is orre
t to within the small errors due to finite sample size.

Figure 2: Steady state expe
tation values of boson number $\langle \hat{n} \rangle$ obtained by gauge simulations (*double triangles*) compared to exact analytic results from Eq. (60) (*solid line*) and positive-P simulations (*circles*) for a wide range of initial coherent states. Size of uncertainty in gauge results due to finite sample size is indicated by vertical extent of 'double-triangle' symbol. Steady state was observed to have been reached in all simulations by $\tau = 7$ or earlier (compare with Fig. 1 and 3), hence this is the time for which the simulation data is plotted. Simulation parameters: 100 000 trajectories; step size $= 0.01$.

C. One- and two-boson absorber

If we now turn on the one-boson decay as well, but still do not have any driving, we expe
t that all states will decay to the vacuum on two time scales 1 and $1/\gamma$. If $\gamma \gg 1$, nothing interesting happens, however if $\gamma \lesssim 1$, we should first see a rapid decay to a mixture of vacuum and one-boson states due to the two-boson pro
ess, and then a slow decay of the one-boson state to the vacuum on a time scale of $\tau \approx 2/\gamma$.

In this case the positive-P equations display different behavior depending on whether γ is above or below the threshold $\gamma = 1$. Below threshold, we have an attractor at $n = (1 - \gamma)/2$, and a repellor at $n = 0$, while above threshold, the attractor is at $n = 0$, and the repellor at $n = -(\gamma - 1)/2$. In either case, there is a singular trajectory along the negative real axis, which can cause boundary-term errors. It turns out that the steady state calculated this way is erroneous while γ < 1, and there are transient boundary-term errors while $\gamma < 2$ [4]. The false steady state below threshold lies at the location of the attractor: $(1 - \gamma)/2$.

Let us try to fix this problem using the same circular gauge (61) as before. The equation for r is now

$$
dr = -r(r - [1 - \gamma]/2)d\tau , \qquad (64)
$$

while the ϕ and Ω evolution is unchanged. So, above threshold we are left with only an attractor at $n = 0$,

Figure 3: Comparison of simulations for system with both single- and double-boson damping. Relative strength $\gamma =$ 0.1; Circles: positive-P simulation. solid line: circular gauge simulation; *dashed line*: exact calculation (truncated numberstate basis). Gauge simulation parameters: 10^5 trajectories; step size varies from 0.0001 to \approx 0.006; initial coherent state $|10\rangle$ with $\langle \hat{n} \rangle = 100$ bosons.

while below threshold we have a repellor at $n = 0$ surrounded by an attracting circle at $r = (1 - \gamma)/2$. This phase spa
e again has no moving singularities.

The results of simulations for the parameter $\gamma = 0.1$ are shown in Fig. 3. The gauge simulation tra
ks the exact results. We have chosen $\gamma \ll 1$ so that a system with two widely differing time scales is tested. The ir
ular gauge avoids the false results of the positive-P simulation. Note also that the gauge simulation remains efficient for a wide range of occupation numbers $-$ from $\langle \hat{n} \rangle \approx 100 \gg 1$, where the positive P is also accurate, to $\langle \hat{n} \rangle \approx 0.1 \ll 1$ where it is totally incorrect.

D. Driven two-boson absorber

The other type of situation to consider is when we have a driving field as well as two-boson damping. In these onsiderations we have set the one-boson damping rate to zero $(\gamma = 0)$, since this process never causes any of the simulation problems anyway, but leaving it out simplifies analysis. Failure of the positive-P representation method has been found in this limit as well [5], and is evident in Fig. 4. The equation for n is no longer stand-alone in this ase, and we must simulate all three omplex variables as in Eq. (53), the Ω equation being the same as in the undriven ase (62).

A treatment of the singular trajectory problem with the same circular gauge (61) leads again to correct results, as seen in Fig. 4.

Figure 4: Driven two-boson absorber with $\varepsilon = 0.05$. Circles: positive P simulation (1000 trajectories); solid line: circular gauge simulation (10⁵ trajectories); *dashed line*: exact calculation (truncated number-state basis). Step size $\Delta t = 0.025$. Initial va
uum state.

$V₁$ THE SINGLE-MODE LASER

Let us now consider the second quantum system for which systematic errors have been seen with the positive-P representation. We will see that the problem here is somewhat different than in the previous case. The differen
e is that for two-boson damping, boundary-term errors occur even when we choose an optimal (i.e., compa
t) initial distribution to represent our starting state, whereas here systematic errors occur only for unreasonably broad initial distributions. Nevertheless, since normally it is assumed that the initial condition can be of arbitrary breadth it is instru
tive to investigate how this problem an be ta
kled with sto
hasti gauge methods.

We have found that sto
hasti gauges an be used to in
rease the allowable breadth to in
lude all reasonable starting conditions, but once one tries to increase the initial spread too mu
h, it be
omes unlikely that any gauge will remove systematic errors, without introducing too mu
h sampling (i.e. random) error instead.

The laser model \mathbf{A} .

Ito stochastic differential equations for a simple photonic or atomic laser model that can be derived from the positive-P distribution are $[5, 6]$

$$
d\tilde{\alpha} = (G - \tilde{\alpha}\tilde{\beta})\tilde{\alpha}d\tau + \sqrt{Q}d\eta,
$$

$$
d\tilde{\beta} = (G - \tilde{\alpha}\tilde{\beta})\tilde{\beta}d\tau + \sqrt{Q}d\eta^*
$$
(65)

in appropriate s
aled variables, with the omplex Gaussian noise $d\eta$ obeying $\langle d\eta d\eta^* \rangle = 2d\tau$. In terms of physical parameters, we have

$$
\tilde{\alpha} = \alpha/\sqrt{\mathcal{N}}
$$

$$
\tilde{\beta} = \beta/\sqrt{\mathcal{N}},
$$
 (66)

where τ is the scaled time, and $\mathcal{N} \gg 1$ is a scaling parameter that equals the number of gain atoms in a simple photoni laser model. Both G the gain parameter and $Q \geq G/N$, the noise parameter, are real and positive.

Since this time we are again interested in the (scaled) boson number $\langle \tilde{n} \rangle = \langle \tilde{\alpha} \tilde{\beta} \rangle = \langle \hat{n} \rangle / \mathcal{N}$, its evolution can be written as a losed equation

$$
d\tilde{n} = -2(\tilde{n} - a)(\tilde{n} - b)d\tau + 2\sqrt{Q\tilde{n}}dW , \qquad (67)
$$

where now the real Gaussian noise obeys $\langle dW dW \rangle$ = $d\tau$, and the deterministic stationary points in the Stratonovich calculus are

$$
a = \frac{1}{2} \left(G + \sqrt{G^2 + 2Q} \right) ,
$$

\n
$$
b = \frac{1}{2} \left(G - \sqrt{G^2 + 2Q} \right) .
$$
 (68)

We find that the stationary point at a is an attractor, and at b we have a repellor. Defining $\Delta = b - \tilde{n}$, we get

$$
d\Delta = 2\Delta(\Delta + \sqrt{G^2 + 2Q}) + \text{noise},\qquad(69)
$$

which shows that we again have a singular trajectory escaping to infinity in finite time along the negative real axis for $\tilde{n} < b$.

\mathbf{R} . Initial conditions

Let us consider the usual case of vacuum initial conditions. A vacuum can be represented by

$$
P^{(+)}(\tilde{\alpha}, \tilde{\beta}) = \delta(\tilde{\alpha})\delta(\tilde{\beta}) , \qquad (70)
$$

but also by Gaussian distributions of any variance σ_0^2 , around the above,

$$
P^{(+)}(\tilde{\alpha}, \tilde{\beta}) = \frac{1}{4\pi^2 \sigma_0^4} \exp\left\{-\frac{|\tilde{\alpha}|^2 + |\tilde{\beta}|^2}{2\sigma_0^2}\right\} \ . \tag{71}
$$

Note: the distribution of \tilde{n} is non-Gaussian, but has a standard deviation of $\sigma_{\tilde{n}} \approx \sqrt{2}\sigma_0^2$ in both the real and imaginary dire
tions.

It has been found by Schack and Schenzle [5] that for the single-mode laser model, a positive-P simulation of pumping from a vacuum will give correct answers if the usual δ -function initial condition (70) is used, but will have systematic errors if the initial condition used has a sufficiently large variance (see Fig. 5). We emphasize here that this is not a real problem in practical cases, as the variance required to cause systematic errors is typically extremely large, on
e the s
aling needed to obtain the usual (approximate) laser model is taken into account.

This can be understood because if we have a sufficiently broad initial distribution, the region of phase space that includes the singular trajectory will be explored by the distribution. Even if initially $\sigma_{\tilde{n}} \ll |b|$, the region $\tilde{n} < b$ may be subsequently explored due to the presen
e of the noise terms.

Apart from the obvious δ -function initial condition, one might want to try the anoni
al distribution of Eq. (10), whi
h is a standard positive-P representation construction $[1]$. It will not cause problems as its variance is $\sigma_0^2 = 1/\mathcal{N}$, which for any realistic case will be very small (i.e., $\sigma_{\tilde{n}} \ll |b|$). Schack and Schenzle discovered anomalous results when they chose $\sigma_0^2 = 1$, due to an erroneous procedure of scaling the equations $-\omega$ while not scaling the canonical initial condition in α . Nevertheless, since any σ_0 is supposed to represent the same state, insight into what an be a
hieved using gauge methods is gained if we analyze the systematic errors for such a relatively large σ_0 .

C. Gauge orre
tions

The Fokker-Plan
k equation orresponding to Eq. (65) is

$$
\frac{\partial P}{\partial \tau} = \left\{ \frac{\partial}{\partial \tilde{\alpha}} [\tilde{n} - G] \tilde{\alpha} + \frac{\partial}{\partial \tilde{\beta}} [\tilde{n} - G] \tilde{\beta} + 2Q \frac{\partial^2}{\partial \tilde{\alpha} \partial \tilde{\beta}} \right\} P \tag{72}
$$

We now introdu
e gauges using the same method as in Sec. II. This leads to the Ito stochastic equations

$$
d\tilde{\alpha} = \tilde{\alpha}(G - \tilde{n})d\tau - \sqrt{Q}(g + i\overline{g})d\tau + \sqrt{Q}d\eta,
$$

\n
$$
d\tilde{\beta} = \tilde{\beta}(G - \tilde{n})d\tau - \sqrt{Q}(g - i\overline{g})d\tau + \sqrt{Q}d\eta^*,
$$

\n
$$
d\Omega = \Omega \left[(g - i\overline{g})d\eta + (g + i\overline{g})d\eta^* \right] / 2.
$$
\n(73)

It is convenient to define a transformed gauge function \tilde{g} , which is also arbitrary, such that

$$
g = \frac{(\tilde{\alpha} + \tilde{\beta})\tilde{g}}{2\sqrt{Q}} ,
$$

$$
\overline{g} = \frac{(\tilde{\alpha} - \tilde{\beta})\tilde{g}}{2i\sqrt{Q}} .
$$
 (74)

Changing to \tilde{n} and $\Theta = \ln(\Omega)$ variables we obtain the Stratonovi
h equation

$$
d\tilde{n} = 2\tilde{n}(G - \tilde{n} - \tilde{g})d\tau + Qd\tau + 2\sqrt{Q\tilde{n}}dW,
$$

$$
d\Theta = -\frac{\tilde{n}\tilde{g}^2}{2Q}d\tau + S_{\Theta}d\tau + \tilde{g}\sqrt{\frac{\tilde{n}}{Q}}dW,
$$
 (75)

with $S_{\Theta}dt$ being the appropriate Stratonovich correction $[g$ iven by the derivative terms in Eq. (44) for a particular gauge function \tilde{g} .

D. Corre
ting for the moving singularities

Consider the deterministic evolution of the real part, \tilde{n}_x , of $\tilde{n} = \tilde{n}_x + i\tilde{n}_y$,

$$
d\tilde{n}_x = -2\tilde{n}_x^2 + 2G\tilde{n}_x + Q + 2\tilde{n}_y^2 - 2\tilde{n}_x \text{Re}[\tilde{g}] + 2\tilde{n}_y \text{Im}[\tilde{g}].
$$
\n(76)

The moving singularity is due to the $-2\tilde{n}_x^2$ leading term for negative values of \tilde{n}_x . We now consider criteria for hoosing the drift gauges as follows.

(1) It is desirable to keep the gauge terms to a minimum because whenever they act the weights of trajectories become more randomized — see Sec. III E1. Thus, let us restrict ourselves to functions \tilde{g} that are only nonzero for $\tilde{n}_x < 0$.

(2) This immediately leads to another restriction on \tilde{g} : To be able to use the efficient numerical algorithms in the Stratonovich calculus, we must be able to calculate the correction term S_{Θ} , which depends on derivatives of $\tilde{g}\sqrt{\tilde{n}/Q}$. This immediately suggests that \tilde{g} must always be continuous, hence, in particular, $\lim_{n_x\to 0}(\tilde{g})=0$. For ease of analysis, let us start with a simple form for the gauge, $\tilde{g} = c - \lambda \tilde{n}_x + \lambda_y \tilde{n}_y$. This restriction immediately implies $c = \lambda_y = 0$, hence

$$
\tilde{g} = \begin{cases}\n-\lambda \text{Re}[\tilde{n}] & \text{if } \text{Re}[\tilde{n}] < 0 \\
0 & \text{if } \text{Re}[\tilde{n}] \ge 0\n\end{cases},\n\tag{77}
$$

and $S_{\Theta} = \lambda (\text{Re}[\tilde{n}] + \tilde{n} + |\tilde{n}|)/2$. when $\text{Re}[\tilde{n}] < 0$, zero otherwise.

(3) The next ne
essary ondition, to remove moving singularities, is that the $-2\tilde{n}_x^2$ term is canceled, hence:

$$
\lambda \ge 1. \tag{78}
$$

(4) Now, if $\lambda = 1$ there are no systematic errors, but the sampling error very qui
kly obs
ures everything be cause n_x still heads to $-\infty$ exponentially due to the $2G\tilde{n}_x$ term. This takes it into regions of everincreasing $|\tilde{g}|$, and weights qui
kly be
ome randomized. For slightly larger parameters λ , the \tilde{n}_x evolution takes trajectories to a point lying far into the negative n_x region where the two leading terms balance. Here the trajectories sit, and quickly accumulate weight noise. It is clear that for an optimum simulation all stationary points of \tilde{n}_x in the nonzero gauge region must be removed. In this system this ondition is

$$
\lambda > 1 + \frac{G^2}{2Q} \,. \tag{79}
$$

An example has been plotted in Fig. 5 where we have parameters $G = 1, Q = 0.25$ (leading to $a \approx 1.1124$ and $b \approx -0.1124$). We are considering an initial condition of $\sigma_0^2 = 0.1$, which is already much larger than the canonical variance for physically likely parameters. Typical values of \tilde{n} initially will be of order $\sigma_{\tilde{n}} \approx 0.14 \gtrsim |b|$ here. A good choice of gauge has $\lambda = 4$. The use of this gauge learly restores the orre
t results.

Figure 5: One-mode laser $G = 1, Q = 0.25$. Dashed line: (correct) positive P simulation with $delta$ -function initial conditions (70) $\sigma_0^2 = 0$ and 10^5 trajectories. Dotted-dashed line: erroneous positive-P simulation with Gaussian initial onditions (71) $\sigma_0^2 = 0.1$ initially, and 10^5 trajectories. Dotted line: positive-P simulation with $\sigma_0^2 = 1$, and 10^4 trajectories. Solid line: gauge calculation for $\sigma_0^2 = 0.1$ with $\lambda = 4$, which corrects the systematic error of the positive P. Only 4000 trajectories, so as not to obs
ure other data. Step size in all ases is 0.005.

E. Non-optimal initial conditions

As we in
rease the spread of the initial distribution beyond $\sigma_{\tilde{n}} \approx |b|$, it becomes increasingly difficult to find a gauge that will give reasonable simulations. (For example we have tried a wide variety of what seemed like promising gauges for $\sigma_0^2 = 0.3$, with the above values of parameters Q and G , and none have come close to success). The problem is that while we can remove systemati errors, large random noise appears and obs
ures whatever we are trying to calculate.

Trajectories that start off at a value of \tilde{n} lying significantly beyond *b* require a lot of modification to their subsequent evolution to (1) stop them from es
aping to $-\infty$ and (2) move them out of the gauged region of phase space so that they do not accumulate excessive weight noise. If there are many of these, the trade-off between the gauge size and length of time spent in the gauged region does not give mu
h benet anymore. Nevertheless, one may be sure that if this is the ase, results will at worst be noisy and unusable, rather than being systemati
ally in
orre
t.

We stress again that this whole matter of non-optimal initial conditions is not a major hurdle to dynamical simulations be
ause a ompa
t starting distribution is generally found very easily.

VI. **CONCLUSIONS**

The positive P representation is well suited to complex quantum me
hani
al problems, su
h as many-body systems, but has been known for about a de
ade to have systematic errors in some cases of its use $-$ due to nonvanishing boundary terms. The gauge P representation, a variant on the usual positive-P representation, an be used to eliminate boundary terms and onsequently all the systematic errors that were encountered previously. It can also reduce sampling error in a simulation, and allows 'imaginary time' calculations of thermal equilibrium states. The fact that correct results are immediately obtained in every ase where systemati errors were found with the positive-P method, is strong evidence that these previous problems were indeed due to boundary terms caused by moving singularities in the analytically continued deterministi equations. Of ourse, boundary terms can occur for other reasons (for example, if the noise term grows too rapidly with radius), so aution is still needed in the gauge hoi
e.

The technique appears to be broadly applicable, and only requires the re
ognition of what instabilities in the stochastic equations could lead to problems. It does not require detailed knowledge of what the boundary terms are, provided instabilities are removed. However, we remark here that the general specification of necessary and sufficient conditions to eliminate boundary terms remains an open problem, and learly requires growth restri
tions on the gauge terms, both in phase spa
e and quantumamplitude space. Care is also required with the choice of the gauge and initial distribution. However, using unsuitable gauges or initial onditions may only lead to large sampling errors, not systemati errors, provided the gauge is hosen to eliminate boundary orre
tions in the first place. Sampling error then allows for a confident assessment of the magnitude of inaccuracies in a simulation, whi
h an be supplemented by numeri
al analysis of the distribution tails.

The main conclusion we come to is that this method does, in the cases studied, provide a complete solution to the problem of simulation of a many-body quantum system in phase spa
e, under onditions where previous direct simulation techniques were not practicable. All known te
hni
al requirements on the path to obtaining a sto
hasti
ally equivalent des
ription to quantum me
hanics, which is applicable to both large and small particle numbers, have been satisfied by this method. For this reason, we believe that gauge simulations an be used to simulate many quantum systems without systemati errors when carrying out more difficult calculations, where no exa
t result is known.

These conclusions must be supplemented by the detailed study of relevant gauges for particular quantum systems. We note, however, that the mathematical techniques employed here for generating sto
hasti gauges, may well be useful for other representations as well as the gauge P representation des
ribed here.

A
knowledgments

Numerical calculations were carried out using open software from the XMDS project [38]. Thanks to the Australian Resear
h Coun
il and the Alexander von Humboldt-Stiftung for providing resear
h support. We thank Damian Pope for suggesting the name sto
hasti gauges.

Appendix A: OTHER EXTENSIONS OF THE POSITIVE-P REPRESENTATION

1. The work of Carusotto, Castin, and Dalibard

Recently, Carusotto, Castin, and Dalibard [7, 8] (CCD) have made related extensions to the positive-P representation. These were derived for the particular case of an intera
ting s
alar Bose gas, and led to a number of onditions for an Ito sto
hasti evolution to be equivalent to a master equation evolution.

It an be shown quite simply that the equations (43) generated by the gauge P representation for this Hamiltonian satisfy the CCD conditions. We conjecture that these provide the most general possible solution to the stochastic problem posed by these authors. In particular, $db = \Pi[g_k dW_k - \bar{N}(\phi_1 dB_2^* + \phi_2^* dB_1)],$ using the above paper's formalism. Our methods an also treat a mu
h larger class of Hamiltonians and master equations than onsidered in the CCD treatment.

In Ref. [7] systematic errors due to boundary terms were not considered. However, evolutions satisfying "exactness" conditions derived using the same procedure can ontain su
h errors.

As an example, following the CCD procedure $[7]$ for a one-mode two-boson absorber master equation, as in Eq. (52) with $\gamma = \varepsilon = 0$, one arrives at the conditions

$$
dB_1 dB_2^* = 0,
$$

\n
$$
dB_a^{*2} = -\phi_a^2,
$$

\n
$$
F_1 = -db dB_1 / \Pi,
$$
\n(A1)

$$
F_2 = -db^* dB_2 / \Pi^* ,
$$

$$
f = \Pi(\bar{N}\phi_1\phi_2^*)^2 , \qquad (A2)
$$

where (referring back to the notation in this present paper),

$$
d\phi_1 = d\alpha/\sqrt{\overline{N}} = F_1 dt + dB_1,
$$

\n
$$
d\phi_2 = d\beta^*/\sqrt{\overline{N}} = F_2 dt + dB_2,
$$

\n
$$
d\Pi = d[\Omega e^{-\phi_1 \phi_2^* \overline{N}}] = f dt + db.
$$
 (A3)

It can be seen that the positive-P equations (54) satisfy these conditions, while producing the erroneous evolution seen in Fig. 1. In summary, the methods of the CCD paper do not obviate the need to hoose gauges that eliminate boundary terms.

2. Noise optimization by Plimak, Olsen, and Collett

In Ref. [9], Plimak, Olsen and Collett have found that for some systems (the Kerr oscillator $\hat{H}~=~\omega_0 \hat{a}^\dagger \hat{a}~+~$ $\kappa \hat{a}^{\dagger 2} \hat{a}^2/2$, in particular), the most obvious (diagonal) choice of noise matrix B may not be the optimal one.

For example, for the above Hamiltonian, one finds that the diffusion matrix (in α , β) variables is

$$
D = i\kappa \begin{bmatrix} -\alpha^2 & 0\\ 0 & \beta^2 \end{bmatrix} = BB^T .
$$
 (A4)

Following the pro
edure in Eq. (30), an equivalent but broader choice of noise matrix B can be any of

$$
B = \sqrt{i\kappa} \begin{bmatrix} i\alpha\cos(g) & i\alpha\sin(g) \\ -\beta\sin(g) & \beta\cos(g) \end{bmatrix}, \quad (A5)
$$

with the usual diagonal decomposition given by $q = 0$.

However, in Ref. $[9]$ it was found that for a positive-P simulation, different decompositions with nonzero constant g gave the lowest sampling error for coherent state initial onditions. In their notation, they introdu
e $\sqrt{A+1} = -\sqrt{2}\cos(g)$, and consider the case of real $A \geq 1$ (i.e., imaginary g) only.

3. Sto
hasti gauges for the Kerr os
illator

In Ref. $[2]$, the sampling error in a Kerr oscillator simulation $-$ equivalent to a one-mode BEC model, apart from linear terms $-$ was reduced substantially by using a representation similar to the gauge P representation formally introduced here. The basic differences were the following.

- 1. Instead of a complex gauge Ω , a phase factor $e^{i\theta}$ with a real θ variable, was used.
- 2. The normalization with respe
t to the behavior of θ was carried out explicitly inside the kernel, rather than post-simulation in the moments as in Eq. (45).

This type of representation is a norm-preserving gauge P representation, as dis
ussed earlier. A parametrized family of gauges led to stable trajectories (as opposed to the large sampling error present with a positive-P simulation). However, some systemati errors were seen due to boundary terms. These boundary terms occurred because of the stochastic growth of the gauge term in Ω space, when θ approached $\pm \pi/2$. With the gauge P representation introdu
ed in this paper, a wide range of gauges do not lead to any systematic errors [33], provided gauge growth is ontrolled.

We note here that the norm-preserving gauges have the property that, in the present notation, $g_k = i[1 - i\Omega'']f_k$. However, while the growth of $Ω'$ is stabilized, there

is growth in the variance of Ω'' . This means that the function f_k must behave as a decreasing function of Ω'' in order to ensure that the distribution is bounded sufficiently in the weight-function space to avoid finite boundary terms. The detailed requirements and onditions for this type of gauge will be treated elsewhere.

- [1] S. Chaturvedi, P. D. Drummond, and D. F. Walls, J. Phys. A 10, L187 (1977); P. D. Drummond and C. W. Gardiner, J. Phys. A 13, 2353 (1980).
- [2] P. Deuar and P. D. Drummond, Comput. Phys. Comm. 142, 442 (2001).
- [3] P. D. Drummond and P. Deuar (To be published).
- [4] A. M. Smith and C. W. Gardiner, Phys. Rev. A $39, 3511$ (1989).
- [5] R. Schack and A. Schenzle, Phys. Rev. A 44, 682 (1991).
- [6] A. Gilchrist, C. W. Gardiner, and P. D. Drummond, Phys. Rev. A 55, 3014 (1997).
- [7] I. Carusotto, Y. Castin, and J. Dalibard, Phys. Rev. A 63, 023606 (2001).
- [8] I. Carusotto and Y. Castin, J. Phys. B 34 , 4589 (2001).
- [9] L. I. Plimak, M. K. Olsen, and M. J. Collett, Phys. Rev. A 64, 025801 (2001).
- [10] K. G. Wilson, Phys. Rev. D 10, 2445 (1974); M. Creutz, Phys. Rev. D 21, 2308 (1980).
- [11] E. L. Pollock and D. M. Ceperley, Phys. Rev. B $30, 2555$ (1984); D. M. Ceperley, Rev. Mod. Phys. 67, 279 (1995); D. M. Ceperley, Rev. Mod. Phys. 71, S438 (1999).
- [12] R. P. Feynman, Int. J. Theor. Phys. 21, 467 (1982).
- [13] The current maximum size is 7 quantum bits in a dissipative, non-s
alable te
hnology, giving a Hilbert-spa
e dimension of 128 without error correction. See $-$ L. M. K. Vandersypen, M. Steffen, G. Breyta, C. S. Yannoni, M. H. Sherwood, and I. L. Chuang, Nature 414, 883 $(2001).$
- [14] J. Von Neumann, Mathematical foundations of quantum mechanics Translated from the German edition by Robert T. Beyer (Prin
eton University Press, Prin
eton, N.J. , 1983).
- [15] E. P. Wigner, Phys. Rev. 40, 749 (1932).
- [16] R. J. Glauber, Phys. Rev. $131, 2766$ (1963).
- [17] E. C. G. Sudarshan, Phys. Rev. Lett. $10, 277$ (1963).
- [18] K. Husimi, Proc. Phys. Math. Soc. Jpn. 22, 264 (1940).
- [19] K. E. Cahill and R. J. Glauber, Phys. Rev. 177, 1882 (1969).
- [20] M. J. Steel, M. K. Olsen, L. I. Plimak, P. D. Drummond, S. M. Tan, M. J. Collett, D. F. Walls, and R. Graham, Phys. Rev. A 58, 4824 (1998).
- [21] S. J. Carter, P. D. Drummond, M. D. Reid, and R. M. Shelby, Phys. Rev. Lett. 58, 1841 (1987);

P. D. Drummond and S. J. Carter, J. Opt. So
. Am. B 4, 1565 (1987); P. D. Drummond, R. M. Shelby, S. R. Friberg, and Y. Yamamoto, Nature 365, 307 (1993).

- [22] S. J. Carter and P. D. Drummond, Phys. Rev. Lett 67, 3757 (1991).
- [23] M. J. Werner, Phys. Rev. A 54 , R2567 (1996).
- $[24]$ P. D. Drummond and J. F. Corney, Phys. Rev. A 60, R2661 (1999).
- [25] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell, Science 269, 198 (1995) . \sim
- [26] C. C. Bradley, C. A. Sackett, J. J. Tollett, and R. G. Hulet, Phys. Rev. Lett. 75, 1687 (1995).
- $[27]$ K. B. Davis, M. -O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle, Phys. Rev. Lett. 75, 3969 (1995).
- [28] J. R. Klauder and E. C. G. Sudarshan, Fundamentals of Quantum Opti
s (Benjamin, New York, 1970).
- [29] L. Arnold, Stochastic Differential Equations: Theory and Applications (Wiley, New York, 1974); C. W. Gardiner, Handbook of Stochastic Methods, 2nd ed. (Springer, Berlin, 1985).
- [30] R. Graham, Z. Phys. B 26, 281 (1977); H. Leschke and M. S
hmutz, Z. Phys. B 27, 85 (1977).
- [31] S. Caratzoulas and P. Pechukas, J. Chem. Phys. 104, 6265 (1996).
- [32] P. D. Drummond and I. K. Mortimer, J. Comput. Phys. 93, 144 (1991).
- [33] P. Deuar and P. D. Drummond (To be published).
- [34] For a recent treatment, see, e.g., A. J. Leggett, Rev. Mod. Phys 73, 307 (2001).
- [35] I. J. D. Craig and K. J. McNeil, Phys. Rev. A 39, 6267 (1989).
- [36] K. J. McNeil and I. J. D. Craig, Phys. Rev. A 41, 4009 (1990).
- [37] C. W. Gardiner, A. Gilchrist, and P. D. Drummond, in Using the Positive P Representation, edited by F. Ehlotzky, Lecture Notes in Physics, Vol. 420 (Springer-Verlag, Berlin, 1993), p. 239.
- [38] G. R. Collecutt and P. D. Drummond http://www. physi
s.uq.edu.au/xmds/