On the coherence verification of the continuous variable state in Fock space

Wang Xiangbin ∗ , Matsumoto Keiji † and Tomita Akihisa ‡

Imai Quantum Computation and Information project, ERATO, Japan Sci. and Tech. Corp.

Daini Hongo White Bldg. 201, 5-28-3, Hongo, Bunkyo, Tokyo 113-0033, Japan

Abstract

Almost all novel observable phenomena in quantum optics are related to the quantum coherence. The coherence here is determined by the relative phase inside a state. Unfortunately, so far all the relevant experimental results in quantum optics are insensitive to the phase information of the coherent state. Lack of phase information may cause serious consequences in many problems in quantum optics. For example, an ensemble of two mode squeezed states is a classical ensemble if the phase in each state is totally random; but it is a non-classical ensemble if the phase in each state is fixed. As a timly application, verification of this type of phase information in an ensemble of two mode squeezed states from the conventional laser is crucial to the validity of the continuous variavable quantum teleportation(CVQT) experiment. Here we give a simple scheme to distinguish two different ensemble of states: the Rudolph-Sanders ensemble, by which each squeezed states emitted has a uniform distribution from $0 - 2\pi$ on the phase value; and the van Enk-Fuchs ensemble, which emmits identical states with a fixed(but unknown) phase for every state. We believe our proposal can help to give a clear picture on whether the existing two mode squeezed states so far are indeed non-classical states which can be used as the entanglement resource.

[∗] email: wang@qci.jst.go.jp

[†] email: keiji@qci.jst.go.jp

[‡] email: a-tomita@az.jp.nec.com

Quantum coherence of different states plays a fundamentally important role in the whole subject of quantum optics. The squeezed states have been widely used to demonstrate various novel non-classical properties in the past. In particular, a two mode squeezed state

$$
|r, \phi\rangle = e^{re^{i\phi}(a_1a_2 - a_1^{\dagger}a_2^{\dagger})}|00\rangle = \sum_{k=0}^{\infty} \lambda^k e^{ik\phi}|00\rangle, \tag{0.1}
$$

where a_i^{\dagger} and a_i are bosonic creation and annihilation operators respectively in the two mode Fock space, $|00\rangle$ is the vaccum state, is hopefully to be used as the entangled resource to carry out various novel tasks in the creteria of quantum information[[1](#page-6-0)] such as the quantum teleportation [\[2–4\]](#page-6-0). Recently, this state has been used as the enatnglement resource to teleport a quantum state between two spatially separated parties [\[4](#page-6-0)]. However, as it is pointed out by Rudolph and Sanders [\[5](#page-6-0)] that the the phase information ϕ here has never been tested, therefore states as defined by the above equation from certain source could have different phase ϕ for each different wavepackets. The lack of the phase information causes the loss of coherence of the state. Although lots of experiments have been done sucessfully to reconstruct the continuous variable states in the past, none of them is related to the phase information. That is to say, a pure state is not the only possible state that is compatible with the observed results. As it is noted [\[5](#page-6-0)] that, a conventional meassurement on optical fields, such as homodyne detection using lasers, involving mixing of defferent incoherent fields and subsquent detection by energy absorption in photodetectors: all such meassurements are completely insensitive to any optical coherence. Actually, the state could be in arbitrary classical probabilistic distribution via the different phases. If we use the random uniform distribution, the observed state is actually a mixed diagonal state in Fock space[[5,6\]](#page-6-0), which is given by

$$
\frac{1}{2\pi} \int_0^{2\pi} |r, \phi \rangle \langle r, \phi | d\phi = \sum_0^{\infty} \lambda^{2n} |nn \rangle \langle nn|.
$$
 (0.2)

Definitely, there is neither quantum coherence nor quantum entanglement in the above state. The state defined by eq(0.2) is a totally classical state. If the phase ϕ of each states from certain source are indeed uniformly distributed from zero to 2π , the so called squeezed states will have little novel properties in the practical use because in such a case all quantum coherence has been lost and all the observable phenomena are just the same as that given by classical optics. For example, we can never really take the advantage of its squeezing property for certain quadrature variable(although a certain quadrature operator ie indeed squeezed on a single wavepacket).

Due to the lack of the phase information, the observable quantum coherence property have never been verified on an ensemble of squeezed states. To answer the question whether the conventional source can produce the nontrivil quantum coherent state we must have a way to detect the phase information ϕ .

A timly application for this type detection is on the validity of the CVQT experiment [\[4](#page-6-0)]. Recently, the phase information of the two mode squeezed states has drawn much attention

of the physicists due to the issue of quantum teleportation of the continuous variable state in Fock space [\[4](#page-6-0)]. Since the tomography result is independent of the phase information ϕ , it is also possible that the the state used as the entanglement source in the quantum teleportation experiment is a mixture of the states with different ϕ , as defined by eq.([0.2\)](#page-1-0). If this is the case, then no quantum state can be teleported by such a separable state. Definitely, an ensemble of N identical copies of pure states as defined by $|r, \phi \rangle \langle r, \phi \rangle^{\otimes N}$ with the unknown ϕ (but all states in the ensemble have the same ϕ) is totally different from the ensemble as defined by eq.([0.2\)](#page-1-0). Unfortunetely, so far there has been no way to distinguish these two totally different cases.

Due to the unclearity of the phase ϕ , there is a very hot discussions on the validity of the entanglement resource used in the CVQT experiment[[5,7–9](#page-6-0)]. The discussions can be sumarrized as the following:

1. Rudolph and Sanders: The phase ϕ in each of the squeezed states from a conventional laser source are uniformly distributed from zero to 2π . The correct form of the quantum state for the ensemble is given by $eq(0.2)$ $eq(0.2)$ $eq(0.2)$

2. van Enk and Fuchs: The traditional formalism is insufficient to describe the meachanism two mode squeezed states produced by the conventional laser source. Using quantum de Finetti theorem one can see that the two mode squeezed states produced by a conventional laser is essentially an ensemble of many coppies of identical two mode squeezed states with a fixd ϕ , though the value of ϕ is unknown.

We may see that the validity of the CVQT experiment is now reduced to which of the above statements correctly describe the property of the source which produses the two mode squeezed states in the CVQT experiment. That is to say, to know the validity of CVQT experiment, we have to distinguish the Rudolph-Sanders source and the van Enk-Fuchs source. In this letter we give a scheme to detect the phase information ϕ in the two mode squeezed state. That is to say, the meassurement result by our scheme is sensitive to the quantum coherence. Using our scheme, the ensemble of states defined by $eq(0.1)$ $eq(0.1)$ or $eq(0.2)$ $eq(0.2)$ can be easily distinguished. Obviously, the scheme has broad potential applications in the whole subject of quantum optics. For example, given many copies of pure squeezed states, we can verify that they are indeed pure states with the same phase ϕ . For another example, given two Fuch sources, each source emmits many identical two mode squeezed states, the phase of the state from each source ϕ_1 and ϕ_2 respectively, they are fixed and unknown, our scheme can be used detect the value $\phi_1 - \phi_2$. As an immediate application, the scheme can be used to judge whether the states used as the entangled resource in a recent quantum teleportation [\[4](#page-6-0)] is the pure state as defined in eq. (0.1) (0.1) or a mixed state defined by eq. (0.2) (0.2) . Consequently, this detection could give a a clear judgement on whether the experiment done in ref.[[4](#page-6-0)] is essentially a quantum teleportation or a classical simulation of quantum teleportation.

Now we show how to distinguish a Roudolph-Sanders source and a van Enk-Fuchs source.

Lets first see what happens to the van Enk-Fuchs source, i.e. a photon source emmits the identical pure squeezed states, all of them have the same but unknown phase ϕ . As it is notedin ref [[10](#page-6-0)] that a two mode squeezed state can be produced with the specifically chosen polarization for each modes as the following

$$
|\psi_1(\phi)\rangle = \sum_{l=0}^{\infty} \lambda^l e^{il\phi} |l\rangle_{ah} |l\rangle_{bv},
$$
\n(0.3)

where the subscripts a, b are for the mode a and b respectively, h and v represents the horizontal and vertical polarizations respectively. Now we consider another wave packet from the same source, of which the mode a and b is exchanged. The quantum state for this wavepacket is $|\psi_2(\phi)\rangle = \sum_{m=0}^{\infty} \lambda^m e^{im\phi} |m\rangle_{av} |m\rangle_{bh}$. The total state is then given by

$$
|\Psi\rangle = |\psi_1\rangle |\psi_2\rangle = \sum_{l=0}^{\infty} \sum_{m=0}^{\infty} \lambda^{l+m} e^{i(l+m)\phi} |l\rangle_{ah} |m\rangle_{av} |l\rangle_{bv} |m\rangle_{bh} . \qquad (0.4)
$$

After taking a measurement on the photon number of each mode, i.e., on the quantity $n = l+m$, the state $|\Psi\rangle$ is collapsed to a specific entangled state. As it has been demonstrated in ref [[10\]](#page-6-0), this type of meassurement cab be carried out by either a quantum non-demolition meas-surement[[11\]](#page-6-0) or by a more feasible way, the destructive photon counting with post selection. Suppose the meassurement result is n for each mode, then the state after the meassurement is $\sum_{m=0}^{n} e^{i[(n-m)\phi+m\phi]}|n-m\rangle_{ah}|m\rangle_{av}|n-m\rangle_{bv}|m\rangle_{bh}$. The phase information is clearly included in the state after the measurement. For simplicity, we consider only the cases of $n = 1$ only. In the real experiment we can set an appropriate value of λ so that we have a significant probability to get the result of $n = 1$. With such a setting, given many coppies of state $|\Psi\rangle$, we can always have a significant number of states with $n = l + m = 1$ after the meassurement. In the case of $n = 1$, the state is

$$
|\Psi_1 \rangle = e^{2i\phi} (|1 \rangle_{ah} |0 \rangle_{av} |1 \rangle_{bv} |0 \rangle_{bh}) + |0 \rangle_{ah} |1 \rangle_{av} |0 \rangle_{bv} |1 \rangle_{bh} . \tag{0.5}
$$

There is only one photon in each mode for the satte defined above. Obviously, the state defined above can be rewritten in the following way

$$
|\Psi_1 \rangle = e^{2i\phi} (|H \rangle_a |V \rangle_b + |V \rangle_a |H \rangle_b).
$$
\n(0.6)

Where the states $|H \rangle$, $|V \rangle$ are for a horizontal or a vertical polarized photon states respectively. For this state, meassurement results for the polarization of the two modes are always different. Clearly, if the states given from the source are indeed pure states, i.e., they are many coppies of $|\Psi_1\rangle$, then we can rotate the polarizers by the same angle to both modes, and in principle we can find certain angle($\pi/4$) by which the meassurement result of the polarization in the two modes are always same ! On the other hand, if the given states from the source are mixed states as defined in eq([0.2\)](#page-1-0), i.e., phase ϕ in each wave packet can be different, we will have a clearly different observation results. In such a case, by the same operation, for all cases we get the meassurement of $l + m = 1$, the state is $\frac{1}{2}(|HV\rangle \langle HV| + |VH\rangle \langle VH|)$. The correlation between the two modes will linearly decreased with $\cos^2 \beta$ when the polarizers are rotated, here β is the angle that is rotated. Consequently, a van Enk-Fuchs source and a Rudolph-Sanders source can be distinguished in the following way:

Initially, in both cases they are totally negatively correlated, i.e., whenever a photon is detected in mode a , there most be no photon detected in mode b , and vice versa. We rotate the polarizers by $\pi/4$, the van Enk-Fuchs source(see eq([0.1\)](#page-1-0)) will give a totally positive correlation for the meassurement result, while the Rudolph-Sanders source(see $eq(0.2)$ $eq(0.2)$ $eq(0.2)$) will give no correlation for the meassurement. Also, in rotating the polarizers, if the correlation does not decrease linearly with $\cos^2 \beta$, the source must be not a Rudolph-Sanders one.

Obviously, the above scheme can be also used to detect the phase difference for two van Enk-Fuchs sources. Now $|\psi_1\rangle$ and $|\psi_2\rangle$ are collected from two different sources whose unknown fixed phase are denoted by ϕ_1 and ϕ_2 respectively. After the meassurement on the photon number basis of each mode is done, the state up to a global phase factor is

$$
|\Psi_1 \rangle = |H \rangle |V \rangle + e^{i(\phi_2 - \phi_1)} |V \rangle |H \rangle. \tag{0.7}
$$

We rotate the polarizers continuously and test the correlations at each stage. There must be certain angle β_0 at which we can observe that the meassurement results of the two modes are totally positively correlated. This angle β_0 determines the phase difference $\phi_2 - \phi_1$.

Note that in a real experiment, we actually do not need to take a meassurement on the basis of of the photon numbers of each mode, i.e. $\sum_{n=0}^{\infty} |n \rangle \langle n|$, where $n = l + m$. The meassure on such a basis could be difficult by our current technology. The only subtle task for us is to collect the wave packet $|\psi_1\rangle$ and $|\psi_2\rangle$ so that they are indistiuguishable. Once we can make $|\psi_1\rangle |\psi_2\rangle$ indistinguishable we can simply carry out the detection scheme by the usual photon counting and obtain the correct result to a good approximation. Suppose we use port A and port B to detect the photons from mode A and B respectively. A polarizer is placed before each port. The polarizers are placed horizontally in the begining. We choose a very small r so that $r^2 \ll 1$ for the photon source. In the photon counting, we only record the data of those events where at least one port detects one photon and no port detects more than one photon. For simplicity, we name the events satisfying this condition as "good events". We will analyse the correlation between the two modes only using the data of the good events. In the whole process, there is only a very small probability that a good event is caused by a state with $l + m \neq 1$, i.e., the wave packet has been collapsed to a state of which the photon number of each mode is not 1. It's easy to see that this type of event can only happen with a small ptobability. First, $l + m = 0$ is impossible, because whenever we observed a good event, we have observed one photon at least in one port, so the photon number of that mode must not be 0, consequently $l + m \neq 0$. Second, $l + m = 2$ or a larger number is possible but the

chance is negligible. We have already set r to be very small. Since $r^2 \ll 1$, the probability that the state $|\Psi\rangle$ collapsed to a state with $l+m=1$ is much larger than that of a state with $l + m > 1$. For example, taking $r = 0.01$, in average, we can have one good event from 10000 events, and the probability that this good event is caused by a stete with $l + m = 1$ is several thousands times larger than that by a state with $l+m \neq 1$. Actually, to a good approximation, we even do not have to require the photon detector to distinguish 1 photon case from many photon case in the detection if r is small enough. Because once the detector detects photons, the probability of 1 photon is much higher than other cases. For clarity, we insist on using the term "1 photon" in the rest parts of the letter. However, whenever the term "one photon" is used, it can be approximately understood as "any number of photons".

Initially, all good evnts must be the events on which 1 photon detected in certain mode and 0 photon is detected in another mode. If the source is the van Enk-Fuchs source, when the polarizers are rotated by $\pi/4$, a photon in each mode must be detected to all good events. If the source is the Rudolph-Sanders source, when the polarizers are rotated by $\pi/4$, to all good events, half of them are $(1_A, 0_B)$ or $(0_A, 1_B)$ and half of them are $(1_A, 1_B)$, where the 0 or 1 represents the number of photons detected, subscripts A, B represent the the port A, B (or the mode a, b respectively.

The phase difference can also be detected for two Enk-Fuchs sources. In this case, we have to rotate the polarizers continuously. The phase difference $\phi_2-\phi_1$ is determined by the angle β_0 we observed 1 photon in each mode to all good events. Thus we see, to a good approximation, our scheme can carried out by the normal photon counting technique.

Acknowledgement: We thank Prof Imai for support.

REFERENCES

- [1] Nielsen M A and Chuang I L, Quantum computation and information,Cambridge university press, 2000.
- [2] C.H. Bennette et al, Teleporting an unknown quantum state via dual classical and EPR channels. Phys. Rev. Lett., 70:1895-1999(1993).
- [3] Bowmeester D et al, Experimental quantum teleportation, Nature, 390, 6660(1997).
- [4] A. Furusawa et al, Unconditional quantum teleportation. Science 282, 706(1998).
- [5] T. Rudolph and Barry C. Sanders., Requirement of optical coherence for continuousvariable quantum teleportation, Phys. Rev. Lett., 87, 077903(2001).
- [6] K. Molmer, Phys. Rev. A 55, 3195(1997).
- [7] S.J. van Enk and C.A. Fuchs, Quantum state of an ideal propagating laser field, Phys. Rev. Lett., 88, 027902(2002).
- [8] S.J. van Enk and C.A. Fuchs, Quantum state of an ideal propagating laser field, arXiv: [quant-ph/0111157.](http://arxiv.org/abs/quant-ph/0111157)
- [9] Rudolph T and Barry C.S., Comment on "Quantum state of an ideal propagating laser field", [arXiv:quant-ph/0112020](http://arxiv.org/abs/quant-ph/0112020).
- [10] Gabirel A Durkin, C. Simon and D. Bouwmeester, Multi-photon Entanglement Concentration and Quantum Cryptography. arXiv: [quant-ph/0109132](http://arxiv.org/abs/quant-ph/0109132)
- [11] Duan L M et al, Entanglement purification of Gaussian continuous variable quantum states, Phys. Rev. Lett., 84, 4002(2000).