## Efficient Simulation of Quantum State Reduction

Dorje C. Brody \* and Lane P. Hughston <sup>†</sup>

\*Blackett Laboratory, Imperial College, London SW7 2BZ, UK

<sup>†</sup>Department of Mathematics, King's College London, The Strand, London WC2R 2LS, UK

(October 29, 2018)

The energy-based stochastic extension of the Schrödinger equation is a rather special nonlinear stochastic differential equation on Hilbert space, involving a single free parameter, that has been shown to be very useful for modelling the phenomenon of quantum state reduction. Here we construct a general closed form solution to this equation, for any given initial condition, in terms of a random variable representing the terminal value of the energy and an independent Brownian motion. The solution is essentially algebraic in character, involving no integration, and is thus suitable as a basis for efficient simulation studies of state reduction in complex systems.

PACS Numbers : 03.65.Ta, 02.50.Cw, 02.50.Ey

The standard energy-based stochastic extension of the Schrödinger equation is given by the following stochastic differential equation:

$$d|\psi_t\rangle = -i\hat{H}|\psi_t\rangle dt - \frac{1}{8}\sigma^2(\hat{H} - H_t)^2|\psi_t\rangle dt + \frac{1}{2}\sigma(\hat{H} - H_t)|\psi_t\rangle dW_t, \qquad (1)$$

with initial condition  $|\psi_0\rangle$ . Here  $|\psi_t\rangle$  is the state vector at time t,  $\hat{H}$  is the Hamiltonian operator,  $W_t$  denotes a one-dimensional Brownian motion, and

$$H_t = \frac{\langle \psi_t | H | \psi_t \rangle}{\langle \psi_t | \psi_t \rangle} \tag{2}$$

is the expectation of  $\hat{H}$  in the state  $|\psi_t\rangle$ . The parameter  $\sigma$ , which has the units  $\sigma \sim [\text{energy}]^{-1}[\text{time}]^{-1/2}$ , governs the characteristic timescale  $\tau_R$  associated with the collapse of the wave function induced by (1). This is given by  $\tau_R = 1/\sigma^2 V_0$ , where  $V_0$  is the initial value of the squared energy uncertainty, which at time t is

$$V_t = \frac{\langle \psi_t | (\hat{H} - H_t)^2 | \psi_t \rangle}{\langle \psi_t | \psi_t \rangle}.$$
(3)

The stochastic equation (1) provides perhaps the simplest known physically plausible model for state vector reduction in quantum mechanics [1,2]. Although its properties have been studied extensively, it has hitherto been necessary to resort to numerical methods to solve (1). The purpose of this article is to present an analytic solution for the dynamics of  $|\psi_t\rangle$ . Apart from its use as a means for generating a general solution to a nonlinear problem in quantum state dynamics, the method we propose also sheds new light on the nature of quantum probability and some of the issues associated with the flow of information when quantum measurements are made.

We begin with a brief overview of the stochastic framework implicit in the extended Schrödinger dynamics given by equation (1). We follow closely here the analysis presented in [3]. Specifically, we introduce first the key notions of filtration, conditional expectation, martingale, and potential. We then demonstrate that the conditional expectation (10) gives rise to the energy expectation process (2). As a consequence, we are led to simple analytic expressions for the energy (15) and the state vector (22) in terms of a pair of underlying state variables. These results open up the possibility of efficiently simulating the reduction process for a variety of models. Finally we illustrate the practical advantages of our method by analysing in some detail the timescale associated with the reduction process in the case of a two-state system.

The dynamics of  $|\psi_t\rangle$  are defined on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  with filtration  $\mathcal{F}_t$   $(0 \leq t < \infty)$ . Here  $\Omega$  is the sample space over which  $\mathcal{F}$  is a  $\sigma$ -field of open sets upon which the probability measure  $\mathbb{P}$  is defined.

The filtration represents the information available at time t. More specifically, a filtration of  $\mathcal{F}$  is a collection  $\{\mathcal{F}_t\}$  of  $\sigma$ -subfields of  $\mathcal{F}$  such that  $s \leq t$  implies  $\mathcal{F}_s \subset \mathcal{F}_t$ . Given a random variable X on  $(\Omega, \mathcal{F}, \mathbb{P})$ for which  $\mathbb{E}[X]$  exists, we write  $\mathbb{E}[X|\mathcal{F}_t]$  for the conditional expectation of X with respect to the  $\sigma$ -subfield  $\mathcal{F}_t \subset \mathcal{F}$ . Intuitively, conditioning with respect to  $\mathcal{F}_t$ means giving the information available up to time t. The nesting  $\mathcal{F}_s \subset \mathcal{F}_t$  for  $s \leq t$  thus gives rise to a notion of causality. For convenience, we use the abbreviation  $\mathbb{E}_t[X] = \mathbb{E}[X|\mathcal{F}_t]$  when the choice of  $\{\mathcal{F}_t\}$  is understood. The conditional expectation  $\mathbb{E}_t[X]$  satisfies: (i) the tower property  $\mathbb{E}_s[\mathbb{E}_t[X]] = \mathbb{E}_s[X]$  for  $s \leq t$ ; and (ii) the law of total probability  $\mathbb{E}[\mathbb{E}_t[X]] = \mathbb{E}[X]$ . If  $\mathbb{E}_t[X] = X$  we say that X is  $\mathcal{F}_t$ -measurable.

The conditional expectation operation allows us to introduce the concept of a martingale, the stochastic analogue of a conserved quantity. A process  $X_t$  is said to be an  $\mathcal{F}_t$ -martingale if  $\mathbb{E}[|X_t|] < \infty$  and  $\mathbb{E}_s[X_t] = X_s$  for all  $0 \le s \le t < \infty$ . In other words,  $X_t$  is an  $\mathcal{F}_t$ -martingale if it is integrable and if its conditional expectation, given information up to time s, is the value  $X_s$  of the process at that time. If the filtration is fixed, then we can simply speak of a martingale without further qualification. There are circumstances, however, where more than one filtration can enter a problem, and then we have to specify with respect to which filtration the martingale property holds.

For a concise mathematical representation of the state reduction process, we also require the concepts of supermartingale and potential. A process  $X_t$  is an  $\mathcal{F}_t$ supermartingale if  $\mathbb{E}[|X_t|] < \infty$  and  $\mathbb{E}_s[X_t] \leq X_s$  for all  $0 \leq s \leq t < \infty$ . Intuitively, a supermartingale is on average a nonincreasing process. A positive supermartingale  $X_t$  with the property  $\mathbb{E}[X_t] \to 0$  as  $t \to \infty$  is called a potential.

The filtration  $\mathcal{F}_t$  with respect to which stochastic extension of the Schrödinger equation (1) is defined is generated in a standard way by the Wiener process  $W_t$ . We signify this by writing  $\mathcal{F}_t = \mathcal{F}_t^W$ . It is straightforward to verify that the Hamiltonian process  $H_t$  is an  $\mathcal{F}_t^W$ -martingale, and that the variance process  $V_t$  is an  $\mathcal{F}_t^W$ -supermartingale. That is to say,  $\mathbb{E}_s[H_t] = H_s$ , and  $\mathbb{E}_s[V_t] \leq V_s$ . These relations can be deduced by applying Ito's lemma to (2) and (3), from which we infer that

$$H_t = H_0 + \sigma \int_0^t V_s \mathrm{d}W_s \tag{4}$$

and

$$V_t = V_0 - \sigma^2 \int_0^t V_s^2 \mathrm{d}s + \sigma \int_0^t \beta_s \mathrm{d}W_s.$$
 (5)

Here  $\beta_t = \langle \psi_t | (\hat{H} - H_t)^3 | \psi_t \rangle / \langle \psi_t | \psi_t \rangle$  is the skewness of the energy distribution at time t. The martingale and the supermartingale relations then follow as a consequence of elementary properties of the stochastic integrals appearing in (4) and (5).

In the case of the ordinary Schrödinger equation with a time-independent Hamiltonian, the energy process (2) is constant. This is usually interpreted as the quantum mechanical expression of an energy conservation law. However, if a system is in an indefinite state of energy then it is not clear a priori what is meant by energy conservation. The martingale condition  $\mathbb{E}_s[H_t] = H_s$  can be interpreted as a generalised energy conservation law applicable in such circumstances. In particular, it implies that once the state reduction has occurred, the probabilistic average of the outcome for the energy must equal the initial expectation.

The supermartingale property satisfied by  $V_t$  on the other hand is the essence of what is meant by a reduction process. In fact, it follows from equation (5) that the asymptotic behaviour of  $V_t$  is given by  $\lim_{t\to\infty} \mathbb{E}[V_t] = 0$ . In other words, the variance process for the energy is a potential. Writing  $H_{\infty} = H_0 + \sigma \int_0^\infty V_t dW_t$  for the random terminal value of the energy, one can prove [3] as a consequence of (4) and (5) that

$$H_t = \mathbb{E}_t[H_\infty] \tag{6}$$

and that

$$V_t = \mathbb{E}_t[(H_\infty - H_t)^2]. \tag{7}$$

That is to say,  $H_t$  and  $V_t$  are respectively the  $\mathcal{F}_t^W$ -conditional mean and variance of  $H_{\infty}$ .

With these facts in hand, we now present a method for obtaining a general solution to the stochastic equation (1). The setup is as follows. We denote by  $E_i$  (i = 1, 2, ...) the eigenvalues of the Hamiltonian of a given quantum system, and write

$$\pi_i = \frac{|\langle \psi_0 | \psi_i \rangle|^2}{\langle \psi_0 | \psi_0 \rangle \langle \psi_i | \psi_i \rangle},\tag{8}$$

for the transition probability from the given initial state  $|\psi_0\rangle$  to the eigenstate  $|\psi_i\rangle$  with energy  $E_i$ . If the spectrum of  $\hat{H}$  is degenerate, then  $|\psi_i\rangle$  denotes the Lüders state, i.e. the projection of  $|\psi_0\rangle$  onto the linear subspace of states corresponding to the eigenvalue  $E_i$ .

Now let the probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  be given, and on it specify a random variable H that takes the values  $E_i$  with probabilities  $\pi_i$ . We also assume that  $(\Omega, \mathcal{F}, \mathbb{P})$ comes equipped with a filtration  $\mathcal{G}_t$  with respect to which a standard Brownian motion  $B_t$  is specified, and that Hand  $B_t$  are *independent*. We assign no *a priori* physical significance to H and  $B_t$ , which are introduced as an ansatz for obtaining a solution for (1).

We now define a random process  $\xi_t$ , which we shall call the signal process, according to the scheme

$$\xi_t = \sigma H t + B_t. \tag{9}$$

Intuitively, one can think of  $\xi_t$  as giving a 'noisy' representation of the information encoded in the random variable H.

We let  $\{\mathcal{F}_t^{\xi}\}$  denote the filtration generated by the process  $\xi_t$ , i.e. the information generated by  $\xi_t$  as time progresses, and consider the conditional expectation

$$H_t = \mathbb{E}\left[H|\mathcal{F}_t^{\xi}\right].$$
 (10)

Clearly,  $\mathcal{F}_t^{\xi} \subset \mathcal{G}_t$  since knowledge of H together with  $\{B_s\}_{0 \leq s \leq t}$  implies knowledge of  $\{\xi_s\}_{0 \leq s \leq t}$ , although the converse is not the case.

The significance of the  $\mathcal{F}_t^{\xi}$ -martingale  $H_t$  is that it represents the 'best estimate' for the value of H given the history of the signal process  $\xi_s$  from time 0 up to time t. More precisely, an  $\mathcal{F}_t^{\xi}$ -measurable random variable  $Y_t$  minimises the expectation of the squared deviation of H from  $Y_t$ , given  $\mathcal{F}_t^{\xi}$ , iff  $Y_t = \mathbb{E}[H|\mathcal{F}_t^{\xi}]$ . This follows, by a variational argument, from the relation  $\mathbb{E}[(H - Y_t)^2|\mathcal{F}_t^{\xi}] = \mathbb{E}[H^2|\mathcal{F}_t^{\xi}] - 2Y_t\mathbb{E}[H|\mathcal{F}_t^{\xi}] + Y_t^2$ .

We proceed to establish the remarkable fact that the process  $H_t$  defined by (10) is statistically indistinguishable from the energy process (2) associated with the stochastic extension of the Schrödinger equation (1).

The argument goes as follows. First, because  $\xi_t$  is a Markov process satisfying  $\lim_{t\to\infty} t^{-1}\xi_t = H$ , we have

$$\mathbb{E}\left[H|\mathcal{F}_t^{\xi}\right] = \mathbb{E}\left[H|\xi_t\right]. \tag{11}$$

In other words, to determine the conditional expectation of H given  $\{\xi_s\}_{0 \le s \le t}$  it suffices to condition on  $\xi_t$  alone.

To calculate  $\mathbb{E}[H|\xi_t]$ , we require a version of the Bayes formula applicable when we consider the probability of a discrete random variable conditioned on the value of a continuous random variable. In particular,

$$\mathbb{P}(H = E_i | \xi_t) = \frac{\pi_i \rho(\xi_t | H = E_i)}{\sum_i \pi_i \rho(\xi_t | H = E_i)},$$
(12)

where  $\pi_i = \mathbb{P}(H = E_i)$ . Here  $\rho(\xi_t | H = E_i)$  denotes the conditional probability density for the continuous random variable  $\xi_t$  given that  $H = E_i$ . Since  $B_t$  is a standard Brownian motion, the conditional density for  $\xi_t$  is

$$\rho(\xi_t | H = E_i) = \frac{1}{\sqrt{2\pi t}} \exp\left(\frac{1}{2t} (\xi_t - \sigma E_i t)^2\right).$$
(13)

It follows from the Bayes law (12) that the conditional probability for the random variable H is

$$\mathbb{P}(H = E_i | \xi_t) = \frac{\pi_i \exp\left(\sigma E_i \xi_t - \frac{1}{2} \sigma^2 E_i^2 t\right)}{\sum_i \pi_i \exp\left(\sigma E_i \xi_t - \frac{1}{2} \sigma^2 E_i^2 t\right)}.$$
 (14)

Therefore, we deduce that the conditional expectation of H given  $\xi_t$  is

$$H_t = \sum_i E_i \mathbb{P}(H = E_i | \xi_t)$$
$$= \frac{\sum_i \pi_i E_i \exp\left(\sigma E_i \xi_t - \frac{1}{2} \sigma^2 E_i^2 t\right)}{\sum_i \pi_i \exp\left(\sigma E_i \xi_t - \frac{1}{2} \sigma^2 E_i^2 t\right)}.$$
(15)

In order to show that  $H_t$  is the energy process of the given quantum system, one further key result is required: namely, that the process  $W_t$  defined by

$$W_t = \xi_t - \sigma \int_0^t H_s \mathrm{d}s \tag{16}$$

is an  $\mathcal{F}_t^{\xi}$ -Brownian motion. To verify this, it suffices, by virtue of Lévy's characterisation of Brownian motion [4], to demonstrate (a) that  $W_t$  is an  $\mathcal{F}_t^{\xi}$ -martingale, and (b) that  $(dW_t)^2 = dt$ . To verify (b) we note that (9) implies  $d\xi_t = \sigma H dt + dB_t$ , and thus  $(d\xi_t)^2 = dt$ . On the other hand, (16) implies that  $dW_t = d\xi_t - \sigma H_t dt$ , and hence  $(dW_t)^2 = (d\xi_t)^2$ . To establish (a), let (15) define a function  $H(\xi, t)$  of two variables such that  $H_t = H(\xi_t, t)$ :

$$H(\xi, t) = \frac{\sum_{i} \pi_{i} E_{i} \exp\left(\sigma E_{i} \xi - \frac{1}{2} \sigma^{2} E_{i}^{2} t\right)}{\sum_{i} \pi_{i} \exp\left(\sigma E_{i} \xi - \frac{1}{2} \sigma^{2} E_{i}^{2} t\right)}.$$
 (17)

Then applying Ito's lemma and using the relation  $(d\xi_t)^2 = dt$ , we obtain

$$dH_t = \left(\partial_t H(\xi_t, t) + \frac{1}{2}\partial_\xi^2 H(\xi_t, t)\right)dt + \partial_\xi H(\xi_t, t)d\xi_t \quad (18)$$

where  $\partial_t H(\xi_t, t)$  denotes  $\partial H(\xi, t)/\partial t$  valued at  $\xi = \xi_t$ , and so on. A short calculation making use of (17) shows that  $\partial_{\xi} H(\xi_t, t) = \sigma V(\xi_t, t)$  and  $\partial_t H(\xi_t, t) + \frac{1}{2} \partial_{\xi}^2 H(\xi_t, t) = -\sigma^2 V(\xi_t, t) H(\xi_t, t)$ , where the function  $V(\xi, t)$  is

$$V(\xi,t) = \frac{\sum_{i} \pi_{i} (E_{i} - H(\xi,t))^{2} \exp\left(\sigma E_{i}\xi - \frac{1}{2}\sigma^{2}E_{i}^{2}t\right)}{\sum_{i} \pi_{i} \exp\left(\sigma E_{i}\xi - \frac{1}{2}\sigma^{2}E_{i}^{2}t\right)}.$$
 (19)

It follows, in particular, that  $H(\xi, t)$  is monotonic in  $\xi$ for any given value of t. Substituting these results into (18), we infer that  $dH_t = \sigma V(\xi_t, t) dW_t$ . However, we know that  $H_t$  is an  $\mathcal{F}_t^{\xi}$ -martingale, and therefore we conclude that  $W_t$  is also an  $\mathcal{F}_t^{\xi}$ -martingale, and that establishes (a). We thus deduce that  $W_t$  is an  $\mathcal{F}_t^{\xi}$ -Brownian motion, as claimed. We call  $W_t$  the innovation process associated with  $H_t$ . The significance of  $W_t$  is that the process  $\xi_t$  defined in (9) satisfies the diffusion equation  $d\xi_t = \sigma H_t dt + dW_t$ , where  $H_t = H(\xi_t, t)$ .

Now let  $|\psi_0\rangle$  be the initial normalised state vector of the quantum system, and let  $\hat{P}_i$  denote for each value of *i* the projection operator onto the Hilbert subspace corresponding to the energy eigenvalue  $E_i$ . We let  $|\psi_i\rangle = \pi_i^{-1/2} \hat{P}_i |\psi_0\rangle$  denote the Lüders state corresponding to  $E_i$ , and write  $\Pi_{it} = \mathbb{P} (H = E_i |\xi_t)$  for the process defined by (14). Then, we can verify that

$$|\psi_t\rangle = \sum_i e^{-iE_i t} \Pi_{it}^{1/2} |\psi_i\rangle$$
(20)

satisfies the stochastic extension of the Schrödinger equation (1) with the given initial condition. In particular, by applying Ito's lemma to (14) and using the diffusion equation satisfied by  $\xi_t$  we obtain

$$d\Pi_{it} = \sigma(E_i - H_t)\Pi_{it}dW_t.$$
 (21)

With another application of Ito's lemma we deduce that  $d\Pi_{it}^{1/2} = -\frac{1}{8}\sigma^2 (E_i - H_t)^2 \Pi_{it}^{1/2} dt + \frac{1}{2}\sigma (E_i - H_t) \Pi_{it}^{1/2} dW_t.$ A short calculation then shows that (20) satisfies (1), and that the expectation of the operator  $\hat{H}$  in the state  $|\psi_t\rangle$ is the process (15).

Summing up, the stochastic equation (1) can be solved as follows. We let H be a random variable taking values  $E_i$  with the probabilities  $\pi_i$  defined by (8), or equivalently  $\pi_i^{1/2} = \langle \psi_0 | \hat{P}_i | \psi_0 \rangle / \langle \psi_0 | \psi_0 \rangle$ . Letting  $B_t$  denote an independent Brownian motion, we set  $\xi_t = \sigma Ht + B_t$ . The solution of (1) is then given by

$$|\psi_t\rangle = \frac{\sum_i \sqrt{\pi_i} \exp\left(-\mathrm{i}E_i t + \frac{1}{2}\sigma E_i \xi_t - \frac{1}{4}\sigma^2 E_i^2 t\right) |\psi_i\rangle}{\left(\sum_i \pi_i \exp\left(\sigma E_i \xi_t - \frac{1}{2}\sigma^2 E_i^2 t\right)\right)^{1/2}}$$
(22)

where the  $\mathcal{F}_t^{\xi}$ -Brownian motion  $W_t$  driving  $|\psi_t\rangle$  in (1) is given by (16). In particular, by use of (22), the expression (15) for  $H_t$  follows at once since  $\langle \psi_i | \hat{H} | \psi_i \rangle = E_i \delta_{ij}$ .

The fact that (15) is indeed a reduction process for the energy can be verified directly as follows. Suppose, in a particular realisation of the process  $H_t$ , the random variable H takes the value  $E_j$  for some choice of the index j. Setting  $\omega_{ij} = E_i - E_j$  and writing  $\xi_t = \sigma E_j t + B_t$ , we have, for the corresponding realisation of  $H_t$ ,

$$H_t = \frac{\sum_i \pi_i E_i \exp\left(\sigma E_i B_t - \frac{1}{2}\sigma^2 E_i(E_i - 2E_j)t\right)}{\sum_i \pi_i \exp\left(\sigma E_i B_t - \frac{1}{2}\sigma^2 E_i\omega_{ij}t\right)}$$
$$= \frac{\pi_j E_j + \sum_i' \pi_i E_i \exp\left(\sigma \omega_{ij} B_t - \frac{1}{2}\sigma^2 \omega_{ij}^2 t\right)}{\pi_j + \sum_i' \pi_i \exp\left(\sigma \omega_{ij} B_t - \frac{1}{2}\sigma^2 \omega_{ij}^2 t\right)}, \quad (23)$$

where  $\sum_{i}' = \sum_{i (\neq j)}$ . However, the exponential martingale  $M_{ijt}$  defined for  $i \neq j$  by

$$M_{ijt} = \exp\left(\sigma\omega_{ij}B_t - \frac{1}{2}\sigma^2\omega_{ij}^2t\right)$$
(24)

that appears in expression (23) has the property:  $\lim_{t\to\infty} \mathbb{P}(M_{ijt} > 0) = 0$ . Hence from

$$H_{t} = \frac{\pi_{j}E_{j} + \sum_{i}'\pi_{i}E_{i}M_{ijt}}{\pi_{j} + \sum_{i}'\pi_{i}M_{ijt}},$$
(25)

we see that  $H_t$  converges to the value  $E_j$  with probability one. A similar argument allows us to verify that if  $H = E_j$  then for each value of i we have  $\lim_{t\to\infty} \prod_{it} = \mathbf{1}_{\{i=j\}}$ , where **1** denotes the indicator function, which shows that  $|\psi_t\rangle$  converges to the Lüders state corresponding to the energy eigenvalue j with probability one [3].

Therefore, we see that the random variable H can be identified with the terminal value  $H_{\infty}$  of the energy process. The fact that H is not  $\mathcal{F}_t^W$ -measurable for  $t < \infty$ indicates that the 'true value' of H is 'hidden' until the reduction process is complete. On a related interpretational point we note that in stochastic models for quantum state reduction it is sometimes assumed that the driving process  $W_t$  is in some way 'external' to the quantum system, representing, e.g., a noisy environmental coupling. This assumption, however, is unnecessary: as far as the flow of information is concerned, we have

$$\mathcal{F}_t^W = \mathcal{F}_t^{\xi} = \mathcal{F}_t^H = \mathcal{F}_t^{|\psi\rangle},\tag{26}$$

and it is thus perfectly consistent to regard the innovation process  $W_t$  as being endogenous.

The advantage of the expressions (15) and (22) is that  $H_t$  and  $|\psi_t\rangle$  are expressed algebraically in terms of the underlying random variable H and the independent Brownian motion  $B_t$ . These can be thought of as representing independent state variables for the reduction dynamics. As a consequence, we are able to investigate properties of the reduction process (1) directly without having to resort to numerical integration. In particular, by use of (22) a numerical simulation of the state reduction of rather complex quantum systems is feasible, including cases for which the Hamiltonian has a nondiscrete spectrum.

In conclusion let us analyse now in detail the timescale associated with the reduction process. For simplicity, we consider a two-state system with energy levels  $E_1$  and  $E_2$ . The initial state is given by  $|\psi_0\rangle$ , and the transition probabilities to the energy eigenstates  $|E_1\rangle$  and  $|E_2\rangle$  are given by  $\pi_1$  and  $\pi_2$ .

Suppose a measurement of the energy is made, and we condition on the outcome of the measurement being  $E_1$ . In that case, according to (25), we have

$$H_t = \frac{\pi_1 E_1 + \pi_2 E_2 M_{21t}}{\pi_1 + \pi_2 M_{21t}},$$
(27)

where  $M_{21t} = \exp(\sigma \omega_{21} B_t - \frac{1}{2} \sigma^2 \omega_{21}^2 t)$ . Writing  $\beta = \frac{1}{4} \sigma^2 \omega_{21}^2$  for the parameter that determines the characteristic rate of reduction, we can work out the probability that  $M_{21t} < e^{-n}$  for some value of n. Since  $B_t$  is normally distributed with zero mean and variance t, we find that

$$\mathbb{P}(M_{21t} < e^{-n}) = \mathbb{P}\left(B_t < \beta^{1/2}t - \frac{1}{2}\beta^{-1/2}n\right)$$
$$= N\left((\beta t)^{1/2} - \frac{1}{2}n(\beta t)^{-1/2}\right) \qquad (28)$$

where N(x) is the standard normal distribution function.

Therefore, for example, we see that provided  $t > 5\tau_R$ , we have  $\mathbb{P}(M_{21t} < e^{-10}) > \frac{1}{2}$ , where  $\tau_R = 1/\beta$ . In particular, as  $H_t$  draws near  $E_1$  we have the relation

$$H_t - E_1 \sim \frac{\pi_2}{\pi_1} (E_2 - E_1) M_{21t}.$$
 (29)

Thus, after only a relatively few multiples of the characteristic reduction timescale, the amount by which  $H_t$ differs from  $E_1$  will typically be reduced to a tiny fraction of the energy difference  $E_2 - E_1$ .

DCB acknowledges support from The Royal Society. LPH acknowledges the Institute for Advanced Study, Princeton, for hospitality while part of this work was carried out.

- L. Diosi, Phys. Lett. A129 419 (1988); N. Gisin, Helv. Phys. Acta 62, 363 (1989); G. C. Ghirardi,
   P. Pearle, and A. Rimini, Phys. Rev. A 42, 78 (1990); A. Barchielli and V. P. Belavkin, J. Phys A. 24, 1495 (1991); I. C. Percival, Proc. R. Soc. London A 447, 189 (1994); H. M. Wiseman and
   L. Diosi, Chem. Phys. 268, 91 (2001).
- L. P. Hughston, Proc. R. Soc. London A 452, 953 (1996); S. L. Adler and L. P. Horwitz, J. Math. Phys. 41, 2485 (2000); S. L. Adler and T. A. Brun, J. Phys. A 34, 4797 (2001); S. L. Adler, J. Phys. A 35, 841 (2002); D. C. Brody and L. P. Hughston, Proc. R. Soc. London A 458, (2002).
- S. L. Adler, D. C. Brody, T. A. Brun and L. P. Hughston, *J. Phys. A* 34, 8795 (2001).
- R. S. Liptser and A. N. Shiryaev, *Statistics of Random Processes* Vols. I and II, 2nd ed. (Springer, Berlin 2000).