

THE COSMIC EVOLUTIONARY THEORY

Ding-Yu Chung

The cosmic evolutionary theory is by means of the evolution of object in four stages: string, membrane, particle, and particle-wave. It is also concurrently by means of the evolution of vacuum in four stages: object vacuum, single vacuum, inclusive vacuum, and exclusive vacuum. Vacuum is a homogeneous space, while object has a definite shape. Object vacuum takes turn to co-exist equally with object at the same location. Single vacuum exists with an object at the same location or different locations in one object to one vacuum relation. Inclusive vacuum exists at the same location with multiple objects. Exclusive vacuum excludes objects at the same location. The universe starts with the pre-universe that is object vacuum-object with a non-zero vacuum energy. It is an equilibrium state between vacuum and the pair of ten-dimensional superstring and anti-superstring. The vacuum fluctuation of object vacuum results in the pre-expanding universe with the boundary 9-brane and 9-antibrane separated by single vacuum with supergravity and anti-supergravity. The collapse and the bounce of the pre-expanding universe result in zero vacuum energy, which induces the fractionalization of one object into multiple low energy objects. The multiple objects require inclusive vacuum each of which associates with multiple objects. Such fractionalization brings about the cosmic expansion that leads to the superluminal inflation to rupture the connection among objects. The gap generated from the rupture is exclusive vacuum that excludes objects. The rupture brings about the big bang, cosmic radiation, elementary particles, dark matter, entropy, force fields, and quantum mechanics. The cosmic evolution involves cyclic universe. The cosmic evolution provides the most comprehensive explanation for the origin of physical laws. The periodic table of elementary particles derived from the cosmic evolution provides the most comprehensive explanation and calculation for the masses of gauge bosons, leptons, quarks, and hadrons in a good agreement with the observed values.

1. Introduction

The cosmic evolutionary theory is by means of the evolution of object in four stages: string, membrane, particle, and particle-wave. It is also concurrently by means of the evolution of vacuum in four stages: object vacuum, single vacuum, inclusive vacuum, and exclusive vacuum. Vacuum is a homogeneous space, while object has a definite shape. Object vacuum takes turn to co-exist equally with object at the same location. Single vacuum exists with an object at the same location or different locations in one object to one vacuum relation. Inclusive vacuum exists at the same location with multiple objects. Exclusive vacuum excludes objects at the same location.

The universe starts with the pre-universe that is object vacuum-object, the equilibrium state between the vacuum and the pairs of ten-dimensional superstring and anti-superstring. The vacuum energy is equal to the energy of the superstring below the Planck energy.

Object vacuum can also undergoes condensation and dilution as vacuum fluctuation. The condensation as the compactification of higher dimensions leads to the formation of an ordinary two-dimensional string-antistring and the empty space as single vacuum, each of which associates with an object. The repulsive force between the string and the antistring keeps single vacuum in between the string and the antistring. The tension of the string or antistring is higher than the single vacuum, which is anti de-Sitter space (AdS). When the string attaches to the adjacent superstring, the attached string provides the eleventh dimension to generate an eleven-dimensional membrane, which has the mass equal to the Planck mass. When the attached string spreads to single vacuum as AdS by the dilution, and wraps around the trapped superstring, the trapped superstring becomes the one-sided Randall-Sundrum [1] 9-brane embedded in the eleven-dimensional bulk with supergravity as Fig. 1.




Fig. 1: the formation of the 9-brane embedded in 11D bulk in the pre-universe

In the same way, the ten-dimensional anti-superstring becomes the one-sided Randall-Sundrum 9-antibrane embedded in the eleven-dimensional bulk with anti-supergravity. The attractive force between the 9-brane and the 9-antibrane forms the pre-expanding universe, where the boundary 9-brane and the boundary 9-antibrane embedded in the eleven-dimensional space-time are separated by single vacuum with supergravity and anti-supergravity as Fig. 2. The force between supergravity and anti-supergravity is repulsive, while the force between the 9-brane and the 9-antibrane is attractive.

This two-brane structure of the pre-expanding universe also appears in the Horava-Witten eleven dimensional supergravity on a manifold with two ten-dimensional boundaries [2], the ekpyrotic universe with the visible and the hidden boundary branes [3], and the brane-antibrane universe [4].

The pre-universe vacuum outside of the pre-expanding universe fills single vacuum, and the formation of brane and antibrane continues. The development of the pre-expanding universe is analogous to the development of a star. In a star, the process of fusion generates heavy elements in the core, that has a strong gravitational attractive force, but the outward explosive thrust of internal fusion reactions prevents the star from collapse to the core. In the pre-expanding universe, the vacuum fluctuation generates the boundary brane and antibrane that attract each other, but it also generates single vacuum and the repulsive supergravity-anti-supergravity that keep the boundary brane and antibrane apart. (In the original ekpyrotic universe model [3], the bulk branes are generated from the boundary branes. In this paper, the boundary branes are generated from the bulk branes.)

At the critical mass of the pre-expanding universe when the rate of the single vacuum formation from the vacuum fluctuation is too slow to keep the 9-brane and the 9-antibrane apart, the pre-expanding universe collapses. During the collapse, the attractive force and the repulsive force cause an inversion that turns the 9-brane and the 9-antibrane inside, and turns supergravity and anti-supergravity outside. (As shown later, electromagnetism is the manifestation of the collapse of the pre-expanding universe.) Eventually, the 9-brane and the 9-antibrane coalesce, and annihilate each other, resulting in the loss of the string property. The result is the generation of the nine space dimensional mixed 9-particle with the Kaluza-Klein structure without identical space dimensions and without fixed number of space dimensions.

At the end of the coalescence, the repulsion between supergravity and anti-supergravity causes a bounce after the collapse. This bounced pre-expanding universe is the mixed pre-expanding universe, consisting of four equal parts: the two mixed 9-particles, supergravity, and anti-supergravity. Single vacuum is absorbed by the mixed 9-particles. The collapse generates a gulf separating the mixed pre-expanding universe and the surrounding pre-universe as Fig. 2.




Fig. 2: from the pre- expanding universe to the mixed pre-expanding universe

The interactions between branes in the form of collision are proposed in various brane models [3, 4, 5]. The cyclic universe model based on the ekpyrotic universe [6] has the collapse-singularity-bounce scheme.

Detached permanently from the pre-universe by the gulf, the mixed pre-expanding universe has zero vacuum energy instead of the non-zero vacuum energy as in the pre-universe. With zero vacuum energy, the mixed 9-particles fractionalize into multiple lower energy mixed particles. The vacuum is transformed from single vacuum associated with only one object to inclusive vacuum associated with multiple objects. It is the beginning of the expanding universe. The transition from non-zero vacuum energy to zero vacuum energy is gradual.

The cosmic expansion mechanism is discussed in Section 2. The two different modes of the expansion are described in Section 3. The derivation of quantum mechanics and gravity is described in Section 4. In Section 5, the ordinary universe consists of the mixed 3-particle as the mixture of leptons and quarks. In Sections 6, 7, and 8, the periodic table of

elementary particles is constructed to account of all leptons, quarks, gauge bosons, and hadrons, and their masses are calculated. In Section 9, the cyclic universe is proposed.

2. *The cosmic expansion mechanism: the dimensional hierarchy*

Single vacuum essentially defines fermion whose space allows only one fermion, and inclusive vacuum essentially defines boson whose space allows multiple bosons. During the collapse, single vacuum is absorbed by the 9-brane as a part of the mixed 9-particle, so during the expansion, single vacuum is linked with fermion, and inclusive vacuum is linked with boson. At the beginning of the expanding universe, the vacuum is transformed from single vacuum associated with only one object to inclusive vacuum associated with multiple objects for fractionalization, and the mixed 9-particles are bosons.

At the same vacuum energy as in supersymmetry, the repeated application of the boson-fermion transformation transforms a boson from one point to the same boson at another point. At the gradually decreasing vacuum energy as in the cosmic expansion, the repeated application of the boson-fermion transformation transforms the mixed 9-particle boson (with the Kaluza-Klein structure) from one point to the mixed 8-particle boson with a lower mass and lower dimension at another point. (In the Kaluza-Klein structure, the space dimensions do not have to be the same, and the number of space dimensions is not fixed.) The stepwise repeated transformations fractionalize the mixed 9-particle into various lower dimensional mixed particles from 8 to 3 with decreasingly lower masses. These transformations bring about the expansion and fractionalization during the cosmic expansion.




Fig. 3: the cosmic expansion through supersymmetry with different vacuum energies

These stepwise transformations constitute the dimensional hierarchy [7,8] where masses decrease with the space-time dimension numbers. In the dimensional hierarchy, each space dimension consists of a dimensional fermion F_D and a dimensional boson B_D . The transformation constant from a fermion into its boson in the adjacent dimension is same as the fine structure constant, α , the probability of a fermion emitting or absorbing a boson. The transformation constant from a boson into its fermion partner in the same dimension is also the fine structure constant, α . This hierarchy is expressed in term of the dimension space-time number, D ,

$$M_{D-1, B} = M_{D, F} \alpha_{D, F}, \quad (1)$$

$$M_{D, F} = M_{D, B} \alpha_{D, B}, \quad (2)$$

where $M_{D, B}$ and $M_{D, F}$ are the masses for a dimensional boson and a dimensional fermion, respectively, and $\alpha_{D, B}$ or $\alpha_{D, F}$ is the fine structure constant, which is the ratio between the energies of a dimensional boson and its dimensional fermionic partner. Assuming α is the same for all dimensional fermions and dimensional bosons, Eq. (3) for the energy relation among the dimensional bosons is derived.

$$M_D = M_P \alpha^{2(11-D)} \quad (3)$$

where M_D is mass of a dimension, D is space-time dimension number, M_P is the Planck mass (the mass of the eleventh dimension), and α is fine structure constant, the probability of a fermion emitting or absorbing a boson. As shown later, all α 's except one is equal to the fine structure constant for electromagnetic field, so Eq. 3 is approximately correct.

3. *The expanding universe: the big band and the big bang*

The expansion and fractionalization process by the repetitive transformation between inclusive vacuum and single vacuum has two different modes: the big band mode for the hidden universe and the big bang mode for the observable universe as Figs. 4 and 5.




Fig. 4: the historical diagram for the formation of the observable and the hidden universes




Fig. 5: the collapse-bounce and the fractionalization process (the big band mode and the big bang mode)

The big band mode is used in the hidden universe. In the big band mode, the mixed 9-particle is fractionalized into the hierarchical mixed particles from 3 to 9, whose masses follow the hierarchical dimensional mass formula as Eq. (3) where p particle has $p + 1$ space-time dimension, D . In the big band mode, the slow rate of expansion (the rate of vacuum energy decrease) brings about the slow sequential stepwise fractionalization from the mixed 9-particle to the mixed 3-particle, resulting in the slow cosmic expansion. Assuming that the pre-universe is elastic, at the end of the cosmic expansion, the pre-universe pushes the hidden universe back, resulting in slow condensation back to the mixed 9-particle. The hidden universe undergoes expansion and contraction, like a big elastic rubber band (big band). Anti-supergravity is not broken. All mixed particles are held together by anti-supergravity, and are adjacent to one another without gap.

The big bang mode is used in the observable universe. In the big bang mode, the expansion is the superluminal inflation that ruptures the supergravity. (The ease of rupture is proportional to the rate of expansion.) The supergravity becomes individualized gravity for each individual object. Consequently, all mixed particles are not held together by supergravity, and are separated by exclusive vacuum as the gap among particles. Unlike inclusive vacuum and single vacuum, exclusive vacuum excludes objects. This superluminal inflation is too fast to allow all mixed particles to become mixed 3-particles, and results in a mixture of mixed particles from 9 to 3.

The mechanism to generate exclusive vacuum is to assign quantized positive and negative charges to the two internal boundary branes within the mixed particle, resulting in the dislocation of energy by the internal annihilation (implosion). The dislocation of energy from the mixed particle forms exclusive vacuum as the empty mixed 3-particle. (The movement of a massless particle leaves exclusive vacuum behind.) The dislocated energy is cosmic radiation, which in turn breaks the entire supergravity. The rupture that causes continuous irreversible disconnection among particles results in largely independent disconnected entities, which constitute entropy for the second law of thermodynamics locally in the observable universe.

The mixed particles that are not annihilated have asymmetrical charges (CP nonconservation), in such way that the mixed particle has two asymmetrical sets (main and auxiliary) of space dimensions from the two boundary branes within the mixed particle. The auxiliary set is dependent on the main set, so the mixed particle appears to have only one set of space dimensions.

During the superluminal inflation, the absorbed single vacuum and the absorbed inclusive vacuum are parts of particles. After the rupture, the space in exclusive vacuum allows single vacuum and inclusive vacuum free to be inside and outside of the original particles, so inclusive vacuum and single vacuum are no longer identified with boson and fermion, respectively, and the repeated boson-fermion transformation for the cosmic expansion stops. In the observable universe, all four forms of vacuum co-exist.

Inclusive vacuum outside of a particle in the big bang mode allows all the mixed particles to have the same number of space dimensions. It is achieved by adding the virtual hierarchical space dimensions to inclusive vacuum surrounding the core mixed particles in the manner of the Kaluza-Klein structure, where the virtual space dimension is a one-dimensional circle associated with every point in the core particle dimension, and every

higher virtual dimension circles the circle of the lower virtual dimension. The virtual space dimensions are the space dimensions in between the core particle space dimension and the gravity (the eleventh space-time dimension). The masses of the hierarchical space dimensions follow the hierarchical dimensional mass formula as Eq. (3).

To provide the virtual hierarchical space dimensions with the same space-time dimension as the core mixed particle, the core mixed particle absorbs the virtual hierarchical dimensions as the scalar fields through Higgs mechanism. (There is no free Higgs boson. So far no free Higgs boson has been detected.) The surrounding virtual hierarchical dimensions are converted into the "dimensional orbitals" with the same space-time dimension as the core mixed particle.

Gravity resided in exclusive vacuum is also a part of the hierarchical dimensional orbitals for the mixed particles. These hierarchical dimensional orbitals are the force and mass fields. The final forms of the mixed particles are the mixed particles from 3 to 9 surrounded by the hierarchical dimensional orbitals as the force and mass fields as Fig. 5.

Since the core mixed particles have two sets (main and auxiliary) of space dimensions, there are also two sets of the dimensional orbitals. For the mixed 3-particle in the big bang mode, there are two sets of the hierarchical seven dimensional orbitals (including gravity) and the non-hierarchical three core particle space dimensions as Figs. 7 and 8. It is the mixture of leptons and quarks.

Exclusive vacuum and cosmic radiation emerge only after the superluminal inflationary emergence of all of the hierarchical mixed particles. Consequently, the superluminal inflationary emergence of the mixed particles followed by the non-inflationary emergence of cosmic radiation constitutes the hybrid inflation [9]. Cosmic radiation, observable elementary particles, and all force fields emerge only at the end of inflation.

The two universes are parallel and incompatible to each other until the quintessence period. This two-universe model appears also in the two-universe (visible and hidden) model in the cyclic universe model based on the ekpyrotic universe [6].

4. *Quantum mechanics and gravity*

After the rupture, inclusive vacuum and single vacuum are no longer identified with boson and fermion, respectively. Inclusive vacuum and single vacuum now exist complementarily in and around all matters, including both boson and fermion. In one inclusive vacuum, there are multiple objects, which can also be multiple states in one inclusive vacuum in one object, so inclusive vacuum fractionalizes an object into multiple states as wave, while single vacuum retains an object as single state as particle. The complementary existence of inclusive vacuum and single vacuum brings about the complementary principle between wave and particle. Since inclusive vacuum relates to boson, and single vacuum relates to fermion, each particle possesses a build-in supersymmetry in vacuum.


Inclusive vacuum overlaps with exclusive vacuum surrounding a particle. In the overlapping space of inclusive vacuum and exclusive vacuum, various parts of the

overlapping space undergo constant random changes from inclusion to exclusion and from exclusion to inclusion. This exclusion-inclusion of an object is essentially equivalent to subjective measurement of an object in Copenhagen interpretation of quantum mechanics. Both the exclusion and subjective measurement change the position and velocity of an object. The rate of change from inclusion to exclusion and from exclusion to inclusion is velocity. The requirement for randomness does not allow constant position and constant velocity (the rate of change) simultaneously. At an exact position, the variation in velocities is large. At an exact velocity, the variation in positions is large. The variation is uncertainty. Therefore, the source of the uncertainty principle is subjective measurement or the objective exclusion. Consequently, an object has the probability to be anywhere in the form of wavefunction.

Exclusive vacuum is the gap between objects, so when two objects entangles (collides), exclusive vacuum disappears. During the entanglement, the original two sets of multiple states no longer exist. By eliminating exclusive vacuum and inclusive vacuum, the entangled single vacuum with the one state, the entangled state, emerges. This is collapse of wavefunction. Therefore, any measurement of wavefunction always ends with one state instead of multiple states.

The cosmic expansion has essentially anti-gravity effect, in which fractionalized particles move away from one another. During the cosmic expansion, the repeated application of the boson-fermion transformation (inclusive vacuum-single vacuum transformation) transforms the mixed p particle from one point to the mixed p-1 particle with one dimension lower at another point. Gravity is the reverse of this cosmic expansion. The coupling from different mass-energy sources through graviton initiates the repeated transformation from inclusive vacuums to single vacuum, and from single vacuum to inclusive vacuum. This repeated transformation generates the D+1 dimensional bulk in which the D particle with D dimensional space-time is embedded. Gravity resides in the D+1 bulk as the one-sided Randall-Sundrum four-dimensional particle embedded in the five-dimensional bulk [1].

The transformations involving other force fields transform D particle into D particle embedded in internal space with various discrete charges. The symmetry in this transformation is gauge symmetry. The summary is described in Fig. 6.






Fig. 6: different forms of vacuum and object in physics

5. *The Ordinary Universe: the mixed 3-particle*

The observable universe consists of the mixed particles from 3 to 9. The ordinary (baryonic) universe in the observable universe consists of the mixed 3-particle, which is the mixture of leptons and quarks. Exotic dark matter in the observable universe consists of the mixed particles from 4 to 9. As shown later, exotic dark matter cannot be seen, but it can be observed by gravity. The ordinary (baryonic) matter is one of the seven mixed particles at equal mass proportions, so the baryonic mass fraction is $1/7$ (0.14). The universal baryonic mass fraction was found to be 0.13 by the observations of primordial deuterium abundance [10]. The calculated value agrees well with the observed value.

For ordinary matter (the mixed 3-particle), there are two sets (main and auxiliary) of the seven dimensional orbitals. The total number of dimensional orbitals is 14 as shown in Fig. 7.




Fig. 7: 14 dimensional orbitals in the mixed 3-particle: 7 main dimensional orbitals (solid line), and 7 auxiliary dimensional orbitals (dot line), D = main dimensional orbital number

As shown in Fig. 7, the auxiliary dimensional orbitals are in the middle of the main dimensional orbitals. The fifth main dimensional orbital is the start of the main dimensional orbitals. To be adjacent to the start of the auxiliary dimensional orbitals, the seventh main dimensional orbitals mixes with the fifth orbitals. Such mixing is manifested by the symmetry mixing in the electroweak interaction between $U(1)$ for the fifth main dimensional orbital and $SU(2)$ for the seventh main dimensional orbitals.

There are dualities between dimensional orbitals and the cosmic evolution. There is duality between gravity resided in $D+1$ bulk and the eleven dimensional supergravity to the ten-dimensional superstring. All other main dimensional orbitals are in the internal space with various discrete charges, and carry various forms of gauge symmetry and space-time symmetry. The auxiliary dimensional orbitals are mostly for quarks.

There is duality between the collapse of the pre-expanding universe and electromagnetism as the fifth main dimensional orbital with $U(1)$ gauge symmetry. The attraction between the 9-brane and the 9-antibrane is manifested as electric force. As the collapse of the pre-expanding universe begins, and the 9-brane and the 9-antibrane start to move, the attractive force and the repulsive force cause an inversion that turns the 9-brane and the 9-antibrane inside, and turns supergravity and the anti-supergravity outside. In the observable universe, there is no repulsive force between supergravity and the anti-supergravity, so the force for the inversion is manifested as the force for the rotation, which is magnetic force, perpendicular to the direction of moving charging particles.

All other dimensional orbitals except gravity are short-range dimensional orbitals. Only the mixed 3-particle (ordinary matter) has the fifth main dimensional orbital, so without the fifth main dimensional orbital, exotic dark matter consists of permanently neutral higher dimensional particles. It cannot emit light, and cannot form atoms.

There is duality between the sixth main dimensional orbital and the hiding of the auxiliary dimensional orbitals. The hiding of the auxiliary dimensional orbitals (quarks) is represented by the hiding of colors in color $SU(3)$ and fractional electric charges for quarks. For the independent existence for a quark composite, color $SU(3)$ with fractional charges has to be converted into colorless $U(1)$ with integer charge and hypercharge, like leptons.

The seventh main dimensional orbital carries the charge for the gauge symmetry within the family of leptons or quarks. The charge is represented by $SU(2)$ for the symmetry between two leptons (e and ν) or between two quarks (u and d). The same set of gauge symmetry groups is assigned to the eighth, the ninth, and the tenth main dimensional orbitals as $U(1)$, $U(1)$, and $SU(2)$, respectively.

Various space-time symmetries reflect the existence of fermions. P nonconservation is required to achieve chiral symmetry for massless leptons (neutrinos), so masses of higher neutrinos in the lepton family are not too large to fit in the lepton family. In order to have a long-term existence of fermions, CP nonconservation is required to distinguish matter from anti-matter. P and CP nonconservations are in pairs of the right and the left. The seventh, the eighth, the ninth, and the tenth main dimensional orbitals are assigned to have P (left), CP (right), CP (left), and P (right) nonconservation, respectively. There is no nonconservation for the fifth and the sixth main dimensional orbitals.

The combination of the gauge symmetry and the space-time symmetry for the main dimensional orbitals from 5 to 10 results in $U(1)$, $SU(3)$ to become $U(1)$, $SU(2)_L$, $U(1)_R$, $U(1)_L$, and $SU(2)_R$, respectively. The eleventh main dimensional orbital is for gravity.

6. *The force fields: the masses of gauge bosons*

The structure of the mixed 3-particle with dimensional orbitals resembles to the structure of atomic orbital. Consequently, the periodic table of elementary particles is constructed to account of all leptons, quarks, gauge bosons, and hadrons.

For the gauge bosons, the seven main dimensional orbitals are arranged as $F_5 B_5 F_6 B_6 F_7 B_7 F_8 B_8 F_9 B_9 F_{10} B_{10} F_{11} B_{11}$, where B and F are boson and fermion in each orbital. The masses of the main dimensional orbital bosons can be derived from Eqs. (1) and (2). Assuming $\alpha_{D,B} = \alpha_{D,F}$, the relation between the bosons in the adjacent main dimensional orbitals, then, can be expressed in terms of the main dimensional orbital number, D, as

$$M_{D-1,B} = M_{D,B} \alpha^2_D, \quad (4)$$

where $D= 6$ to 11 , and $E_{5,B}$ and $E_{11,B}$ are the energies for the main dimensional orbital five and the main dimensional orbital eleven, respectively. The lowest energy is the Coulombic field,

$$E_{5,B} = \alpha M_{6,F} = \alpha M_e \quad (5)$$

The bosons generated are the main dimensional orbital bosons or B_D . Using only α_e , the mass of electron, the mass of Z^0 , and the number (seven) of dimensional orbitals, the masses of B_D as the gauge boson can be calculated as shown in Table 1.

Table 1. The Masses of the main dimensional orbital bosons:
 $\alpha = \alpha_e$, D = main dimensional orbital number

B_D	M_D	GeV (calculated)	gauge boson	Interaction, symmetry
B_5	$M_e \alpha$	3.7×10^{-6} (given)	A	electromagnetic, U(1)
B_6	M_e / α	7×10^{-2}	$\pi_{1/2}$	strong, SU(3) \longrightarrow U(1)
B_7	$M_e / \alpha_w^2 \cos \theta_w$	91.177 (given)	Z_L^0	weak (left), SU(2) _L
B_8	M_7 / α^2	1.7×10^6	X_R	CP (right) nonconservation, U(1) _R
B_9	M_8 / α^2	3.2×10^{10}	X_L	CP (left) nonconservation, U(1) _L
B_{10}	M_9 / α^2	6.0×10^{14}	Z_R^0	weak (right), SU(2) _R
B_{11}	M_{10} / α^2	1.1×10^{19}	G	gravity, D particle embedded in D+1 bulk

In Table 1, $\alpha = \alpha_e$ (the fine structure constant for electromagnetic field), and α_w is not same as α of the rest, because there is a mixing between B_5 and B_7 as the symmetry mixing between U(1) and SU(2) in the standard theory of the electroweak interaction, and $\sin \theta_w$ is not equal to 1. As shown later, B_5 , B_6 , B_7 , B_8 , B_9 , and B_{10} are A (massless photon), $\pi_{1/2}$, Z_L^0 , X_R , X_L , and Z_R^0 , respectively, responsible for the electromagnetic field, the strong interaction, the weak (left handed) interaction, the CP (right handed) nonconservation, the CP (left handed) nonconservation, and the P (right handed) nonconservation, respectively. The calculated value for θ_w is 29.69° in good agreement with 28.7° for the observed value of θ_w

[11]. The calculated energy for B_{11} is 1.1×10^{19} GeV in good agreement with the Planck mass, 1.2×10^{19} GeV.

As in the standard theory to the electroweak interaction, the boson mixing of $U(1)$ and $SU(2)_L$ is to create electric charge and to generate the bosons for leptons and quarks by combining isospin and hypercharge. The hypercharges for both e^+ and ν are 1, while for both u and d quarks, they are 1/3 [12]. The electric charges for e^+ and ν are 1 and 0, respectively, while for u and d quarks, they are 2/3 and -1/3, respectively.

There is no strong interaction from the internal symmetry for the dimensional bosons, originally. The strong interaction is generated by "leptonization" [13], which means quarks have to behave like leptons. Leptons have integer electric charges and hypercharges, while quarks have fractional electric charges and hypercharges. The leptonization is to make quarks behave like leptons in terms of "apparent" integer electric charges and hypercharges. Obviously, the result of such a leptonization is to create the strong interaction, which binds quarks together in order to make quarks to have the same "apparent" integer electric charges and hypercharges as leptons. There are two parts for this strong interaction: the first part is for the charge, and the second part is for the hypercharge. The first part of the strong interaction allows the combination of a quark and an antiquark in a particle, so there is no fractional electric charge. It involves the conversion of $\pi_{1/2}$ boson in B_6 into the electrically chargeless meson field by combining two $\pi_{1/2}$, analogous to the combination of e^+ and e^- fields, so the meson field becomes chargeless. (The mass of π , 135 MeV, is twice of the mass of a half-pion boson, 70 MeV, minus the binding energy. $\pi_{1/2}$ becomes pseudoscalar up or down quark in pion.) In the meson field, no fractional charge of quark can appear. The second part of the strong interaction is to combine three quarks in a particle, so there is no fractional hypercharge. It involves the conversion of B_5 ($\pi_{1/2}$) into the gluon field with three colors. The number of colors (three) in the gluon field is equal to the ratio between the lepton hypercharge and quark hypercharge. There are three $\pi_{1/2}$ in the gluon field, and at any time, only one of the three colors appears in a quark. Quarks appear only when there is a combination of all three colors or color-anticolor. In the gluon field, no fractional hypercharge of quarks can appear. By combining both of the meson field and the gluon field, the strong interaction is the three-color gluon field based on the chargeless vector meson field from the combination of two $\pi_{1/2}$'s. The total number of $\pi_{1/2}$ is 6, so the fine structure constant, α_s , for the strong force is

$$\begin{aligned}\alpha_s &= 6\alpha e^1 \\ &= 0.119\end{aligned}\tag{6}$$

which is in a good agreement with the observed value, 0.124 [14].

The dimensional boson, B_8 , is a CP violating boson, because B_8 is assumed to have the CP-violating $U(1)_R$ symmetry. The ratio of the force constants between the CP-invariant W_L in B_8 and the CP-violating X_R in B_8 is

$$\begin{aligned}\frac{G_8}{G_7} &= \frac{\alpha E_7^2 \cos^2 \Theta_W}{\alpha_W E_8^2} \\ &= 5.3 \times 10^{-10},\end{aligned}\quad (7)$$

which is in the same order as the ratio of the force constants between the CP-invariant weak interaction and the CP-violating interaction with $|\Delta S| = 2$.

The dimensional boson, $B_9 (X_L)$, has the CP-violating $U(1)_L$ symmetry. The lepton (l_9) and the quark (q_9) are outside of the three families for leptons and quarks, so baryons in dimension nine do not have to have the baryon number conservation. The baryon that does not conserve baryon number has the baryon number of zero. The combination of the CP-nonconservation and the baryon number of zero leads to the baryon, \bar{p}_9^- , with the baryon number of zero and the baryon, p_9^+ , with the baryon number of 1. The decay of \bar{p}_9^- is as follows.

$$\bar{p}_9^- \rightarrow l_9^- \bar{l}_9^{\circ}$$

The combination this \bar{p}_9^- and p_9^+ as well as leptons, $l_9^{\circ} \bar{l}_9^{\circ}$ results in

$$\begin{aligned}p_9^+ \bar{p}_9^- + l_9^{\circ} \bar{l}_9^{\circ} &\rightarrow p_9^+ + l_9^- \bar{l}_9^{\circ} + l_9^{\circ} \bar{l}_9^{\circ} \\ &\rightarrow p_9^+ + l_9^- + \text{radiation} + l_9^{\circ} \\ &\rightarrow n_9^{\circ} + \text{radiation}\end{aligned}$$

Consequently, excess baryons, n_9° , are generated in the dimension nine. These excess baryons, n_9° , become the predecessors of excess neutrons in the low energy level.

The ratio of force constants between X_R with CP-independent baryon number and X_L with CP-dependent baryon number is

$$\begin{aligned}\frac{G_9}{G_8} &= \frac{\alpha E_8^2}{\alpha E_9^2} \\ &= 2.8 \times 10^{-9},\end{aligned}\quad (8)$$

which is in the same order as the observed ratio of the numbers between the left handed baryons (proton or neutron) and photons in the universe.

7. *The periodic table of elementary particles: the masses of leptons and quarks*

The two sets of seven dimensional orbitals result in 14 dimensional orbitals (Fig. 8) for gauge bosons, leptons, and quarks. The periodic table for elementary particles is shown in Table 2.




Fig. 8. leptons and quarks in the dimensional orbits
D = main dimensional number, a = auxiliary dimensional number

Table 2. The Periodic Table of elementary particles

D = main dimensional orbital number, a = auxiliary dimensional orbital number

D	a = 0	1	2	a = 0	1	2	3	4	5
	<u>Lepton</u>			<u>Quark</u>					<u>Boson</u>
5	$l_5 = \nu_e$			$q_5 = u_5 = 3\nu_e$					$B_5 = A$
6	$l_6 = e$			$q_6 = d_6 = 3e$					$B_6 = \pi_{1/2}$
7	$l_7 = \nu_\mu$	μ_7	τ_7	$q_7 = 3\mu$	u_7/d_7	s_7	c_7	b_7	$B_7 = Z_L^0$
8	$l_8 = \nu_\tau$	μ_8		$q_8 = \mu'$	b_8	t_8			$B_8 = X_R$
9	l_9			q_9					$B_9 = X_L$
10	f_{10}								$B_{10} = Z_R^0$
11	f_{11}								$B_{11} = G$

D is the dimensional orbital number for the seven main dimensional orbitals. The auxiliary dimensional orbital number, a, is for the seven extra auxiliary dimensional orbitals. All gauge bosons, leptons, and quarks are located on the seven main dimensional orbits and seven auxiliary orbits. ν_e , e , ν_μ , and ν_τ are main dimensional fermions for main dimension 5, 6, 7, and 8, respectively. All neutrinos have zero mass because of chiral symmetry.

The dimensional fermions for $D = 5$ and 6 are neutrino (l_5) and electron (l_6), respectively. To generate a quark whose mass is higher than the lepton is to add the lepton to the boson from the combined lepton-antilepton, so the mass of the quark is three times of the mass of the corresponding lepton in the same main dimension. The equation for the quark mass is

$$M_{q_D} = 3M_{l_D} \quad (9)$$

A heavy lepton (μ or τ) is the combination of the dimensional leptons and the auxiliary dimensional leptons. In the same way, a heavy quark is the combination of the

dimensional quarks from Eq.(9) and the auxiliary quarks. The mass of the auxiliary dimensional fermion (AF for both heavy lepton and heavy quark) is generated from the corresponding main dimensional boson as follows.

$$M_{AF_{D,a}} = \frac{M_{B_{D-1,0}}}{\alpha_a} \sum_{a=0}^a a^4, \quad (10)$$

where α_a = auxiliary dimensional fine structure constant, and a = auxiliary dimension number = 0 or integer. The first term, $\frac{M_{B_{D-1,0}}}{\alpha_a}$, of the mass formula (Eq.(10)) for the auxiliary dimensional fermions is derived from the mass equation, Eq. (1), for the dimensional fermions and bosons. The second term, $\sum_{a=0}^a a^4$, of the mass formula is for Bohr-Sommerfeld quantization for a charge - dipole interaction in a circular orbit as described by A. Barut [14]. $1/\alpha_a$ is $3/2$. The coefficient, $3/2$, is to convert the main dimensional boson mass to the mass of the auxiliary dimensional fermion in the higher dimension by adding the boson mass to its fermion mass which is one-half of the boson mass. The formula for the mass of auxiliary dimensional fermions (AF) becomes

$$\begin{aligned} M_{AF_{D,a}} &= \frac{M_{B_{D-1,0}}}{\alpha_a} \sum_{a=0}^a a^4 \\ &= \frac{3}{2} M_{B_{D-1,0}} \sum_{a=0}^a a^4 \\ &= \frac{3}{2} M_{F_{D,0}} \alpha_D \sum_{a=0}^a a^4 \end{aligned} \quad (11)$$

When the mass of this auxiliary dimensional fermion is added to the sum of masses from the corresponding main dimensional fermions (zero auxiliary dimension number) with the same electric charge and the same main dimension, the fermion mass formula for heavy leptons and quarks is derived as follows.

$$\begin{aligned} M_{F_{D,a}} &= \sum M_{F_{D,0}} + M_{AF_{D,a}} \\ &= \sum M_{F_{D,0}} + \frac{3}{2} M_{B_{D-1,0}} \sum_{a=0}^a a^4 \\ &= \sum M_{F_{D,0}} + \frac{3}{2} M_{F_{D,0}} \alpha_D \sum_{a=0}^a a^4 \end{aligned} \quad (12)$$

Each fermion can be defined by main dimensional numbers (D's) and auxiliary dimensional numbers (a's). Heavy leptons and quarks consist of one or more D's and a's. The compositions and calculated masses of leptons and quarks are listed in Table 3.

Table 3. The Compositions and the Constituent Masses of Leptons and Quarks
D = main dimensional orbital number and a = auxiliary dimensional orbital number

	D_a	Composition	Calc. Mass
<u>Leptons</u>			
	<u>D_a for leptons</u>		
ν_e	5_0	ν_e	0
e	6_0	e	0.51 MeV (given)
ν_μ	7_0	ν_μ	0
ν_τ	8_0	ν_τ	0
μ	$6_0 + 7_0 + 7_1$	$e + \nu_\mu + \mu_7$	105.6 MeV
τ	$6_0 + 7_0 + 7_2$	$e + \nu_\mu + \tau_7$	1786 MeV
<u>Quarks</u>			
	<u>D_a for quarks</u>		
u	$5_0 + 7_0 + 7_1$	$u_5 + q_7 + u_7$	330.8 MeV
d	$6_0 + 7_0 + 7_1$	$d_6 + q_7 + d_7$	332.3 MeV
s	$6_0 + 7_0 + 7_2$	$d_6 + q_7 + s_7$	558 MeV
c	$5_0 + 7_0 + 7_3$	$u_5 + q_7 + c_7$	1701 MeV
b	$6_0 + 7_0 + 7_4$	$d_6 + q_7 + b_7$	5318 MeV
t	$5_0 + 7_0 + 7_5 + 8_0 + 8_2$	$u_5 + q_7 + t_7 + q_8 + t_8$	176.5 GeV

The lepton for main dimension five is ν_e , and the quark for the same main dimension is u_5 , whose mass is equal to $3 M\nu_e$ from Eq. (9). The lepton for the main dimension six is e, and the quark for this main dimension is d_6 . u_5 and d_6 represent the “light quarks” or “current quarks” which have low masses. The main dimensional lepton for the dimensions seven is ν_μ . All ν 's become massless by the chiral symmetry to preserve chirality. The auxiliary dimensional leptons (Al) in the main dimension seven are μ_7 and τ_7 whose masses can be calculated by Eq. (13) derived from Eq. (11).

$$\begin{aligned}
M_{Al7,a} &= \frac{3}{2} M_{B6,0} \sum_{a=0}^a a^4 \\
&= \frac{3}{2} M_{\pi_{1/2}} \sum_{a=0}^a a^4 \quad ,
\end{aligned} \tag{13}$$

where $a = 1, 2$ for μ_7 and τ_7 , respectively.

For heavy quarks, q_7 (the main dimensional fermion, F_7 , for quarks in the main dimension seven) is 3μ instead of massless 3ν as in Eq. (9). According to the mass formula, Eq. (11), of the auxiliary fermion, the mass formula for the auxiliary quarks, $Aq_{7,a}$, is as follows.

$$\begin{aligned}
M_{Aq_{7a}} &= \frac{3}{2} M_{q_7} \alpha_7 \sum_{a=0}^a a^4 \\
&= \frac{3}{2} (3 M_\mu) \alpha_w \sum_{a=0}^a a^4 \quad ,
\end{aligned} \tag{14}$$

where $\alpha_7 = \alpha_w$, and $a = 1, 2, 3, 4,$ and 5 for $u_7/d_7, s_7, c_7, b_7,$ and $t_7,$ respectively.

The main dimensional lepton for the main dimension eight is ν_τ , whose mass is zero to preserve chirality. The heavy lepton for the main dimensional eight is μ' as the sum of $e, \mu,$ and μ_8 (auxiliary dimensional lepton). Because there are only three families for leptons, μ' is the extra lepton, which is "hidden". μ' can appear only as $\mu + \text{photon}$. The pairing of $\mu + \mu$ from the hidden μ' and regular μ may account for the occurrence of same sign dilepton in the high energy level [15]. For the main dimension eight, q_8 (the F_8 for quarks) is μ' instead of $3\mu'$, because the hiding of μ' allows q_8 to be μ' . The hiding of μ' for leptons is balanced by the hiding of b_8 for quarks. The calculated masses are in good agreement with the observed constituent masses of leptons and quarks [16,17]. The observed mass of the top quark is $174.3 \pm 5.1 \text{ GeV}$ [17] in a good agreement with the calculated value, 176.5 GeV .

The masses of leptons, quarks, gauge bosons, and hadrons are calculated with only four known constants: the number of spatial dimensions, the mass of electron, the mass of Z° , and α_e . Most importantly, the calculation shows that exactly seven main and seven auxiliary dimensional orbitals are needed for all fundamental interactions, leptons, and quarks.

8. *The masses of hadrons*

As molecules are the composites of atoms, hadrons are the composites of elementary particles. Hadron can be represented by elementary particles in many different ways. One way to represent hadron is through the nonrelativistic constituent quark model where the mass of a hadron is the sum of the masses of quarks plus a relatively small binding energy. $\pi_{1/2}$ (u or d pseudoscalar quark), u, d, s, c, b, and t quarks mentioned above are nonrelativistic constituent quarks. On the other hand, except proton and neutron, all hadrons are unstable, and decay eventually into low-mass quarks and leptons. Is a hadron represented by all nonrelativistic quarks or by low-mass quarks and leptons? This paper proposes a dual formula: the full quark formula for all nonrelativistic constituent quarks ($\pi_{1/2}$, u, d, s, c, and b) and the basic fermion formula for the lowest- mass quark ($\pi_{1/2}$ and u) and lepton (e). The calculation of the masses of hadrons requires both formulas. The full quark formula sets the initial mass for a hadron. This initial mass is matched by the mass resulted from the combination of various particles in the basic fermion formula. The mass of a hadron is the mass of the basic fermion formula plus the binding energy (negative energy).

$$\begin{aligned}
M_{\text{full quark formula}} &\approx M_{\text{basic fermion formula}} \\
M_{\text{hadron}} &= M_{\text{basic fermion formula}} + \text{binding energy}
\end{aligned} \tag{15}$$

This dual representation of hadrons by the full quark formula and the basic quark formula is seen in the treatment of nucleons [18] as a mixture of the quark model and the vector meson dominance model which can be represented by $\pi_{1/2}$ and u.

The full quark formula consists of the vector quarks (u, d, s, c, and b) and pseudoscalar quark, $\pi_{1/2}$ (70.03 MeV), which is B_6 (dimension boson in the dimension six). The strong interaction converts B_6 ($\pi_{1/2}$) into pseudoscalar u and d quark in pseudoscalar π meson. The combination of the pseudoscalar quark ($\pi_{1/2}$) and vector quarks (q) results in hybrid quarks (q') whose mass is the average mass of pseudoscalar quark and vector quark.

$$Mq' = 1/2 (Mq + M\pi_{1/2}) \quad (16)$$

Hybrid quarks include u', d', and s' whose masses are 200.398, 201.164, and 314.148 MeV, respectively. For baryons other than n and p, the full quark formula is the combination of vector quarks, hybrid quarks (u' and d'), and pseudoscalar quark ($\pi_{1/2}$). For example, Λ° (uds) is u'd's3, where 3 denotes 3 $\pi_{1/2}$. For pseudoscalar mesons (J = 0), the full quark formula is the combination of $\pi_{1/2}$ and q' (u', d' and s') or $\pi_{1/2}$ alone. For vector mesons (J > 0), the full quark formula is the combination of vector quarks (u, d, s, c, and b) and $\pi_{1/2}$. For examples, π° is 2, η (1295, J=0) is u'u'd'd'8, and K_1 (1400, J=1) is ds8. The compositions of hadrons from the particles of the full quark formula are listed in Table 4.

Table 4. Particles for the full quark formula

	$\pi_{1/2}$	u', d'	s'	u, d	s	c, b
mass (MeV) (from Tables 1 and 3 and Eq. (16))	70.025	200.40, 201.16	314.15	330.8, 332.3	558	1701, 5318
n and p				√		
baryons other than n and p	√	√			√	√
mesons (J = 0) except $c\bar{c}$ and $b\bar{b}$	√	√	√			√
mesons (J > 0) and $c\bar{c}$ and $b\bar{b}$	√			√	√	√

Different energy states in hadron spectroscopy are the results of differences in the numbers of $\pi_{1/2}$. The higher the energy state, the higher the number of $\pi_{1/2}$. The number of $\pi_{1/2}$ attached to q or q' in the full quark formula is restricted. The number of $\pi_{1/2}$ can be 0, ± 1 , ± 3 , ± 4 , ± 7 , and $\pm 3n$ ($n > 2$). This is the “ $\pi_{1/2}$ series.” The number, 3, is indicative of a baryon-like (3 quarks in a baryon) number. The number, 4, is the combination of 1 and 3,

while the number, 7, is the combination of 3 and 4. Since $\pi_{1/2}$ is essentially pseudoscalar u and d quarks, the $\pi_{1/2}$ series is closely related to u and d quarks. Since s is close to u and d, the $\pi_{1/2}$ series is also related to s quark. Each presence of u, d, and s associates with one single $\pi_{1/2}$ series. The single $\pi_{1/2}$ series associates with baryons and the mesons with $u \bar{u}$, $d \bar{d}$, $s \bar{s}$, $d \bar{c}$, $u \bar{c}$, $d \bar{b}$, $u \bar{b}$, $s \bar{c}$, and $s \bar{b}$. The combination of two single $\pi_{1/2}$ series is the double $\pi_{1/2}$ series (0, ± 2 , ± 6 , ± 8 , ± 14 , and $\pm 6n$) associating with $u \bar{d}$ and $d \bar{s}$ mesons. For the mesons ($c \bar{c}$ and $b \bar{b}$) without u, d, or s, the numbers of $\pi_{1/2}$ attached to quarks can be from the single $\pi_{1/2}$ series or the double $\pi_{1/2}$ series.

The basic fermion formula is similar to M. H. MacGregor's light quark model [19], whose calculated masses and the predicted properties of hadrons are in very good agreement with observations. In the light quark model, the mass building blocks are the "spinor" (S with mass 330.4 MeV) and the mass quantum (mass = 70MeV). For the basic fermion formula, S is u, the quark with the lowest mass (330.77 MeV), and the basic quantum is $\pi_{1/2}$ (mass = 70.025 MeV). For examples, in the basic fermion formula, neutron is SSS, and K_1 (1400, J=1) is $S_2 11$, where S_2 denotes two S, and 11 denotes 11 $\pi_{1/2}$'s.

In additional to S and $\pi_{1/2}$, the basic fermion formula includes P (positive charge) and N (neutral charge) with the masses of proton and neutron. As in the light quark model, the mass associated with positive or negative charge is the electromagnetic mass, 4.599 MeV, which is nine times the mass of electron. This mass (nine times the mass of electron) is derived from the baryon-like electron that represents three quarks in a baryon and three electron in d_6 quark as in Table 2. This electromagnetic mass is observed in the mass difference between π^0 (2) and π^+ (2+) where + denotes positive charge. The calculated mass different is one electromagnetic mass, 4.599 MeV, in good agreement with the observed mass difference, 4.594 MeV, between π^0 and π^+ . (The values for observed masses are taken from "Particle Physics Summary "[17].) The particles in the basic fermion formula are listed in Table 5.

Table 5. Particles in the basic fermion formula

	S	$\pi_{1/2}$	N	P	electromagnetic mass
mass (MeV)	330.77	70.0254	939.565	938.272	4.599

Hadrons are the composites of quarks as molecules composing of atoms. As atoms are bounded together by chemical bonds, hadrons are bounded by "hadronic bonds," connecting the particles in the basic fermion formula. These hadronic bonds are similar to the hadronic bonds in the light quark model.

The hadronic bonds are the overlappings of the auxiliary dimensional orbits. As in Eq (11), the energy derived from the auxiliary orbit for S (u quark) is

$$\begin{aligned}
 E_a &= (3/2) (3 M_\mu) \alpha_w \\
 &= 14.122 \text{ MeV}
 \end{aligned}
 \tag{17}$$

The auxiliary orbit is a charge - dipole interaction in a circular orbit as described by A. Barut [14], so a fermion for the circular orbit and an electron for the charge are embedded in this hadronic bond. The fermion for the circular orbit is the supersymmetrical fermion for the auxiliary dimensional orbit according to Eq (2).

$$M_f = E_a \alpha_w \quad (18)$$

The binding energy (negative energy) for the bond (S , S) between two S's is twice of 14.122 MeV minus the masses of the supersymmetrical fermion and electron.

$$\begin{aligned} E_{S-S} &= -2 (E_a - M_f - M_e) \\ &= -26.384 \text{ MeV} \end{aligned} \quad (19)$$

It is similar to the binding energy (-26 MeV) in the light quark model. An example of S , S bond is in neutron (S – S – S) which has two S – S bonds. The mass of neutron can be calculated as follows.

$$\begin{aligned} M_n &= 3M_S + 2E_{S-S} \\ &= 939.54 \text{ MeV}, \end{aligned} \quad (20)$$

which is in excellent agreement with the observed mass, 939.57 MeV. The mass of proton is the mass of neutron minus the mass difference (three times of electron mass = M_{3e}) between u and d quark as shown in Table 2. Proton is represented as S – S – (S -3e). The calculation of the mass of proton is as follows.

$$\begin{aligned} E_a \text{ for } (S-3e) &= (3/2) (3 (M_\mu - M_{3e})) \alpha_w \\ M_f &= E_a \alpha_w \\ M_p &= 2 M_S + M_{(S-3e)} + E_{S,S} + E_{S,(S-3e)} \\ &= 938.21 \text{ MeV} \end{aligned} \quad (21)$$

The calculated mass is in a good agreement with the observed mass, 938.27 MeV.

The binding energy for $\pi_{1/2} - \pi_{1/2}$ bond can be derived in the same way as Eqs (17), (18), and (19).

$$\begin{aligned} E_a &= (3/2) M\pi_{1/2} \alpha_w \\ M_f &= E_a \alpha_w \\ E_{\pi_{1/2} - \pi_{1/2}} &= -2 (E_a - M_f - M_e) \\ &= -5.0387 \text{ MeV} \end{aligned} \quad (22)$$

It is similar to the binding energy (-5 MeV) in the light quark model. An example for the binding energy of $\pi_{1/2} - \pi_{1/2}$ bond is in π^0 . The mass of π^0 can be calculated as follows.

$$M_{\pi^0} = 2 M\pi_{1/2} + E_{\pi_{1/2} - \pi_{1/2}}$$

$$= 135.01 \text{ MeV.} \quad (23)$$

The calculated mass of π^0 is in excellent agreement with the observed value, 134.98 MeV. There is one $\pi_{1/2} - \pi_{1/2}$ bond per pair of $\pi_{1/2}$'s, so there are two $\pi_{1/2} - \pi_{1/2}$ bonds for 4 $\pi_{1/2}$'s, and three $\pi_{1/2} - \pi_{1/2}$ bonds for 6 $\pi_{1/2}$'s.

Another bond is $N - \pi_{1/2}$ or $P - \pi_{1/2}$, the bond between neutron or proton and $\pi_{1/2}$. Since N is SSS, $N - \pi_{1/2}$ bond is derived from $S - \pi_{1/2}$. The binding energy of $S - \pi_{1/2}$ is the average between $S-S$ and $\pi_{1/2} - \pi_{1/2}$.

$$E_{S - \pi_{1/2}} = 1/2 (E_{S-S} + E_{\pi_{1/2} - \pi_{1/2}}) \quad (24)$$

An additional dipole (e^+e^-) is needed to connected $S - \pi_{1/2}$ to neutron.

$$\begin{aligned} E_{N - \pi_{1/2}} &= E_{S - \pi_{1/2}} + 2 M_e \\ &= -14.689 \text{ MeV.} \end{aligned} \quad (25)$$

It is similar to -15 MeV in the light quark model. An example for $P - \pi_{1/2}$ is Σ^+ which is represented by P4 whose structure is $2 - P - 2$. The 4 $\pi_{1/2}$'s are connected to P with two $P - \pi_{1/2}$ bonds. The mass of Σ^+ is as follows.

$$\begin{aligned} M_{\Sigma^+} &= M_P + 4 M_{\pi_{1/2}} + 2 E_{N - \pi_{1/2}} \\ &= 1189.0 \text{ MeV.} \end{aligned} \quad (26)$$

The calculated mass is in a good agreement with the observed mass, $1189.4 \pm 0.07 \text{ MeV}$.

There is $N - N$ hadronic bond between two N's. N has the structure of $S - S - S$. $N - N$ has a hexagonal structure shown in Fig. 9.




Fig. 9. The structure of $N - N$

There are two additional $S - S$ for each N. The total number of $S - S$ bonds between two N's is 4. An example is Ω_c^0 (ssc) which has the basic fermion formula of $N_2 S_9$ with the structure of $3 - N - N - 6 S$. The mass of Ω_c^0 can be calculated as follows.

$$\begin{aligned} M_{\Omega_c^0} &= 2M_N + M_S + 9M_{\pi_{1/2}} + 4 E_{S-S} + 2 E_{N - \pi_{1/2}} \\ &= 2705.3 \text{ MeV} \end{aligned} \quad (27)$$

The calculated mass for Ω_c° is in a good agreement with the observed mass (2704 ± 4 MeV). For baryons other than p and n, there are two or three N - $\pi_{1/2}$ or P - $\pi_{1/2}$ bonds per baryon. It is used to distinguish $J = 1/2$ and $J \geq 3/2$. For $J = 1/2$ which have asymmetrical spins (two up and one down), there are two bonds for three S in a baryon to represent the asymmetrical spins. For $J \geq 3/2$, there are three bonds for three S in a baryon except if there is only one $\pi_{1/2}$, only two bonds exists for a baryon.

Among the particles in the basic fermion formula, there are hadronic bonds, but not all particles have hadronic bonds. In the basic fermion formula, hadronic bonds appear only among the particles that relate to the particles in the corresponding full quark formula. The related particles are the “core” particles that have hadronic bonds, and the unrelated particles are “filler” particles that have no hadronic bonds. In the basic fermion formula, for baryons other than p and n, the core particles are P, N, and $\pi_{1/2}$. For the mesons consisting of u and d quarks, the core particles are $\pi_{1/2}$, S, and N. For the mesons containing one u, d, or s along with s, c, or b, the core particles are S and N, and no hadronic bond exist among $\pi_{1/2}$'s, which are the filler particles. For the mesons ($c \bar{c}$ and $b \bar{b}$), the only hadronic bond is N - N. The occurrences of hadronic bonds are listed in Table 6.

Table 6. Hadronic bonds in hadrons

	S - S	$\pi_{1/2} - \pi_{1/2}$	N(P) - $\pi_{1/2}$	N - N
binding energy (MeV)	-26.384	-5.0387	-14.6894	2 S-S per N
baryons other than n and p	√		√	√
mesons with u and d only	√	√		√
mesons containing one u, d, or s along with s, c, or b	√			√
$c \bar{c}$ or $b \bar{b}$ mesons				√

An example is the difference between π and f_0 . The decay modes of f_0 include the mesons of s quarks from K meson. Consequently, there is no $\pi_{1/2} - \pi_{1/2}$ for f_0 . The basic fermion formula for f_0 is 14. The mass of f_0 is as follows.

$$\begin{aligned} M &= 14M\pi_{1/2} \\ &= 980.4 \text{ MeV} \end{aligned} \tag{28}$$

The observed mass is 980 ± 10 MeV.

In addition to the binding energies for hadronic bonds, hadrons have Coulomb energy (-1.2 MeV) between positive and negative charges and magnetic binding energy (± 2 MeV per interaction) for S - S from the light quark model [19]. In the light quark

model, the dipole moment of a hadron can be calculated from the magnetic binding energy. Since in the basic fermion formula, magnetic binding energy becomes a part of hadronic binding energy as shown in Eq (19), magnetic binding energy for other baryons is the difference in magnetic binding energy between a baryon and n or p. If a baryon has a similar dipole moment as p or n, there is no magnetic binding energy for the baryon. An example for Coulomb energy and magnetic binding energy is Λ (uds, J=1/2) whose formula is P3- with the structure of 2 – P – 1- where "-" denotes negative charge. The dipole moment of Λ is $-6.13 \mu_N$, while the dipole moment of proton (P) is $2.79 \mu_N$ [19]. According to the light quark model, this difference in dipole moment represents -6 MeV magnetic binding energy. The Coulomb energy between the positive charge P and the negative charge 1- is -1.2 MeV. The electromagnetic mass for 1- is 4.599 MeV. The mass of Λ is calculated as follows.

$$\begin{aligned} M_{\Lambda} &= M_p + 3M_{\pi_{1/2}} + M_{e.m.} + 2 E_{N-\pi_{1/2}} + E_{mag} + E_{coul} \\ &= 1116.4 \text{ MeV} \end{aligned} \quad (29)$$

The observed mass is 1115.7 ± 0.0006 MeV.

An example of the dual representation of the full quark formula and the basic fermion formula as expressed by Eq. (15) is Λ (uds). The full quark formula for Λ is u'd's'3 with mass of 1169.9 MeV. This mass is matched by the mass of the basic fermion formula, P3- with the mass of 1152.9 MeV. The final mass of Λ is 1152.9 MeV plus the various binding energies.

Table 7 is the results of calculation for the masses of baryons selected from Ref. [17]. The hadrons selected are the hadrons with precise observed masses and precise known quantum states such as isospin, spin, and decay mode.

Table 7. The masses of baryons

Baryons	$I(J^P)$	Full quark formula	Basic fermion formula	Calculated mass	Observed mass	Difference
<u>n and p</u>						
n	$1/2(1/2^+)$	udd	SSS	939.54	939.57	-0.03
p	$1/2(1/2^+)$	uud	SSS-3e	938.21	938.27	-0.06
<u>uds, uus, dds</u>						
Λ°	$0(1/2^+)$	u'd's3	P3-	1116.4	1115.7	0.7
Σ^+	$1(1/2^+)$	u'u's4	P4	1189.0	1189.4	-0.4
Σ°	$1(1/2^+)$	u'd's4	P4-	1192.4	1192.6	-0.2
Σ^-	$1(1/2^+)$	d'd's4	N4-	1194.9	1197.4	-2.5
Σ^+	$1(3/2^+)$	u'u's7	P7	1384.4	1382.8	1.6
Σ°	$1(3/2^+)$	u'd's7	P7-	1381.8	1383.7	-1.9
Σ^-	$1(3/2^+)$	d'd's7	N7-	1390.3	1387.2	3.1
$\Lambda^\circ (s_{01})$	$0(1/2^-)$	u'd's7	P7-	1402.5	1406.0	-3.5
$\Lambda^\circ (D_{03})$	$0(3/2^-)$	u'd's9	N9	1525.7	1519.5	6.2
<u>uss, dss</u>						
Ξ°	$1/2(1/2^+)$	u'ss	NS1	1311.0	1314.9	-3.9
Ξ^-	$1/2(1/2^+)$	d'ss	NS1-	1315.6	1321.3	-5.7
Ξ°	$1/2(3/2^+)$	u'ss4	N9+-	1533.7	1531.8	1.9
Ξ^-	$1/2(3/2^+)$	d'ss4	N10-	1530.3	1535.0	-4.7
Ξ°	$1/2(3/2^-)$	u'ss9	$N_2 1+-$	1822.2	1823.0	-0.8
<u>udc, ddc, uuc</u>						
Λ_c^+	$0(1/2^+)$	u'd'c3	PS15	2290.0	2284.9	5.1
Σ_c^0	$1(1/2^+)$	d'd'c7	$N_2 10$	2452.5	2452.2	0.3
Σ_c^{++}	$1(1/2^+)$	u'u'c7	$N_2 10^{++}$	2452.5	2452.8	-0.3
Σ_c^+	$1(1/2^-)$	u'd'c7	$N_2 10^+$	2449.1	2453.6	-4.5
Λ_c^+	$0(1/2^-)$	u'd'c9	$N_2 12^+$	2589.1	2593.9	-4.8
<u>usc, dsc</u>						
Ξ_c^+	$1/2(1/2^+)$	u'sc1	NS ₃ 8+	2467.3	2466.3	1.0
Ξ_c°	$1/2(1/2^+)$	d'sc1	$N S_3 8^{+-}$	2470.7	2471.8	-1.1
<u>sss</u>						
Ω^-	$0(3/2^+)$	sss	NS6-	1665.7	1672.5	-6.8
<u>ssc</u>						
Ω_c°	$0(1/2^+)$	ssc	$N_2 S_9$	2705.2	2704.0	1.2
<u>udb</u>						
Λ_b°	$0(1/2^+)$	u'd'b1	$N_2 S_9 13$	5631.5	5624.0	7.5

The full quark formula for baryons other than p and n involves hybrid quarks (u' and d'), vector quarks (s, c, and b), and pseudoscalar quark ($\pi_{1/2}$). The whole spectrum of baryons (other than p and n) follows the single $\pi_{1/2}$ series with 0, 1, 3, 4, 7, and 3n. $\pi_{1/2}$ is

closely related to u and d, so the lowest energy state baryons (Λ and Σ) that contain low mass quark and high number of u or d have high number of $\pi_{1/2}$. . On the contrary, the lowest energy state baryons (Ω , Ω_c , and Λ_b) that contain high mass quarks and no or low number of u and d have low number of $\pi_{1/2}$ or no $\pi_{1/2}$. $\pi_{1/2}$'s are added to the lowest energy state baryons to form the high-energy state baryons.

The basic fermion formula includes P, N, S, and $\pi_{1/2}$. If the dominant decay mode includes p, the basic fermion formula includes P. If the dominant decay mode includes n, the basic fermion formula includes N. If the decay mode does not include either n or p directly, the baryon with the dipole moment similar to p or n has the basic fermion formula with P or N, respectively. If the dipole moment is not known, the designation of P or N for the higher energy state baryon follows the designation of P or N for the corresponding lower energy state baryon. Only N is used in the multiple nucleons such as N_2 (two N's).

The result of calculated masses for light unflavored mesons is listed in Table 8. Light unflavored mesons have zero strange, charm, and bottom numbers.

Table 8. Light unflavored mesons

Meson	$I(J^{PC})$	Full quark formula	Basic fermion formula	Calculated mass.	Observed mass	Difference
<u>J = 0, only u, d, or lepton in decay mode</u>						
π^0	1 (0 ⁺⁺)	2	2	135.01	134.98	0.04
π^\pm	1 (0 ⁻)	2	2 \pm	139.61	139.57	0.04
η	0 (0 ⁺⁺)	8	8+-	548.0	547.3	0.7
η'	0 (0 ⁺⁺)	14	14+-	958.1	957.8	0.3
η	0 (0 ⁺⁺)	u'u'u'd'd'd'2	S ₃ 5+-	1292.6	1297.0	-4.4
f ₀	0 (0 ⁺⁺)	u'u'u'd'd'd'6	S ₄ 4	1514.0	1500.0	14.0
<u>J > 0, only u, d, or lepton in decay modes</u>						
ρ	1 (1 ⁻)	ud2	S ₂ 2	770.2	769.3	0.9
ω	0 (1 ⁻)	ud2	S ₂ 2+-	785.4	788.6	-3.2
h ₁	0 (1 ⁺)	ud8	S ₂ 8	1176.2	1170.0	6.2
ω	0 (1 ⁻)	uudd2	S ₃ 7+-	1415.6	1419.0	-3.4
ω	0 (1 ⁻)	uudd6	S ₄ 6	1650.0	1649.0	1.0
ω_3	0 (3 ⁻)	uudd6	S ₄ 6	1662.0	1667.0	-5.0
<u>J = 0, u and d in major decay modes, s in minor decay modes</u>						
f ₀	0 (0 ⁺⁺)	u'd's'4	14	980.4	980.0	0.4
a ₀	1 (0 ⁺⁺)	u'd's'4	14	985.4	984.8	0.6
<u>J > 0, u and d in major decay modes, s in minor decay mode</u>						
a ₁	1 (1 ⁺⁺)	uds	S13	1232.7	1230.0	2.7
b ₁	1 (1 ⁺)	uds	S13	1232.7	1239.5	-6.8
f ₂	0 (2 ⁺⁺)	uds1	S ₂ 9	1278.4	1275.4	3.0
f ₁	0 (1 ⁺⁺)	uds1	S ₂ 9+-	1278.4	1281.9	-3.5
a ₂	1 (2 ⁺⁺)	uds1	S14	1316.2	1318.1	-1.9
f ₁	0 (1 ⁺⁺)	uds4	S ₃ 7	1429.7	1426.3	3.4
ρ	1 (1 ⁻)	uds4	S ₂ 12	1475.5	1465.0	10.5
π_2	1 (2 ⁺)	uds7	S ₂ 15	1685.5	1670.0	15.5
ρ_3	1 (3 ⁻)	uds7	S ₂ 15	1693.5	1691.0	2.5
ρ	1 (1 ⁻)	uds7	S ₂ 15+-	1698.6	1700.0	-1.4
f ₄	0 (4 ⁺⁺)	uds12	S ₄ 7	2038.5	2034.0	4.5
<u>s in major decay modes</u>						
Φ	0 (1 ⁻)	ss-1	S ₃ +-	1017.6	1019.4	-1.8
f ₂ '	0 (2 ⁺⁺)	ss7	S ₄ 4+-	1532.0	1525.0	7.0
Φ	0 (1 ⁻)	ss9	S ₄ 6+-	1672.1	1680.0	-7.9
Φ_3	0 (3 ⁻)	ss12	S ₆	1852.7	1854.0	-1.3
f ₂	0 (2 ⁺⁺)	ss15	S ₆ 2	2000.7	2011.0	-10.3
f ₂	0 (2 ⁺⁺)	ss18	N S ₃ 6	2299.3	2297.0	2.3
f ₂	0 (2 ⁺⁺)	ss18	N S ₄ 2+-	2331.5	2339.0	-7.5

The full quark formulas for light unflavored mesons are different for different decay modes. There are three different types of the full quark formula for light unflavored mesons. Firstly, when the decay mode is all leptons or all mesons with u and d quarks, the full quark formula consists of $\pi_{1/2}$, u', d', u, and d quarks. Secondly, when u, d, and leptons are in the major decay modes, and s is in the minor decay modes, the full quark formula consists of $\pi_{1/2}$, u', d', s', u, d, and s quarks. The most common full quark formula for such mesons is uds, which is essentially $\frac{1}{2}(u \bar{u} d \bar{d} s \bar{s})$. Finally, when s quarks are in the major or all decay mode, the full quark formula consists of $\pi_{1/2}$'s and s quarks. When $J = 0$, the full quark formula includes $\pi_{1/2}$'s and hybrid quarks (u', d', and s'). When $J > 0$, the full quark formula includes $\pi_{1/2}$'s and vector quarks (u, d, and s). Since there are double presence of u and d quarks for the full quark formula with u and d quark, it follows the double $\pi_{1/2}$ series: 0, 2, 6, 8, 14, and $6n$. The full quark formula with odd presence of u, d, and s follows the single $\pi_{1/2}$ series: 0, ± 1 , ± 3 , ± 4 , ± 7 , and $\pm 3n$.

The basic fermion formula includes $\pi_{1/2}$'s, S, and N. $\pi_{1/2} - \pi_{1/2}$ hadronic bond exists in the full quark formula with u and d quarks, and does not exist in the full quark formula with s quark. S – S bond exists in all formula. Light unflavored mesons decay into symmetrical low mass mesons, such as 2γ , $3\pi^0$, and $\pi^0 2\gamma$ from η , or asymmetrical high and low mass mesons, such as asymmetrical $\pi^+\pi\eta$, $\rho^0\gamma$, and $\pi^0\pi^0\eta$ from η' . To distinguish the asymmetrical decay modes from the symmetrical decay modes, one “counter $\pi_{1/2} - \pi_{1/2}$ hadronic bond” is introduced in η' [19]. The binding energy for the counter $\pi_{1/2} - \pi_{1/2}$ hadronic bond is 5.04 MeV, directly opposite to -5.04 MeV for the $\pi_{1/2} - \pi_{1/2}$ hadronic bond. All light unflavored mesons with asymmetrical decay modes include this counter $\pi_{1/2} - \pi_{1/2}$ hadronic bonds. For an example, the mass of η' ($14+-$) is calculated as follows.

$$\begin{aligned} M_{\eta'} &= 14 M_{\pi_{1/2}} + 2 M_{e.m.} + E_{e.m.} + 7E_{\pi_{1/2} - \pi_{1/2}} - E_{\pi_{1/2} - \pi_{1/2}} \\ &= 958.2 \text{ MeV} \end{aligned} \quad (30)$$

The observed mass is 957.8 ± 0.14 MeV.

The result of the mass calculation for the mesons consisting of u, d, or s with s, c, or b is listed in Table 9.

Table 9. Mesons with s, c, and b

Meson	J^{PC}	Full quark formula	Basic fermion formula	Calculated mass.	Observed mass	Difference
<u>Light strange mesons</u>						
K^\pm	0^-	7	7_\pm	494.8	493.7	1.1
K^0	0^-	7	7_{+-}	498.2	497.7	0.5
K^*	0^+	d's'14	$S_2 11_{+-}$	1413.4	1412.0	1.4
K^*	1^-	us	$S8_\pm$	895.6	891.6	4.0
K^*	1^-	ds	S8	899.0	896.1	2.9
K_1	1^+	ds6	$S_2 9$	1273.4	1273.0	0.4
K_1	1^+	ds8	$S_2 11$	1405.4	1402.0	3.4
K^*	1^-	ds8	$S_2 11_{+-}$	1413.4	1414.0	-0.6
$K^{*\pm}$	2^+	us8	$N7_\pm$	1434.3	1425.6	8.7
K^*	2^+	ds8	$N7_{+-}$	1437.7	1432.4	5.3
K^*	1^-	ddss	$S_3 11_{+-}$	1717.8	1717.0	0.8
K_2	2^-	ddss	N12	1779.9	1773.0	6.9
K_3^*	3^-	ddss	N12	1779.9	1776.0	3.9
K_2	2^-	ds14	$S_4 8_{+-}$	1812.1	1816.0	-3.9
K_4^*	4^+	ds18	$S_5 7_{+-}$	2046.5	2045.0	1.5
<u>Charmed mesons</u>						
D^0	0^-	u'c1	S_6_{+-}	1860.7	1864.5	-3.8
D^\pm	0^-	d'c1	S_6_\pm	1857.3	1869.3	-12.0
D^{*0}	1^-	uc1	$S_6 2_{+-}$	2000.7	2006.7	-6.0
$D^{*\pm}$	1^-	dc1	$S_6 2_\pm$	1997.4	2010.0	-12.6
D_1	1^+	uc7	$S_6 8_{+-}$	2420.9	2422.2	-1.3
D_2^*	2^+	uc7	$N S_4 4$	2463.6	2458.9	4.7
$D_2^{*\pm}$	2^+	dc7	$N S_4 4_\pm$	2468.2	2459.0	9.2
<u>Charmed strange mesons</u>						
D_s	0^-	s'c1	$S_5 6$	1968.5	1968.6	0.1
D_{s1}	1^+	sc4	NS18	2535.4	2535.4	0.0
<u>Bottom mesons</u>						
B^\pm	0^-	u'b	$N_4 S20_\pm$	5283.1	5279.0	4.1
B^0	0^-	d'b	$N_4 S 20$	5278.5	5279.4	-0.9
B^*	1^-	db	N_6	5320.8	5325.0	-4.2
B_s	0^-	s'b	$N_5 S_3$	5373.5	5369.6	3.9

The full quark formula for these mesons contains $\pi_{1/2}$'s, u', d', s', u, d, s, c, and b. The full quark formula for the mesons with $J = 0$ contains $\pi_{1/2}$, hybrid quarks (u', d', and s'), c, and b quarks. The full quark formula with $J > 0$ contains $\pi_{1/2}$'s and vector quarks (u, d, s, c, and b). Strange mesons have double presence of d and s, so it follows the double

$\pi_{1/2}$ series. All other mesons have single presence of u, d, or s, so they follow the single $\pi_{1/2}$ series. The basic fermion formula consists of $\pi_{1/2}$'s, S, and N. There is no $\pi_{1/2} - \pi_{1/2}$ bond. There are S – S and N – N bonds. For example, the mass of B_s ($N_5 S_3$) with the observed mass as 5369.6 ± 2 MeV is calculated as follows.

$$\begin{aligned} M &= 5M_N + 3M_S + (10+ 2) M_{S-S} \\ &= 5373.5 \text{ MeV} \end{aligned} \quad (31)$$

There are 10 S – S bonds for N_5 (two S – S bonds for each N) and 2 S – S bonds for S_3 . The result of mass calculation for c \bar{c} and b \bar{b} mesons is listed in Table 10.

Table 10. c \bar{c} and b \bar{b} mesons

Meson	J ^{PC}	Full quark formula	Basic fermion formula	Calculated mass.	Observed mass	Difference
<u>c \bar{c} mesons</u>						
η_c (1s)	0 ⁺	cc-6	N ₃ S ₃ 15	2982.3	2979.8	2.5
J/ ψ	1 ⁻	cc-3	N ₂ S ₄	3096.7	3096.9	-0.2
χ_c (1p)	0 ⁺⁺	cc2	N ₂ S ₂ 14	3415.5	3415.0	0.5
χ_c' (1p)	1 ⁺⁺	cc3	N ₂ S ₄ 6	3516.8	3510.5	6.5
χ_{c2} (1p)	2 ⁺⁺	cc4	N ₂ S ₂ 16	3555.5	3556.2	-0.7
U _{2s}	1 ⁻	cc6	N ₃ S ₁ 10	3691.4	3686.0	5.4
ψ	1 ⁻	cc7	N ₃ S ₁ 11	3761.4	3769.9	-8.5
ψ	1 ⁻	cc12	N ₄ 7	4037.4	4040.0	-2.6
ψ	1 ⁻	cc14	N ₄ S ₄	4158.1	4159.0	-0.9
ψ	1 ⁻	cc18	N ₄ S ₂ 3	4418.8	4415.0	3.8
<u>b \bar{b} mesons</u>						
Υ (1s)	1 ⁻	bb-12	N ₇ S ₆ 18	9452.7	9460.3	-7.6
χ_b (1p)	0 ⁺⁺	bb-4	N ₁₀ S ₃	9860.3	9859.9	0.4
χ_{b1} (1p)	1 ⁺⁺	bb-3	N ₁₀ S ₁ 10	9899.0	9892.7	6.3
χ_{b2} (1p)	2 ⁺⁺	bb-3	N ₁₀ 15	9918.4	9912.6	5.8
Υ (2s)	1 ⁻	bb-1	N ₁₀ S ₂ 7	10019.7	10023.3	-3.6
χ_{b0} (1p)	0 ⁺⁺	bb2	N ₁₀ S ₂ 10	10229.8	10232.1	-2.3
χ_{b1} (2p)	1 ⁺⁺	bb2	N ₁₀ S ₄ 1	10261.1	10255.2	5.9
χ_{b2} (2p)	2 ⁺⁺	bb2	N ₁₀ S ₄ 1	10261.1	10268.5	-7.4
Υ (3s)	1 ⁻	bb3	N ₁₀ S ₃ 7	10350.5	10355.3	-4.8
Υ (4s)	1 ⁻	bb7	N ₁₁ S ₇	10575.7	10580.0	-4.3
Υ	1 ⁻	bb12	N ₁₂ 3	10851.6	10865.0	-13.4
Υ	1 ⁻	bb14	N ₁₁ S ₃ 4	11027.2	11019.0	8.2

The full quark formula contains c or b. Since it contains no u, d, or s related to $\pi_{1/2}$, it follows a mixed $\pi_{1/2}$ series from both the single $\pi_{1/2}$ series and the double $\pi_{1/2}$ series. Since c

and b are high mass quarks unlike the low mass u and d quarks that relate to $\pi_{1/2}$, the $\pi_{1/2}$ series actually starts from a negative $\pi_{1/2}$. The basic fermion formula contains $\pi_{1/2}$'s, S, and N. The only hadronic bond is N – N. An example is J/ ψ ($N_2 S_4$) whose mass (3096.9 ± 0.04 MeV) is calculated as follows.

$$\begin{aligned} M &= 2 M_N + 4 M_S + 4 M_{S-S} \\ &= 3096.7 \text{ MeV} \end{aligned} \tag{32}$$

9. *The Cyclic Universe*

The hidden universe and the observable universe are parallel to each other. The hidden universe is hidden from the observable universe until the quintessence transition [20], when the hidden universe fractionalizes into mixed 3-particle, compatible with the empty mixed 3-particle (exclusive vacuum) overlapping with inclusive vacuum in the observable universe. The empty mixed 3-particle with negative tension from the observable universe and the mixed 3-particle with positive tension from the hidden universe form the five dimensional bulk as in the two-sided Randall - Sundrum model [1] as Fig. 10.




Fig. 10: Quintessence

The five-dimensional bulk provides space for quintessence as the scalar field from the hidden universe. When the hidden universe is in the fractionalization (expansion) phase from the mixed 4-particles to the mixed 3-particles, quintessence accelerates expansion in the observable universe. After a certain period in the hidden universe, the mixed 3-particle starts to condense into the mixed 4-particle, inducing contraction in the observable universe. When all of the hidden universe mixed 3-particles are converted into the mixed 4-particles, quintessence ceases to exist.

Quintessence controls the rate of expansion and contraction for the observable universe during the quintessence period. It makes the hidden and the observable universes to have the same size and to contract at the same rate, so eventually both the observable universe and the hidden universe end at the same time.

At the end of the contraction, the hidden universe becomes the hidden mixed 9-particle, and the observable universe becomes a cosmic black hole without exclusive vacuum. Without exclusive vacuum, the cosmic black hole is converted back into the observable mixed 9-particle as in the beginning of the mixed pre-expanding universe. At the end of the contraction, the universe again becomes the mixed pre-expanding universe

with the 9-mixed particles. This mixed pre-expanding universe then starts another cycle of the universe as Fig. 11.




Fig. 11: the cyclic universe

10. Conclusion

The cosmic evolutionary theory is by means of the evolution of object in four stages: string, membrane, particle, and particle-wave. It is also concurrently by means of the evolution of vacuum in four stages: object vacuum, single vacuum, inclusive vacuum, and exclusive vacuum.

The universe starts with the pre-universe as object vacuum-object, where object and vacuum take turn to exist equally at the same location. It is an equilibrium state between vacuum and the pair of ten-dimensional superstring and anti-superstring with a non-zero vacuum energy. The vacuum fluctuation of object vacuum results in the pre-expanding universe with the boundary 9-brane and 9-antibrane separated by single vacuum each of which exists with object in one object to one vacuum relation. The collapse and the bounce of the pre-expanding universe result in zero vacuum energy, which induces the fractionalization of one object into multiple low energy objects. The multiple objects require inclusive vacuum each of which associates with multiple objects. Such fractionalization brings about the cosmic expansion that leads to the superluminal inflation to rupture the connection among objects. The gap generated from the rupture is exclusive vacuum that excludes objects. The rupture brings about the big bang, cosmic radiation, elementary particles, dark matter, entropy, force fields, and quantum mechanics.

The cosmic evolution involves cyclic universe: the pre-universe, the pre-expanding universe, the mixed pre-expanding universe, the expanding universe, quintessence, the contracting universe, and back to the mixed pre-expanding universe as Fig. 11. The cosmic evolution provides the most comprehensive explanation for the origin of physical laws.

The periodic table of elementary particles (Fig. 8 and Table 2) derived from the cosmic evolution can account for all leptons, quarks, and gauge bosons. The calculated masses (Tables 1 and 3) of elementary particles derived from the periodic table are in good agreement with the observed values. For example, the calculated mass (176.5 GeV) of the top quark has an excellent agreement with the observed mass (174.3 ± 5.1 GeV).

A hadron can be represented by the full quark formula (Table 4) and the basic quark formula (Table 5). The full quark formula consists of all quarks, pseudoscalar quark in pion, and the hybrid quarks. The basic quark formula consists of the lowest mass quarks. The relation between these two formulas is expressed by Eq. (15). As a molecule is the composite of atoms with chemical bonds, a hadron is the composite of elementary particles with "hadronic bonds" (Table 6) which are the overlappings of the auxiliary dimensional orbits. The calculated masses (Tables 7, 8, 9, and 10) of hadrons are in good agreement with the observed values. For examples, the calculated masses for neutron and pion are 939.54 and 135.01 MeV in excellent agreement with the observed masses, 939.57 and 134.98 MeV, respectively. The overall standard deviation between the calculated masses and the observed masses for all hadrons (108 hadrons) is 5.1 MeV, comparing to ± 11.4 MeV, the average observed error for the masses of the hadrons.

The masses of gauge bosons, leptons, quarks, and hadrons can be calculated using only four known constants: the number of the extra spatial dimensions in the eleven dimensional membrane, the mass of electron, the mass of Z^0 , and α_e . The periodic table of elementary particles derived from the cosmic evolution provides the most comprehensive explanation and calculation for the masses of elementary particles and hadrons.

References

- [1] L. Randall and R. Sundrum, Nucl. Phys. **B557** (1999) 79; Phys. Rev. Lett. **83** (1999) 3370 ; Phys. Rev. Lett. **83** (1999) 4690
- [2] P. Horava and E. Witten, Nucl. Phys. **B475** (1996) 94 [hep-th/9603142].
- [3] J. Houry, B. A. Ovrut, P. J. Steinhardt, and N. Turok, hep-th/0103239; R. Kallosh, L. Kofman, and A. Linde, hep-th/0104073; S. Rasanen, hep-th/0111279.
- [4] G. Dvali, Q. Shafi, and S. Solganik, hep-th/0105203; C. P. Burgess, M. Majumdar, D. Nolte, F. Quevedo, G. Rajesh, and R. J. Zhang, hep-th/0105204.
- [5] A. Mazumdar, Phys.Lett. **B469** (1999) 55 [hep-ph/9902381]; A. Riotto, Phys. Rev. D **61** (2000) 123506; J. M. Cline, Phys. Rev. D **61** (2000) 02313; G. Dvali and S. H. H. Tye, Phys. Lett. **B450** (1999) 72; G. Shiu and S. H. H. Tye, hep-th/0106274; M. Bucher, hep-th/0107148; A. Nornov, hep-th/0109090; M.F. Parry, D. A. Steer, hep-th/0109207, D. Langlois, K. Maeda, D. Wards, gr-qc/0111013; J. Garriga, T. Tanaka, hep-th/0112028.
- [6] J. Houry, B. A. Ovrut, N. Seiberg, P. J. Steinhardt, and N. Turok, hep-th/0108187; P. J. Steinhardt, and N. Turok, hep-th/0111098; P. J. Steinhardt, and N. Turok, astro-

- ph/0112537; J. Martin, P. Peter, N. Pinto-Neto, and D. Schwarz, hep-th/0112128; G. Felder, A. Frolov, L. Kofman, and A. Linde; P. Steinhardt and N. Turok, astro-ph/0204479
- [7] D. Chung, hep-ph/0003237, D. Chung, *Speculations in Science and Technology* 20 (1997) 259; *Speculations in Science and Technology* 21(1999) 277
 - [8] D. Chung, hep-ph/0201115
 - [9] A. Linde, *Phys. Rev D***49** (1994) 748 [hep-ph/9307002], G. Felder, J. Garcia-Bellido, P. B. Greene, L. Kofman, A. Linde, I. Tkachev, *Phys. Rev. Lett.* **87** (2001) 011601 [hep-ph/0012142]; C. Herdeiro, S. Hirano, and R. Kallosh, hep-th/0111147.
 - [10] D. Tytler, S. Burles, L. Lu, X-M, Fan, A. Wolfe, and B. Savage, *AJ*, **117** (1999) 63; F. C. van den Bosch, A. Burkert, and R. A. Swaters, astro-ph/0105082
 - [11] P. Langacher, M. Luo, and A. Mann, *Rev. Mod. Phys.* **64** (1992) 87.
 - [12] E. Elbaz, J. Meyer, and R. Nahabetian, *Phys. Rev. D*23. (1981)1170
 - [13] A. O. Barut and D. Chung, *Lett. Nuovo Cimento* 38 (1983) 225
 - [14] A.O. Barut, *Phys. Rev. Lett.* 42.(1979) 1251
 - [15] L. Hall, R. Jaffe, J. Rosen1985, *Phys. Rep.* 125 (1985) 105
 - [16] C.P. Singh, *Phys. Rev. D*24 (1981) 2481; D. B. Lichtenberg *Phys. Rev. D*40 (1989) 3675
 - [17] D.E. Groom et al, *The European Physical Journal* **C15** (2000) 1
 - [18] G.E. Brown, M. Rho, V. Vento, *Phys Lett* 97B (1980); G.E. Brown, *Nucl. Phys.* A374 (1982) 630
 - [19] M.H. MacGregor, *Phys. Rev. D*9 (1974) 1259; *Phys. Rev. D*10 (1974) 850; *The Nature of the Elementary Particles* (Springer-Verlag, New York, 1978)
 - [20] N. Balcall, J.P. Ostriker, S. Perlmutter, and P.J. Steinhardt, *Science* **284**, 1481-1488, (1999); C. Arnedariz-Picon, V. Mukhanov, and P.J. Steinhardt, *Phys.Rev.Lett.* **85**, 4438 (2000); K. Maeda, astro-ph/0012313; V. Sahni, astro-ph/0202076; D. Huterer, astro-ph/0202256.