Experimentally obtaining the Likeness of Two unknown qubits on an NMR Quantum Information Processor

XUE Fei,^{1,*} DU JiangFeng,¹ ZHOU XianYi,¹ HAN RongDian,¹ and WU JiHui²

¹Department of Modern Physics, University of Science and Technology of China, Hefei, People's Republic of China, 230027

²Laboratory of Structure Biology, University of Science and Technology of China, Hefei, People's Republic of China, 230027

(Dated: January 7, 2019)

Recently quantum states discrimination has been frequently studied. In this paper we study them from the other way round, the likeness of two quantum states. The fidelity is used to describe the likeness of two quantum states. Then we presented a scheme to obtain the fidelity of two unknown qubits directly from the integral area of the spectra of the assistant qubit(spin) on an NMR Quantum Information Processor. Finally we demonstrated the scheme on a three-qubit quantum information processor. The experimental data are consistent with the theoretical expectation with an average error of 0.05, which confirms the scheme.

PACS numbers: 03.67.Lx, 82.56.-b

Quantum Information Processing(QIP) has been the subject of much recent interest, not only because it has great advantages in efficient algorithms and secure communications, but also because quantum information differs from classical information in several fundamental ways. One important difference is that qubits can hold superposition states while classical bits can only hold either 0 or 1 at the same time. So unlike two classical bits whose relationship is either the same or inverse, the relationship of two qubits is more complex. The complication comes from the superposition principle of quantum mechanics. Recently quantum state discrimination has been studied frequently. [1, 2, 3, 4, 5] In this paper we study quantum states from the other way round, i.e., considering the likeness of two quantum states. We focus on such questions that whether two unknown quantum states are the same or not, morever, the extent to which two states are alike, which is useful in quantum encryption and quantum states comparison.

Since Gershenfeld and Chuang realized quantum computation with NMR technique in 1997[6] a lot of jobs have been done with Nuclear Magnetic Resonance Quantum Information Processor(NMR QIP). NMR QIP has successfully demonstrated some efficient algorithms, such as Deutsch-Jozsa algorithm[7], searching algorithm[8, 9], Bernstein-Vazirani parity problem[10], and Shor's quantum factoring algorithm[11]; and some fundamental ideas in quantum information, such as Dense Coding[12], error correction[13], quantum games[14], creation of Greenberger Horne Zeilinger states[15] and approximate quantum cloning[16]. In this paper we will enlarge the list.

In the paper, first we give a brief review of the concept of the fidelity which originated from quantitative measures of the accuracy of transmission in communication theory, then we present a scheme to obtain the quantitative likeness(the fidelity) of two unknown qubits on an NMR QIP. The fidelity comes from the integral area of the spectra of the assistant qubit(spin), rather than from the tomography of the spin system. This makes it convenient to implement the scheme by NMR QIP. Finally we demonstrated the scheme on a three-qubit quantum information processor by obtaining the fidelity of two unknown qubits.

Before explaining the scheme, let us first give a review of the problem to be considered. The question is: given two unknown quantum states $|\psi_1\rangle$ and $|\psi_2\rangle$, what is the relationship of them or how similar they are. In order to quantitively describe the likeness some functions are needed.

Naturally we want that the function has the following characters:

a. $0 \leq F(\psi_1, \psi_2) \leq 1$ and $F(\psi_1, \psi_2) = 1$, if and only if $|\psi_1\rangle = |\psi_2\rangle$.

b. $F(\psi_1, \psi_2) = F(\psi_2, \psi_1).$

c. $F(\psi_1, \psi_2)$ is invariant under any unitary transformations on both states.

The fidelity appearing in the communication theory is a good candidate for it. The origin of the fidelity is a quantitative measure of the accuracy of transmission. It has desired properties and thus is a sensible choice as the quantitative measure of the likeness of two unknown quantum states. The fidelity of two quantum states is defined as [17]

$$F(\psi_1, \psi_2) = |\langle \psi_1 | \psi_2 \rangle|^2.$$
 (1)

Now we have the function of the quantitative measure of the likeness of two quantum states, then the question turns to be obtaining the fidelity $F(\psi_1, \psi_2)$.

In Quantum Fingerprinting[18], Harry Buhrman have presented a quantum circuit (Fig.1). It was used to test $|\psi_1\rangle = |\psi_2\rangle$ or $|\langle\psi_1|\psi_2\rangle| \leq \delta$ in that paper. We found that this quantum circuit can be further used to obtain fidelity $F(\psi_1, \psi_2)$ of the two unknown quantum states.

$$(H \otimes I)(G_{Fredkin})(H \otimes I)|0\rangle|\psi_1\rangle|\psi_2\rangle \tag{2}$$

where H is the Hadamard gate, which maps $|b\rangle \rightarrow \frac{1}{\sqrt{2}}(|0\rangle + (-1)^{b}|1\rangle)$. $G_{Fredkin}$ is the Fredkin gate which

^{*}Electronic address: Feixue@mail.ustc.edu.cn



FIG. 1: Quantum circuit to obtain the likeness of two unknown quantum states $|\psi_1\rangle$ and $|\psi_2\rangle$.

is also called contolled-swap gate, the first qubit is the controlling qubit, the controlled qubits can be generalized to quantum states. Tracing through the execution of the circuit, the final state is

$$\frac{1}{2}|0\rangle(|\psi_1\rangle|\psi_2\rangle+|\psi_2\rangle|\psi_1\rangle)+\frac{1}{2}|1\rangle(|\psi_1\rangle|\psi_2\rangle-|\psi_2\rangle|\psi_1\rangle) \quad (3)$$

After tracing out the other two qubits, the density matrix of the first qubit of this final state is reduced to the form,

$$\begin{pmatrix} \frac{1}{2}(1+|\langle\psi_1|\psi_2\rangle|^2) & 0\\ 0 & \frac{1}{2}(1-|\langle\psi_1|\psi_2\rangle|^2) \end{pmatrix}$$
(4)

When this state is projected to $|0\rangle$ and $|1\rangle$, the difference of the two probability is $|\langle \psi_1 | \psi_2 \rangle|^2$, which is just the fidelity $F(\psi_1, \psi_2)$. Then the question is to obtain the difference of two outcome probability.

The outcome probabilities can not be obtained from few experiments for a general quantum computer. But for NMR QIP, which is a Bulk Quantum computer[6], the situation is very different. In NMR QIP pseudo-pure states which are deviation density matrices of the spin system were proposed as the initial states, in stead of genuine pure states[6, 25]. The system of the spins in NMR may be convenient described by the product operator notation[19]. The state of the first qubit can sure be expressed in the notation of the deviation density matrix and the notation of product operator,

$$\rho_{1} = \begin{pmatrix} \rho_{00} & \rho_{01} \\ \rho_{10} & \rho_{11} \end{pmatrix}$$

$$= (\rho_{00} - \rho_{11})I_{z}^{1} + (\rho_{01} + \rho_{10})I_{x}^{1}$$

$$+ (\rho_{01} - \rho_{10})iI_{y}^{1} + \frac{1}{2}(\rho_{00} + \rho_{11})I$$
(5)
(6)

where $I_a^i = \frac{1}{2}\sigma_a^i$, a = x, y, z, and I is the identity matrix. So if the coefficient of I_z^1 could be observed, then the difference of the two probabilities—the fidelity is obtained. Of course it can be done by constructing the density matrix of the spin system. However the fidelity can be gotten in a more simple method. The fidelity—the coefficient of I_z^1 can be gotten by the following operations(Fig.2 part 3). First apply a gradient pulse to remove the non-diagonal part of the density matrix if there are any, then get rid of the part that correspond to the spins beside the first spin, finally apply a 90_y pulse on the first qubit. After performing these operations the state of the spin system becomes

$$(\rho_{00} - \rho_{11})I_x^1 + \frac{1}{2}(\rho_{00} + \rho_{11})I.$$
 (7)



FIG. 2: The pulse sequence for the network of obtaining the likeness of two qubits and the reading pulse sequence. The symbols on the indicate the phases in which the pulses are applied. All experiments were done on Bruker Avance DMX400 spectrometer at temperature 300K.

The identity matrix in NMR is not observable, so the signal from the first spin, the integral area of peaks, now corresponds to $\rho_{00} - \rho_{11}$, which is proportional to $|\langle \psi_1 | \psi_2 \rangle|^2$ -the fidelity $F(\psi_1, \psi_2)$.

On a three-qubit alanine(in D_2O) NMR QIP [22], we demonstrated the scheme. Spin- C_1 serves as qubit-1(the assistant spin), Spin- C_2 serves as qubit-2, which holds $|\psi_1\rangle$, and Spin- C_3 serves as qubit-3, which holds $|\psi_2\rangle$. We let $|\psi_1\rangle = \cos\frac{\theta_1}{2}|0\rangle - e^{i\varphi_1}\sin\frac{\theta_1}{2}|1\rangle$ and $|\psi_2\rangle =$ $\cos\frac{\theta_2}{2}|0\rangle - e^{i\varphi_2}\sin\frac{\theta_2}{2}|1\rangle$. Specifically we studied two situations systemically: $|\psi_2\rangle = |0\rangle$, changing $|\psi_1\rangle$ and $|\psi_1\rangle = |0\rangle$, changing $|\psi_2\rangle$. In each situation we have collected 20 experimental data, where θ_i varies from 0 to 180 degree, with a step of 45 degree, and for each θ_i , φ_i varies as 0, 90, 180 and 270 degree.

The pseudo-pure state is implemented by temporal averaging of three separate experiments, see Ref.[20] for the detail. The pulse sequence of the scheme is shown in Fig.2. It is applied after the preparation pulses for the pseudo-pure state. It includes three parts: 1. Prepare $|\psi_1\rangle = \cos \frac{\theta_1}{2}|0\rangle - e^{i\varphi_1} \sin \frac{\theta_1}{2}|1\rangle$ or $|\psi_2\rangle = \cos \frac{\theta_2}{2}|0\rangle - e^{i\varphi_2} \sin \frac{\theta_2}{2}|1\rangle$, it is implemented by selective pulses in certain phase on Spin- C_2 or Spin- C_3 . 2. The sequence for network in Fig.1. The two Hadamard gates are implemented by $R_y^1(90)$ and $R_{-y}^1(90)$, the Fredkin gate is implemented with three transition pulses TP1,TP2,TP3, see Ref.[21, 22] for the detail. 3. Reading sequence. Processing of tracing out Spin- C_2 and Spin- C_3 is done by integrating the entire multiplet of the Spin- C_1 . The errors in the experiment are estimated by $Err = |F_{exp} - F_{theory}|$.

The experimental data each of which is the sum of three experimental data, after the normalization, are plotted in Fig.3. Though the biggest error in the experimental data reaches about 0.11. The experimental data are consistent with the theoretical expectation with a average error of 0.05. It is sufficient to demonstrate the scheme. We have also tried the experiments that both $|\psi_1\rangle$ and $|\psi_2\rangle$ were changing. It is hard to make sure that both φ_1 and φ_2 are what we specify, because that the spins are precessing in different frequency, and the temporal averaging technique is also increase the difficulty. If these are taken into considering and be emen-



FIG. 3: Experiment results. Arrows are normalized experimental data, directions of them indicate φ_i . (a) corresponds $|\psi_2\rangle = |0\rangle$; (b) corresponds $|\psi_1\rangle = |0\rangle$. The theoretical fidelity $F_{theory}(\psi_1, \psi_2) = \cos^2(\frac{\theta_i}{2})$.

dated the scheme can still be confirmed. There are two

- L. S. Phillips, S. M. Barnett and D. T. Pegg 1998 Phys. Rev. A 58 3259.
- [2] Zhang Shengyu, Feng Yuan, Sun Xiaoming, et al. 2001 Phys. Rev. A 64 062103.
- [3] S. M. Barnett 2001 Phys. Rev. A 64 030303.
- [4] A. Chefles 2001 Phys. Rev. A 64 062305.
- [5] J. Walgate, A. J. Short, L. Hardy, et al. 2000 Phys. Rev. Lett. 85 4972.
- [6] N. Gershenfeld and I. Chuang, 1997 Science 275 350.
- [7] J. Kim, Jae-Seung Lee, and S. Lee, 2000 Phys. Rev. A 62 022312.
- [8] I. L. Chuang, N. Gershenfeld, and M. Kubinec 1998 Phys. Rev. Lett. 80 3408.
- [9] Li Xiao, G. L. Long, Hai-Yang Yan, et al. 2002 The Journal of Chemical Physics, 117, 3310.
- [10] Du Jiangfeng, Shi Mingjun, Zhou Xianyi, *et al.* 2001 Phys. Rev. A **64** 042306.
- [11] L. M. K. Vandersypen, M. Steffen, G. Breyta, et al. 2001 Nature 414 863.
- [12] Fang Ximing, Zhu Xiwen, Feng Mang, et al. 2000 Phys. Rev. A 61 022307.
- [13] E. Knill, R. Lafflamme, R. Martinez, *et al.* arXiv:quant-ph/0101034.
- [14] Du Jiangfeng, Li Hui, Xu Xiaodong, et al. 2002 Phys.

3

main sources of the errors. One is the imperfections of the pulses in the experiments, which bring about some errors in the implementation of the circuit and can be reduced by refining the pulse sequence. The other is the effect of the relaxation times. The length of the pulse sequences reaches 0.3s, while T_2 of the alanine is about 0.893s in the experiment. So the effect of the relaxation is not neglectable and will influence the final spectra. Morever, different terms of the spin system, such as I_z and I_x , have different relaxation behaviors and different relaxation speeds, hence these errors are harder to be reduced.

To summarize, we introduced the fidelity as a quantitative measure of the likeness of two unknown quantum states. Then we presented a scheme to obtain the fidelity on an NMR Quantum Information Processor, showed that the fidelity can be set proportional to the intensity of the signal from the assistant qubit(spin). Finally we experimentally demonstrated the scheme on a threequbit NMR QIP which is implemented by the solution of alanine. The network(Fig.1) with slight difference was studied in Ekert's work[23, 24]. In a sense our work in this paper gives an experimentally demonstration of some ideas in the paper[23]. Besides on NMR QIP, any quantum computer based on bulk spins can use the scheme to obtain the fidelity of two unknown qubits.

This work was supported by the National Nature Science Foundation of China (Grants No. 10075041, No. 10075044 and No. 10104014), and the National Fundamental Research Program(Grant No. 2001CB309300).

Rev. Lett. 88 137902.

- [15] R. Laflamme, E. Knill, W. H. Zurex, et al. 1998 Phil. Trans. R. Soc. Lond. A 356 1941.
- [16] H. K. Cummins, C. Jones, A. Furze, et al. 2002 Phys. Rev. Lett. 88 187901.
- [17] R. Jozsa 1994 J. Modern Optics 54 2315.
- [18] H. Buhrman, R. Cleve, J. Watrous, et al. 2001 Phys. Rev. Lett. 87 167902.
- [19] O. W. Sørensen, G W. Eich, M. H. Levitt, et al. 1983 Prog. Nucl. Magn. Reson. Spectrosc 16 163.
- [20] Xue Fei, Du JiangFeng, Fan YangMei, et al. 2002 Acta Physica Sinica 51 763.
- [21] Du JiangFeng, Shi MingJun, Wu JiHui, et al. 2001 Phys. Rev. A Vol. 63 042302.
- [22] Xue Fei, Shi MingJun, Du JiangFeng, et al. 2002 Chin. Phys. Lett. 19 1048.
- [23] A. K. Ekert, C. M. Alves, D. K. L. Oi, *et al.* 2002 Phys. Rev. Lett. 88 217901.
- [24] P. Horodecki and A. Ekert, 2002 Phys. Rev. Lett. 89, 127902.
- [25] E. Knill, I. Chuang, and R. Laflamme, 1998 Phys. Rev. A 57, 3348.