Bell's Theorem Without Inequalities for two Maximally Entangled Particles

W. LiMing* and Z. L. Tang
Dept. of Physics, South-China Normal University, Guangzhou 510631, P. R. China

C. Liao

School for Information and Optoelectronic Science and Engineering, South-China Normal University, Guangzhou 510631, P. R. China (Dated: October 28, 2018)

A proof of Bell's theorem without inequalities for two maximally entangled particles is proposed using the technique of quantum teleportation. It follows Hardy's arguments for a non-maximally entangled state with the help of two auxiliary particles without correlation. The present proof can be tested by measurements with 100% probability.

PACS numbers: 03.65.Ud,03.67.-a, 03.65.Ta

Bell's theorem claims that quantum mechanics cannot be reproduced with the hidden variable local model[1]. Recently, Bell's theorem appeared in forms without inequalities [2, 3, 4, 5], exhibiting greater contradiction between the local model and quantum mechanics. Greenberger et al (GHZ) proposed a proof for three entangled particles, thus three observers are needed[2]. Hardy proposed a proof for two non-maximally entan-Cabello proposed a GHZ-like and gled particles[3]. more delicate proof for two pairs of maximally entangled particles [4, 5]. It seems that non-locality exists only in a maximally entangled state of three or more particles or in a non-maximally entangled state of two particles. The paradox of Einstein, Podolsky, and Rosen (EPR), however, argues the non-locality of two maximally entangled particles[6]. The question is whether it is possible to demonstrate Bell's theorem without using inequalities by two maximally entangled particles.

In this paper I propose a proof of Bell's theorem without using inequalities for two maximally entangled particles, using the technique of quantum teleportation. It follows Hardy's arguments for the contradiction between the hidden variable local model and the non-locality of quantum mechanics.

Consider two spin-1/2 particles, 1 and 2, in a maximally entangled state, i.e., a spin singlet:

$$|\Psi\rangle_{12} = \frac{1}{\sqrt{2}}(|+\rangle_1|-\rangle_2 - |-\rangle_1|+\rangle_2) \tag{1}$$

where "+" and "-" denote, respectively, spin up and down. These two particles are transmitted in opposite directions to two observers, Alice and Bob, with a space-like separation. Both Alice and Bob have prepared an auxiliary particle, A and B, respectively, with the following spin states

$$|A\rangle = |+\rangle$$
 , $|B\rangle = (|+\rangle + |-\rangle)/\sqrt{2}$, (2)

which are the eigenstates of σ_z and σ_x , respectively. These two auxiliary particles are not correlating with each other and never transmitted between the two observers, thus are considered as part of the apparatus. The total spin state of this four-particle system is given by

$$|\Psi\rangle = |A\rangle|\Psi\rangle_{12}|B\rangle \tag{3}$$

In order to use the technique of quantum teleportation this state is expanded in the Bell basis of Alice's particles A and 1 as follows:

$$|\Psi\rangle = -\frac{1}{2} \left[|\Psi^{-}\rangle_{A1}| + \rangle_{2} + |\Psi^{+}\rangle_{A1}| + \rangle_{2} - |\Phi^{-}\rangle_{A1}| - \rangle_{2} - |\Phi^{+}\rangle_{A1})| - \rangle_{2} \right]|B\rangle$$
(4)

where the Bell states are given by

$$|\Psi^{\pm}\rangle = \frac{1}{\sqrt{2}}(|+\rangle|-\rangle \pm |-\rangle|+\rangle) \tag{5}$$

$$|\Phi^{\pm}\rangle = \frac{1}{\sqrt{2}}(|+\rangle|+\rangle \pm |-\rangle|-\rangle)$$
 (6)

 $|\Psi\rangle$ can also be expanded in the Bell basis of Bob's particles 2 and B as follows:

$$|\Psi\rangle = -\frac{1}{2\sqrt{2}}|A\rangle [(|+\rangle_1 + |-\rangle_1)|\Psi^-\rangle_{2B} + (|+\rangle_1 - |-\rangle_1)|\Psi^+\rangle_{2B} - (|+\rangle_1 + |-\rangle_1)|\Phi^-\rangle_{2B} - (-|+\rangle_1 + |-\rangle_1)|\Phi^+\rangle_{2B}]$$
(7)

According to the principle of quantum teleportation, if Alice measures one of the Bell states of her two particles, A and 1, particle 2 will collapse to the corresponding quantum state, see (4). For example, if Alice measures the Bell state, $|\Psi^-\rangle_{A1}$, particle 2 collapses to state $|+\rangle$. In the same way, if Bob measures the state, $|\Psi^-\rangle_{2B}$, particle 1 collapses to state $(|+\rangle + |-\rangle)/\sqrt{2}$, see (7). It is seen that the state of particle A of Alice is copied into

^{*}Electronic address: wliming@scnu.edu.cn

the state of particle 2 of Bob without transmitting any physical information between Alice and Bob[7, 8]. The above two measurements correspond to the following two projecting operators:

$$\hat{D}_1 = |\Psi^-\rangle_{A1}\langle\Psi^-|
\hat{D}_2 = |\Psi^-\rangle_{2B}\langle\Psi^-|$$
(8)

Including another two operators \hat{U}_1 and \hat{U}_2 ,

$$\hat{U}_1 = |+\rangle_1 \langle +| \quad , \quad \hat{U}_2 = |+\rangle_2 \langle +| \tag{9}$$

one has four physical observable quantities, $\hat{D}_1, \hat{D}_2, \hat{U}_1, \hat{U}_2$. They take values 0 or 1 corresponding to their eigenvalues, denoted as D_1, D_2, U_1, U_2 .

Now if Alice measures \hat{U}_1 on particle 1, and Bob measures \hat{U}_2 on particle 2, they have

$$U_1 U_2 = 0. (10)$$

This is because, since particles 1 and 2 are in a maximally entangled state, (1), their spins are always opposite to each other.

From (4), if Alice measures \hat{D}_1 on her particles A and 1, and Bob measures \hat{U}_2 on his particle 2, they have

if
$$D_1 = 1$$
 then $U_2 = 1$ (11)

From (7), if Bob measures \hat{D}_2 on his particles 2 and B, and Alice measures \hat{U}_1 on his particle 1, they have

if
$$D_2 = 1$$
 then $U_1 = 1$ (12)

The most important fact here is $[\hat{D}_1, \hat{D}_2] = 0$, indicating that \hat{D}_1 and \hat{D}_2 can be measured simultaneously. This allows Alice and Bob to make a joint measurement for these two observable quantities on the system. The probability of the system with $D_1 = D_2 = 1$ is given by

$$P = \langle \Psi | \hat{D}_1 \hat{D}_2 | \Psi \rangle = \frac{1}{16}$$
 for $D_1 = D_2 = 1$ (13)

Now Hardy's arguments can be applied to deduce a contradiction between the non-locality of the entangled state, $|\Psi\rangle$, and the hidden variable local model[3]. The latter claims that Alice's choice of measurement cannot influence the outcome of any measurement of Bob, since there is a space-like space-time separation between Alice and Bob. For example, for a run that Alice and Bob obtain $D_1 = U_2 = 1$, if Alice had measured something else, say \hat{U}_1 for particle 1, instead of \hat{D}_1 , he would not affect the outcome of Bob's measurement $U_2 = 1$, vice versa. In another word, due to (11), if Alice obtained $D_1 = 1$, she can predict without any uncertainty the outcome of Bob's measurement, $U_2 = 1$. According to EPR's argument, $U_2 = 1$ is a physical reality element of particle 2.

Consider a run of a joint measurement that Alice and Bob found $D_1 = D_2 = 1$. This run does exist since the

probability exists, see (13). In the hidden variable local model, we have the following deductions:

Deduction 1: From the fact that we have $D_1 = 1$ it follows from (11) that if \hat{U}_2 had been measured we would obtain $U_2 = 1$. In another word, $U_2 = 1$ is a physical reality element of particle 2.

Deduction 2: Since Alice's choice of measurement does not affect the outcome of Bob's measurement, even if \hat{U}_1 had been measured on particle 1 instead of \hat{D}_1 we would still have $U_2 = 1$.

Deduction 3: By a similar argument we can deduce from the fact $D_2 = 1$ and (12) that $U_1 = 1$. In another word, $U_1 = 1$ is a physical reality element of particle 1.

Deduction 4: Thus, for this run, we have $U_1U_2 = 1$. Hence, if we had measured \hat{U}_1 and \hat{U}_2 instead of \hat{D}_1 and \hat{D}_2 , we would have obtained $U_1U_2 = 1$, which, however, contradicts (10).

It is seen that from the hidden variable local model we arrive at a contradiction and therefore, an entangled state of quantum mechanics must be nonlocal. Physical reality element does not exist.

To demonstrate this proof experimentally, one needs first to verify (10, 11,12), and then measure \hat{D}_1 and \hat{D}_2 . If $D_1 = D_2 = 1$ is observed then non-locality of quantum mechanics is verified.

In the above formalism, only one pair of Bell states, i.e., $|\Psi^{-}\rangle_{A1}$ and $|\Psi^{-}\rangle_{2B}$, are considered. In fact, operators \hat{D}_{1} and \hat{D}_{2} can be made up of each pair of Bell states, such as

$$\begin{array}{l} \hat{D}_1 = |\Psi^-\rangle_{A1}\langle\Psi^-| \text{ and } \hat{D}_2 = |\Psi^+\rangle_{2B}\langle\Psi^+|,.....\\ \hat{D}_1 = |\Psi^+\rangle_{A1}\langle\Psi^+| \text{ and } \hat{D}_2 = |\Phi^-\rangle_{2B}\langle\Phi^-|,...... \end{array}$$

Totally, one has $4 \times 4 = 16$ different pairs of \hat{D}_1 and \hat{D}_2 . This is obviously true because the four Bell states compose a complete basis. For each pair of operators \hat{D}_1 and \hat{D}_2 one defines corresponding operators \hat{U}_1 and \hat{U}_2 . Using each group of operators \hat{D}_1 , \hat{D}_2 , \hat{U}_1 , \hat{U}_2 one can deduce the above nonlocal property of entangled states. Therefore, from (13), the total probability to test the Bell theorem is $16 \times 1/16 = 100\%$.

In summary, Bell's theorem without inequalities is proved for a maximally entangled state based on the technique of quantum teleportation. Auxiliary particles A and B have not been transmitted between Alice and Bob and are not correlating to each other, thus are taken as part of the apparatus. The non-locality proved here belongs to the maximally entangled state of particles 1 and 2. This proof can be tested by measurements with 100% probability.

Acknowledgments

This work was supported by the National Fundamental Research Program under Grant No 2001CD309300.

- [1] J. S. Bell, Physics (Long Island City, N.Y.) 1, 195(1964).
- [2] D. M. Greenberger, M. A. Horne, A. shimony, A. Zeilinger, Am. J. Phys. 58, 1131(1990).
- [3] L. Hardy, Phys. Rev. Lett. 71, 1665(1993).
- [4] A. Cabello, Phys. Rev. Lett. 86, 1911(2001).
- [5] A. Cabello, Phys. Rev. Lett. 87, 10403(2001).
- [6] A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. 47,
- 777(1935); D. Bohm, $\it Quantum\ Theory\ (Prentice\ Hall,\ Englewood\ Cliffs,\ NJ,\ 1951).$
- [7] D. Bouwnmeester, J. W. Pan, K. Mattle, M. Eibl, H. Weinfurter and A. Zeilinger, Nature 390, 575(1997).
- [8] C. H. Bennet, et al., Phys. Rev. Lett. 70, 1895(1993).