

# On the communication cost of entanglement transformations

Patrick Hayden\*

*Institute for Quantum Information, Caltech 107-81, Pasadena, CA 91125, USA*

Andreas Winter†

*Department of Computer Science, University of Bristol,  
Merchant Venturers Building, Woodland Road, Bristol BS8 1UB, United Kingdom*

(Dated: 31<sup>st</sup> May 2002)

We study the amount of communication needed for two parties to transform some given joint pure state into another one, either exactly or with some fidelity. Specifically, we present a method to lower bound this communication cost even when the amount of entanglement does not increase. Moreover, the bound applies even if the initial state is supplemented with unlimited entanglement in the form of EPR (Einstein–Podolsky–Rosen) pairs and the communication is allowed to be quantum mechanical.

We then apply the method to the determination of the communication cost of asymptotic entanglement concentration and dilution. While concentration is known to require no communication whatsoever, the best known protocol for dilution, discovered by Lo and Popescu [Phys. Rev. Lett. 83(7):1459–1462, 1999], requires a number of bits to be exchanged which is of the order of the square root of the number of EPR pairs. Here we prove a matching lower bound of the same asymptotic order, demonstrating the optimality of the Lo–Popescu protocol up to a constant factor and establishing the existence of a fundamental asymmetry between the concentration and dilution tasks.

We also discuss states for which the minimal communication cost is proportional to their entanglement, such as the states recently introduced in the context of “embezzling entanglement” [W. van Dam and P. Hayden, quant-ph/0201041].

PACS numbers: 03.65.Ta, 03.67.Hk

Keywords: Entanglement transformations, entanglement dilution, communication, Rényi entropy

## I. PURE STATE ENTANGLEMENT TRANSFORMATIONS

The quantification of entanglement began with the study of the following question: assume that two parties, generically referred to as *Alice* and *Bob*, share  $n$  copies of a bipartite pure state  $|\phi_{AB}\rangle$  which by local operations and classical communication (LOCC), they would like to convert into a state that has high fidelity to  $k$  copies of the target state  $|\psi_{AB}\rangle$ , with  $k$  as large as possible. The basic question is then, what is  $\lim k/n$  as  $n \rightarrow \infty$  and the fidelity goes to one?

It turns out [5] that this optimal asymptotic ratio is equal to  $E(\phi)/E(\psi)$ , where

$$E(\phi) = S(\text{Tr}_B |\phi\rangle\langle\phi|) = -\text{Tr} (\text{Tr}_B |\phi\rangle\langle\phi| \log \text{Tr}_B |\phi\rangle\langle\phi|)$$

is the von Neumann entropy of Alice’s reduced state. For this reason,  $E$  is often called the *entropy of entanglement*. One consequence of this result is that pure state entanglement can be interconverted *asymptotically losslessly* between its different forms, justifying the introduction of the *ebit* as a resource quantity, with its ubiquitous “incarnation”, the two-qubit EPR pair state, which, up to

a local change of basis, can be written as

$$|\phi_2^+\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle).$$

For the sake of quantifying entanglement, however, not only local actions by Alice and Bob but *classical communication* was considered unlimited. It is precisely these communication requirements that we study in the present paper, in which we follow Lo’s [16] suggestion to study the communication complexity of distributed quantum information processing.

This goal notwithstanding, our point of departure will not be the theory of asymptotically faithful transformations but, rather, its finite (and more refined) variant of transformations from  $|\phi_{AB}\rangle$  to  $|\psi_{AB}\rangle$  up to fidelity  $1 - \epsilon$ , as laid out in [23], building on previous work [19], [12], [18] for the zero-error case.

Up to local unitaries, pure entangled states are uniquely defined by the spectrum of their reduced states (either at Alice’s or Bob’s side), the eigenvalues known as the *Schmidt coefficients*  $\lambda_j$ . Indeed, it is possible to choose bases in the entangled system such that

$$|\phi_{AB}\rangle = \sum_j \sqrt{\lambda_j} |i\rangle_A \otimes |i\rangle_B.$$

The theory relates the feasibility of such an LOCC transformation to the *majorisation order* of the Schmidt coefficients ( $\lambda$ ) of  $|\phi\rangle$  and ( $\mu$ ) of  $|\psi\rangle$ , both vectors arranged

---

\*Electronic address: patrick@cs.caltech.edu

†Electronic address: winter@cs.bris.ac.uk

in nonincreasing order:

$$|\phi\rangle \xrightarrow{\text{LOCC}} |\psi\rangle \quad \text{iff} \quad (\lambda) \prec (\mu),$$

where  $(\lambda) \prec (\mu)$  is defined to mean

$$\forall k \quad \sum_{j=1}^k \lambda_j \leq \sum_{j=1}^k \mu_j,$$

which can be shown to be equivalent to the existence of a doubly stochastic matrix  $M$  such that  $(\lambda) = M \cdot (\mu)$ . By the results of [12] and [15], any such allowed transformation can always be achieved by one-way communication, say from Alice to Bob, of  $2 \log \text{rank} \text{Tr}_B |\phi\rangle\langle\phi|$  classical bits.

The organisation of the paper is as follows. In section II we will explain the mathematical model of approximate pure state transformations and derive our main result, a lower bound on the communication cost of state transformations which holds even if the initial state is supplemented by an unlimited number of EPR pairs, and even if the communication is quantum mechanical. To our knowledge this is the first *quantitative* statement of its kind. (The need for some nonzero amount of communication in certain transformations was pointed out in [18].) We then apply the result in section III to the asymptotic transformations mentioned in the introduction, proving a lower bound of  $\Omega(\sqrt{n})$  on the communication necessary for entanglement dilution, which, up to a constant factor, matches the  $O(\sqrt{n})$  construction of Lo and Popescu [17] for this task. In section IV we analyse a class of states that require for their creation from EPR pairs communication of the same order as their entanglement, before ending with a discussion of some open problems.

## II. A LOWER BOUND ON THE COMMUNICATION COST

Assume that initially Alice and Bob share the state  $|\phi\rangle$ , then execute several rounds of local actions and classical communication, and finally end up with some joint state  $\tilde{\rho}$  that has high fidelity to  $|\psi\rangle$ . Allowing the use of *quantum bits* to communicate, we give Alice and Bob even more power, thereby potentially reducing the communication cost, while at the same time simplifying the appearance of the protocol: because each of the local actions can be implemented using ancillae and unitary transformations, the whole process can be reduced to a series of exchanges of quantum systems of certain dimensions  $d_i$  between Alice and Bob, with a final tracing out (discarding) of part of Alice's and part of Bob's system. The total communication cost of such a procedure is just  $C = \sum_{i=1}^N \log d_i$  qubits.

The idea of the lower bound is very simple, and is explained most straightforwardly for *exact* state transformations, when  $\tilde{\rho} = |\psi\rangle\langle\psi|$ . During the process of transformation we monitor a certain quantity  $\Delta$  associated to




FIG. 1: In round  $i$  Alice (Bob) performs some unitary  $U_i$  on her (his) system, which separates into a residual system and a  $d_i$ -dimensional system that is sent to Bob (Alice). In the last,  $N^{\text{th}}$ , round, the receiver of the message may perform a unitary on his/her system, and then Alice and Bob trace out subsystems  $A'$  and  $B'$ .

Alice's reduced density operator, showing that, for each qubit communicated, it can only increase by a constant, and then observe that the final partial trace never increases  $\Delta$  at all. The difference between the initial and the final  $\Delta$  then provides a lower bound on the communication.

Specifically we shall consider, for  $\rho = \text{Tr}_B |\phi\rangle\langle\phi|$ ,

$$\Delta(\rho) := S_0(\rho) - S_\infty(\rho), \quad (1)$$

where  $S_\alpha$  are the *Rényi entropies* [21] of order  $\alpha$ :

$$S_\alpha(\rho) := \frac{1}{1-\alpha} \log \text{Tr}(\rho^\alpha).$$

For  $\alpha = 0, 1, \infty$  the Rényi entropies are defined by continuous extension, with resulting formulas

$$\begin{aligned} S_0(\rho) &= \log \text{rank } \rho, \\ S_1(\rho) &= -\text{Tr}(\rho \log \rho), \\ S_\infty(\rho) &= -\log \|\rho\|_\infty, \end{aligned}$$

where  $\|\cdot\|_\infty$  is the supremum norm: for selfadjoint operators it is the largest absolute value of an eigenvalue. (Throughout the paper,  $\log$  and  $\exp$  are understood to be base 2.) Note that  $\Delta(\rho) \geq 0$  since  $S_\alpha(\rho)$  is non-increasing in  $\alpha$  [21], or by inspection of the definition. Furthermore, if all the non-zero eigenvalues of  $\rho$  are the same then  $S_0(\rho) = S_\infty(\rho)$  so that  $\Delta(\rho) = 0$ . Otherwise,  $\Delta(\rho)$  will be strictly greater than zero. Therefore,  $\Delta(\rho)$  can be interpreted as a measure of the variation in the eigenvalues of  $\rho$ .

The key observation is that, in communication round  $i$ , the Rényi entropy of Alice's reduced state, whose spectrum characterises the entanglement, cannot change too much. To see this, we assume without loss of generality that it is Alice's turn to perform a unitary, rotating her reduced state to  $\rho_{AA'}$ . This step, obviously, does not change the Rényi entropy at all. Next, she gives Bob the  $d_i$ -dimensional system  $A'$ , leaving her with the new reduced state  $\rho_A$ , for which we have the relation (see [11])

$$S_\alpha(\rho_{AA'}) - \log d_i \leq S_\alpha(\rho_A) \leq S_\alpha(\rho_{AA'}) + \log d_i, \quad (2)$$

which implies (inserting  $\alpha = 0, \infty$ )

$$\Delta(\rho_A) \leq \Delta(\rho_{AA'}) + 2 \log d_i. \quad (3)$$

Thus, the quantity  $\Delta$  can increase (or decrease) by at most  $2 \log d_i$  in step  $i$ . After the last round of communication has taken place, the joint state is  $|\tilde{\psi}_{ABA'B'}\rangle = |\psi_{AB}\rangle \otimes |\iota_{A'B'}\rangle$ . (Note that if this were not a product state,  $\tilde{\rho}$  would necessarily not be pure.) Hence, by induction over the number of rounds, summing over the the eqs. (3) yields

$$\Delta(\text{Tr}_{BB'}|\tilde{\psi}_{ABA'B'}\rangle\langle\tilde{\psi}_{ABA'B'}|) \leq \Delta(\text{Tr}_B|\phi\rangle\langle\phi|) + 2C. \quad (4)$$

The effect on  $\Delta$  of the final partial trace over the primed system is easy to understand: because the Rényi entropies are additive under tensor products, i.e.

$$S_\alpha(\rho \otimes \sigma) = S_\alpha(\rho) + S_\alpha(\sigma),$$

we obtain

$$\begin{aligned} \Delta(\text{Tr}_B|\psi\rangle\langle\psi| \otimes \text{Tr}_{B'}|\iota\rangle\langle\iota|) \\ = \Delta(\text{Tr}_B|\psi\rangle\langle\psi|) + \Delta(\text{Tr}_{B'}|\iota\rangle\langle\iota|), \end{aligned} \quad (5)$$

and the rightmost term is nonnegative. This proves

**Theorem 1** *A (deterministic) pure state transformation of  $|\phi_{AB}\rangle$  into  $|\psi_{AB}\rangle$  requires at least*

$$C \geq \frac{1}{2} \left( \Delta(\text{Tr}_B|\psi\rangle\langle\psi|) - \Delta(\text{Tr}_B|\phi\rangle\langle\phi|) \right)$$

*bits of communication, even if quantum communication is allowed.*  $\square$

We note that in [11] the analogous theorem for the bare Rényi entropies  $S_\alpha$  was used to prove bounds on the communication required to perform entanglement transformations in an approximate setting. There, changes in  $S_\alpha$  reflected changes in the amount of entanglement present in the system. The advantage of using  $\Delta$  is precisely that it does not measure entanglement but, rather, variation in the Schmidt coefficients.

**Remark 2** *Obviously, a similar result holds for*

$$\Delta^{\alpha\beta}(\rho) := S_\alpha(\rho) - S_\beta(\rho),$$

*with arbitrary  $0 \leq \alpha < \beta \leq \infty$ . Even though  $\Delta^{\alpha\beta}(\rho) \leq \Delta(\rho)$ , for some  $\alpha$  and  $\beta$  the increase of the former quantity in an entanglement transformation may exceed the increase of the latter.*

**Remark 3** *As an example of a nontrivial consequence of theorem 1 we may observe that it puts severe restrictions on the entanglement transformations possible without any communication: none of the  $\Delta^{\alpha\beta}$  must increase.*

*For example, from a maximally entangled state only other maximally entangled states (with possibly smaller Schmidt rank) may be obtained. If the Schmidt rank of the target divides that of the initial state this is clearly possible, while inspection of eq. (5) shows that this is also necessary.*

For the case of high-fidelity transformations this approach turns out to be too simple: neither  $S_0$  nor  $S_\infty$  can be well controlled if we switch from a state to one close by. For example, for the dilution task, which consists of the creation of  $(\alpha|00\rangle + \beta|11\rangle)^{\otimes n}$  from EPR pairs, theorem 1 implies a lower bound of  $\Omega(n)$ , while we know from [17] that arbitrarily high fidelity can be achieved with  $O(\sqrt{n})$  bits of communication.

Instead, we invent robust versions of  $S_0$ ,  $S_\infty$  and  $\Delta$ : let the eigenvalues of  $\rho$  be denoted  $r_j$  and then define, for  $0 \leq \epsilon < 1$ ,

$$S_{0,\epsilon}(\rho) := \log \min \left\{ |J| : \sum_{j \in J} r_j \geq 1 - \epsilon \right\}, \quad (6)$$

$$S_{\infty,\epsilon}(\rho) := -\log \min \left\{ \max_{j \in J} r_j : \sum_{j \in J} r_j \geq 1 - \epsilon \right\}, \quad (7)$$

$$\Delta_\epsilon(\rho) := \log \min \left\{ |J| \left( \max_{j \in J} r_j \right) : \sum_{j \in J} r_j \geq 1 - \epsilon \right\}, \quad (8)$$

all the minimisations are understood to be over subsets  $J$  of the eigenvalue indices  $j$ . Note that

$$\Delta_\epsilon(\rho) \geq S_{0,\epsilon}(\rho) - S_{\infty,\epsilon}(\rho), \quad (9)$$

with equality generally only if  $\epsilon = 0$ , in which case these quantities reduce to the above  $S_0$ ,  $S_\infty$  and  $\Delta$ .

**Remark 4** *Note that  $\Delta_\epsilon$  has the following ‘‘high-fidelity’’ relation to  $\Delta_0$ :*

$$\Delta_\epsilon(\rho) = \min \{ \Delta_0(P\rho P) : \text{Tr}(\rho P) \geq 1 - \epsilon \},$$

*where the minimisation is over all projections  $P$  commuting with  $\rho$ , extending the definition of  $\Delta_0$  to sub-normalised density operators. The operators  $P\rho P$  can be interpreted as post-measurement states after the event ‘‘ $P$ ’’ has occurred, normalised to the event probability.*

*More generally, we could allow any  $0 \leq B \leq \mathbb{1}$  in the above minimisation, such that  $\text{Tr}(\rho B) \geq 1 - \epsilon$ , and substituting the post-measurement states  $\sqrt{B}\rho\sqrt{B}$ . (By a result in [24] this operator has high fidelity to the state  $\rho$ .) It is easy to see that the resulting quantity is within a distance of  $\log(1 - \epsilon)$  from  $\Delta_\epsilon$ .*

We now prove a few lemmas which will together comprise our method of estimating the communication cost, by providing the tools to estimate  $\Delta_\epsilon$  for the appropriate reduced states. We begin with the simple observation that for all states  $\rho$  and  $\epsilon' < \epsilon < 1$ :

$$\Delta_\epsilon(\rho) \geq \log(1 - \epsilon), \quad (10)$$

$$\Delta_\epsilon(\rho) \leq \Delta_{\epsilon'}(\rho). \quad (11)$$

**Lemma 5** *If for two states  $\rho$  and  $\sigma$ ,  $\|\rho - \sigma\|_1 \leq \epsilon$ , then*

$$\Delta_0(\rho) \geq \Delta_{\sqrt{\epsilon}}(\sigma) + \log(1 - \sqrt{\epsilon}).$$

(Where  $\|\cdot\|_1$  is the trace norm, for selfadjoint operators given by the sum of the absolute values of all eigenvalues, counting multiplicities.)

*Proof.* To begin, denote the eigenvalue lists of  $\rho$  and  $\sigma$  by  $(r)$  and  $(s)$ , respectively, in nonincreasing order. Then, because (see [20])

$$\|(r) - (s)\|_1 \leq \|\rho - \sigma\|_1 \leq \epsilon,$$

we may concentrate on the eigenvalues only. Define, for  $\delta = \sqrt{\epsilon}$ ,

$$J := \{j : (1 - \delta)s_j \leq r_j \leq (1 + \delta)s_j\}.$$

Then, for the complement  $J^c$  of  $J$ ,

$$\delta s(J^c) = \sum_{j \notin J} \delta s_j \leq \sum_{j \notin J} |r_j - s_j| \leq \epsilon,$$

implying

$$\sum_{j \in J} s_j \geq 1 - \sqrt{\epsilon}.$$

We may clearly assume that  $s$  is nonzero on  $J$ , otherwise shrinking  $J$  without affecting the last inequality.

Thus, by the definition of  $\Delta_{\sqrt{\epsilon}}$ ,

$$\log\left(|J| \max_{j \in J} s_j\right) \geq \Delta_{\sqrt{\epsilon}}(\sigma).$$

On the other hand, by the definition of  $J$ ,

$$j \in J \implies r_j \neq 0,$$

which implies that  $\text{rank } \rho \geq |J|$ . Similarly,

$$j \in J \implies r_j \geq (1 - \sqrt{\epsilon}) s_j,$$

implies  $\max_j r_j \geq (1 - \sqrt{\epsilon}) \max_{j \in J} s_j$ . Comparing the last two observations to the definition of  $\Delta_0(\rho)$  finishes the proof of the claim.  $\square$

**Lemma 6** *For any two states  $\tau$  and  $\omega$ , and  $\epsilon < 1$ ,*

$$\Delta_\epsilon(\tau \otimes \omega) \geq \Delta_{\sqrt{\epsilon}}(\tau) + \log(1 - \sqrt{\epsilon}).$$

*Proof.* Denote the eigenvalues of  $\tau$  and  $\omega$  by  $t_i$  and  $w_k$ , respectively. Let  $J$  be a set of indices  $i$  and  $k$  such that

$$\Delta_\epsilon(\tau \otimes \omega) = \log\left(|J| \max_{ik \in J} t_i w_k\right),$$

and

$$(t \otimes w)(J) = \sum_{ik \in J} t_i w_k \geq 1 - \epsilon. \quad (12)$$

We shall be interested, for certain  $k$ , in the sections

$$S_k := \{i : ik \in J\}$$

of  $J$  along  $k$ , in particular in the set

$$K := \left\{k : t(S_k) = \sum_{i \in S_k} t_i \geq 1 - \sqrt{\epsilon}\right\}.$$

It follows from the definition of  $K$  and the constraint of eq. (12) that

$$w(K) = \sum_{k \in K} w_k \geq 1 - \sqrt{\epsilon}. \quad (13)$$

The proof is a standard Markov inequality argument: observe that we can rewrite eq. (12) using the sections:

$$1 - \epsilon \leq (t \otimes w)(J) = \sum_k w_k t(S_k).$$

Now the right hand side is a probability average over the values  $t(S_k)$ , taken with probability  $w_k$ . We decompose the sum into two contributions which we estimate separately:

$$\begin{aligned} 1 - \epsilon &\leq \sum_{k \in K} w_k t(S_k) + \sum_{k \notin K} w_k t(S_k) \\ &\leq w(K) + (1 - w(K))(1 - \sqrt{\epsilon}). \end{aligned}$$

Hence  $(1 - w(K))\sqrt{\epsilon} \leq \epsilon$ , which is our claim.

Now define

$$J' := \bigcup_{k \in K} S_k \times \{k\},$$

and successively estimate

$$\begin{aligned} |J| \max_{ik \in J} (t_i w_k) &\geq |J'| \max_{ik \in J'} (t_i w_k) \\ &= \sum_{l \in K} |S_l| \max_{ik \in J'} (t_i w_k) \\ &\geq \sum_{k \in K} |S_k| \max_{i \in S_k} (t_i w_k) \\ &= \sum_{k \in K} w_k \left( |S_k| \max_{i \in S_k} t_i \right) \\ &\geq \sum_{k \in K} w_k \exp(\Delta_{\sqrt{\epsilon}}(\tau)) \\ &\geq (1 - \sqrt{\epsilon}) \exp(\Delta_{\sqrt{\epsilon}}(\tau)), \end{aligned}$$

the second last line because of  $t(S_k) \geq 1 - \sqrt{\epsilon}$ , the last line by eq. (13), which proves the lemma.  $\square$

**Remark 7** We do not know if a symmetric version of this lemma holds, with an additional term to the right analogous to the one for  $\tau$ :

$$\Delta_\epsilon(\tau \otimes \omega) \stackrel{?}{\geq} (1 - \epsilon')\Delta_{\epsilon'}(\tau) + (1 - \epsilon')\Delta_{\epsilon'}(\omega) + \epsilon'',$$

with  $\epsilon', \epsilon''$  functions of  $\epsilon$  which vanish for  $\epsilon \rightarrow 0$ .

This would constitute a form of “quasi-additivity” for  $\Delta$ , since the validity of the analogous reverse inequality

$$\Delta_{2\epsilon}(\tau \otimes \omega) \leq \Delta_\epsilon(\tau) + \Delta_\epsilon(\omega)$$

is quite easy to see. While it may not be useful to improve on our present results, confirmation of the “quasi-additivity” would be of conceptual interest.

We are now ready to state our central result, which applies whenever the output state has high Uhlmann fidelity  $F(\sigma, \omega) = \left(\text{Tr}\sqrt{\sigma^{1/2}\omega\sigma^{1/2}}\right)^2$  [14, 22] with the desired state, even if the output is mixed:

**Theorem 8** Consider a state transformation protocol that takes  $|\phi_{AB}\rangle$  to  $|\psi_{AB}\rangle$  with fidelity  $1 - \epsilon$ , exchanging a total of  $C$  qubits in the process. Then, with  $\delta = \sqrt[8]{4\epsilon}$ ,

$$2C \geq \Delta_\delta(\text{Tr}_B|\psi\rangle\langle\psi|) - \Delta_0(\text{Tr}_B|\phi\rangle\langle\phi|) + 2\log(1 - \delta).$$

*Proof.* Like in the zero-error case, we follow the increase of  $\Delta_0$  over the course of the protocol: after the last communication has taken place, the joint state is  $|\tilde{\psi}_{ABA'B'}\rangle$ , and we have (compare eq. (4))

$$2C \geq \Delta_0(|\tilde{\psi}_{AA'}\rangle) - \Delta_0(\text{Tr}_B|\phi\rangle\langle\phi|),$$

where  $|\tilde{\psi}_{AA'}\rangle = \text{Tr}_{BB'}|\tilde{\psi}\rangle\langle\tilde{\psi}|$ .

Now, since  $\text{Tr}_{A'B'}|\tilde{\psi}\rangle\langle\tilde{\psi}|$  has fidelity  $1 - \epsilon$  to  $|\psi_{AB}\rangle$ , we can choose a pure state  $|\iota_{A'B'}\rangle$  such that

$$F(|\tilde{\psi}_{ABA'B'}\rangle, |\psi_{AB}\rangle \otimes |\iota_{A'B'}\rangle) \geq 1 - \epsilon.$$

Introducing  $\psi_A = \text{Tr}_B|\psi\rangle\langle\psi|$  and  $\iota_{A'} = \text{Tr}_{B'}|\iota\rangle\langle\iota|$ , we infer, from the monotonicity of the fidelity, that

$$F(|\tilde{\psi}_{AA'}\rangle, \psi_A \otimes \iota_{A'}) \geq 1 - \epsilon,$$

from which it follows by standard inequalities [20] that

$$\|\tilde{\psi}_{AA'} - \psi_A \otimes \iota_{A'}\|_1 \leq \sqrt{4\epsilon}.$$

Now we can use lemma 5 to lower bound  $\Delta_0(|\tilde{\psi}_{AA'}\rangle)$  in terms of  $\Delta_{\sqrt[4]{4\epsilon}}(\psi_A \otimes \iota_{A'})$ , which is bounded in turn, using lemma 6, by  $\Delta_{\sqrt[8]{4\epsilon}}(\psi_A)$ , which proves the theorem.  $\square$

Using the additivity of the Rényi entropies, and that  $S_\alpha(\frac{1}{2}\mathbb{1}) = 1$  for all  $\alpha$ , we observe that

$$\Delta_0\left(\rho \otimes \frac{1}{2}\mathbb{1}\right) = \Delta_0(\rho).$$

This implies

**Corollary 9** The lower bound on  $C$  of theorem 8 continues to hold even if the starting state  $|\phi_{AB}\rangle$  is supplemented by unlimited numbers of EPR pairs.  $\square$

Now suppose that  $|\phi_{AB}\rangle$  can be converted into a high-fidelity copy of  $|\psi_{AB}\rangle$  using an LOCC protocol in which only  $C$  bits are exchanged between Alice and Bob. By consuming EPR pairs for superdense coding [6], this protocol can be converted into a protocol requiring only  $C/2$  qubits of communication. Since the lower bound of the corollary applies to the modified protocol, we conclude that for classical communication our bound can be improved by a factor of two.

**Corollary 10** If the state transformation  $|\phi_{AB}\rangle$  to  $|\psi_{AB}\rangle$  can be accomplished with fidelity  $1 - \epsilon$  by exchanging a total of  $C$  classical bits then, with  $\delta = \sqrt[8]{4\epsilon}$ ,

$$C \geq \Delta_\delta(\text{Tr}_B|\psi\rangle\langle\psi|) - \Delta_0(\text{Tr}_B|\phi\rangle\langle\phi|) + 2\log(1 - \delta).$$

$\square$

**Remark 11** Sometimes, direct application of these results can give an overly conservative lower bound because  $\Delta_0(\text{Tr}_B|\phi\rangle\langle\phi|)$  can be much larger than the corresponding  $\Delta_\epsilon(\text{Tr}_B|\phi\rangle\langle\phi|)$ .

Here we note that a lower bound on  $C$  in terms of  $\Delta_\epsilon$  of both the initial and the final state exists: simply observe that changing the initial state  $|\phi\rangle$  into some state  $|\phi'\rangle$  with fidelity  $1 - \epsilon_0$ , the protocol results in a state  $\rho'$  that has fidelity  $1 - \epsilon_0$  to  $\rho$  (because the fidelity does not decrease under completely positive trace preserving maps), which in turn has fidelity  $1 - \epsilon$  to  $|\psi\rangle$ . By a result of [3] this implies that the transformation from  $|\phi'\rangle$  to  $|\psi\rangle$  has fidelity  $1 - \epsilon'$ , with some universal function  $\epsilon'$  of  $\epsilon$  and  $\epsilon_0$ . We may then apply theorem 8 to this transformation.

**Remark 12** Of course one can also define a robust version of our previous  $\Delta^{\alpha\beta}$  (see remark 2):

$$\Delta_\epsilon^{\alpha\beta}(\rho) := \min \left\{ \frac{\log\left(\sum_{j \in J} r_j^\alpha\right)}{1 - \alpha} - \frac{\log\left(\sum_{j \in J} r_j^\beta\right)}{1 - \beta} \right\},$$

again with minimisation over all subsets of indices  $J$  such that  $\sum_{j \in J} r_j \geq 1 - \epsilon$ . Unsurprisingly, a variant of theorem 8 also holds for this quantity:

Consider an entanglement transformation from  $|\phi\rangle$  to  $|\psi\rangle$  with fidelity  $1 - \epsilon$  and a total communication cost of  $C$  qubits. Then, for  $0 \leq \alpha < 1 < \beta \leq \infty$ ,

$$2C \geq \Delta_\delta^{\alpha\beta}(\text{Tr}_B|\psi\rangle\langle\psi|) - \Delta_0^{\alpha\beta}(\text{Tr}_B|\phi\rangle\langle\phi|) + \delta',$$

with  $\delta = \sqrt[8]{4\epsilon}$  and  $\delta' = \left(\frac{2\alpha}{1-\alpha} + \frac{2\beta}{\beta-1}\right)\log(1 - \sqrt{\delta})$ .

The proof is slightly more cumbersome version of the proof for the  $\Delta_\delta = \Delta_\delta^{0\infty}$  case.

### III. ENTANGLEMENT CONCENTRATION AND DILUTION

In [5] it was shown that, using only local operations, Alice and Bob can convert a state  $|\psi_{AB}\rangle^{\otimes n}$  to a high fidelity approximation of  $|\phi_2^+\rangle^{\otimes nE(\psi)-O(\sqrt{n})}$ . We reproduce the argument here, as the relevant concepts are used again in the dilution protocol and our lower bound.

Diagonalise  $\rho_A = \text{Tr}_B |\psi\rangle\langle\psi| = \sum_{i=1}^d r_i |e_i\rangle\langle e_i|$ . For a distribution  $P$  on  $\{1, \dots, d\}$  we can introduce the *type classes* of sequences  $i^n = i_1 \dots i_n$ :

$$\mathcal{T}_P^n := \{i^n : \forall i \ N(i|i^n) = nP(i)\},$$

where  $N(i|i^n)$  counts the number of occurrences of  $i$  in  $i^n$ . The number of non-empty type classes is  $\binom{n+d-1}{d-1} \leq (n+1)^d$ , and the corresponding  $P$  are called *n-types*.

For  $\delta > 0$  we have the set of *typical sequences*

$$\mathcal{T}_{r,\delta}^n := \bigcup \left\{ \mathcal{T}_P^n : P \text{ s.t. } \forall i \ |P_i - r_i| \leq \frac{\delta \sqrt{r_i(1-r_i)}}{\sqrt{n}} \right\}.$$

Standard facts about these concepts are to be found in [25] (see also [7]):

$$r^{\otimes n}(\mathcal{T}_{r,\delta}^n) \geq 1 - \frac{d}{\delta^2}, \quad (14)$$

$$\forall i^n \in \mathcal{T}_P^n \quad r^{\otimes n}(i^n) = \exp(-n(D(P||r) + H(P))), \quad (15)$$

with the relative entropy (or entropy divergence)  $D(P||r) = \sum_i P_i \log \frac{P_i}{r_i}$ . Furthermore,

$$|\mathcal{T}_{r,\delta}^n| \leq \exp(nH(r) + Kd\delta\sqrt{n}), \quad (16)$$

$$|\mathcal{T}_P^n| \leq \exp(nH(P)), \quad (17)$$

$$|\mathcal{T}_P^n| \geq (n+1)^{-d} \exp(nH(P)), \quad (18)$$

$$|\mathcal{T}_P^n| \geq \exp(nH(r) - Kd\delta\sqrt{n}) \quad \text{if } P \text{ typical}, \quad (19)$$

for an absolute constant  $K > 0$ . These sets allow for the definition of corresponding projectors  $\Pi_P^n := \sum_{i^n \in \mathcal{T}_P^n} |e_{i^n}\rangle\langle e_{i^n}|$ , and similarly  $\Pi_{\rho,\delta}^n$ , with probability and trace relations identical to eqs. (14–19). Note that  $H(r) = S(\rho)$ , by definition.

The concentration protocol only requires Alice and Bob to each independently perform the projective measurement  $(\Pi_P^n)_P$  *n-type*. (Without loss of generality  $|\psi\rangle$  is in Schmidt diagonal form, and the bases with respect to which the projectors are defined are *identical* eigenbases of the reduced states.) The result  $P$  will be the same for Alice and Bob, and by eq. (14) it will be typical with probability  $\geq 1 - \epsilon$  (choosing  $\delta$  large enough). Moreover, by eq. (19) the resulting states  $|\phi_P\rangle$  are maximally entangled states of Schmidt rank  $\geq \exp(nH(r) - Kd\delta\sqrt{n})$ . Local measurements, corresponding to a partition of  $\mathcal{T}_P^n$  into blocks of size  $2^m$  (and a remainder of smaller size), for  $m = \lfloor nH(r) - Kd\delta\sqrt{n} + \log \epsilon \rfloor$ , project this further down to a state isomorphic to  $|\phi_2^+\rangle^{\otimes m}$ , with probability  $1 - \epsilon$ . This shows that  $|\psi\rangle^{\otimes n}$  can be converted by

local operations into  $m$  EPR pairs, with fidelity  $1 - 2\epsilon$ , establishing that asymptotically  $|\psi\rangle$  is worth  $E(\psi)$  EPR pairs.

In the same work it was demonstrated that the reverse is true as well: using LOCC,  $|\phi_2^+\rangle^{\otimes nE(\psi)+O(\sqrt{n})}$  can be converted to a high fidelity approximation of  $|\psi_{AB}\rangle^{\otimes n}$ .

Alice simply prepares the state  $\Pi_{\rho,\delta}^n |\psi\rangle^{\otimes n}$  (properly normalized) locally. By eq. (16) it has Schmidt rank  $\leq \exp(nH(r) + Kd\delta\sqrt{n})$ , enabling Alice to teleport [4] the half intended for Bob using  $nH(r) + Kd\delta\sqrt{n}$  EPR pairs.

Note that this method requires communication of  $2nE(\psi) + O(\sqrt{n})$  classical bits from Alice to Bob, which is of the order of the entanglement manipulated. Whether this amount can be reduced is, therefore, a legitimate and interesting question. In [17] it was shown that, indeed, communication of  $O(\sqrt{n})$  classical bits are sufficient, by the following method:

They demonstrated that there exists a state  $|\chi\rangle$  entangling  $O(\sqrt{n})$  qubits, and local unitaries  $U_A$  and  $U_B$  such that

$$F \left( (U_A \otimes U_B) |\psi\rangle^{\otimes n}, |\phi_2^+\rangle^{\otimes nE-O(\sqrt{n})} \otimes |\chi\rangle \right) \geq 1 - \epsilon. \quad (20)$$

This state arises naturally by looking at what was done in the concentration procedure above, in a reversible setting. Applying the same dilution procedure as before but to the smaller state  $|\chi\rangle$ , that is, local preparation by Alice followed by teleportation of Bob's share, then only consumes  $O(\sqrt{n})$  ebits and twice that amount of classical communication (as Lo [16] has shown this factor can be reduced to 1, i.e., a state of Schmidt rank  $d$  can be prepared using  $\log d$  bits of entanglement and communicating  $\log d$  classical bits).

Let us now apply our main result to show that any protocol to create  $|\psi\rangle^{\otimes n}$  up to fidelity  $1 - \epsilon$  from EPR pairs must use  $\Omega(\sqrt{n})$  bits of communication:

Noting first that EPR pairs have  $\Delta_0 = 0$ , we have only to lower bound  $\Delta_\delta(\psi_A^{\otimes n})$  in order to make use of theorem 8. This we do by using eq. (9). First, we show that

$$S_{\infty,\delta}(\psi_A^{\otimes n}) \leq nE(\psi) - D(\epsilon)\sqrt{n} + o(\sqrt{n}),$$

with a constant  $D(\epsilon) > 0$  (for  $\delta = \sqrt[8]{4\epsilon} < 1/2$ ):

Observe that  $S_{\infty,\delta}$  is particularly easy to understand; it is the negative logarithm of the largest eigenvalue such that the sum of the eigenvalues exceeding this one is bounded by  $\delta$ .

Define the independent and identically distributed (i.i.d.) random variables  $X_j$ ,  $j = 1, \dots, n$ , by letting

$$\Pr\{X_j = -\log r_i\} = r_i,$$

where  $(r_i)$  are the Schmidt coefficients of  $|\psi\rangle$ . Note that their expectation  $\mathbb{E}X_j$  equals  $E(\psi)$ , and that they are nonconstant, unless  $|\psi\rangle$  is maximally entangled, so that the variance  $\sigma^2$  is nonzero.

Hence we can apply the central limit theorem:

$$\Pr \left\{ \sum_{j=1}^n X_j \leq nE(\psi) + x\sigma\sqrt{n} \right\} \rightarrow \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt.$$

This implies that the sum of the largest eigenvalues, from  $\exp(-nE(\psi) + D(\epsilon)\sqrt{n} + o(\sqrt{n}))$  up (including multiplicities), is bounded from below by  $\delta$ , and our claim follows.

Next, we lower bound  $S_{0,\delta}(\psi_A^{\otimes n})$ . An optimal set  $J$  in the definition, eq. (6), must consist of the indices of the  $|J|$  largest eigenvalues such that their sum is barely above  $1 - \epsilon$ .

Once more invoking the central limit theorem, the sum of the smallest eigenvalues (including multiplicities) of  $\psi_A^{\otimes n}$  up to  $\exp(-nE(\psi) - D(\epsilon)\sqrt{n} + o(\sqrt{n}))$  is at least  $\delta$ .

We exhibit now a large type class inside the set corresponding to larger eigenvalues, which by the preceding is a subset of  $J$ : there exists an  $n$ -type  $P$  such that  $|P_i - r_i| \leq 1/n$ , for all  $i$ . This entails that for  $i^n \in \mathcal{T}_P^n$

$$\begin{aligned} \log r^{\otimes n}(i^n) &= n \sum_i P_i \log r_i \\ &= n \sum_i \left( r_i \pm \frac{1}{n} \right) \log r_i \\ &= -nH(r) \pm \sum_i |\log r_i| \\ &= -nE(\psi) \pm C. \end{aligned}$$

Thus,  $\mathcal{T}_P^n \subset J$  as soon as  $D(\epsilon) > 0$  and  $n$  is large enough.

On the other hand, because  $\|P - r\|_1 \leq d/n$ , we have (using a well-known estimate for Shannon entropies, see [7]) that

$$|H(P) - H(r)| \leq \frac{d}{n} \log n,$$

and we conclude, by eq. (18), that

$$\begin{aligned} |J| &\geq |\mathcal{T}_P^n| \\ &\geq (n+1)^{-d} \exp(nH(P)) \\ &\geq \exp(nH(r) - d \log n - d \log(n+1)). \end{aligned}$$

It follows that  $S_{0,\delta}(\psi_A^{\otimes n}) \geq nE(\psi) - O(\log n)$ .

Combining the estimates of  $S_{0,\delta}$  and  $S_{\infty,\delta}$ , we obtain

**Theorem 13** *For every bipartite pure state  $|\psi_{AB}\rangle$  that is neither separable nor maximally entangled and every sufficiently small  $\epsilon$  there exists a positive constant  $D(\epsilon)$  such that the communication cost of creating  $|\psi\rangle^{\otimes n}$  up to fidelity  $1 - \epsilon$  from EPR pairs is at least  $C \geq D(\epsilon)\sqrt{n} - o(\sqrt{n})$ .  $\square$*

**Remark 14** *Recently, secret shared randomness has been proposed as a “classical analogue of entanglement” [9], partly to increase intuition on entanglement transformations, and partly to be able to distinguish the*

*quantum effects of entanglement from those that are statistically explainable.*

*Specifically, pure state entanglement was paralleled to classical perfect correlation: Alice and Bob share a joint random variable  $(X, Y)$ , where  $X$  belongs to Alice,  $Y$  to Bob and  $X = Y$  with probability 1. Entanglement transformations by LOCC have their analogue in transformations of these random variables by local (classical) actions and public discussion, which can be listened to by an eavesdropper. The analogue of EPR pairs are shared random bits:  $\Pr\{X = Y = 0\} = \Pr\{X = Y = 1\} = 1/2$ .*

*Now it is an easy result of the theory of shared randomness (see [1] for definitions) that in an i.i.d. setting  $(X, Y)$  can be asymptotically converted into the Shannon entropy  $H(X)$  of  $X$  many shared secret bits and, inversely, this amount of shared randomness can be used to generate  $(X, Y)$ : more precisely, both transformations can be performed with asymptotically vanishing total variational distance of the distributions. These operations are the classical analogues of entanglement concentration and dilution.*

*What is remarkable is that in this setting both the concentration and dilution processes require no public discussion whatsoever. Thus, our  $\Omega(\sqrt{n})$  lower bound is a purely quantum phenomenon that has no counterpart in the “classical analogue”.*

#### IV. STATES WITH LARGE COMMUNICATION COST

In [10], the states

$$|\mu(n)\rangle = \frac{1}{\sqrt{H_n}} \sum_{i=1}^n \frac{1}{\sqrt{i}} |i\rangle|i\rangle,$$

with the harmonic sum  $H_n = \sum_{i=1}^n \frac{1}{i}$ , were introduced to show that the concept of “approximate pure state transformations with unlimited catalysis” allows *any* state transformation (this was dubbed “embezzling entanglement” in [10]). In particular it was shown that for every pure state  $|\phi\rangle$  of Schmidt rank  $m$  there are local isometries  $U_A$  and  $U_B$  such that

$$F(|\mu(n)\rangle \otimes |\phi\rangle, (U_A \otimes U_B)|\mu(n)\rangle) \geq 1 - \frac{\log m}{\log n}.$$

It is straightforward to verify that the entanglement of  $|\mu(n)\rangle$  is asymptotically  $\frac{1}{2} \log n$ , and we shall demonstrate here that the communication cost to produce it from EPR pairs is of the same order:

Theorem 8 asks us to lower bound  $\Delta_\delta$  of Alice’s reduced state

$$\rho_A = \frac{1}{H_n} \sum_{i=1}^n \frac{1}{i} |i\rangle\langle i|,$$

which we do using eq. (9):

$$S_{0,\delta}(\rho_A) = \log \min \left\{ k : \sum_{i=k+1}^n \frac{1}{iH_n} \leq \delta \right\}, \quad (21)$$

$$S_{\infty,\delta}(\rho_A) = \log H_n + \log \max \left\{ k : \sum_{i=1}^{k-1} \frac{1}{iH_n} \leq \delta \right\}. \quad (22)$$

Now, asymptotically  $(\log n) - 1 \leq H_n \leq \log(n+1)$ , and eqs. (21) and (22) allow us to estimate

$$\Delta_\delta(\rho_A) \geq ((1-2\delta) \log n) - 4 - \log \log(n+1), \quad (23)$$

resulting in a lower bound

$$C \geq \left( \frac{1}{2} - \sqrt[8]{4\epsilon} \right) \log n - O(\log \log n)$$

for the communication cost to create  $|\mu(n)\rangle$  up to fidelity  $1-\epsilon$  from EPR pairs. In fact, the classical communication cost is, by corollary 10, lower bounded by  $(1-o(1)) \log n$ , asymptotically matching the upper bound  $\log n$  from Lo's earlier mentioned state preparation method in [16].

Other states with entanglement being of the same order as the communication necessary to create them are the  $|\chi\rangle$  of eq. (20): their entanglement is at most  $O(\sqrt{n})$  while theorem 13 implies a lower bound of  $\Omega(\sqrt{n})$  on the communication resources.

## V. CONCLUSION

We have exhibited the first quantitative lower bound on the communication cost of general entanglement transformations. It is good enough to prove that the Lo/Popescu protocol of entanglement dilution is within a constant factor of being optimal, requiring  $\Theta(\sqrt{n})$  bits of communication. Also, it can be used to show that there exist states whose communication cost for creation from EPR pairs is of the same order as their entanglement, making local preparation and teleportation essentially the optimal strategy.

It is unknown to us how tight our lower bound can be made or if there is an upper bound involving similar

quantities, so we leave these questions open for future research. On a different note, it has repeatedly been speculated (such as in [17]) that the classical communication cost is related to the *loss of entanglement* in a transformation. Observe that this seems to fit perfectly for concentration and dilution, and it might be that in an appropriate model the entanglement loss in a pure state transformation provides an upper bound on the minimal communication cost required to perform it.

Other applications may include the study of quantum communication complexity, where a technique for lower bounding the communication exists [2, 8, 11] that requires estimation of the communication cost of certain pure state entanglement transformations. In the cited works this cost was lower bounded by observing that some measure of entanglement has increased. Our method could be useful as it gives nontrivial lower bounds even when the entanglement remains constant or decreases, and continues to be effective in the presence of unlimited numbers of EPR pairs.

After the present paper was finished, the independent work of Harrow and Lo [13] came to our attention, which proves the  $\Omega(\sqrt{n})$  lower bound on entanglement dilution by a different method (though there are similarities) that simultaneously provides a lower bound on the entanglement loss.

## Acknowledgments

We thank Wim van Dam for his suggestion to also consider general Rényi entropies, and Karol and Michal Horodecki for posing the problem solved in remark 3. We want to thank Aram Harrow and Hoi-Kwong Lo for making their draft of [13] available to us and for stimulating discussions.

P.H. was supported by US National Science Foundation grant no. EIA-0086038 and a Sherman Fairchild Fellowship. A.W. is supported by the U.K. Engineering and Physical Sciences Research Council. This work was carried out during the second author's visit to the Institute of Quantum Information, Caltech, in January 2002.

- 
- [1] R. Ahlswede, I. Csiszár, "Common Randomness in Information Theory and Cryptography. I. Secret Sharing", IEEE Trans. Inf. Theory, vol. 39, no. 4, pp. 1121-1132, 1993.
- [2] A. Ambainis, L. Schulman, A. Ta-Shma, U. Vazirani, A. Wigderson, "The quantum communication complexity of sampling", in: Proceedings of 39<sup>th</sup> FOCS, pp. 342-351, 1998.
- [3] H. Barnum, C. A. Fuchs, R. Jozsa, B. Schumacher, "General fidelity limit for quantum channels", Phys. Rev. A,

vol. 54, no. 6, pp. 4707-4711, 1996.

- [4] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, W. K. Wootters, "Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels", Phys. Rev. Letters, vol. 70, no. 13, pp. 1895-1899, 1993.
- [5] C. H. Bennett, H. J. Bernstein, S. Popescu, B. W. Schumacher, "Concentrating partial entanglement by local operations", Phys. Rev. A, vol. 53, no. 4, pp. 2046-2052, 1996.



- [6] C. H. Bennett, S. Wiesner, “Communication via one- and two-particle operators on Einstein-Podolsky-Rosen states”, *Phys. Rev. Letters*, vol. 69, no. 20, pp. 2881-2884, 1992.
- [7] I. Csiszar, J. Körner, *Information Theory: Coding Theorems for Discrete Memoryless Systems*, Academic Press, London 1981.
- [8] R. Cleve, W. van Dam, M. A. Nielsen, A. Tapp, “Quantum Entanglement and the Communication Complexity of the Inner Product Function”, in: *Proceedings of 1<sup>st</sup> NASA International Conference on Quantum Computing and Quantum Communications*, Colin P. Williams (Ed.), Springer Verlag, LNCS 1509, pp. 61-74, 1998.
- [9] D. Collins, S. Popescu, “A classical analogue of entanglement”, *Phys. Rev. A*, vol. 65, 032321, 2002.
- [10] W. van Dam, P. Hayden, “Embezzling Entangled Quantum States”, e-print [quant-ph/0201041](#), 2002.
- [11] W. van Dam, P. Hayden, “Communication Complexity of Quantum State Transformations”, e-print [quant-ph/0204093](#), 2002.
- [12] L. Hardy, “Method of areas for manipulating the entanglement properties of one copy of a two-particle pure entangled state”, *Phys. Rev. A*, vol. 60, no. 3, pp. 1912-1923, 1999.
- [13] A. Harrow, H.-K. Lo, “A tight lower bound on the amount of classical communication necessary for entanglement dilution”, e-print [quant-ph/0204096](#), 2002.
- [14] R. Jozsa, “Fidelity for mixed quantum states”, *J. Mod. Opt.*, vol. 41, pp. 2315-2323, 1994.
- [15] J. G. Jensen, R. Schack, “A Simple Algorithm for Local Conversion of Pure States”, *Phys. Rev. A*, vol. 63, 062303, 2001.
- [16] H.-K. Lo, “Classical-communication cost in distributed quantum-information processing: A generalization of quantum-communication complexity”, *Phys. Rev. A*, vol. 62, 012313, 2000.
- [17] H.-K. Lo, S. Popescu, “Classical Communication Cost of Entanglement Manipulation: Is Entanglement an Interconvertible Resource?”, *Phys. Rev. Letters*, vol. 83, no. 7, pp. 1459-1462, 1999.
- [18] H.-K. Lo, S. Popescu, “Concentrating entanglement by local actions: Beyond mean values”, *Phys. Rev. A*, vol. 63, 022301, 2001.
- [19] M. A. Nielsen, “Conditions for a Class of Entanglement Transformations”, *Phys. Rev. Letters*, vol. 83, no. 2, pp. 436-439, 1999.
- [20] M. A. Nielsen, I. L. Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, Cambridge (U.K.), 2000.
- [21] A. Rényi, “On measures of entropy and information”, *Proc. 4<sup>th</sup> Berkeley Sympos. Math. Statist. and Prob.*, Vol. I, pp. 547-561, Univ. California Press, Berkeley, CA, 1961.
- [22] A. Uhlmann, “The ‘transition probability’ in the state space of a \*-algebra”, *Rep. Math. Phys.*, vol. 9, pp. 273-279, 1976.
- [23] G. Vidal, D. Jonathan, M. A. Nielsen, “Approximate transformations and robust manipulation of bipartite pure-state entanglement”, *Phys. Rev. A*, vol. 62, 012304, 2000.
- [24] A. Winter, “Coding theorem and strong converse for quantum channels”, *IEEE Trans. Inf. Theory*, vol. 45, no. 7, pp. 2481-2485, 1999.
- [25] J. Wolfowitz, *Coding Theorems of Information Theory*, 2nd edition, Springer Verlag, Berlin 1964.