

Quantum Dynamics as a Stochastic Process

M. S. Torres Jr and J. M. A. Figueiredo¹
Universidade Federal de Minas Gerais - Dept. de Física
Caixa Postal 702 - Belo Horizonte - Brazil - 30.123-970
(October 28, 2018)

We study the classical motion of a particle subject to a stochastic force. We then present a perturbative schema for the associated Fokker-Planck equation where, in the limit of a vanishingly small noise source, a consistent dynamical model is obtained. The resulting theory is similar to Quantum Mechanics, having the same field equations for probability measures, the same operator structure and symmetric ordering of operators. The model is valid for general electromagnetic interaction as well as many body systems with mutual interactions of general nature.

Since their introduction in Physics stochastic processes became an indispensable tool in the analysis of random phenomena. In many common physical situations this mathematical theory arises as a natural description language to such an extent that, rigorously speaking, we can't discard noise effects from any realistic description of natural phenomena. On the other hand Quantum Mechanics introduced probability concepts to physics without explicit reference to any random phenomena, although it is believed that this theory must generate a random vacuum field in order to explain more subtle quantum processes. Thus it was quite natural that efforts in seeking a stochastic nature to quantum phenomena have been presented, the pioneering one being featured by Nelson [2]. In the very beginning of the Quantum Theory Planck [3], Einstein and Hopf [4] tried to understand blackbody radiation in a statistical basis. Subsequently Einstein and Stern [5] developed a stochastic theory based also on a random vacuum field in order to explain low temperature corrections to specific heat of the solids.

The advent of the Quantum Theory solved the intriguing problems issued by the enormous sequence of experiments that opened the mysteries of the microscopic world in the first two decades of the last century. The price paid was the astonishing set of obscure epistemological foundations presented by Quantum Theory. At that time such structural problems were a scenario for deep discussions and controversies between Einstein and Bohr. It is in EPR and Bell-like [6] experiments that researchers usually look for evidence of a nontrivial indication that Quantum Mechanics is in fact a fundamental theory. The successful predictions of Quantum Mechanics have wakened physicists' epistemological demands in justifying its basic axioms, and in many cases some authors were amazed by its intriguing (epistemological) mysteries, which are believed to be inevitable and denying any possibility of scientific reasoning on alternative paths.

Rescuing the original phenomenological view of the scientific thought of Planck and Einstein, as presented in their first works, Boyer [7] showed that relativistic invariance of the Einstein-Hopf vacuum leads to a vacuum spectral density that is compatible with blackbody radiation law. This may be an indication that a fluctuating vacuum field, if real, may explain some observable phenomena only understood within the scope of the Quantum Theory. Unfortunately Boyer's work and its subsequent consequences proved to have no advantage over Quantum Electrodynamics yet presenting some additional difficulties on many calculations. Nevertheless, the idea of a real fluctuating vacuum everywhere in the Universe should not be discarded since it provides a possible empirical basis for a phenomenological theory compatible to Quantum Mechanics without its epistemological inconsistencies and hopefully providing additional, predictable phenomena.

In this work we present a theory that points toward this line. We study the motion of a particle subject to a stochastic force of Wiener type [8], generating a specific kind of Brownian motion described by its associated Fokker-Planck equation. The main ideas displayed here were presented on an earlier work [9] where a deeper justification for the stochastic process we consider is found as well as some consequences not discussed this time. On the other hand we do make a generalization of that work featuring a) the tridimensional case b) motion is subject to electric and magnetic external fields and c) the many-particle case. As a result we show that it is possible to obtain a fully consistent physical theory that is a proper generalization of Quantum Mechanics, based on phase space dynamics, with full classical content. In addition the perturbative series of the Fokker-Planck equation assumed here seems to be very efficient, since it provides the correct quantum mechanical symmetric ordering of operators coming from a full classical interpretation, without the need to calculate redundant combinations of operators. In consequence a new term in the expected values of fourth order (and above) correlations shows up. A careful calculation using Quantum Mechanics demonstrates the existence of this extra term, never considered in conventional quantum mechanical calculations.

As stressed in the above considerations our picture is classical on its epistemological nature. Nontrivial effects of noise is the main ingredient on the kind of dynamical model we introduce now. Lets consider the motion of a particle of mass μ subject to electric and magnetic fields. A stochastic force is added to the associated Hamilton equations of motion which are then given by

$$\begin{aligned} d\mathbf{x}(t) &= \nabla_p \mathcal{H} dt \equiv \frac{1}{\mu} \left(\mathbf{p}(t) - \frac{e}{c} \mathbf{A}(\mathbf{x}(t), t) \right) dt \\ d\mathbf{p}(t) &= -\nabla_x \mathcal{H} dt + \sqrt{2\mu P} d\mathbf{W}(t) \\ \mathcal{H}(\mathbf{p}, \mathbf{x}, t) &= \frac{1}{2\mu} \left[\mathbf{p} - \frac{e}{c} \mathbf{A}(\mathbf{x}, t) \right]^2 + V(\mathbf{x}, t) \end{aligned} \quad (1)$$

where the subscript in the ∇ operator stands for derivatives in the corresponding variable, P is a "vacuum power" and $d\mathbf{W}(t)$ is a vectorial Wiener-type stochastic variable satisfying $d\mathbf{W}(t)^2 = dt$. As we will see the particle may gain energy from the vacuum at a rate P . This apparently unphysical situation will be carefully removed later when a proper perturbation theory will be developed, leaving a consistent theory at the end of our reasoning. However we must be aware of the fact that a real noise may exist. For example, since Universe is expanding against a event horizon having a specific gravity, some energy is necessary to change its curvature. Quantum mechanical calculations of this effect leads to an effective "vacuum temperature" known as the Gibbons-Hawking effect [10]. We may also invert the argument and claim that vacuum effects supply the energy for the expanding Universe and simultaneously supports quantum phenomena. Therefore it would not be a surprise if, eventually, experiments come to confirm that, in fact, every massive particle absorbs some energy from vacuum. This has in fact has been considered by some cosmological models in order to explain the accelerated expansion [11].

The motion in an electromagnetic field described by eqn(1) demands the inclusion of a radiation reaction term that gives back to vacuum part of the received power thus minimizing the effects of direct observation of these fluctuations. We leave for a future work the inclusion of this term since we believe it should not contribute to the main conclusions of the present work. The pertinent point is that classical orbits are very sensitive to stochastic perturbations *no matter* the value of vacuum power. This means that even for infinitesimally small value of P the above equations admit a probabilistic description, resulting in a Fokker-Planck equation for the associated probability distribution. The procedure to get this equation from a Wiener source is standard, essentially using general properties of Ito calculus [8] along with statistical properties of the noise source. We naturally arrive at a phase space description where classical trajectory concept is lost and the associated phase space distribution function $\Phi(\mathbf{x}, \mathbf{p}, t)$ satisfies

$$\frac{\partial \Phi}{\partial t} = -(\nabla_p \mathcal{H}) \cdot (\nabla_x \Phi) + (\nabla_x \mathcal{H}) \cdot (\nabla_p \Phi) + \mu P \nabla_p^2 \Phi \quad (2)$$

Now \mathbf{x} and \mathbf{p} are no longer functions of time, meaning that we can't assign a given trajectory to any point in phase space. All sample trajectories are possible and, although real, they are not differentiable [8], so a deterministic relationship between position and momentum is not possible anymore. In short, the concept of a classical deterministic motion is lost although particle reality, as well as its trajectory, is not.

One general feature of this equation is that it is norm preserving. In fact the time derivative of the norm is

$$\frac{d}{dt} \int \Phi(\mathbf{x}, \mathbf{p}, t) d\Omega = \int [-(\nabla_p \mathcal{H}) \cdot (\nabla_x \Phi) + (\nabla_x \mathcal{H}) \cdot (\nabla_p \Phi) + \mu P \nabla_p^2 \Phi] d\Omega$$

where $d\Omega = d^3\mathbf{x}d^3\mathbf{p}$ is phase space differential volume. Direct integration by parts gives

$$\frac{d}{dt} \int \Phi(\mathbf{x}, \mathbf{p}, t) d\Omega = \int [(\nabla_x \cdot \nabla_p \mathcal{H}) \Phi - (\nabla_p \cdot \nabla_x \mathcal{H}) \Phi] d\Omega = 0$$

This remarkable property is general for the kind of stochastic process we consider here, being valid for any type of classical motion, regardless its specific dynamics, since it is Hamiltonian-independent. This will be considered later as a fundamental condition for defining the structure of the probability measure supporting our theory. The following function, with domain in phase space

$$H(\mathbf{x}, \mathbf{p}, t) \equiv \frac{1}{2m} \left(\mathbf{p} - \frac{e}{c} \mathbf{A}(\mathbf{x}, t) \right)^2 + V(\mathbf{x}, t)$$

cannot be confused with the (classical) particle's energy due to the breakdown of trajectory reality but its average is an observable that changes at a rate P [9]

$$\frac{d}{dt} \langle H \rangle = \frac{d}{dt} \int \Phi(\mathbf{x}, \mathbf{p}, t) H(\mathbf{x}, \mathbf{p}, t) d\Omega = P$$

and should be interpreted as the mean particle energy. Thus, as anticipated, the particle gains energy at a rate P that is proportional to noise intensity. The (small) value of vacuum power would, if a real phenomenon, be a thermodynamic property of the Universe, common to all existing elementary particles. In this case, its measurable consequences can probably exist at cosmological scales, but we will show that its stochastic effects are dominant at microscopic ones, influencing the motion of small bodies and giving rise to a dynamical process that, as we will see, resembles quantum dynamics. This is the kind of motion we want to describe here. It shares many common features with Quantum Mechanics but it is neither a Quantum Theory nor a tentative task to give it some stochastic justification but a full thermodynamic theory showing results that promises to be as precise as Quantum Mechanics while still having the desired clearness of a thermodynamic process.

Note that motion follows in mean classical laws and do not depend on vacuum power. To see this observe that, although particle's velocity has only a stochastic meaning, its average is a well defined time-dependent observable given by $\mu \langle \mathbf{v} \rangle \equiv \langle \mathbf{p} - \frac{e}{c} \mathbf{A} \rangle$ whose time derivative is given by

$$\mu \frac{d}{dt} \langle \mathbf{v} \rangle \equiv \frac{d}{dt} \langle \mathbf{p} - \frac{e}{c} \mathbf{A} \rangle = \int \left[\left(\mathbf{p} - \frac{e}{c} \mathbf{A} \right) \frac{\partial}{\partial t} \Phi(\mathbf{x}, \mathbf{p}, t) - \frac{e}{c} \frac{\partial \mathbf{A}}{\partial t} \Phi(\mathbf{x}, \mathbf{p}, t) \right] d\Omega$$

Direct manipulation of the Fokker-Planck equation (2) show that the particle follows in mean Newton's Law where a Lorentz force term drives the motion. The result is

$$\mu \frac{d}{dt} \langle \mathbf{v} \rangle = \int \left[-\nabla_x V - \frac{e}{c} \frac{\partial \mathbf{A}}{\partial t} + \left(\mathbf{p} - \frac{e}{c} \mathbf{A} \right) \times (\nabla \times \mathbf{A}) \right] \Phi d\Omega = \langle e\mathbf{E} + \left(\mathbf{p} - \frac{e}{c} \mathbf{A} \right) \times \mathbf{B} \rangle$$

and is the stochastic version of Ehrenfest's theorem. This shows how vacuum stochastic effects would be visible even for a vanishingly small value of the vacuum power. Since motion suffers from random perturbations that smear phase space trajectories no matter the value of noise intensity and since the resulting average dynamics do not depend explicitly on its value a nontrivial theory must survive even in the limit of zero power. This "ghost" effect is a consequence of the known singular small-noise perturbation expansion of the Fokker-Planck equation [8]. To solve eqn(2) we look for a perturbative schema that captures explicitly these features.

The Fokker-Planck equation we have obtained is linear but does not have constant coefficients implying that linear operations on it may not be simple. This is particularly true for the spatial variable dependence due to the arbitrariness of the external fields but the momentum variable has a simpler linear dependence, allowing a tractable procedure in the Fourier space. In this case we did arrive to a closed form of the transformed equation where a very important feature turns out: due to its specific dependency on the momentum variable, the resulting equation (in Fourier space) is nonlinear on the scale conversion between adjoint spaces. Thus we can't simply rescale and incorporate units in the transformed variables because the physics changes. This means that only one value of the conversion factor is compatible with experimental results. In the context of the present work that's the way Planck's constant becomes important in defining scales for microscopic phenomena. More explicitly, consider the adjoint space of the momentum variable defined by

$$\varphi(\mathbf{x}, \mathbf{y}, t) = \int \Phi(\mathbf{x}, \mathbf{p}, t) \exp\left(-i \frac{\mathbf{p} \cdot \mathbf{y}}{\hbar}\right) d^3 \mathbf{p}$$

where \hbar is Planck's constant. Here the y -space hosts modes of the probability distribution associated to momentum which are waves having wavelength of value $y \simeq \hbar/p$. Direct substitution of this definition in eqn(2) results in the following equation

$$\frac{\partial \varphi}{\partial t} = \frac{\hbar}{i\mu} \frac{\partial^2 \varphi}{\partial y_j \partial x_j} - \frac{1}{i\hbar} \frac{\partial U}{\partial x_j} y_j \varphi + \frac{e}{\mu c} \frac{\partial}{\partial x_j} (A_j \varphi) + \frac{e}{\mu c} y_l \frac{\partial A_j}{\partial x_l} \frac{\partial \varphi}{\partial y_j} - \frac{\mu P}{\hbar^2} y_j y_j \varphi \quad (3)$$

where $U = V(\mathbf{x}, t) + \frac{e^2}{2\mu c^2} A_j A_j$. As anticipated this equation is nonlinear in the constant \hbar a fact present even for zero vacuum power. In what follows, we expand in a Taylor series in the variable \mathbf{y} around the origin in order to get a perturbation theory for this equation. This corresponds to a high kinetic energy perturbation series. In [9] we did this (in one dimension) by explicitly including a vacuum characteristic mode $k_v \equiv \sqrt{P/(\hbar c^2)}$ as a scaling factor in the expansion which allows a non-singular phase space reconstruction. Since we intend to take the limit $P \rightarrow 0$ later in the present work, a simpler but rather singular approach (in phase space) will be developed here. We just

approximate the Gaussian kernel used in [9] by the unity, albeit still preserving the diffusion term in eqn(3). In one dimension, the limit of $k_v = 0$ of the ref [9] obviously coincides with ours. We believe that inclusion of vacuum modes explicitly would be necessary if radiation reaction terms were included in the Fokker-Planck equation in order to get a full thermodynamic theory. Our task here is to show that a theory exists at sufficiently high energy where direct vacuum effects have minor contribution to dynamics in such a way the limit $P \rightarrow 0$ makes sense. In the present representation in adjoint space we have developed an expansion close to the origin in the variable \mathbf{y} ($\mathbf{p} \rightarrow \infty$) written as

$$\varphi(\mathbf{x}, \mathbf{y}, t) = \sum_{l,m,n} \varphi_{l,m,n}(\mathbf{x}, t) y_1^l y_2^m y_3^n \quad (4)$$

where the subscripts in \mathbf{y} refer to its Cartesian components. When inserted in eqn(3) this results in a recursive series for the coefficients $\varphi_{l,m,n}$, with only one of them unknown since that equation involves only first order derivatives in \mathbf{y} . At these high (kinetic) energies, vacuum modes have minor influence and only appear explicitly after terms of order two. However due to the existence of non-constant coefficients in eqn(3), even low order terms have a nonlinear dependence on Planck's constant, leaving the expansion with nontrivial physics. Explicitly, we have

$$\begin{aligned} \frac{\partial \varphi_{l,m,n}}{\partial t} = & \frac{\hbar}{i\mu} \left[(l+1) \frac{\partial \varphi_{l+1,m,n}}{\partial x_1} + (m+1) \frac{\partial \varphi_{l,m+1,n}}{\partial x_2} + (n+1) \frac{\partial \varphi_{l,m,n+1}}{\partial x_3} \right] - \\ & - \frac{1}{i\hbar} \left[\frac{\partial U}{\partial x_1} \varphi_{l-1,m,n} + \frac{\partial U}{\partial x_2} \varphi_{l,m-1,n} + \frac{\partial U}{\partial x_3} \varphi_{l,m,n-1} \right] + \\ & + \frac{e}{\mu c} \left(l \frac{\partial A_1}{\partial x_1} + m \frac{\partial A_2}{\partial x_2} + n \frac{\partial A_3}{\partial x_3} \right) \varphi_{l,m,n} + \\ & + \frac{e}{\mu c} (l+1) \left(\frac{\partial A_1}{\partial x_2} \varphi_{l+1,m-1,n} + \frac{\partial A_1}{\partial x_3} \varphi_{l+1,m,n-1} \right) + \\ & + \frac{e}{\mu c} (m+1) \left(\frac{\partial A_2}{\partial x_1} \varphi_{l-1,m+1,n} + \frac{\partial A_2}{\partial x_3} \varphi_{l,m+1,n-1} \right) + \\ & + \frac{e}{\mu c} (n+1) \left(\frac{\partial A_3}{\partial x_1} \varphi_{l-1,m,n+1} + \frac{\partial A_3}{\partial x_2} \varphi_{l,m-1,n+1} \right) + \\ & + \frac{e}{\mu c} \nabla \cdot (\mathbf{A} \varphi_{l,m,n}) - \frac{\mu P}{\hbar^2} [\varphi_{l-2,m,n} + \varphi_{l,m-2,n} + \varphi_{l,m,n-2}] \end{aligned} \quad (5)$$

being implicit in this equation that the terms exist only for nonnegative indices. We will show below that a consistent dynamical theory comes out for $P = 0$, having properties common to Quantum Dynamics. For this, we collect the following first two terms in the above equation which are independent of the vacuum power P

$$\frac{\partial \varphi_{0,0,0}}{\partial t} = \frac{\hbar}{i\mu} \nabla \cdot \mathbf{J} + \frac{e}{\mu c} \nabla \cdot (\mathbf{A} \varphi_{0,0,0}) \quad (6a)$$

$$\frac{\partial J_l}{\partial t} = \frac{\hbar}{i\mu} \frac{\partial T_{l,j}}{\partial x_j} - \frac{1}{i\hbar} \varphi_{0,0,0} \frac{\partial U}{\partial x_l} + \frac{e}{\mu c} \frac{\partial}{\partial x_j} (A_j J_l) + \frac{e}{\mu c} \frac{\partial A_j}{\partial x_l} J_j \quad (6b)$$

where

$$\begin{aligned} \mathbf{J} &\equiv (\varphi_{1,0,0}, \varphi_{0,1,0}, \varphi_{0,0,1}) \\ \mathbf{T} &\equiv \begin{bmatrix} 2\varphi_{2,0,0} & \varphi_{1,1,0} & \varphi_{1,0,1} \\ \varphi_{1,1,0} & 2\varphi_{0,2,0} & \varphi_{0,1,1} \\ \varphi_{1,0,1} & \varphi_{0,1,1} & 2\varphi_{0,0,2} \end{bmatrix} \end{aligned}$$

are the probability current and the correlation tensor respectively. Another set of equations involving $\varphi_{1,1,0}$ and its cyclic permutations as well as the $\varphi_{1,1,1}$ term are also P -independent. Existence of these P -independent coefficients is consequence of the high kinetic energy character of the perturbative expansion. All coefficients are linked by a recursive chain, so if we find a consistent way to calculate $\varphi_{0,0,0}$ the entire series is solved. The first P -dependent term involves time derivative of $\varphi_{2,0,0}$ (and its cyclic permutations) and spatial derivatives involving order three terms. Thus it appears that $\varphi_{0,0,0}$ and \mathbf{J} may be evaluated in a truly P -independent way representing the ghost effect referred to above. This means that, even for a very small vacuum source, the stochastic character of this dynamical problem is not lost, thus preserving its probability foundation. Detectable effects must exist and we have proved in [9] that

(if real) they take account of quantum phenomena of an elementary particle in its nonrelativistic limit. Calculations made here give an improvement on the mathematical structure of the theory, show the consistency in the interaction with electromagnetic fields as well as the consistence in the many particle case.

We initiate our analysis noting that $\varphi_{0,0,0}(\mathbf{x}, t) = \int \Phi(\mathbf{x}, \mathbf{p}, t) d^3\mathbf{p}$, meaning the first coefficient is always real and its spatial integral equals the time-independent norm in phase space. We get this result immediately by integrating eqn(6a) in the whole space. We introduce the probability amplitude $\Psi(\mathbf{x}, t)$ as $\varphi_{0,0,0}(\mathbf{x}, t) \equiv |\Psi(\mathbf{x}, t)|^2$. The last argument demands that $\Psi(\mathbf{x}, t)$ belongs to a particular \mathcal{L}^2 space \mathbf{H} having the property

$$\frac{d}{dt} \int |\Psi(\mathbf{x}, t)|^2 d^3\mathbf{x} = \int \left(\Psi^* \frac{\partial \Psi}{\partial t} + \Psi \frac{\partial \Psi^*}{\partial t} \right) d^3\mathbf{x} = 0 \quad (7)$$

which means $(\Psi, i\partial_t \Psi) = (i\partial_t \Psi, \Psi)$ for all $\Psi \in \mathbf{H}$ where (\cdot, \cdot) stands for internal product in \mathcal{L}^2 . It may be proved [12] that there is an Hermitean operator $\tilde{H}(\mathbf{x}, t)$ such that

$$i\hbar \frac{\partial \Psi}{\partial t} = \tilde{H} \Psi \quad (8)$$

for every Hilbert space vector satisfying eqn(7). Defining the (Hermitean) Hilbert space operators

$$\begin{aligned} \mathcal{M} &\equiv \frac{1}{2\mu} \left(\pi_j - \frac{e}{c} A_j(\mathbf{x}, t) \right) \left(\pi_j - \frac{e}{c} A_j(\mathbf{x}, t) \right) + V(\mathbf{x}, t) \\ \pi &\equiv \frac{\hbar}{i} \nabla \end{aligned}$$

it is not difficult to prove that for any Ψ we have

$$\Psi^* [\mathcal{M}, \pi_j] \Psi = -\frac{\hbar}{i} \frac{\partial U}{\partial x_j} |\Psi|^2 - \frac{e\hbar^2}{2\mu c} \frac{\partial^2 A_l}{\partial x_l \partial x_j} |\Psi|^2 - \frac{e\hbar^2}{\mu c} \Psi^* \frac{\partial A_l}{\partial x_j} \frac{\partial \Psi}{\partial x_l}$$

which upon integration in the whole space gives

$$\int \Psi^* [\mathcal{M}, \pi_j] \Psi d^3\mathbf{x} = -\frac{\hbar}{i} \int \frac{\partial U}{\partial x_j} |\Psi|^2 d^3\mathbf{x} - \frac{e\hbar^2}{2\mu c} \int \frac{\partial A_l}{\partial x_j} \left[\Psi^* \frac{\partial \Psi}{\partial x_l} - \Psi \frac{\partial \Psi^*}{\partial x_l} \right] d^3\mathbf{x} \quad (9)$$

We use this result in the space integration of eqn(6b) in order to get, after substitution for the potential energy term,

$$\hbar^2 \frac{d}{dt} \int J_l d^3\mathbf{x} = \int \Psi^* [\mathcal{M}, \pi_l] \Psi d^3\mathbf{x} + \frac{e\hbar^2}{\mu c} \int \frac{\partial A_l}{\partial x_j} \left[J_l - \frac{1}{2} \left(\Psi \frac{\partial \Psi^*}{\partial x_l} - \Psi^* \frac{\partial \Psi}{\partial x_l} \right) \right] d^3\mathbf{x} \quad (10)$$

The freedom in defining Ψ allows us to choose a specific functional dependence for the probability amplitude. A natural choice would simplify eqn(10) in order to generate an equation for Ψ and consequently closing the whole series. If we assume

$$\mathbf{J} \equiv \frac{1}{2} (\Psi \nabla \Psi^* - \Psi^* \nabla \Psi) \quad (11)$$

we obtain a considerable simplification of eqn(10). In addition the left side of this equation may be rewritten using eqn(8) in order to get a form similar to the right side. The result is

$$\hbar^2 \frac{d}{dt} \int \mathbf{J} d^3\mathbf{x} = \frac{\hbar^2}{2} \frac{d}{dt} \int (\Psi \nabla \Psi^* - \Psi^* \nabla \Psi) d^3\mathbf{x} = \int \Psi^* [\tilde{H}, \pi] \Psi d^3\mathbf{x}$$

in such a way that eqn(10) admits the (Hilbert space) solution $\tilde{H} = \mathcal{M}$. Using eqn(8) a field equation for the probability amplitude is also obtained

$$i\hbar \frac{\partial \Psi}{\partial t} = \frac{1}{2\mu} \left(\frac{\hbar}{i} \nabla - \frac{e}{c} \mathbf{A} \right)^2 \Psi + V(\mathbf{x}, t) \Psi$$

a condition that simultaneously satisfy eqn(6a) completing the self consistency of the proposed solution. The current vector \mathbf{J} corresponds to the (classical) momentum of the particle averaged over the momentum sector of the phase space

$$i\hbar\mathbf{J}(\mathbf{x}, t) = \int \mathbf{p}\Phi(\mathbf{x}, \mathbf{p}, t) d^3\mathbf{p}$$

so that all classical observables may be effectively calculated using a similar rule [9]. Since the spatial average of $i\hbar\mathbf{J}$ is equal to the classical average of the momentum we have

$$\langle \mathbf{p} \rangle \equiv \int \mathbf{p}\Phi(\mathbf{x}, \mathbf{p}, t) d^3\mathbf{p}d^3\mathbf{x} = \int \Psi^* \left(\frac{\hbar}{i} \nabla \right) \Psi d^3\mathbf{x}$$

We have obtained a clean formulation of Quantum Mechanics with all axioms and rules in a nice and phenomenological way as demanded by the traditional scientific reasoning. In addition to the tridimensional formulation presented here, the interaction with electromagnetic fields had the effect of determining in a clever way the functional form of the probability current because the vector potential explicitly couples to this current in the perturbative expansion, a fact not possible in the one dimensional case. Time derivative of eqn(11), along with Schrödinger equation, can be used in eqn(6b) in order to get a explicit form of the correlation tensor. The result is

$$T_{j,l} = \frac{1}{4} \left(\Psi^* \frac{\partial^2 \Psi}{\partial x_j \partial x_l} + \Psi \frac{\partial^2 \Psi^*}{\partial x_j \partial x_l} - \frac{\partial \Psi^*}{\partial x_j} \frac{\partial \Psi}{\partial x_l} - \frac{\partial \Psi}{\partial x_j} \frac{\partial \Psi^*}{\partial x_l} \right) \quad (12)$$

and completes the determination of the two first terms in the series that do not explicitly depend on the noise. Everything would work very well if vacuum fluctuations are a real, detectable phenomena. In this case the presumably small value of the vacuum power P prevents us of its detection using present experimental resolution, but since the two first terms in the expansion used here for developing the theory are independent of its value, a consistent dynamical model exists in the $P \rightarrow 0$ limit. On the other hand, usual observables are at most second order on the dynamical variables. Their averages calculated by the method presented here depends only on the first two terms of the perturbative series and coincide with those predicted by Quantum Mechanics [9] for any value of P . Consequently their values are not affected by the vacuum power itself even if it is real. In other words, no difference is noticeable between Quantum Mechanics with all its axioms and mysteries and the thermodynamic theory presented in this work in what concern a) field dynamics for Ψ (the Schrödinger equation) and b) up to second order observables. The reason, as anticipated, is the high kinetic energy limit used which prevents a explicit dependence on vacuum effects although the stochastic nature of the problem cannot be discarded even for zero vacuum power.

The very existence of other terms in the expansion show that the Fokker-Planck equation represents a dynamical theory richer than Quantum Mechanics at least in the nonrelativistic limit case. It is noticeable that the probability distribution function $\Phi(\mathbf{x}, \mathbf{p}, t)$ cannot be confused with Wigner's function which, differently of our case, does not satisfy the Fokker-Planck equation in the $P \rightarrow 0$ limit where this equation assumes a Liouville-like form. Since our theory is purely classical, expected values are calculated using classical variables that are automatically converted in quantum-like operators by the perturbative procedure we developed. In this aspect, Wigner's function shares some commons properties with the distribution function given by the Fokker-Planck equation. The particular case of the momentum angular operators is interesting. We have

$$L_l = \varepsilon_{jkl} x_j p_k \Rightarrow \langle L_l \rangle = \varepsilon_{jkl} \int x_j p_k \Phi d^3\mathbf{x} d^3\mathbf{p} = \int \Psi^* \tilde{L}_l \Psi d^3\mathbf{x}$$

where $\tilde{\mathbf{L}}$ is the quantum mechanical angular momentum operator. Thus each component of the averaged angular momentum coincides with the quantum mechanical value. However the expected value of the squared angular momentum differs from the usual quantum mechanical value. In fact we have

$$\langle \mathbf{L}^2 \rangle = \int (x_j x_j p_k p_k - x_j x_k p_j p_k) \Phi d^3\mathbf{x} d^3\mathbf{p} = -\hbar^2 \int (x_j x_j T_{kk} - x_j x_k T_{jk}) d^3\mathbf{x} \quad (13)$$

so by explicit calculation using eqn(10) we get

$$\langle \mathbf{L}^2 \rangle = \int \Psi^* \tilde{L}^2 \Psi d^3\mathbf{x} + \frac{3}{2} \hbar^2 \quad (14)$$

that is, the particle has an additional "zero point" squared angular momentum equal to $\frac{1}{2}\hbar^2$ per rotational degree of freedom. The case of a free particle is sufficient to understand this result within the framework of the present theory: while each component of the angular momentum must be zero for that particle, stochastic effects prevent it to describe a perfectly linear motion. Consequently, some motion in the plane must exist, which demands the existence

of a fluctuation in the total angular momentum. By the same argument a rotor will also present a zero point kinetic energy because $\langle E_c \rangle = \frac{1}{2I} \langle \mathbf{L}^2 \rangle$, where I is the moment of inertia of the particle. Consider now a three dimensional harmonic oscillator; in its fundamental state, the energy is $E_0 = \frac{3}{2}\hbar\omega$ (both in Quantum Mechanics and in the present stochastic model) and the total angular momentum is zero, but our model predicts that a kinetic energy equal to $\frac{1}{2I} \langle \mathbf{L}^2 \rangle$ with $I = 2\mu r^2$ and $r \equiv \sqrt{\hbar/(2\mu\omega)}$ is present. It appears that the stochastic model retains full consistence in the interpretation of quantum phenomena.

The discrepancy to direct calculation using quantum theory displayed in the eqn(14) is removed when a fully symmetric quantum operator is used. Consider the (classical) fourth order position-moment correlation function $\chi^4_{ijkl} \equiv x_i x_j p_l p_k$. There are 24 possible permutations of the associated quantum operators and many of them are redundant permutations of commuting ones. Thus the complete full symmetric quantum representation of χ^4 is

$$\begin{aligned} \widetilde{\chi^4}_{ijkl} &= \frac{1}{6} (\widetilde{x}_i \widetilde{x}_j \widetilde{p}_l \widetilde{p}_k + \widetilde{p}_l \widetilde{p}_k \widetilde{x}_i \widetilde{x}_j) + \\ &\frac{1}{12} (\widetilde{x}_i \widetilde{p}_l \widetilde{p}_k \widetilde{x}_j + \widetilde{x}_j \widetilde{p}_l \widetilde{p}_k \widetilde{x}_i + \widetilde{p}_l \widetilde{x}_i \widetilde{x}_j \widetilde{p}_k + \widetilde{p}_k \widetilde{x}_i \widetilde{x}_j \widetilde{p}_l) + \\ &\frac{1}{24} (\widetilde{x}_i \widetilde{p}_l \widetilde{x}_j \widetilde{p}_k + \widetilde{x}_j \widetilde{p}_l \widetilde{x}_i \widetilde{p}_k + \widetilde{x}_j \widetilde{p}_k \widetilde{x}_i \widetilde{p}_l + \widetilde{x}_i \widetilde{p}_k \widetilde{x}_j \widetilde{p}_l) + \\ &\frac{1}{24} (\widetilde{p}_l \widetilde{x}_i \widetilde{p}_k \widetilde{x}_j + \widetilde{p}_l \widetilde{x}_j \widetilde{p}_k \widetilde{x}_i + \widetilde{p}_k \widetilde{x}_j \widetilde{p}_l \widetilde{x}_i + \widetilde{p}_k \widetilde{x}_i \widetilde{p}_l \widetilde{x}_j) \end{aligned}$$

in such a way that after a simple but long calculation we get

$$\begin{aligned} \widetilde{\chi^4}_{ijkl} \Psi &= \widetilde{x}_i \widetilde{x}_j \widetilde{p}_l \widetilde{p}_k \Psi + \frac{\hbar}{2i} (\delta_{j,l} x_i + \delta_{i,l} x_j) \widetilde{p}_k \Psi + \\ &\frac{\hbar}{2i} (\delta_{j,k} x_i + \delta_{i,k} x_j) \widetilde{p}_l \Psi - \frac{\hbar^2}{4} (\delta_{i,l} \delta_{j,k} + \delta_{i,k} \delta_{j,l}) \Psi \end{aligned} \quad (15)$$

Since $\mathbf{L}^2 = \sum_{l,k} (\chi^4_{kkll} - \chi^4_{klkl})$ the corresponding symmetric operator is

$$\widetilde{\mathbf{L}}_N^2 \equiv \sum_{l,k} \left(\widetilde{\chi^4}_{kkll} - \widetilde{\chi^4}_{klkl} \right) = \sum_{l,k} (\widetilde{x}_k \widetilde{x}_k \widetilde{p}_l \widetilde{p}_l - \widetilde{x}_k \widetilde{x}_l \widetilde{p}_k \widetilde{p}_l) - \frac{2\hbar}{i} \sum_l \widetilde{x}_l \widetilde{p}_l + \frac{3\hbar^2}{2}$$

Expected value of $\widetilde{\mathbf{L}}_N^2$ is given by $\int \Psi^* \widetilde{\mathbf{L}}_N^2 \Psi d\mathbf{x}$, which is exactly the result displayed in eqn(13). This shows clearly that a symmetric-ordered squared angular momentum operator also presents zero point fluctuations as predicted by eqn(14) which has a purely classical interpretation. It appears that the perturbation expansion given by eqn(4) gives automatically the symmetric-ordering correlation functions from the classical observables. This means that quantum calculations using a classical construction without axioms or rules is possible. This opens conditions for perturbative analysis using higher order terms, which may come much easier with the above formalism.

The many body problem can also be handled inside the present formalism. The kind of noise considered here is uncorrelated, which means that the Wiener process for this problem is just a set of Wiener variables for each particle. The resulting Fokker-Planck equation has the same appearance shown in eqn(2), the difference being the existence of an additional index to label particles. We have found no differences on field dynamics for Ψ to Quantum Theory and, as above, all basic axioms and rules come naturally. As expected correlation functions above fourth order present differences involving vacuum terms. At least in the fourth order a symmetric-ordering is naturally obtained in a similar way worked out for the one particle case.

The Fokker-Planck equation for the many particle case is

$$\frac{\partial \Phi(\mathbf{x}, \mathbf{y}, t)}{\partial t} = \sum_{\alpha=1}^N \left[-(\nabla_{\alpha p} \mathcal{H}) \cdot (\nabla_{\alpha x} \Phi) + (\nabla_{\alpha x} \mathcal{H}) \cdot (\nabla_{\alpha p} \Phi) + m P \nabla_{\alpha p}^2 \Phi \right]$$

where greek symbols stands for particle labeling. Also \mathbf{x} and \mathbf{y} stands for $\{\mathbf{x}_\alpha\}$ and $\{\mathbf{p}_\alpha\}$. Interactions include not only external fields but mutual forces as well. The perturbative expansion analogous to eqn(4) is written as

$$\varphi(\mathbf{x}, \mathbf{y}, t) = \rho(\mathbf{x}, t) + \sum_{\alpha} J_{\alpha i}(\mathbf{x}, t) y_{\alpha}^i + \frac{1}{2} \sum_{\alpha\beta} T_{\alpha\beta ij} y_{\alpha}^i y_{\beta}^j + \frac{1}{6} \sum_{\alpha\beta\gamma} C_{\alpha\beta\gamma ijk} y_{\alpha}^i y_{\beta}^j y_{\gamma}^k + \dots$$

and its coefficients satisfy equations similar to those found above. In particular the correlation tensor $T_{\alpha\beta ij}$ has the following form

$$T_{\alpha\beta ij} = \frac{1}{4} \left(\Psi^* \frac{\partial^2 \Psi}{\partial x_\alpha^i \partial x_\beta^j} + \Psi \frac{\partial^2 \Psi^*}{\partial x_\alpha^i \partial x_\beta^j} - \frac{\partial \Psi^*}{\partial x_\alpha^i} \frac{\partial \Psi}{\partial x_\beta^j} - \frac{\partial \Psi}{\partial x_\alpha^i} \frac{\partial \Psi^*}{\partial x_\beta^j} \right)$$

resulting in the angular momentum the expression

$$\langle \mathbf{L}^2 \rangle = \sum_\alpha \int \Psi^* \widetilde{L}_\alpha^2 \Psi d^3 \mathbf{x} + N \frac{\hbar^2}{2} \frac{D(D-1)}{2} \quad (16)$$

where D is the spatial dimensionality of the problem. As before, the additional vacuum term is common to particle-particle correlations above fourth order and can be understood if symmetric-ordering of the associated quantum operators is assumed. In this case, the generalization of the fourth order operator displayed in eqn(15) is

$$\begin{aligned} \widetilde{\chi}_{\alpha\beta\gamma\delta ijkl}^4 \Psi &= \widetilde{x}_{\alpha i} \widetilde{x}_{\beta j} \widetilde{p}_{\gamma l} \widetilde{p}_{\delta k} \Psi + \frac{\hbar}{2i} (\delta_{\beta\delta} \delta_{j,l} x_i + \delta_{\alpha\delta} \delta_{i,l} x_j) \widetilde{p}_{\gamma k} \Psi + \\ &\frac{\hbar}{2i} (\delta_{\alpha\gamma} \delta_{j,k} x_i + \delta_{\beta\delta} \delta_{i,k} x_j) \widetilde{p}_{\delta l} \Psi - \frac{\hbar^2}{4} (\delta_{\alpha\gamma} \delta_{\beta\delta} \delta_{i,l} \delta_{j,k} + \delta_{\alpha\delta} \delta_{\beta\gamma} \delta_{i,k} \delta_{j,l}) \Psi \end{aligned}$$

leading to an expression of the squared angular momentum equal to

$$\begin{aligned} \widetilde{\mathbf{L}}_N^2 &\equiv \sum_{l,k} (\widetilde{\chi}_{kkl}^4 - \widetilde{\chi}_{klk}^4) = \sum_{\alpha,\beta,i,j} (\widetilde{x}_{\alpha i} \widetilde{x}_{\beta i} \widetilde{p}_{\alpha j} \widetilde{p}_{\beta j} - \widetilde{x}_{\alpha i} \widetilde{x}_{\beta j} \widetilde{p}_{\alpha j} \widetilde{p}_{\beta i}) - \\ &-\frac{\hbar}{i} (D-1) \sum_{\alpha,i} \widetilde{x}_{\alpha i} \widetilde{p}_{\alpha i} + N \frac{\hbar^2}{2} \frac{D(D-1)}{2} \end{aligned}$$

whose expected value coincides with the classical result shown in eqn(16).

We see that a stochastic theory exists able to explain the results of Quantum Mechanics. It is based on a phenomenological reasoning demanding that vacuum fluctuations be real. We think that a phenomenological theory has advantages over a purely axiomatic one because its foundations can experimentally be tested and eventually enlarged. An axiomatic theory can be extended owing to its results and by feeling only. Furthermore, a rational interpretation of quantum phenomena is provided by the stochastic theory enabling a comfortable epistemological basis. No new phenomena was predicted within the limits of the nonrelativistic formulation presented here besides additional terms in the correlation functions. However an explicit calculation of Bell-like correlations may present new limits to quantum measurements as well as shed some light on interpretation of entangled states within the classical framework, possible in principle using the stochastic approach.

Concluding we have shown that the limit $P \rightarrow 0$ of the Fokker-Planck equation associated to a Wiener process describing the motion of a charged particle in a electromagnetic field represents a well defined stochastic process presenting all ingredients of Quantum Mechanics. This means that, at least in the mathematical sense, there exists a stochastic model that is exactly equivalent to Quantum Mechanics.

We have also shown that fourth order (classical) particle correlations are compatible with symmetric ordering of quantum operators, opening possibilities to get, in a simple way, higher order perturbation expansions of quantum phenomena. Another advantage is the natural epistemological scenario where all of the strange axioms of Quantum Mechanics come quite naturally from a well defined phenomenological theory having a strong thermodynamic appeal. This phenomenological foundation may be, at least in principle, experimentally tested for vanishingly small vacuum power since, in this case, we predict small differences to Quantum Theory. These differences were shown in eqn(5), where there are additional terms involving the vacuum power P that are expected to give rise to correction terms in observables involving powers greater than two of the particle's conjugate momentum. Anyway, we believe the present model opens possibilities to a still better understanding of Nature inside our classic rational approach to scientific reasoning.

[1] email: josef@fisica.ufmg.br, marzojr@fisica.ufmg.br

[2] E. Nelson, Phys. Rev. **150**, 1079 (1966).

- [3] M. Planck, Ann. d. Phys. **37**, 642 (1912).
- [4] A. Einstein and L. Hopf, Ann. d. Phys. **33**, 1105 (1910).
- [5] A. Einstein and O. Stern, Ann. d. Phys. **40**, 551 (1913).
- [6] R. I. G. Hughes, *The Structure and Interpretation of Quantum Mechanics* (Harvard University Press, Cambridge, 1989).
- [7] T. H. Boyer, Phys. Rev. **182**, 1374 (1969).
- [8] C. W. Gardiner, *Handbook of Stochastic Methods* (Springer-Verlag, Berlin, 1990).
- [9] J. M. A. Figueiredo, submitted to Journal of Physics A.
- [10] G. W. Gibbons and S. W. Hawking, Phys. Rev. D **15**, 2738 (1977).
- [11] M. S. Turner, astro-ph/0108103 (2001).
- [12] N. Wheeler, Reed College Physics Department/Spring 2000 (notes)