

Pseudo-Hermiticity, weak pseudo-Hermiticity and η -orthogonality condition

B. Bagchi ^{a,*}, C. Quesne ^{b,†}

^a *Department of Applied Mathematics, University of Calcutta,
92 Acharya Prafulla Chandra Road, Calcutta 700 009, India*

^b *Physique Nucléaire Théorique et Physique Mathématique, Université Libre de Bruxelles,
Campus de la Plaine CP229, Boulevard du Triomphe, B-1050 Brussels, Belgium*

Abstract

We discuss certain features of pseudo-Hermiticity and weak pseudo-Hermiticity conditions and point out that, contrary to a recent claim, there is no inconsistency if the correct orthogonality condition is used for the class of pseudo-Hermitian, PT-symmetric Hamiltonians of the type $H_\beta = [p + i\beta\nu(x)]^2/2m + V(x)$.

PACS: 03.65.Ca

Keywords: Non-Hermitian Hamiltonians; PT Symmetry; Pseudo-Hermiticity; Supersymmetric Quantum Mechanics

Corresponding author: C. Quesne, Physique Nucléaire Théorique et Physique Mathématique, Université Libre de Bruxelles, Campus de la Plaine CP229, Boulevard du Triomphe, B-1050 Brussels, Belgium

Telephone: 32-2-6505559

Fax: 32-2-6505045

E-mail: cquesne@ulb.ac.be

*E-mail: bbagchi@cucc.ernet.in

†Directeur de recherches FNRS; E-mail: cquesne@ulb.ac.be

In recent times it has been stressed that neither Hermiticity nor PT symmetry serves as a necessary condition for a quantum Hamiltonian to preserve the reality of its bound-state eigenvalues [1, 2, 3, 4, 5]. In fact, it has been realized [5] that the existence of real eigenvalues can be associated with a non-Hermitian Hamiltonian provided it is η -pseudo-Hermitian:

$$\eta H = H^\dagger \eta, \quad (1)$$

where η is a Hermitian linear automorphism and, assuming $\hbar = 2m = 1$,

$$H = p^2 + V(x) \quad (2)$$

for $V(x) \in \mathbb{C}$ and $p = -i\partial_x$. Then, in such a case, the spectrum of a diagonalizable H is real if there exists a linear invertible operator O such that $\eta = (OO^\dagger)^{-1}$. Moreover, one can relax H to be only weak pseudo-Hermitian [6] by not restricting η to be Hermitian.

The purpose of this Letter is to establish the following results:

- (i) The twin concepts of pseudo-Hermiticity and weak pseudo-Hermiticity are complementary to one another.
- (ii) For a first-order differential realization, η may be anti-Hermitian but for the second-order case, η is necessarily Hermitian. For both cases, we make connections to the same PT-symmetric Scarf II Hamiltonian (having normalizable eigenfunctions) to show that the choice of η is not unique in ascertaining the character of the Hamiltonian.
- (iii) For the class of η -pseudo-Hermitian, PT-symmetric Hamiltonians described by [7, 8] ¹

$$H_\beta = [p + i\beta\nu(x)]^2 + V(x), \quad \beta \in \mathbb{R}, \quad (3)$$

where the odd function $\nu(x) \in \mathbb{R}$, $V(x)$ is PT-symmetric, and

$$\eta = \exp \left[-2\beta \int^x \nu(y) dy \right], \quad (4)$$

our earlier derivation [9] of the generalized continuity equation for Hamiltonians of the form (2) [with $V(x)$ PT-symmetric] can be extended to H_β as well. The resulting η -orthogonality condition needs to be implemented judiciously.

¹Note that in Ref. [8], it is assumed that $\hbar = m = 1$ instead of $\hbar = 2m = 1$.

We begin by addressing to the point (i) above. Consider some non-Hermitian η that is subject to the condition (1). Taking Hermitian conjugate, we obtain, on adding and subtracting, the following combinations

$$\eta_+ H = H^\dagger \eta_+, \quad \eta_- H = H^\dagger \eta_-, \quad (5)$$

where $\eta_\pm = \eta \pm \eta^\dagger$. While the first one of (5) corresponds to strict pseudo-Hermiticity, the second one points to weak pseudo-Hermiticity with a new anti-Hermitian operator η_- . Note that η_+ is Hermitian. It is thus clear that weak pseudo-Hermiticity is not more general than pseudo-Hermiticity but works complementary to it.

We now turn to (ii). Decomposing $V(x)$ and η as

$$\begin{aligned} V(x) &= V_R(x) + iV_I(x), \\ \eta &= \frac{d}{dx} + f(x) + ig(x), \end{aligned} \quad (6)$$

where $V_R, V_I, f, g \in \mathbb{R}$, we get, on inserting (6) into (1), the relations

$$\begin{aligned} V_I &= i(f' + ig'), \\ V_R' + iV_I' &= -(f'' + ig'') - 2iV_I(f + ig). \end{aligned} \quad (7)$$

In (7) the primes denote the order of differentiations with respect to the variable x . We are then led to the conditions

$$V_R' = -2gg', \quad cg' = 0, \quad c \in \mathbb{R}, \quad (8)$$

which imply the existence of two solutions corresponding to $c = 0$ and $g' = 0$, respectively. In the following we concentrate on the case $c = 0$ because $g' = 0$ yields a trivial result that corresponds to a real constant potential with no normalizable eigenfunction.

For the choice $c = 0$, it turns out that

$$f = 0, \quad V_R = -g^2 + k, \quad V_I = -g', \quad (9)$$

where $k \in \mathbb{R}$. In consequence, we have the results

$$\begin{aligned} V(x) &= -g^2(x) + k - ig'(x), \\ \eta &= \frac{d}{dx} + ig(x). \end{aligned} \quad (10)$$

The above form of $V(x)$ shows that, in the framework of supersymmetric quantum mechanics, we can associate to it an imaginary superpotential $W(x) = ig(x)$, its partner being the complex conjugate potential. We also observe that, for even g functions, the representation of η makes it anti-Hermitian in character. Let us consider the following specific example for $g = d \operatorname{sech} x$, $d \in \mathbb{R}$. We get from (10)

$$\begin{aligned} V(x) &= -d^2 \operatorname{sech}^2 x + k + id \operatorname{sech} x \tanh x, \\ \eta &= \frac{d}{dx} + id \operatorname{sech} x. \end{aligned} \quad (11)$$

It is obvious that $V(x)$ is a particular case of the generalized PT-symmetric Scarf II potential investigated previously by us [10] in connection with the complex algebra $\mathfrak{sl}(2, \mathbb{C})$. A comparison with the results obtained there shows that, in the present case, we have a single series of real eigenvalues with normalizable eigenfunctions provided $d > \frac{1}{2}$. The corresponding Hamiltonian is both P-pseudo-Hermitian and η -weak-pseudo-Hermitian with η given by (11). Our example confirms the assertion [11] that, for a given non-Hermitian Hamiltonian, there could be infinitely many η satisfying the weak-pseudo-Hermiticity or the pseudo-Hermiticity condition.

We next attend to a second-order differential representation of η :

$$\eta = \frac{d^2}{dx^2} - 2p(x) \frac{d}{dx} + b(x), \quad (12)$$

where $p, b \in \mathbb{C}$. Substituting (12) into the condition (1), we obtain the constraints

$$\begin{aligned} b &= -p' + p^2 - \frac{p''}{2p} + \left(\frac{p'}{2p}\right)^2 + \frac{\gamma}{4p^2}, \\ V &= 2p' + p^2 + \frac{p''}{2p} - \left(\frac{p'}{2p}\right)^2 - \frac{\gamma}{4p^2} - \delta, \\ V^* &= -2p' + p^2 + \frac{p''}{2p} - \left(\frac{p'}{2p}\right)^2 - \frac{\gamma}{4p^2} - \delta, \end{aligned} \quad (13)$$

where $\gamma, \delta \in \mathbb{R}$. From the last two relations in (13), it is clear that $p(x)$ must be pure imaginary,

$$p(x) = ia(x), \quad (14)$$

where $a(x) \in \mathbb{R}$. As such $V(x)$ and η acquire the forms

$$\begin{aligned} V(x) &= 2ia' - a^2 + \frac{a''}{2a} - \left(\frac{a'}{2a}\right)^2 + \frac{\gamma}{4a^2} - \delta, \\ \eta &= \frac{d^2}{dx^2} - 2ia(x)\frac{d}{dx} + b(x), \end{aligned} \quad (15)$$

with $b(x) = -V(x) + ia' - 2a^2 - \delta$. In (15), η can be easily recognized to be a Hermitian operator since it can be written in the form $\eta = -\tilde{O}^\dagger \tilde{O}$, where $\tilde{O} = \frac{d}{dx} + r - ia$, $\tilde{O}^\dagger = -\frac{d}{dx} + r + ia$, and $r^2 - r' = \frac{a''}{2a} - \left(\frac{a'}{2a}\right)^2 + \frac{\gamma}{4a^2}$. In Ref. [12], such a decomposition of η was assumed, a priori, to arrive at some non-Hermitian Hamiltonians with real spectra.

Let us, however, confine ourselves to the following choice

$$a(x) = -\frac{1}{2}B(2A + 1) \operatorname{sech} x, \quad \gamma = 0, \quad \delta = \frac{1}{4}, \quad (16)$$

where $A + \frac{1}{2} > 0$, $B > 0$, and $A - B + \frac{1}{2}$ is not an integer. We are again led to the PT-symmetric Scarf II potential having a more general form than obtained with the first-order differential realization of η :

$$V(x) = -V_1 \operatorname{sech}^2 x - iV_2 \operatorname{sech} x \tanh x, \quad (17)$$

where $V_1 = \frac{1}{4}[B^2(2A + 1)^2 + 3] > 0$ and $V_2 = -B(2A + 1) \neq 0$. According to Refs. [13, 14], the condition for real eigenvalues for the Hamiltonian corresponding to (17) is $|V_2| \leq V_1 + \frac{1}{4}$. Here it amounts to $[B(2A + 1) - 2]^2 \geq 0$, which is always met.

Of particular interest is the special case $B = 1$:

$$V(x) = -(A^2 + A + 1) \operatorname{sech}^2 x + i(2A + 1) \operatorname{sech} x \tanh x. \quad (18)$$

On setting $A + \frac{1}{2} = -\lambda$ ($\lambda < 0$), Eq. (18) can be seen to reduce to the potential $V^{(1)} - \frac{1}{4}$ of Ref. [15] for $\mu = 1$. The associated energy levels of (18) are [10]: $E_n^{(-\lambda)} = -(\lambda + n + \frac{1}{2})^2$ and coincide with $E_n^{(2)} - \frac{1}{4}$ of [15]. Note that there is, in general, a doubling of energy levels in transiting from the real to the PT-symmetric Scarf II potential. In fact, the second algebra of $\mathfrak{sl}(2, \mathbb{C})$ leads to an additional energy level $E_0^{(1)} = -\frac{1}{4}$ that is consistent with the zero-energy state of [15].

Finally, we take up a general derivation of the continuity equation for the class of Hamiltonians H_β given by (3). The associated Schrödinger equation reads

$$i\frac{\partial\psi(x,t)}{\partial t} = -\left(-\frac{\partial}{\partial x} + \beta\nu(x)\right)^2 \psi(x,t) + V(x)\psi(x,t). \quad (19)$$

From this it follows that the function $\psi^*(-x,t)$ satisfies

$$-i\frac{\partial\psi^*(-x,t)}{\partial t} = -\left(-\frac{\partial}{\partial x} + \beta\nu(x)\right)^2 \psi^*(-x,t) + V(x)\psi^*(-x,t). \quad (20)$$

On considering Eq. (19) for some solution $\psi_1(x,t)$ and Eq. (20) for some other solution $\psi_2(x,t)$ and then multiplying (19) and (20) by $\exp[-2\beta\int^x\nu(y)dy]\psi_2^*(-x,t)$ and $\exp[-2\beta\int^x\nu(y)dy]\psi_1(x,t)$, respectively, we obtain, on subtracting, a natural generalization of the continuity equation for PT-symmetric quantum mechanics to its η -pseudo-Hermitian extension, namely

$$\frac{\partial P_\eta(x,t)}{\partial t} + \frac{\partial J_\eta(x,t)}{\partial x} = 0, \quad (21)$$

where

$$\begin{aligned} P_\eta(x,t) &= \eta\psi_2^*(-x,t)\psi_1(x,t), \\ J_\eta(x,t) &= \frac{\eta}{i}\left[\psi_2^*(-x,t)\frac{\partial\psi_1(x,t)}{\partial x} - \psi_1(x,t)\frac{\partial\psi_2^*(-x,t)}{\partial x}\right], \end{aligned} \quad (22)$$

and η is defined in (4). If $\psi_1(x,t) \rightarrow 0$ and $\psi_2(x,t) \rightarrow 0$ as $x \rightarrow \pm\infty$, as is normally expected for bound-state wave functions, then integration of (21) over the entire real line gives the conservation law

$$\frac{\partial}{\partial t} \int_{-\infty}^{\infty} dx \eta \psi_2^*(-x,t)\psi_1(x,t) = 0. \quad (23)$$

In the case of energy eigenfunctions

$$\psi_1(x,t) = u_1(x)e^{-iE_1t}, \quad \psi_2(x,t) = u_2(x)e^{-iE_2t}, \quad (24)$$

corresponding to the eigenvalues E_1 and E_2 respectively, Eq. (23) reduces to

$$(E_1 - E_2^*) \int_{-\infty}^{\infty} dx \eta u_2^*(-x)u_1(x) = 0. \quad (25)$$

Equation (25) represents the η -orthogonality condition [5]. Obviously it transforms to the PT-orthogonality [9, 16]

$$(E_1 - E_2^*) \int_{-\infty}^{\infty} dx u_2^*(-x) u_1(x) = 0 \quad (26)$$

for $\nu(x) = 0$. Indeed Eq. (25) can be derived from (26) by effecting a gauge transformation on the wave functions u of H in a manner $u \rightarrow \exp[-\int^x \beta\nu(y)dy]u$, $V(x)$ being PT-symmetric. As such H_β may be looked upon as a gauge-transformed version of H . However, it needs to be emphasized that care should be taken to correctly implement the normalization conditions deriving from (25) and (26) and which are appropriate to the Hamiltonians H_β and H , respectively. Thus although PT-symmetric, the form of the η -pseudo-Hermitian Hamiltonian H_β at once suggests that the normalization condition related to (25) is to be used rather than that connected with (26), a point overlooked in Ref. [8].

In summary, we have shown that η -pseudo-Hermiticity and weak pseudo-Hermiticity are essentially complementary concepts. We have provided an explicit example of PT-symmetric Scarf II model to demonstrate that η does not necessarily have a unique representation to determine the character of the associated non-Hermitian Hamiltonian. We have also pointed out the correct use of the η -orthogonality condition when dealing with a pseudo-Hermitian gauge-transformed Hamiltonian.

References

- [1] D. Bessis, unpublished (1992).
- [2] C.M. Bender, S. Boettcher, Phys. Rev. Lett. 80 (1998) 5243.
- [3] C.M. Bender, S. Boettcher, P.N. Meisinger, J. Math. Phys. 40 (1999) 2201.
- [4] P. Dorey, C. Dunning, R. Tateo, J. Phys. A 34 (2001) 5679.
- [5] A. Mostafazadeh, J. Math. Phys. 43 (2002) 205; 43 (2002) 2814.
- [6] L. Solombrino, Weak pseudo-Hermiticity and antilinear commutant, Preprint quant-ph/0203101.
- [7] A. Mostafazadeh, On the pseudo-Hermiticity of general PT-symmetric standard Hamiltonians in one dimension, Preprint math-ph/0204013.
- [8] Z. Ahmed, Phys. Lett. A 294 (2002) 287.
- [9] B. Bagchi, C. Quesne, M. Znojil, Mod. Phys. Lett. A 16 (2001) 2047.
- [10] B. Bagchi, C. Quesne, Phys. Lett. A 273 (2000) 285.
- [11] A. Mostafazadeh, Pseudo-supersymmetric quantum mechanics and isospectral pseudo-Hermitian Hamiltonians, Preprint math-ph/0203041.
- [12] T.V. Fityo, A new class of non-Hermitian Hamiltonians with real spectra, Preprint quant-ph/0204029.
- [13] Z. Ahmed, Phys. Lett. A 282 (2001) 343; 287 (2001) 295.
- [14] B. Bagchi, C. Quesne, Non-Hermitian Hamiltonians with real and complex eigenvalues in a Lie-algebraic framework, Preprint math-ph/0205002, to be published in Phys. Lett. A.
- [15] B. Bagchi, R. Roychoudhury, J. Phys. A 33 (2000) L1.
- [16] G.S. Japaridze, J. Phys. A 35 (2002) 1709.