## What's Wrong with these Observables?

Asher Peres

Department of Physics, Technion—Israel Institute of Technology, 32000 Haifa, Israel

An imprecise measurement of a dynamical variable (such as a spin component) does not, in general, give the value of another dynamical variable (such as a spin component along a slightly different direction). The result of the measurement cannot be interpreted as the value of any observable that has a classical analogue.

# 1. MEMORABILIA

I was never compelled to learn solid state physics, and my first encounter with David Mermin was his paper [1] "Is the moon there when nobody looks? Reality and the quantum theory". That paper gave a wonderful proof of Bell's theorem. I wrote a comment [2] saying "Mermin gives a remarkably simple illustration of Bell's theorem, but leaves the impression that something mysterious is implied. The situation is much simpler: the pair of photons is a single, nonlocal, indivisible entity . . . It is only because we force upon the photon pair the description of two separate particles that we get the paradox of Einstein, Podolsky and Rosen." David answered to me, privately, "one person's mystery is another person's explanation."

The following year, we met in a conference that Danny Greenberger had organized in the World Trade Center in New York City [3] and it was friendship at first sight. I gave a talk "When is a quantum measurement?" after which Abner Shimony came to me and said, with a big friendly smile "Asher, you understand nothing! You speak just like Niels Bohr!" One person's insult is another person's compliment.

Over the years, David and I collaborated many times, but I never had the honor of being his co-author. Many of our discussions revolved around the Kochen-Specker (KS) theorem [4]. In 1990, David relayed to me information that Conway and Kochen had found a new proof of that theorem using only 33 rays, instead of 117 in the original proof. The only clue I had was that they used a cubic lattice for constructing their rays. This led me to find a much simpler proof, also with 33 rays, but this one using an irrational lattice. I wrote to Simon Kochen, asking him to inform me when he publishes his proof, so that I could publish mine immediately afterwards. Kochen answered that my proof was indeed very simple, but meanwhile he had found one with only 31 rays. I was both happy and disappointed. However, soon afterwards, I found a proof with only 24 rays, in four dimensions. I wrote to Kochen that 24 < 31 and that I would publish my results, while mentioning the existence of their proof with 31 rays [5]. Conway and Kochen then never bothered to publish the latter.

In my paper [5] I also stated that 24 was the minimal number number in four dimensions. David was not convinced by my argument. We exchanged several letters on this issue but could not resolve it. Meanwhile I was curious to see the 31 rays in three dimensions. I guessed they were a subset of the same cubic lattice and I did the search with a computer program. The algorithm appears in my book [6], where there is also an exercise: write a computer program for the four-dimensional case, and check that 24 rays are the minimum number. I never bothered to solve that boring exercise, but two students took up the challenge and found that it was possible to remove any one of the 24 rays, and still have a KS set. Michael Kernaghan, in Canada, found a KS set with 20 rays [7] and then Adán Cabello, together with José Manuel Estebaranz and Guillermo García Alcaine in Madrid, found a set of 18 rays [8]. They still hold the world record (probably for ever).

Soon after I found my 33 + 24 rays, there was the Gulf war and Saddam Hussein sent numerous scud missiles on Israeli targets. Each time there was a raid alert, we had to go into an air-tight room and don gas masks (fortunately they were never needed). Each alert lasted about half an hour, until the scud debris were examined by civil defense experts. To help time pass, I read Boojums [9] through the goggles of the mask. This was sometimes difficult, because the goggles were fogged by my breath, but this was always enjoyable and instructive.

It is a pleasure to dedicate this article to David Mermin, for his birthday and many more birthdays.

### 2. WHAT CAN BE OBSERVED?

Standard texbooks on quantum mechanics tell you that observable quantities are represented by Hermitian operators, their possible values are the eigenvalues of these operators, and that the probability of detecting eigenvalue  $\lambda_n$ , corresponding to eigenvector  $u_n$ , is  $|\langle u_n|\psi\rangle|^2$ , where  $\psi$  is the (pure) state of the quantum system that is observed. With a bit more sophistication to include mixed states, the probability can be written in a general way  $\langle u_n|\rho|u_n\rangle$ . Really bad books also claim that the state of the physical system after the measurement collapses into the corresponding  $u_n$ . This is sheer nonsense. (Finding appropriate references is left as an exercise for the reader.)

The simple and obvious truth is that quantum phenomena do not occur in a Hilbert space. They occur

in a laboratory. If you visit a real laboratory, you will never find there Hermitian operators. All you can see are emitters (lasers, ion guns, synchrotrons and the like) and detectors. The experimenter controls the emission process and observes detection events. The theorist's problem is to predict the probability of response of this or that detector, for a given emission procedure. Quantum mechanics tells us that whatever comes from the emitter is represented by a state  $\rho$  (a positive operator, usually normalized to 1). Detectors are represented by positive operators  $E_{\mu}$ , where  $\mu$  is an arbitrary label whose sole role is to identify the detector. The probability that detector  $\mu$  be excited is  $\operatorname{tr}(\rho E_{\mu})$ . A complete set of  $E_{\mu}$ , including the possibility of no detection, sums up to the unit matrix and is called a positive operator valued measure (POVM) [6].

The various  $E_{\mu}$  do not in general commute, and therefore a detection event does not correspond to what is commonly called the "measurement of an observable." Still, the activation of a particular detector is a macroscopic, objective phenomenon. There is no uncertainty as to which detector actually clicked.

### 3. IMPRECISE MEASUREMENTS

There has recently been considerable controversy about the possibility of testing the physical implications of the KS theorem or even "nullifying" the latter [10]. The "observables" that are usually considered are spin components of a spin-1 particle, which is the paradigm of the KS theorem. Opinions are varied [11] but more often than not assume that the result of an imprecise measurement of a spin component  $\mathbf{n} \cdot \mathbf{J}$  is the value of another spin component, along some direction  $\mathbf{n}'$  which is close to the correct  $\mathbf{n}$ . As shown below, this assumption is generally unfounded (except in the trivial case of spin- $\frac{1}{2}$  particles).

When I first learnt of this "nullification," my gut feeling was that the claim was not even wrong. However, finding a decisive argument that would convince the authors [10] proved as elusive as giving an experimental proof of a violation of Bell's inequality. It is always possible to find loopholes. All I can do is to challenge them to find a loophole in the following argument.

Whatever the causes of imprecision are, if we don't measure exactly  $\mathbf{n} \cdot \mathbf{J}$ , we have to give a physical interpretation to the result of our experimental procedure. Do we measure some other nearby operator,

$$A = \mathbf{n} \cdot \mathbf{J} + \sum_{m,n} a_{mn} J_m J_n / \hbar,$$

where the unknown dimensionless coefficients  $a_{mn} = a_{nm}^*$  are small? If the array  $a_{mn}$  is antisymmetric, and therefore imaginary, we have  $A = \mathbf{m} \cdot \mathbf{J}$ , where  $\mathbf{m} \approx \mathbf{n}$  (but  $\mathbf{m}$  need not be a unit vector). However, in general, the operator A defined above also contains bilinear terms which

cannot be reduced to another component of J. These terms have no classical analogue and no name in our vocabulary. Quantum mechanics has a much richer choice of dynamical variables than classical mechanics.

Actually, the result of the measurement procedure may not even be the value of any dynamical variable that looks like A. Traditional concepts such as "measuring Hermitian operators," that were borrowed or adapted from classical physics, are not appropriate in the quantum world. In the latter, as explained above, we have emitters and detectors, and calculations are performed by means of POVMs.

Any lack of precision in the experimental setup simply means that the positive operator  $E_{\mu}$  which is actually implemented in the laboratory only approximates the original  $E_{\mu}$  that the experimenter intended to use. That  $E_{\mu}$  may have been associated with a classical interpretation such as the value of a component of angular momentum. The actual  $E_{\mu}$  may have no classical interpretation at all. In particular, there is no reason to expect it to correspond to a component of  ${\bf J}$  along a slightly different direction.

What does this imply for attempts to test experimentally the KS theorem? That theorem is only a statement about Euclidean geometry. Any purported experimental test should be analyzed as explained above: particles are emitted according to a specified procedure, and quantum theory is used to predict the probability of excitation of various detectors. After we take into account known imperfections of the emission and detection processes, any discrepancy would imply either a poor understanding of the laboratory equipment, or a failure of quantum theory. The KS theorem itself is not involved. Its only role is to restrict our freedom of concocting realistic noncontextual theories that would replace quantum mechanics.

### Acknowledgment

This work was supported by the Gerard Swope Fund and the Fund for Encouragement of Research.

- [1] N. D. Mermin, *Physics Today* **38** (4) 38 (1985).
- [2] S. J. Feingold and A. Peres, Physics Today 38 (11) 15 (1985).
- [3] D. M. Greenberger, editor, "New Techniques and Ideas in Quantum Measurement Theory" Ann. New York Acad. Sc. 480 (1986).
- [4] S. Kochen and E. Specker, J. Math. Mech. 17, 59 (1967).
- [5] A. Peres, J. Phys. A 24, L175 (1991).
- [6] A. Peres, Quantum Theory: Concepts and Methods (Kluwer, Dordrecht, 1993).

- [7] M. Kernaghan, J. Phys. A  $\bf 27$ , L829 (1994).
- [8] A. Cabello, J. M. Estebaranz, and G. García Alcaine, Phys. Lett. A 212, 183 (1996).
- [9] N. D. Mermin, Boojums All the Way Through: Communicating Science in a Prosaic Age (Cambridge Univ. Press, Cambridge, 1990).
- [10] D. A. Meyer, Phys. Rev. Lett. 83, 3751 (1999); A. Kent,
   ibid. 83, 3755 (1999); R. K. Clifton and A. Kent, Proc.
   Roy. Soc. Lond. A 456, 2101 (2000).
- [11] H. Havlicek, G. Krenn, J. Summhammer, and K. Svozil, J. Phys. A 34, 3071 (2001); C. Simon, Č. Brukner, and A. Zeilinger, Phys. Rev. Lett. 86, 4427 (2001); T. Breuer, ibid. 88, 24042 (2002); D. M. Appleby, Phys. Rev. A 65, 022105 (2002); A. Cabello, ibid. 65, 052101 (2002); J.-Å. Larsson, Europhys. Lett. 58, 799 (2002).
  See also numerous articles in the quant-ph electronic archive.