## Comments on Struyve and Baere's paper on experiments to distinguish Bohmian mechanics from quantum mechanics

## Partha Ghose

S. N. Bose National Centre for Basic Sciences, Block JD, Sector III, Salt Lake, Kolkata 700 098

It is shown in detail why the arguments put forward by Struyve and Baere (quant-ph/0108038) against my conclusions are incorrect.

There are several strands to the arguments I used that seem to have been overlooked by the authors. Let me take them up one by one. But right in the beginning I would like to emphasize that two aspects of the discussion must be clearly separated from each other, the question of (a) the theoretical incompatibility between dBB and quantum mechanics, a question that can be settled with the help of *gedanken* experiments alone, and (b) the feasibility of real experiments to test this incompatibility.

Quantum Equilibrium Hypothesis

Without loss of generality let us restrict the discussion to the two-particle case. In dBB particle trajectories  $\mathbf{x}_1(t)$  and  $\mathbf{x}_2(t)$  are introduced through the *guidance condition* 

$$\mathbf{p}_i = \nabla_{\mathbf{x}_i} S(\mathbf{x}_1, \mathbf{x}_2, t) \tag{1}$$

where  $S(\mathbf{x}_1, \mathbf{x}_2, t)$  is the phase of the wave function  $\psi(\mathbf{x}_1, \mathbf{x}_2, t)$  in quantum mechanics. (I use the notation  $\mathbf{x} = (x, y, z)$ .) Having introduced the trajectories, one then postulates that

$$\rho(\mathbf{x}_1(t_0), \mathbf{x}_2(t_0)) = |\psi(\mathbf{x}_1, \mathbf{x}_2, t_0)|^2$$
(2)

at some time time  $t_0$ . Then the continuity equation guarantees that this relation is valid at all times t:

$$\rho(\mathbf{x}_1(t), \mathbf{x}_2(t)) = |\psi(\mathbf{x}_1, \mathbf{x}_2, t)|^2$$
(3)

This is the only hypothesis in dBB that may be called the quantum equilibrium hypothesis (QEH). Writing it in the form

$$\rho(\mathbf{x}_1, \mathbf{x}_2, t) = |\psi(\mathbf{x}_1, \mathbf{x}_2, t)|^2 \tag{4}$$

is misleading because this relation is a mathematical identity without any physical content. The guidance condition being the only input in dBB that is additional to quantum mechanics, it goes without saying that any analysis that does not take this condition into account and merely uses the identity (4) and the continuity equation will trivially reproduce the same results as quantum mechanics in every case.

Note that  $\mathbf{x}_1$  and  $\mathbf{x}_2$  are variables that cover the entire support of the wave function  $\psi$ , and hence the trajectories  $\mathbf{x}_1(t)$  and  $\mathbf{x}_2(t)$ ) must also cover the same support. Therefore, to realize the distribution  $\rho(\mathbf{x}_1(t), \mathbf{x}_2(t))$  one must have an infinitely large number of particles of types 1 and 2 distributed in the prescribed manner  $\rho$  at time t. One then has a full ensemble.

If we take any observable  $\mathcal{O}$ , its (space) average in dBB is then given by

$$\bar{O} = \int_{t} \mathcal{O}\rho(\mathbf{x}_{1}(t), \mathbf{x}_{2}(t)) d\mathbf{x}_{1}(t) d\mathbf{x}_{2}(t)$$
(5)

This tells us that one has to integrate over all possible trajectories  $\mathbf{x}_1(t)$  and  $\mathbf{x}_2(t)$  at a fixed time t in order to compute the average in dBB. This is the ensemble average.

Now consider a theoretically admissible situation in which one has particle 1 at a definite position  $\mathbf{X}_1(t)$  and particle 2 at  $\mathbf{X}_2(t)$  at some time t and everywhere else the wave function (in dBB) is empty. Since, by hypothesis, empty parts of the wave function do not fire detectors, only two trajectories out of all the possible ones contribute to detections in this case. If one repeats the situation over and over again with identical copies of the system at different times  $(t_1, ..., t_N)$ , the particles will be in different positions  $\mathbf{X}_1(t_i)$  and  $\mathbf{X}_2(t_i)$  each time with probabilities determined by the distribution function  $\rho(\mathbf{X}_1(t_i), \mathbf{X}_2(t_i))$ . If one collects all the results together and computes the time average of the joint detection probability, one would obtain (for two detectors  $D_1$  and  $D_2$  placed with their x and z coordinates the same and fixed)

$$P_{12}^* = \lim_{N \to \infty} \frac{1}{N} \sum_{t_1}^{t_N} \int_{D_1, D_2, t_i} \rho(\mathbf{x}_1(t_i), \mathbf{x}_2(t_i)) \, \delta(\mathbf{x}_1(t_i) - \mathbf{X}_1(t_i)) \, \delta(\mathbf{x}_2(t_i) - \mathbf{X}_2(t_i)) \, dy_1(t_i) \, dy_2(t_i)$$
(6)

$$= \lim_{N \to \infty} \frac{1}{N} \sum_{t_1}^{t_N} \rho(\mathbf{X}_1(t_i), \mathbf{X}_2(t_i)) = \bar{P}_{12}$$
 (7)

where  $\bar{P}_{12}$  is the space average. This result is, however, true only provided the guidance condition does not imply a constraint like  $Y_1(t_i) + Y_2(t_i) = 0$  on the trajectories for every  $t_i$ .

If there is a constraint like  $\delta(Y_1(t_i) + Y_2(t_i))$  on the trajectories due to some symmetry, it will make the system non-ergodic. One would then obtain

$$P_{12}^* = \lim_{N \to \infty} \frac{1}{N} \sum_{t_1}^{t_N} \frac{1}{\delta(0)} \int_{D_1, D_2, t_i} \rho(\mathbf{x}_1(t_i), \mathbf{x}_2(t_i)) \, \delta(\mathbf{x}_1(t_i) - \mathbf{X}_1(t_i)) \, \delta(\mathbf{x}_2(t_i) - \mathbf{X}_2(t_i))$$
(8)

$$. \delta(Y_1(t_i) + Y_2(t_i)) dy_1(t_i) dy_2(t_i) \neq \bar{P}_{12}$$
(9)

Nevertheless, if one computes the time average of an observable other than the joint detection probability, one would still obtain

$$O_{12}^* = \lim_{N \to \infty} \frac{1}{N} \sum_{t_1}^{t_N} \frac{1}{\delta(0)} \int_{t_i} \mathcal{O}\rho(\mathbf{x}_1(t_i), \mathbf{x}_2(t_i)) \, \delta(\mathbf{x}_1(t_i) - \mathbf{X}_1(t_i)) \, \delta(\mathbf{x}_2(t_i) - \mathbf{X}_2(t_i))$$
(10)

$$\delta(Y_1(t_i) + Y_2(t_i)) \, dy_1(t_i) \, dy_2(t_i) = \bar{O}$$
(11)

because the constraint, being a consequence of the guidance condition, is necessarily consistent with the distribution function  $\rho$  by hypothesis, and most importantly the restrictions on the integration variables coming from the supports of the detectors is removed.

All this shows that QEH is applicable to the full ensemble but not to individual processes that make up this ensemble. There could be information regarding individual processes determined by Bohmian dynamics (through the guidance condition) that are hidden in the full ensemble. The symmetric trajectories in a double-slit experiment with single particles is a clear example of how the information that the trajectories are symmetrical about the line of symmetry and do not cross, is masked in the distribution function  $\rho(\mathbf{x}(t))$ .

With this background let us now consider a gedanken experiment in which two particles are incident at a time on a screen S with two point-like slits (the points being assumed to be limits of spheres). The source of the particles is assumed to be such that it produces entangled pairs of particles, one pair at a time, and it is so placed that out of all the pairs it emits, a significant number is produced such that one particle passes through each slit. The rest of the pairs are blocked by the screen S. On the far side of the slit two spherical waves will emerge from the two slits. I have shown by an explicit analytical calculation (quant-ph/0103126) that the trajectories of the particles will be symmetrical about the line of symmetry between the slits. The system is therefore non-ergodic. Let two detectors be placed at a fixed distance X along the x-axis asymmetrically about the line of symmetry x = 0. It follows from eqn. (8) that  $P_{12}^* = 0$  in this case. Since  $\bar{P}_{12} \neq 0$ , this establishes the incompatibility.

If one widens the slits somewhat and uses Gaussian profiles, then, as Struyve and Baere have shown,

$$\sigma(t) \equiv Y_1(t) + Y_2(t) = \sigma(0)\sqrt{1 + (\hbar/2m\sigma_0^2)^2 t^2}$$
(12)

This spreading can be made as small as one desires by requiring that

$$(\hbar/2m\sigma_0^2)^2 t^2 << 1 \tag{13}$$

There is nothing in the theory that precludes such a condition. Since this spreading is calculable in any case, it can be taken care of in the design of the experiment.

Ergodicity

First of all, let me state that the definition (and proof) of ergodicity of quantum mechanics that I gave in quantph/0103126 is not due to me but can be found in standard text books such as that by Toda et al [1]. Furthermore, the cross terms have not been ignored in the proof, as the authors seem to imply, but they can actually be shown to vanish in the limit of large time. In fact, this is the most crucial part of the proof supplied by mathematicians in the most general case. Having said that, let me also state that the proof of ergodicity of dBB given by Struyve and Baere is flawed by the trivial use of QEH in the form (4) as explained above.

Finally, let me comment on their equation (26),

$$\int_{M} f^* d\mu = \bar{f} \tag{14}$$

What the authors have not stated is the condition under which this theorem is valid, namely, that the flow be such that there is no invariant subspace of the full phase space M, i.e.,  $\{\phi_t(E) = E \subseteq M \Rightarrow E \equiv M \text{ or } \emptyset\}$ . In other words, this theorem is valid for ergodic systems only. Therefore there is no inconsistency in my arguments.

## Double pendulum

In this case it is obvious that the Bohmian motion of the individual pendulums remains periodic (as in the classical case which is non-ergodic if the two eigenfrequencies are commensurate), and therefore the motion does not cover the entire phase space torus That is sufficient to prove non-ergodicity for commensurate frequencies. Since there is a general theorem that states that there must be at least one observable in a non-ergodic system whose space and time averages are different, the incompatibility with quantum mechanics is established.

[1] M. Toda, R. Kubo and N. Saitô, Statistical Physics I, second edition, Springer, 1995, p.222.