

$so(2, 1)$ scattering via Euclidean Connection for the Natanzon Hypergeometric Potentials

Hermann Albrecht¹ and Sebastián Salamó^{2, 3}

Universidad Simón Bolívar, Departamento de Física,
A. P. 89000, Valle de Sartenejas, Miranda 1080-A, Venezuela.

Abstract

The $so(2, 1)$ Euclidean Connection [1] formalism is used to calculate the S matrix for the Hypergeometric Natanzon Potentials (HNP) [2].

1 Introduction

Algebraic scattering has been an active research field for many years. Pioneer works in the late 1960's and 1970's by Zwanziger [3] and Barut, et al. [4] dealing with the Coulomb system founded the basic ideas of this area. More recently, Alhassid et al. [1] developed an algebraic formalism, known as the Euclidean connection, to calculate the S matrix for systems whose Hamiltonian may be written as a function of the Casimir operator of an $so(2, 1)$ algebra.

In this work we study the application of the Euclidean connection to obtain in a straightforward manner the S matrix for the HNP , by extending the treatment developed in [5].

We start with a brief review of [5] and then introduce the basic ideas of the Euclidean connection for $so(2, 1)$ [1]. Then we study its application to the scattering sector of the HNP , obtaining the S matrix for this family of potentials.

2 Bound states

As is well known, the Hypergeometric Natanzon Potentials (HNP) [2] are given by:

$$V(z) = \frac{f z^2 - (h_0 - h_1 + f) z + h_0 + 1}{R(z)} + \left(a + \frac{a + (c_1 - c_0)(2z - 1)}{z(z - 1)} - \frac{5}{4} \frac{\Delta}{R(z)} \right) \frac{z^2 (1 - z)^2}{R(z)^2} \quad (1)$$

where

$$R(z) = a z^2 + \tau z + c_0 \quad \Delta = \tau^2 - 4 a c_0 \quad \tau = c_1 - c_0 - a \quad (2)$$

¹Email: halbrech@fis.usb.ve

²Email: ssalamo@usb.ve

³Accepted for publication in Rev. Mex. Física, 2002.

where f , h_0 , h_1 , a , c_0 , c_1 are the *Natanzon Parameters* and $z(r)$ is a real arbitrary function that satisfies

$$\frac{dz}{dr} \equiv z' = \frac{2z(z-1)}{\sqrt{R(z)}} \quad z(r) \in [0, 1] \quad (3)$$

and the transformation $r(z)$ is assumed to take $r \rightarrow \infty$ to $z = 1$. In [5] the bound state sector of the *HNP* was studied by means of an $so(2, 1)$ algebra, obtaining an algebraic description of it. The realization used is

$$\begin{aligned} J_0 &= -i\partial_\phi \\ J_\pm &= e^{\pm i\phi} \left\{ \pm \frac{\sqrt{z}(z-1)}{z'} \partial_r - \frac{1}{2} \frac{i(z+1)}{\sqrt{z}} \partial_\phi \right\} \\ &\quad \pm e^{\pm i\phi} \left\{ \frac{1}{2}(z-1) \left[\frac{1 \mp p_\nu}{\sqrt{z}} - \frac{z''\sqrt{z}}{z'^2} \right] \right\} \\ Q &= \frac{z(z-1)^2}{z'^2} \partial_r^2 + \frac{1}{4} \frac{(z-1)^2}{4z} \partial_\phi^2 + \frac{1}{2} \frac{i p_\nu (z^2-1)}{z} \partial_\phi \\ &\quad + \frac{(z-1)^2}{4} \frac{[z^2(2z'''z' - 3z'') - z'^4(p_\nu^2 - 1)]}{z z'^4} \end{aligned} \quad (4)$$

where p is a parameter, which allows the use of the $so(2, 1)$ algebra to completely describe the *HNP*. The algebraic treatment makes use of

$$(Q - q)\Psi(r, \phi) = \mathcal{G}(r)(E - H)\Psi(r, \phi) \quad (5)$$

where H is the standard radial Hamiltonian, q is the eigenvalue of the Casimir operator Q and $\mathcal{G}(r)$ is a function of r . $\Psi(r, \phi)$ is simultaneously an eigenfunction of Q and J_0 as well as of H , and is given by $\Psi(r, \phi) = e^{im\phi}\Phi(r)$. For the bound states the $\mathcal{D}^{(+)}$ representation is used, therefore the compact generator J_0 has eigenvalues [6]

$$m = \nu + \frac{1}{2} + \sqrt{q_\nu + \frac{1}{4}} \quad \nu = 0, 1, 2, \dots, \nu_{max} \quad (6)$$

The energy spectrum is obtained from [5]:

$$2\nu + 1 = \alpha_\nu - \beta_\nu - \delta_\nu \quad (7)$$

where

$$\begin{aligned} \alpha_\nu &\equiv \sqrt{-aE_\nu + f + 1} = p_\nu + m_\nu \\ \beta_\nu &\equiv \sqrt{-c_0E_\nu + h_0 + 1} = p_\nu - m_\nu \\ \delta_\nu &\equiv \sqrt{-c_1E_\nu + h_1 + 1} = \sqrt{4q_\nu + 1} \end{aligned}$$

This are the basic equations of the algebraization derived in [5].

3 Euclidean connection

The main motivation for Euclidean connection [1] is that the asymptotic states (regular solution) are written as [7]:

$$|\phi\rangle^{(\infty)} = \mathcal{A}e^{ikr} + \mathcal{B}e^{-ikr} \quad (8)$$

where \mathcal{A} and \mathcal{B} are the Jost functions and $e^{\pm ikr}$ are eigenfunctions of an Euclidean algebra $e(2)$:

$$[L_z, P_x] = iP_y \quad [L_z, P_y] = -iP_x \quad [P_x, P_y] = 0 \quad (9)$$

A realization in polar coordinates of the ladder and Casimir operators is as follows:

$$\begin{aligned} P_{\pm} &= e^{\pm i\phi} \left[-i\partial_r + \frac{1}{r} \left(\pm i\partial_{\phi} + \frac{1}{2} \right) \right] \\ \mathbf{P}^2 &= -\partial_r^2 - \frac{1}{r^2} \left(\partial_{\phi}^2 + \frac{1}{4} \right) \end{aligned} \quad (10)$$

where $P_{\pm} \equiv P_x \pm iP_y$ and $\mathbf{P}^2 \equiv P_+P_-$. Taking the limit $r \rightarrow \infty$, the asymptotic generators are obtained:

$$\begin{aligned} P_{\pm}^{(\infty)} &= -ie^{\pm i\phi} \partial_r & L_z^{(\infty)} &= -i\partial_{\phi} \\ \mathbf{P}^2^{(\infty)} &= -\partial_r^2 \end{aligned} \quad (11)$$

that also closes an $e(2)$ algebra. The eigenstate basis $\{|\pm k, m\rangle^{(\infty)}\}$ is

$$|\pm k, m\rangle^{(\infty)} = e^{\pm ikr} e^{im\phi} \quad (12)$$

and the action of the asymptotic generators upon it is given by:

$$\begin{aligned} P_{\pm}^{(\infty)}|\pm k, m\rangle^{(\infty)} &= (\pm k)|\pm k, m \pm 1\rangle^{(\infty)} \\ \mathbf{P}^2|\pm k, m\rangle^{(\infty)} &= k^2|\pm k, m\rangle^{(\infty)} \end{aligned} \quad (13)$$

As is well known, we can expand the ladder operators of $so(2, 1)$ in terms of the $e(2)$ generators [8]. Therefore, given an $so(2, 1)$ algebrization of the bound states, it is natural to study its extension to the scattering sector by means of the mentioned expansion. The expanded ladder operator $J_+^{(\infty)}$ of $so(2, 1)^{(\infty)}$ in the continuous representation (where $j = -\frac{1}{2} + if(k)$) obtained in [1] is:

$$J_+^{(\infty)}(\pm k) = \frac{e^{i\gamma_{\pm}(k)}}{\pm k} \left[\left(-\frac{1}{2} \mp if(k) \right) P_+^{(\infty)} + L_z^{(\infty)} P_+^{(\infty)} \right] \quad (14)$$

where $f(k)$ is defined by:

$$k^2 = h(-f^2(k)) \quad E_j = h \left[q + \frac{1}{4} \right] \quad (15)$$

where $h(\eta)$ is the function that connects the Energy E_j with the Casimir eigenvalue q or, equivalently, the Hamiltonian with the Casimir operator Q .

A recurrence relation may be calculated and solved for the coefficients A_m and B_m by applying (14) directly to (8), with $|\phi\rangle^{(\infty)} = e^{-im\phi}|j, m\rangle^{(\infty)}$, and equating it to the action of $J_+^{(\infty)}$ upon $|j, m\rangle^{(\infty)}$. Since $S_m(k) = A_m/B_m$, then:

$$S_m(k) = e^{im[\gamma_+ + \gamma_-]} \frac{\Gamma\left[m + \frac{1}{2} - if(k)\right]}{\Gamma\left[m + \frac{1}{2} + if(k)\right]} \Delta(k) \quad (16)$$

where $\Delta(k)$ is a constant factor to be determined.

4 S matrix for the HNP

Starting from (4), the asymptotic algebra is readily obtained by taking the limit $r \rightarrow \infty$ (or equivalently $z = 1$).

$$\begin{aligned} J_0^{(\infty)} &= -i \partial_\phi \\ J_\pm^{(\infty)} &= e^{\pm i\phi} \left[\mp \frac{\sqrt{c_1}}{2} \partial_r - i \partial_\phi \pm \frac{1}{2} \right] \\ Q^{(\infty)} &= \frac{c_1}{4} \partial_r^2 - \frac{1}{4} \end{aligned} \quad (17)$$

Since these generators form also an $so(2, 1)$ algebra, the Euclidean connection may be applied directly. Equation (17) corresponds to the $so(2, 1)^{(\infty)}$ algebra mentioned before. We only need to determine the expansion coefficients $f(k)$ and $\gamma_\pm(k)$. This is accomplished by equating the action of (14) with the one of (17), obtaining

$$\begin{aligned} \gamma_\pm(k) = 0 \quad f(k) &= \frac{k\sqrt{c_1}}{2} \\ \Rightarrow h'(\eta) &= -\frac{4}{c_1} \eta \end{aligned} \quad (18)$$

From (7) we know that

$$\begin{aligned} E_j = -\frac{4q - h_1}{c_1} &= h \left[q + \frac{1}{4} \right] \\ \Rightarrow h(\eta) &\equiv -\frac{4}{c_1} \eta + \frac{h_1 + 1}{c_1} \end{aligned} \quad (19)$$

Since we are dealing with the scattering sector and $V_{PNH} \propto h_1 + 1$ as $r \rightarrow \infty$ [2], $h_1 = -1$. Therefore $h'(\eta) = h(\eta)$ and the S matrix is given by (16):

$$S_m(k) = \frac{\Gamma\left(m + \frac{1}{2} - \frac{1}{2} ik\sqrt{c_1}\right)}{\Gamma\left(m + \frac{1}{2} + \frac{1}{2} ik\sqrt{c_1}\right)} \Delta_m(k) \quad (20)$$

where $\Delta_m(k)$ is a regular function of k derived by Natanzon [2]. It can easily be verified that the poles are in agreement with the bound state energy spectrum.

5 Conclusions

We have successfully extended the algebraic treatment for the bound states developed in [5] to the scattering sector via the $so(2,1)$ Euclidean connection, therefore obtaining a purely algebraic description of the HNP using an $so(2,1)$ algebra. This can also be accomplished with a methodology analogous to the one used by Frank, et al. [9]. This was done in detail and it is to be published soon [10]. Meanwhile, a short review may be found in this volume [11]. Even though Alhassid, et al. had obtained an algebraic description of the HNP in [12] by means of an $so(2,2)$ algebra, it is important to notice that our treatment is simpler.

Acknowledgments

This work has been partially supported by grants USB-61D30 and FONACIT No. 6-2001000712.

References

- [1] Alhassid, Y., Gürsey, F., Iachello, F. *Ann. Phys.* **167**, (1986) 181.
- [2] Natanzon, G. A., *Vestnik Leningradskogo Universiteta*, **10**, (1971) 22-8, physics/9907032; *Theor. Math. Phys.*, **38**, (1979) 146.
- [3] Zwanziger, D., *J. Math. Phys.*, **8**, (1967) 1858.
- [4] Barut, A. O., Rasmussen, W., *J. Phys. B: Atom. Molec.*, **6**, (1973) 1965; Barut, A. O., Rasmussen, W., Salamó, S., *Phys. Rev. D*, **10**, (1974) 622; *Phys. Rev. D*, **10**, (1974) 630; Rasmussen, W., Salamó, S., *J. Math. Phys.*, **20**, (1979) 1064.
- [5] Cordero, P., Salamó, S., *Found. Phys.*, **23**, 675 (1993). *J. Math. Phys.*, **35**, (1994) 3301.
- [6] Barut, A.O., Fronsdal, C. *Proc. R. Soc. London, Ser. A*, **287**, (1965) 532.
- [7] Taylor, J. R., *Scattering Theory: The Quantum Theory on Nonrelativistic collisions*, John Wiley & Sons., 1972.
- [8] Gilmore, R. *Lie Groups, Lie algebras, and some of their applications*, J. Wiley & Sons., 1974.
- [9] Frank, A., Wolf, Kurt B., *Phys. Rev. Lett.*, **52**(20), 1737 (1984).
- [10] Albrecht, H., Salamó, S., *to be published*.
- [11] Salamó, S., to appear in *Rev. Mex. Fís.*, (2002).
- [12] Alhassid, Y., Wu, J., Gürsey, F., *Ann. Phys.*, **196** (1989) 163-191.