

Towards quantum superpositions of a mirror

William Marshall^{1,2}, Christoph Simon¹, Roger Penrose^{3,4} and Dik Bouwmeester^{1,2}

¹ *Department of Physics, University of Oxford, Oxford OX1 3PU, United Kingdom*

² *Department of Physics, University of California, Santa Barbara, CA 93106*

³ *Center for Gravitational Physics and Geometry,*

The Pennsylvania State University, University Park, PA 16802

⁴ *Department of Mathematics, University of Oxford, Oxford OX1 3LB, United Kingdom*

(Dated: February 1, 2008)

We propose a scheme for creating quantum superposition states involving of order 10^{14} atoms via the interaction of a single photon with a tiny mirror. This mirror, mounted on a high-quality mechanical oscillator, is part of a high-finesse optical cavity which forms one arm of a Michelson interferometer. By observing the interference of the photon only, one can study the creation and decoherence of superpositions involving the mirror. All experimental requirements appear to be within reach of current technology.

In 1935 Schrödinger pointed out that according to quantum mechanics even macroscopic systems can be in superposition states [1]. The quantum interference effects are expected to be hard to detect due to environment induced decoherence [2]. Nevertheless there have been proposals on how to create and observe macroscopic superpositions in various systems [3–5], and experiments demonstrating superposition states of superconducting devices [6] and fullerene molecules [7]. One long-term motivation for this kind of experiment is the question of whether unconventional decoherence processes such as gravitationally induced decoherence or spontaneous wave-function collapse [8, 9] occur.

Here we present a scheme that is close in spirit to Schrödinger’s original discussion. A small quantum system (a photon) is coupled to a large system (a mirror) in order to create a macroscopic superposition. The basic principle of the experiment as described in Ref. [9] grew out of discussions in 1997 [10]. It consists of a Michelson interferometer in which one arm has a tiny moveable mirror. The radiation pressure of a single photon displaces the tiny mirror. The initial superposition of the photon being in either arm causes the system to evolve into a superposition of states corresponding to two distinct locations of the mirror. In the present proposal a high-finesse cavity is used to enhance the interaction between the photon and the mirror. The observed interference of the photon allows one to study the creation of coherent superposition states periodic with the motion of the mirror. We perform a detailed analysis of the requirements for such an experiment.

The proposed setup is shown in Fig. 1. In the cavity in arm A one of the mirrors is very small and attached to a micromechanical oscillator, similar to the cantilevers in atomic force microscopes. While the photon is in the cavity, it exerts radiation pressure on the mirror. Under the conditions that the oscillator period is much longer than the photon roundtrip time, and the amplitude of the mirror’s motion is small compared to the cavity length,




FIG. 1: The proposed setup: a Michelson interferometer for a single photon, where in each arm there is a high-finesse cavity. The cavity in arm A has a very small end mirror mounted on a micro-mechanical oscillator. The single photon comes in through I. If the photon is in arm A, the motion of the small mirror is affected by its radiation pressure. The photon later leaks out of either cavity and is detected at D1 or D2.

the system can be described by a Hamiltonian [11]

$$H = \hbar\omega_c a^\dagger a + \hbar\omega_m b^\dagger b - \hbar g a^\dagger a (b + b^\dagger), \quad (1)$$

where ω_c is the frequency of the photon in the cavity for the (empty) cavity length L , a is the creation operator for the photon, ω_m and b are the frequency and phonon creation operator for the center of mass motion of the mirror, and the coupling constant is $g = \frac{\omega_c}{L} \sqrt{\frac{\hbar}{2M\omega_m}}$, where M is the mass of the mirror. To start with, let us suppose that initially the photon is in a superposition of being in either arm A or B, and the mirror is in a coherent state $|\beta\rangle = e^{-|\beta|^2/2} \sum_{n=0}^{\infty} \frac{\beta^n}{\sqrt{n!}} |n\rangle$, where $|n\rangle$ are the eigenstates of the harmonic oscillator. Then the initial state is $|\psi(0)\rangle = \frac{1}{\sqrt{2}}(|0\rangle_A |1\rangle_B + |1\rangle_A |0\rangle_B)|\beta\rangle$. After a time t the state of the system will be given by [12]

$$|\psi(t)\rangle = \frac{1}{\sqrt{2}} e^{-i\omega_c t} (|0\rangle_A |1\rangle_B |\beta e^{-i\omega_m t}\rangle + e^{i\kappa^2(\omega_m t - \sin \omega_m t)} |1\rangle_A |0\rangle_B |\beta e^{-i\omega_m t} + \kappa(1 - e^{-i\omega_m t})\rangle), \quad (2)$$

where $\kappa = g/\omega_m$. In the second term on the r.h.s. the motion of the mirror is altered by the radiation pressure of the photon in cavity A . The parameter κ quantifies the displacement of the mirror in units of the size of the coherent state wavepacket. In the presence of the photon the mirror oscillates around a new equilibrium position determined by the driving force.

The maximum possible interference visibility for the photon is given by twice the modulus of the off-diagonal element of the photon's reduced density matrix. By tracing over the mirror one finds from Eq. (2) that the off-diagonal element has the form

$$\frac{1}{2}e^{-\kappa^2(1-\cos\omega_m t)}e^{i\kappa^2(\omega_m t - \sin\omega_m t) + i\kappa\text{Im}[\beta(1-e^{i\omega_m t})]} \quad (3)$$

where Im denotes the imaginary part. The first factor is the modulus, reaching a minimum after half a period at $t = \pi/\omega_m$, when the mirror is at its maximum displacement. The second factor gives the phase, which is identical to that obtained classically due to the varying length of the cavity. For general t the phase depends on β , i.e. the initial conditions of the mirror. However, the effect of β averages out after every full period.

In the absence of decoherence, after a full period, the system is in the state $\frac{1}{\sqrt{2}}(|0\rangle_A|1\rangle_B + e^{i\kappa^2 2\pi}|1\rangle_A|0\rangle_B)|\beta\rangle$, such that the mirror is again disentangled from the photon. Full interference can be observed if the photon is detected at this time. If the environment of the mirror "remembers" that the mirror has moved, then, even after a full period, the photon will still be entangled with the mirror's environment, and thus the interference for the photon will be reduced. Therefore the setup can be used to measure the decoherence of the mirror.

In practice the mirror will be in a thermal state, which can be written as a mixture of coherent states $|\beta\rangle$ with a Gaussian probability distribution $(1/\pi\bar{n})e^{-|\beta|^2/\bar{n}}$, where $\bar{n} = 1/(e^{\hbar\omega_m/kT} - 1)$ is the mean thermal number of excitations. In order to determine the expected interference visibility of the photon at a time t for an initial mirror state which is thermal, one has to average the off-diagonal element Eq. (3) over β with this distribution. The result is

$$\frac{1}{2}e^{-\kappa^2(2\bar{n}+1)(1-\cos\omega_m t)}e^{i\kappa^2(\omega_m t - \sin\omega_m t)}. \quad (4)$$

As a consequence of the averaging of the β -dependent phase in Eq. (3), the off-diagonal element now decays on a timescale $1/(\kappa\omega_m\sqrt{\bar{n}})$ after $t = 0$, i.e. very fast for the experimentally relevant case of $\kappa \sim 1$ and large \bar{n} . However, remarkably, it still exhibits a revival [4] at $t = 2\pi/\omega_m$, when photon and mirror become disentangled and the phase in Eq. (3) is independent of β , such that the phase averaging does not reduce the visibility. This behaviour is shown in Fig. 2. The magnitude of the revival is reduced by any decoherence of the mirror.




FIG. 2: Time evolution of the interference visibility of the photon over one period of the mirror's motion for $\kappa = 1$, $T = 1\text{mK}$ and $T = 60\mu\text{K}$. The time is given in units of $1/\omega_m$. The visibility decays very fast after $t = 0$, but in the absence of decoherence there is a sharp revival of the visibility after a full period. The width of each peak scales like $1/\sqrt{T}$.

The revival demonstrates the coherence of the superposition state that exists at intermediate times. For $\kappa^2 \gtrsim 1$ the state of the system is a superposition involving two distinct mirror positions. More precisely, for a thermal mirror, the state of the system is a mixture of such superpositions. However, this does not affect the fundamentally non-classical character of the state. We now discuss the experimental requirements for achieving such a superposition and observing its recoherence at $t = 2\pi/\omega_m$.

Firstly, we require $\kappa^2 \gtrsim 1$, which physically means that the momentum kick imparted by the photon has to be larger than the initial quantum uncertainty of the mirror's momentum. Let N denote the number of roundtrips of the photon in the cavity during one period of the mirror's motion, such that $2NL/c = 2\pi/\omega_m$. This allows us to rewrite the condition $\kappa^2 \gtrsim 1$ as

$$\frac{2\hbar N^3 L}{\pi c M \lambda^2} \gtrsim 1, \quad (5)$$

where λ is the wavelength of the light. The factors entering Eq. (5) are not all independent. The achievable N , which is determined by the quality of the mirrors, and the minimum mirror size (and hence M) both depend strongly on λ . The mirror's lateral dimension should be an order of magnitude larger than λ to limit diffraction losses. Its thickness in order to achieve sufficiently high reflectivity depends on λ as well.

Eq. (5) allows one to compare the viability of different wavelength ranges. While the highest values for N are achievable for microwaves (up to 10^{10}), this is counteracted by their cm wavelengths. On the other hand there are no good mirrors for highly energetic photons. The optical regime seems optimal. Here we propose an experiment with λ around 630 nm.

The cavity mode needs to have a very narrow focus on the tiny mirror, which requires the other cavity end mirror to be large due to beam divergence. The maximum cavity length is therefore limited by the difficulty of making large high quality mirrors. We propose a cavity length of 5 cm, and a small mirror size of $10 \times 10 \times 10$

microns, leading to a mass of order 5×10^{-12} kg.

Such a small mirror on a mechanical oscillator can be fabricated by coating a silicon cantilever with alternating layers of SiO_2 and a metal oxide. The best current mirrors are made in this way. A larger silicon oscillator has been coated with $\text{SiO}_2/\text{Ta}_2\text{O}_5$ and used as part of a high-finesse cavity in Ref. [13].

For the above dimensions the condition (5) is satisfied for $N = 5.6 \times 10^6$. Correspondingly, we are aiming for a photon loss per reflection not larger than 3×10^{-7} , about a factor of 4 below the best reported values for such mirrors [14], and for a transmission of 10^{-7} , consistent with the $10\mu\text{m}$ mirror thickness [15]. For these values, about 1% of the photons are still left in the cavity after a full period of the mirror. For the above values of N and L one obtains a frequency $\omega_m = 2\pi \times 500$ Hz. This leads to a quantum uncertainty for the mirror position of order 10^{-13} m, which corresponds to the necessary displacement in the superposition in order to achieve $\kappa^2 \sim 1$.

The fact that a relatively large L is needed to satisfy Eq. (5) implies that the creation of superpositions following the related proposal of Ref. [4], which relies on a micro-cavity, imposes requirements that are probably beyond the reach of current technology. A large L is helpful because, for a given N , it allows one to use a lower frequency ω_m , and thus a more weakly bound mirror that is easier to displace by the photon.

Secondly, the requirement of observing the revival puts a bound on the acceptable environmental decoherence. To estimate the expected decoherence we model the mirror's environment by an (Ohmic) bath of harmonic oscillators. The effect of this environment can approximately be described by a decoherence rate $\gamma_D = \gamma_m k T M (\Delta x)^2 / \hbar^2$ governing the decay of off-diagonal elements between different mirror positions [2]. Here γ_m is the damping rate for the mechanical oscillator, T is the temperature of the environment, which is constituted mainly by the internal degrees of freedom of the mirror and cantilever, and Δx is the separation of two coherent states that are originally in a superposition. This approximation is strictly valid only for times much longer than $2\pi/\omega_m$ and for Δx large compared to the width of the individual wavepackets. A more rigorous description would follow Ref. [16] and take into account the movement of the mirror. Our analysis indicates that the order of magnitude of the decoherence is well captured by γ_D . Assuming that the experiment achieves $\kappa^2 \gtrsim 1$, i.e. a separation by the size of a coherent state wavepacket, $\Delta x \sim \sqrt{\frac{\hbar}{M\omega_m}}$, the condition $\gamma_D \lesssim \omega_m$ can be cast in the form

$$Q \gtrsim \frac{kT}{\hbar\omega_m} = \bar{n}, \quad (6)$$

where $Q = \omega_m/\gamma_m$ is the quality factor of the mechanical oscillator. Bearing in mind that $Q \gtrsim 10^5$ has been

achieved [17] for silicon cantilevers of approximately the right dimensions and frequency, this implies that the temperature has to be 3 mK or less.

Thirdly, the stability requirements for the experiment are very strict. The phase of the interferometer has to be stable over the whole measurement time. This means that the distance between the large cavity end mirror and the equilibrium position [18] of the small mirror has to be stable to of order $\lambda/20N = 0.6 \times 10^{-14}$ m. The required measurement time can be determined in the following way. A single run of the experiment starts by sending a weak pulse into the interferometer, such that on average 0.1 photons go into either cavity. This probabilistically prepares a single-photon state as required to a good approximation. The two-photon contribution has to be kept low because it causes noise in the interferometer. From Eq. (4) the width of the revival peak is $2/\kappa\omega_m\sqrt{\bar{n}}$. This implies that only a fraction $\sim 1/\pi\sqrt{\bar{n}}$ of the remaining photons will leak out in the time interval of the revival. It is therefore important to work at the lowest possible temperature. A temperature of 60 μK has been achieved with a nuclear demagnetization cryostat [19].

Considering the required low value of ω_m and the fact that approximately 1% of the photons remain after a full period for the assumed loss, this implies a detection rate of approximately 100 photons per hour in the revival interval, given a detection efficiency of 70 %. Thus a measurement time of order 10 minutes should give statistically significant results. After one such measurement period the interferometer can be readjusted, and the experiment can be repeated. Stability of order 10^{-13} m/min for an STM inside at 8 K was achieved with a rather simple suspension [20]. Gravitational wave observatories using interferometers also require very high stability in order to have a length sensitivity of 10^{-19} m over timescales of a ms or greater [21], for arm lengths of order 1 km.

The experiment requires ultra-high vacuum conditions in order to ensure that events where an atom hits the cantilever are sufficiently rare not to cause significant errors, which is at the level of about 5/s. Background gas particle densities of order $100/\text{cm}^3$ have been achieved [22] and are sufficient for our purposes.

After every single run of the experiment the mirror has to be damped to reset it to its initial (thermal) state. This could be done by electric or magnetic fields, e.g. following Ref. [23], where a Nickel sphere was attached to the cantilever, whose Q could then be changed by three orders of magnitude by applying a magnetic field.

There are several ways in which the total measurement time could be decreased, thus relaxing the stability requirements. Switchable cavity mirrors would allow coupling the photon into and out of the cavity at any desired time, instead of relying on postselection. This allows the data collection in the revival interval to be increased by three orders of magnitude. The transmission of a distributed Bragg reflector (DBR) can be changed

by altering the refractive index of one of the two materials. Highly reflective mirrors have been fabricated from alternating layers of GaAs and AlAs [24]. The refractive index of GaAs can be changed dramatically by optical pumping [25]. A change from 3.6 to 3.15 on a ps timescale was observed at a probe wavelength of 830nm for pump intensities in the range 1 kJ/m^2 (just below the damage threshold of the material). Such a large change in refractive index would allow efficient optical switching [26] and is a promising avenue that deserves experimental investigation. The main open question is whether the absorption of the materials [27] can be made low enough to match our very strict demands.

Since the width of the revival peak scales like $1/\sqrt{T}$, the required measurement time can also be decreased by decreasing the temperature below $60 \mu\text{K}$. Passive cooling techniques may be improved. In addition, active cooling of mirror oscillators has been proposed [28], and even implemented experimentally for a large mirror [29]. The mirror's movement is sensed via the output light of an optical cavity, and a feedback loop is used to damp the motion. Applying the theory of Ref. [28] to our mirror, one finds that cooling even to the ground state of the center of mass motion is conceivable. This would reduce the required measurement time, and thus the stability requirements, by a factor of 50.

In principle the proposed setup has the potential to test wave function reduction models, in particular the one of Ref. [9]. Based on [9] and [16], we estimate that an improvement of the ratio Q/T by about five orders of magnitude from the values discussed in this paper ($Q = 10^5$ and $T = 60\mu\text{K}$) would make the predicted wavefunction decoherence rate comparable to the environmental decoherence rate. Improvements in Q are certainly conceivable. In particular, Q is known to increase with decreasing temperature [17]. Active cooling methods for the mirror's center of mass motion as described above could in principle also be used to indirectly cool the mirror's internal degrees of freedom. Whether such sympathetic cooling is a realistic avenue is in itself a fascinating subject for further study.

We have performed a detailed study of the experimental requirements for the creation and observation of quantum superposition states of a mirror consisting of 10^{14} atoms, approximately nine orders of magnitude more massive than any superposition observed to date. Our analysis suggests that, while very demanding, this goal appears to be in reach of current technology. It is remarkable that a tabletop experiment has the potential to test quantum mechanics in an entirely new regime. Preliminary experiments on components of the proposal are currently under way.

This work was supported by the E.U. (IST-1999 - 10033). W.M. is supported by EPSRC (award no. 00309297). C.S. is supported by a Marie Curie Fellowship of the E.U. (no. HPMF-CT-2001-01205). R.P. thanks

the NSF for support under contract 00-90091 and the Leverhulme Foundation for an Emeritus Fellowship. We would like to thank S. Bose, M. Davies, R. Epstein, T. Knuuttila, R. Lalezari, J. Pethica, and J. Roberts for useful discussions.

-
- [1] E. Schrödinger, *Die Naturwissenschaften* **23**, 807 (1935).
 - [2] D. Giulini *et al.*, *Decoherence and the Appearance of a Classical World in Quantum Theory* (Springer, Berlin, 1996); W.H. Zurek, *Phys. Today* **44**, 36 (1991).
 - [3] J. Ruostekoski, M. J. Collett, R. Graham, and D. F. Walls, *Phys. Rev. A* **57**, 511 (1998); J. I. Cirac, M. Lewenstein, K. Molmer, and P. Zoller, *Phys. Rev. A* **57**, 1208 (1998).
 - [4] S. Bose, K. Jacobs, and P.L. Knight, *Phys. Rev. A* **59**, 3204 (1999).
 - [5] A.D. Armour, M.P. Blencowe, and K.C. Schwab, *Phys. Rev. Lett.* **88**, 148301 (2002).
 - [6] C.H. van der Wal *et al.*, *Science* **290**, 773 (2000); J.R. Friedman, V. Patel, W. Chen, S.K. Tolpygo, and J.E. Lukens, *Nature* **406**, 43 (2000).
 - [7] M. Arndt *et al.*, *Nature* **401**, 680 (1999).
 - [8] G.C. Ghirardi, A. Rimini and T. Weber, *Phys. Rev. D* **34**, 470 (1986); G.C. Ghirardi, P. Pearle, and A. Rimini, *Phys. Rev. A* **42**, 78 (1990); I.C. Percival, *Proc. Roy. Soc. London, Ser. A* **447**, 189 (1994); D.I. Fivel, *Phys. Rev. A* **56**, 146 (1997); L. Diosi, *Phys. Rev. A* **40**, 1165 (1989).
 - [9] R. Penrose, in A. Fokas *et al.* (Eds.), *Mathematical Physics 2000* (Imperial College, London, 2000).
 - [10] J. Schmiedmayer, R. Penrose, D. Bouwmeester, J. Daprich, H. Weinfurter, A. Zeilinger, private communication (1997). See also D. Bouwmeester *et al.*, in N. Dadhich and J. Narlikar (Eds.), *Gravitation and Relativity: At the turn of the Millennium* (IUCAA, Pune, 1998).
 - [11] C.K. Law, *Phys. Rev. A* **51**, 2537 (1994); C.K. Law, *Phys. Rev. A* **49**, 433 (1993).
 - [12] S. Mancini, V.I. Man'ko, and P. Tombesi, *Phys. Rev. A* **55**, 3042 (1997); S. Bose, K. Jacobs, and P.L. Knight, *Phys. Rev. A* **56**, 4175 (1997).
 - [13] I. Tittonen *et al.*, *Phys. Rev. A* **59**, 1038 (1999).
 - [14] G. Rempe, R.J. Thompson, H.J. Kimble, and R. Lalezari, *Opt. Lett.* **17**, 363 (1992).
 - [15] C.J. Hood, H.J. Kimble, and J. Ye, *Phys. Rev. A* **64**, 033804 (2001).
 - [16] W.T. Strunz and F. Haake, quant-ph/0205108 (2002)
 - [17] H.J. Mamin and D. Rugar, *Appl. Phys. Lett.* **79**, 3358 (2001).
 - [18] As we have seen above, the thermal fluctuations of the small mirror around this equilibrium position (which are of order 10^{-12} m for $60 \mu\text{K}$) lead to a narrowing of the visibility peaks, but do not destroy the revival.
 - [19] W. Yao *et al.*, *J. Low Temp. Phys.* **120**, 121 (2000).
 - [20] B.C. Stipe, M.A. Rezaei and W. Ho., *Rev. Sci. Inst.* **70**, 137 (1999).
 - [21] S. Rowan and J. Hough, *Living Review in Relativity*, **3**, 2000-3 (2000),
 - [22] G. Gabrielse *et al.*, *Phys. Rev. Lett.* **65**, 1317 (1990).
 - [23] K. Wago, D. Botkin, C.S. Yannoni, and D. Rugar, *Appl. Phys. Lett.* **72**, 2757 (1998).
 - [24] T.E. Sale, *Vertical Cavity Surface Emitting Lasers* (Re-

- search Studies Press, Baldock, 1995).
- [25] L. Huang, J.P. Callan, E.N. Glezer, and E. Mazur, Phys. Rev. Lett. **80**, 185 (1998).
- [26] M. Scalora, J.P. Dowling, C.M. Bowden, and M.J. Bloemer, Phys. Rev. Lett. **73**, 1368 (1993).
- [27] M.D. Sturge, Phys. Rev. **127**, 768 (1962).
- [28] S. Mancini, D. Vitali, and P. Tombesi, Phys. Rev. Lett. **80**, 688 (1998).
- [29] P.F. Cohadon, A. Heidmann, and M. Pinard, Phys. Rev. Lett. **83**, 3174 (1999).