

# From a Mechanical Lagrangian to the Schrödinger Equation. A Modified Version of the Quantum Newton's Law

A. Bouda\*

Laboratoire de Physique Théorique, Université de Béjaïa,  
Route Targa Ouazemour, 06000 Béjaïa, Algeria

October 30, 2018

## Abstract

In the one-dimensional stationary case, we construct a mechanical Lagrangian describing the quantum motion of a non-relativistic spinless system. This Lagrangian is written as a difference between a function  $T$ , which represents the quantum generalization of the kinetic energy and which depends on the coordinate  $x$  and the temporal derivatives of  $x$  up to the third order, and the classical potential  $V(x)$ . The Hamiltonian is then constructed and the corresponding canonical equations are deduced. The function  $T$  is first assumed arbitrary. The development of  $T$  in a power series together with the dimensional analysis allow us to fix univocally the series coefficients by requiring that the well-known quantum stationary Hamilton-Jacobi equation be reproduced. As a consequence of this approach, we formulate the law of the quantum motion representing a new version of the quantum Newton's law. We also analytically establish the famous Bohm's relation  $\mu\dot{x} = \partial S_0/\partial x$  outside of the framework of the hydrodynamical approach and show that the well-known quantum potential, although it is a part of the kinetic term, it plays really a role of an additional potential as assumed by Bohm.

PACS: 03.65. Ca; 03.65. Ta

Key words: Lagrangian, Hamiltonian, Conjugate momentum, Hamilton's principal function, Quantum Hamilton-Jacobi equation.

---

\*Electronic address: bouda\_a@yahoo.fr

## 1 Introduction

In the one-dimensional space, the quantum stationary Hamilton-Jacobi equation (QSHJE) for a non-relativistic spinless particle of mass  $\mu$  and energy  $E$  is [1]

$$\frac{1}{2\mu} \left( \frac{\partial S_0}{\partial x} \right)^2 + V(x) - E = \frac{\hbar^2}{4\mu} \left[ \frac{3}{2} \left( \frac{\partial S_0}{\partial x} \right)^{-2} \left( \frac{\partial^2 S_0}{\partial x^2} \right)^2 - \left( \frac{\partial S_0}{\partial x} \right)^{-1} \left( \frac{\partial^3 S_0}{\partial x^3} \right) \right], \quad (1)$$

where  $S_0$  and  $V(x)$  are respectively the reduced action and the external potential. The solution of this equation, investigated in [2, 3, 4, 5], is written in [6] as

$$S_0 = \hbar \arctan \left( a \frac{\phi_1}{\phi_2} + b \right) + \hbar \kappa, \quad (2)$$

where  $(\phi_1, \phi_2)$  is a real set of independent solutions of the Schrödinger equation (SE) and  $(a, b, \kappa)$  are real integration constants satisfying the condition  $a \neq 0$ .

In Bohm's theory [7], Eq. (1) can be obtained from the SE by writing the wave function in the form

$$\phi(x) = A(x) \exp \left( \frac{i}{\hbar} S_0(x) \right). \quad (3)$$

It is shown that the real function  $A(x)$  is proportional to  $(\partial S_0 / \partial x)^{-1/2}$  [1, 5].

In one dimension, the well-known quantum potential, represented by the term proportional to  $\hbar^2$  in (1), is written in terms of higher derivatives of  $S_0$ . Thus, it seems not correct to consider in a dynamical equation this term as a potential but it may be a quantum correction of the kinetic part represented by the first term in (1). However, we will show in Section 5 that this term plays indeed a role of an additional potential.

An unsatisfactory feature of the form (3) of the wave function is the fact that for bound states, described by real wave functions up to a constant phase factor,  $S_0$  must be constant. Floyd remarked this problem and proposed to use a trigonometric representation in the real wave function cases [8, 9]. He also proposed that quantum trajectories were obtained by using Jacobi's theorem [9, 10],

$$t - t_0 = \frac{\partial S_0}{\partial E}, \quad (4)$$

as in classical mechanics. In Ref. [11], it is shown that the resulting trajectories depend on the choice of the couple of solutions of the SE used in the expression of  $S_0$ . This represents an unsatisfactory feature since the mathematical choices affect the physical results. Furthermore, in Ref. [12], it is shown that the time delay in tunneling through a potential barrier, calculated by using Jacobi's theorem as proposed by Floyd, manifests also some ambiguities.

From an equivalence postulate, Faraggi and Matone [3, 4] derived quantum mechanics. They deduced that the wave function must be written in the unified form

$$\phi(x) = \left( \frac{\partial S_0}{\partial x} \right)^{-1/2} \left[ \alpha \exp \left( \frac{i}{\hbar} S_0 \right) + \beta \exp \left( -\frac{i}{\hbar} S_0 \right) \right] \quad (5)$$

both for bound and unbound states,  $\alpha$  and  $\beta$  being complex constants. In the case where the wave function  $\phi$  is real, we have  $|\alpha| = |\beta|$  but never  $S_0 = cte$ . This

result is also reproduced in Ref. [5] outside of the framework of the equivalence postulate.

Recently, by appealing to the quantum transformation [4, 13] which allows to write the QSHJE in the classical form, the relation

$$\dot{x} \frac{\partial S_0}{\partial x} = 2[(E - V(x))] \quad (6)$$

is derived in Ref. [6]. It leads to a third order differential equation which appears as a quantum generalization of the first integral of Newton's law [6]. The relativistic version of this law is also obtained for a spinless particle in Ref. [14]. When the quantum coordinate [4, 13] is used to apply Jacobi's theorem [6] or to express the Lagrangian in order to obtain the equation of motion [11], the formalism does not seem to suffer from any mathematical ambiguity. Furthermore, the fundamental result is reproduced with many formulations [11]. However, some unsatisfactory features are appeared. First, at the turning points (where  $V(x) = E$ ), since  $\partial S_0/\partial x$  never has a vanishing value, relation (6) indicates that  $\dot{x} = 0$ . Then, from (6) we can show that all the higher temporal derivatives of  $x$  take a vanishing value at the turning points:  $\dot{x} = 0$ ,  $\ddot{x} = 0$ ,  $\dddot{x} = 0$ , ... This means that when the particle gets to these points, it will stand still forever. Another feature which seems to be unsatisfactory is the extreme rapid divergence of the velocity in the classically forbidden regions which is manifested for the three different potentials considered in [12]. Other comments about relation (6) are given by Floyd in [15].

In this paper, we present a new version of the quantum law of motion free from the previous criticisms. In Section 2, we propose a generalization of the classical mechanics to the cases where the Lagrangian is depending on  $x$ ,  $\dot{x}$ ,  $\ddot{x}$  and  $\ddot{\dot{x}}$ . In Section 3, we will use a dimensional analysis to construct a quantum Lagrangian for a spinless particle. We will show in Section 4 that the well-known QSHJE can be reproduced from the previous Lagrangian with a judicious choice of some parameters. We, therefore, present the resulting quantum law of motion and apply our result in the free particle case.

## 2 The Generalized Classical Mechanics

Let us consider any Lagrangian,  $L$ , depending on  $(x, \dot{x}, \ddot{x}, \ddot{\dot{x}}, t)$ . In the next section we will justify the choice of this set of variables. Let us define the Hamilton's principal function as

$$S = \int L(x, \dot{x}, \ddot{x}, \ddot{\dot{x}}, t) dt . \quad (7)$$

The least action principle leads to the generalized Euler-Lagrange equation

$$\frac{d^3}{dt^3} \frac{\partial L}{\partial \ddot{\dot{x}}} - \frac{d^2}{dt^2} \frac{\partial L}{\partial \ddot{x}} + \frac{d}{dt} \frac{\partial L}{\partial \dot{x}} - \frac{\partial L}{\partial x} = 0 . \quad (8)$$

The corresponding Hamiltonian is constructed in Ref. [11]. Its expression is given by

$$H = \left( \frac{\partial L}{\partial \dot{x}} - \frac{d}{dt} \frac{\partial L}{\partial \ddot{x}} + \frac{d^2}{dt^2} \frac{\partial L}{\partial \ddot{\dot{x}}} \right) \dot{x} + \left( \frac{\partial L}{\partial \ddot{x}} - \frac{d}{dt} \frac{\partial L}{\partial \ddot{\dot{x}}} \right) \ddot{x} + \frac{\partial L}{\partial \ddot{\dot{x}}} \ddot{\dot{x}} - L , \quad (9)$$

so that if we require that  $\partial L/\partial t = 0$ , with the use of (8), we obtain  $dH/dt = 0$ . This means that  $H$  is a constant of motion when the time is considered uniform.

The conjugate momentum,  $P$ , must be defined so that if we require that the space be homogeneous,  $P$  must be a constant of motion. In other words, if we require that  $\partial L/\partial x = 0$ , we must obtain  $\dot{P} \equiv dP/dt = 0$ . With the use of Eq. (8), it follows that

$$P = \frac{\partial L}{\partial \dot{x}} - \frac{d}{dt} \frac{\partial L}{\partial \ddot{x}} + \frac{d^2}{dt^2} \frac{\partial L}{\partial \dot{\ddot{x}}}. \quad (10)$$

Thus, the generalized Euler-Lagrange equation takes the form

$$\dot{P} = \frac{\partial L}{\partial x}. \quad (11)$$

We will call  $P$  the principal conjugate momentum. The form (9) of the Hamiltonian suggests that we define two secondary conjugate momentums

$$\Pi = \frac{\partial L}{\partial \ddot{x}} - \frac{d}{dt} \frac{\partial L}{\partial \dot{\ddot{x}}} \quad (12)$$

and

$$\Xi = \frac{\partial L}{\partial \dot{\ddot{x}}}, \quad (13)$$

so that

$$H = P\dot{x} + \Pi\ddot{x} + \Xi\dot{\ddot{x}} - L. \quad (14)$$

By taking into account relations (10), (11), (12) and (13), from (14) we deduce that

$$dH = -\dot{P}dx + \dot{x}dP - \dot{\Pi}d\dot{x} + \dot{\ddot{x}}d\Pi - \dot{\Xi}d\dot{\ddot{x}} + \dot{\ddot{x}}d\Xi - \frac{\partial L}{\partial t}dt. \quad (15)$$

If we suppose that the Hamiltonian can be written as follows

$$H = H(x, P; \dot{x}, \Pi; \ddot{x}, \Xi; t), \quad (16)$$

we therefore deduce that

$$dH = \frac{\partial H}{\partial x}dx + \frac{\partial H}{\partial P}dP + \frac{\partial H}{\partial \dot{x}}d\dot{x} + \frac{\partial H}{\partial \Pi}d\Pi + \frac{\partial H}{\partial \ddot{x}}d\ddot{x} + \frac{\partial H}{\partial \Xi}d\Xi + \frac{\partial H}{\partial t}dt. \quad (17)$$

Identifying (15) and (17), we obtain the generalized canonical equations

$$\dot{x} = \frac{\partial H}{\partial P}, \quad (18)$$

$$\dot{P} = -\frac{\partial H}{\partial x}, \quad (19)$$

$$\dot{\ddot{x}} = \frac{\partial H}{\partial \Pi}, \quad (20)$$

$$\dot{\Pi} = -\frac{\partial H}{\partial \dot{\ddot{x}}}, \quad (21)$$

$$\dot{\dot{\ddot{x}}} = \frac{\partial H}{\partial \Xi}, \quad (22)$$

$$\dot{\Xi} = -\frac{\partial H}{\partial \ddot{x}}, \quad (23)$$

$$\frac{\partial L}{\partial t} = -\frac{\partial H}{\partial t}. \quad (24)$$

We would like to indicate that the analogue of these equations for a Lagrangian depending only on  $(x, \dot{x}, \ddot{x})$  is obtained in Ref. [16].

We stress to call the reader attention to a contradiction which seems to appear when we compare the Lagrangian formulation and the Hamiltonian one. Generally, relation (8) is a six order differential equation while the last set of equations seems to lead to a fourth order equation. In fact, since  $H$  is a function of  $(x, P, \dot{x}, \Pi, \ddot{x}, \Xi, t)$ , by combining (18), (20) and (22), we would be able to express  $P$ ,  $\Pi$  and  $\Xi$  in terms of  $x$ ,  $\dot{x}$ ,  $\ddot{x}$ ,  $\dot{\ddot{x}}$  and  $t$ . Then, Eq. (19) would be written as a fourth order differential equation, expressed only in terms of  $x(t)$  and its temporal derivatives. However, as we will see in Appendix II where a concrete case is examined, Eqs. (18) and (20) are identities and can not be combined with (22) in order to express  $(P, \Pi, \Xi)$  in terms of  $(x, \dot{x}, \ddot{x}, \dot{\ddot{x}}, t)$ . So, there is no contradiction. We would like to add that a similar false contradiction appears also when we consider a Lagrangian depending only on  $(x, \dot{x}, \ddot{x}, t)$ .

### 3 The Quantum Lagrangian

Our goal is to construct in the stationary case a Lagrangian from which we can start to get to the well-known QSHJE, which in turn, leads to the SE. The higher derivatives of  $S_0$  appearing in (1) suggest that our Lagrangian depends on  $x$ ,  $\dot{x}$  and higher temporal derivatives of  $x$ . Since the only parameters which appear in the SE as well as in the QSHJE are the mass  $\mu$  and the Planck constant  $\hbar$ , the Lagrangian must also depend on  $\mu$  and  $\hbar$

$$L = L(\mu, \hbar, x, \dot{x}, \ddot{x}, \dots).$$

Of course, we have not taken into account the possibility of the dependence on other physical parameters through the external potential. We will see that this fact is not important for our construction.

The reduced action, given in (2), depends on the energy  $E$  through the functions  $\phi_1$  and  $\phi_2$  and on two non-additive integration constants  $a$  and  $b$ . This indicates that the fundamental law describing the quantum motion is a fourth order differential equation. It follows that  $x(t)$  will contain four integration constants as it is in the earlier formulations of trajectory representation of quantum mechanics [6, 10, 17]. From the mathematical point of view, Eq. (8) indicates that it is sufficient to write the Lagrangian as a function of  $(x, \dot{x}, \ddot{x})$  to obtain a fourth order differential equation. Nevertheless, from the physical point of view, the set  $(x, \dot{x}, \ddot{x})$  is not sufficient to define the “mechanical state” of the system. In fact, the knowledge of this set at any time does not allow to determine the future positions of the system since the fundamental law of motion must be a fourth order one. However, the knowledge at any time of the set  $(x, \dot{x}, \ddot{x}, \dot{\ddot{x}})$  is sufficient to predict the future motion since it must allow to determine the four integration constants. Thus, the “mechanical state” is defined by this last set and the Lagrangian must be written as

$$L = L(\mu, \hbar, x, \dot{x}, \ddot{x}, \dot{\ddot{x}}). \quad (25)$$

Note that the analogue of this reasoning for classical mechanics was proposed by Landau-Lifchitz [18].

Now, one mathematical difficulty appears since Eq. (8) indicates that the Lagrangian (25) often leads to a six order differential equation. In what follows,

we will see how to overcome this problem. For the moment, let us write the Lagrangian in a natural form

$$L = T(\mu, \hbar, x, \dot{x}, \ddot{x}, \dot{\dot{x}}) - V(x) . \quad (26)$$

where  $T$  is the quantum generalization of the kinetic energy whose form is assumed independent on the external potential  $V(x)$ . Let us suppose that  $T$  is a regular function so that we can develop it in a power series with respect to  $\hbar$  as follows

$$T = \sum_n \hbar^n T_n(\mu, x, \dot{x}, \ddot{x}, \dot{\dot{x}}) . \quad (27)$$

In the limit  $\hbar \rightarrow 0$ ,  $T$  must not diverge. Thus, we impose the condition

$$T_n = 0, \quad \forall n < 0 . \quad (28)$$

Before going further, it is interesting to remark that in order to simplify our investigation, there are two physical conditions that we can impose:

- in the limit  $\hbar \rightarrow 0$ , the function  $T$  goes to the classical expression  $\mu\dot{x}^2/2$ ;
- in the absence of the external potential, the space must be homogeneous, and then the condition  $\partial L/\partial x = 0$  implies that  $\partial T/\partial x = 0$ , meaning that we can suppress the dependence on  $x$  of  $T$  in (26) and (27).

However, in order to persist in the most general construction for  $T$  and  $L$ , we do not impose these two conditions. In what follows, we will see that the homogeneity of the space when  $V(x) = 0$  and the relation  $\lim_{\hbar \rightarrow 0} T = \mu\dot{x}^2/2$  are consequences of the well-known QSHJE.

Because all the terms of the series (27) must have a dimension of the energy, the unit of measurement of the function  $T_n$  is

$$[T_n] = kg^{-n+1}m^{-2n+2}s^{n-2} . \quad (29)$$

In relation (27), we see that the only physical parameter which may appear in the expression of  $T_n$  is the mass  $\mu$ . Then, relation (29) indicates that  $\forall n \geq 0$  the function  $T_n$  takes the form

$$T_n = \frac{1}{\mu^{n-1}} f_n(x, \dot{x}, \ddot{x}, \dot{\dot{x}}) . \quad (30)$$

By taking into account relations (28) and (30), (27) turns out to be

$$T = \sum_{n=0}^{\infty} \frac{\hbar^n}{\mu^{n-1}} f_n(x, \dot{x}, \ddot{x}, \dot{\dot{x}}) , \quad (31)$$

where the unit measurement of the function  $f_n$  is

$$[f_n] = m^{-2n+2}s^{n-2} . \quad (32)$$

As indicated above, the dependence on  $\dot{\dot{x}}$  will induce a differential equation of sixth order. The unique manner to avoid this difficulty is to assume that this dependence is linear. Then, the most general form for  $f_n$  is

$$f_n(x, \dot{x}, \ddot{x}, \dot{\dot{x}}) = u_n(x, \dot{x}, \ddot{x}) + \dot{\dot{x}} v_n(x, \dot{x}, \ddot{x}) , \quad (33)$$

where  $u_n$  and  $v_n$  are two functions depending only on  $x$ ,  $\dot{x}$  and  $\ddot{x}$ . We mention that the term  $\ddot{x}v_n(x, \dot{x}, \ddot{x})$  will induce two fifth order terms when we apply (8). However, these terms cancel each other out.

By taking into account relation (32), the unit measurements of  $u_n$  and  $v_n$  are

$$[u_n] = m^{-2n+2} s^{n-2}, \quad (34)$$

$$[v_n] = m^{-2n+1} s^{n+1}. \quad (35)$$

Assuming  $u_n$  and  $v_n$  as regular functions, we can develop them in a power series as follows

$$u_n = \sum_{ijk} \alpha_{ijk}^{(n)} x^k \dot{x}^i \ddot{x}^j, \quad (36)$$

$$v_n = \sum_{ijk} \beta_{ijk}^{(n)} x^k \dot{x}^i \ddot{x}^j, \quad (37)$$

where  $\alpha_{ijk}^{(n)}$  and  $\beta_{ijk}^{(n)}$  are dimensionless real numbers since  $u_n$  and  $v_n$  do not depend on any physical parameter.

Relations (34) and (36) imply that

$$i + j + k = -2n + 2, \quad -i - 2j = n - 2,$$

leading to  $i = -3n - 2k + 2$  and  $j = n + k$ . It follows that all the numbers  $\alpha_{ijk}^{(n)}$  for which  $i \neq -3n - 2k + 2$  and  $j \neq n + k$  must take a vanishing value. So, by defining

$$\alpha_{nk} \equiv \alpha_{-3n-2k+2, n+k, k}^{(n)}, \quad (38)$$

we have

$$u_n = \sum_k \alpha_{nk} \frac{x^k \ddot{x}^{n+k}}{\dot{x}^{3n+2k-2}}. \quad (39)$$

Relations (35) and (37) imply that

$$i + j + k = -2n + 1, \quad -i - 2j = n + 1,$$

leading to  $i = -3n - 2k + 3$  and  $j = n + k - 2$ . It follows that all the numbers  $\beta_{ijk}^{(n)}$  for which  $i \neq -3n - 2k + 3$  and  $j \neq n + k - 2$  must take a vanishing value. So, by defining

$$\beta_{nk} \equiv \beta_{-3n-2k+3, n+k-2, k}^{(n)}, \quad (40)$$

we have

$$v_n = \sum_k \beta_{nk} \frac{x^k \ddot{x}^{n+k-2}}{\dot{x}^{3n+2k-3}}. \quad (41)$$

From (39) and (41), we see that for  $k < 0$  the functions  $u_n$  and  $v_n$  diverge at  $x = 0$ . In order to avoid these divergences,  $\alpha_{nk}$  and  $\beta_{nk}$  must take a vanishing value for  $k < 0$ . Concerning the possible divergences which will appear in the case where  $\dot{x}$  or  $\ddot{x}$  take a vanishing value, in the following Sections, we will see that  $\dot{x}$  never reaches a vanishing value and  $\ddot{x}$  is never present in the denominator. Thus, by taking into account relations (33), (39) and (41), expression (31) turns out to be

$$T = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n}{\mu^{n-1}} \left[ \alpha_{nk} \frac{x^k \ddot{x}^{n+k}}{\dot{x}^{3n+2k-2}} + \beta_{nk} \frac{x^k \ddot{x}^{n+k-2} \dot{x}}{\dot{x}^{3n+2k-3}} \right]. \quad (42)$$

In this relation,  $k$  is considered as an integer number meaning that  $u_n$  and  $v_n$  are assumed infinitely differentiable at  $x = 0$ . If it is not the case, in order to keep  $k$  integer, it is sufficient to substitute in the above relations  $x$  by  $(x - x_0)$ ,  $x_0$  being a point chosen in such a way as to have  $u_n$  and  $v_n$  infinitely differentiable at this point.

However, with the form (42) of  $T$ , we have a problem concerning the Hamiltonian formulation. In fact, by applying (13), we get an expression for  $\Xi$  only in terms of  $x$ ,  $\dot{x}$  and  $\ddot{x}$ . This means that  $\Xi$ ,  $x$ ,  $\dot{x}$  and  $\ddot{x}$  can not be considered as independent variables in the Hamiltonian approach. In addition, if we apply (12) and (20), we see that there is a contradiction between the obtained results unless we put  $\beta_{nk} = 0$  for every  $n$  and  $k$ . This forces us to lose the dependence on  $\ddot{x}$ . However, as explained at the beginning of this Section, in order to obtain a Lagrangian describing the ‘‘mechanical state’’ of the system, the presence of  $\ddot{x}$  is required. We stress to indicate that it is not the generalized classical mechanics presented in Section 2 which is ambiguous, but the linear terms pose also problems in the formulation where we have a Lagrangian of classical type. As shown in Appendix I, it is instructive to consider a Lagrangian depending only on  $x$  and  $\dot{x}$  as in classical mechanics, to see why linear terms induce mathematical ambiguities and how to overcome these difficulties. The solution consists in adding to the Lagrangian a quadratic term in  $\dot{x}$ , proportional to a constant  $\lambda$ , and in taking at the end the limit  $\lambda \rightarrow 0$ . That’s what we will do in the quantum case. We will add a quadratic term in  $\dot{x}$  and, with the use of (26) and (42), we write the quantum Lagrangian as

$$L = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n}{\mu^{n-1}} \left[ \alpha_{nk} \frac{x^k \ddot{x}^{n+k}}{\dot{x}^{3n+2k-2}} + \beta_{nk} \frac{x^k \ddot{x}^{n+k-2} \dot{x}}{\dot{x}^{3n+2k-3}} \right] + \frac{1}{2} \lambda \dot{x}^2 - V(x), \quad (43)$$

$\lambda$  being a constant extremely small so that, after having obtained the equation of motion, we can take the limit  $\lambda \rightarrow 0$ . We stress that  $\lambda$  is independent on  $\mu$  and  $\hbar$ . Of course, the Lagrangian (43) leads to a six order differential equation. However, as we will see in Section 4, when we take the limit  $\lambda \rightarrow 0$ , we obtain a fourth order equation of motion. We emphasize that this additional term is useless in the Lagrangian approach but it is necessary for a coherent formulation of the Hamiltonian approach.

Now, let us calculate the conjugate momentums by applying (10), (12) and (13). We get

$$P = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n}{\mu^{n-1}} \left\{ \begin{aligned} & \left[ (3n+2k-2)(n+k-1)\alpha_{nk} + (3n+2k-2)(3n+2k-3)\beta_{nk} \right. \\ & \quad - (k+1)(n+k+1)\alpha_{n,k+1} - (k+1)(6n+4k-3)\beta_{n,k+1} \\ & \quad \left. + (k+1)(k+2)\beta_{n,k+2} \right] \frac{x^k \ddot{x}^{n+k}}{\dot{x}^{3n+2k-1}} \\ & + \left[ - (n+k)(n+k-1)\alpha_{nk} - (n+k)(3n+2k-3)\beta_{nk} \right. \\ & \quad \left. + (k+1)(n+k-1)\beta_{n,k+1} \right] \frac{x^k \ddot{x}^{n+k-2} \dot{x}}{\dot{x}^{3n+2k-2}} \end{aligned} \right\}$$



$$+ \lambda \ddot{x} , \quad (44)$$

$$\begin{aligned} \Pi = & \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n}{\mu^{n-1}} \left[ (n+k)\alpha_{nk} \right. \\ & \left. + (3n+2k-3)\beta_{nk} - (k+1)\beta_{n,k+1} \right] \frac{x^k \ddot{x}^{n+k-1}}{\dot{x}^{3n+2k-2}} - \lambda \ddot{x} , \quad (45) \end{aligned}$$

$$\Xi = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n \beta_{nk}}{\mu^{n-1}} \frac{x^k \ddot{x}^{n+k-2}}{\dot{x}^{3n+2k-3}} + \lambda \dot{x} . \quad (46)$$

Since we see in these three relations nine variables ( $P, \Pi, \Xi, x, \dot{x}, \ddot{x}, \dots, \ddot{\ddot{x}}$ ), it is clear that in the set  $(x, P, \dot{x}, \Pi, \ddot{x}, \Xi)$ , all the six variables can be considered independent. Then, by using (14), (43) and (46), the Hamiltonian takes the form

$$\begin{aligned} H(x, P, \dot{x}, \Pi, \ddot{x}, \Xi) = & P \dot{x} + \Pi \ddot{x} - \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n \alpha_{nk}}{\mu^{n-1}} \frac{x^k \ddot{x}^{n+k}}{\dot{x}^{3n+2k-2}} \\ & + \frac{1}{2\lambda} \left[ \Xi - \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n \beta_{nk}}{\mu^{n-1}} \frac{x^k \ddot{x}^{n+k-2}}{\dot{x}^{3n+2k-3}} \right]^2 + V(x) . \quad (47) \end{aligned}$$

In Appendix II, the resulting canonical equations are deduced and it is shown that they allow to reproduce the same expressions (44), (45) and (46) for the conjugate momenta  $P$ ,  $\Pi$  and  $\Xi$  obtained from the Lagrangian (43). In particular, it is also shown that the Hamiltonian formulation is equivalent to the Lagrangian one and both of them lead to the same law of motion for any  $\lambda$ . This law can be easily obtained by using (44) in the expression of  $\dot{P}$  given in Appendix II and which we obtain by applying (19). Calculating the temporal derivative of (44), we see that the expression of  $\dot{P}$  has no term containing  $\ddot{\ddot{x}}$ . The term containing  $\ddot{\ddot{x}}$  is proportional to  $\lambda$ . In conclusion, we obtain in the Lagrangian formulation as well as in the Hamiltonian one the same fourth order equation of motion when  $\lambda \rightarrow 0$ .

#### 4 Toward the Quantum Stationary Hamilton-Jacobi Equation

Now, our task consists in finding the numerical values of the dimensionless parameters  $\alpha_{nk}$  and  $\beta_{nk}$  with which we can reproduce the well-known QSHJE. In the stationary case, the Hamilton's principal function is related to the reduced action by  $S = S_0 - Et$ . Then, we write

$$dS = dS_0 - E dt . \quad (48)$$

On the other hand, by using (7) and (14), we can deduce that

$$dS = L dt = P dx + \Pi d\dot{x} + \Xi d\ddot{x} - E dt , \quad (49)$$

where the Hamiltonian  $H$  is substituted by  $E$ . Comparing (48) and (49), we obtain

$$dS_0 = P dx + \Pi d\dot{x} + \Xi d\ddot{x} , \quad (50)$$

from which we deduce

$$\begin{aligned} \frac{dS_0}{dx} &= P + \Pi \frac{d\dot{x}}{dx} + \Xi \frac{d\ddot{x}}{dx} \\ &= P + \Pi \frac{\ddot{x}}{\dot{x}} + \Xi \frac{\dot{\ddot{x}}}{\dot{x}} . \quad (51) \end{aligned}$$

We indicate that the left hand side in this last equation represents the partial derivative  $\partial S_0/\partial x$  appearing in (1) and in the abstract. In fact, in (1),  $S_0$  depends on  $x$  and not on any other variable. Furthermore, from the solution (2) of (1), we see that  $S_0$  is a function only of  $x$  and some integrations constants.

Thus, the usual conjugate momentum [2, 4, 5, 6, 10], represented here by the left hand side of (51), differs from the principal conjugate momentum  $P$  that we have defined in (10). This difference can be explicitly calculated from (51).

Using expression (44), (45) and (46) in the limit  $\lambda \rightarrow 0$ , relation (51) turns out to be

$$\frac{dS_0}{dx} = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n}{\mu^{n-1}} \left[ A_{nk} \frac{x^k \ddot{x}^{n+k}}{\dot{x}^{3n+2k-1}} + B_{nk} \frac{x^k \ddot{x}^{n+k-2} \dot{x}}{\dot{x}^{3n+2k-2}} \right], \quad (52)$$

where

$$\begin{aligned} A_{nk} = & (3n^2 + 2k^2 + 5nk - 4n - 3k + 2)\alpha_{nk} \\ & + (3n + 2k - 1)(3n + 2k - 3)\beta_{nk} - (k + 1)(n + k + 1)\alpha_{n,k+1} \\ & - 2(k + 1)(3n + 2k - 1)\beta_{n,k+1} + (k + 1)(k + 2)\beta_{n,k+2}, \end{aligned} \quad (53)$$

and

$$\begin{aligned} B_{nk} = & -(n + k)(n + k - 1)\alpha_{nk} - (3n^2 + 2k^2 + 5nk \\ & - 3n - 3k - 1)\beta_{nk} + (k + 1)(n + k - 1)\beta_{n,k+1}. \end{aligned} \quad (54)$$

With the use of (51), Eq. (14) turns out to be

$$H = \dot{x} \frac{dS_0}{dx} - L. \quad (55)$$

Substituting in this last relation  $H$  by  $E$  and using (43) in the limit  $\lambda \rightarrow 0$ , we deduce that

$$E - V(x) = \dot{x} \frac{dS_0}{dx} - \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n}{\mu^{n-1}} \left[ \alpha_{nk} \frac{x^k \ddot{x}^{n+k}}{\dot{x}^{3n+2k-2}} + \beta_{nk} \frac{x^k \ddot{x}^{n+k-2} \dot{x}}{\dot{x}^{3n+2k-3}} \right]. \quad (56)$$

By using this last expression, we can rewrite Eq. (1) in the form

$$\begin{aligned} & \dot{x} \left( \frac{dS_0}{dx} \right)^3 - \frac{1}{2\mu} \left( \frac{dS_0}{dx} \right)^4 + \frac{\hbar^2}{4\mu} \left[ \frac{3}{2} \left( \frac{d^2 S_0}{dx^2} \right)^2 - \frac{dS_0}{dx} \frac{d^3 S_0}{dx^3} \right] \\ & = \left( \frac{dS_0}{dx} \right)^2 \left[ \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n}{\mu^{n-1}} \left( \alpha_{nk} \frac{x^k \ddot{x}^{n+k}}{\dot{x}^{3n+2k-2}} + \beta_{nk} \frac{x^k \ddot{x}^{n+k-2} \dot{x}}{\dot{x}^{3n+2k-3}} \right) \right]. \end{aligned} \quad (57)$$

With the use of (52) in Appendix III, it is shown that the unique physical solution for  $\alpha_{nk}$  and  $\beta_{nk}$  satisfying this last relation is

$$\alpha_{00} = \frac{1}{2}, \quad \alpha_{20} = \frac{5}{8}, \quad \beta_{20} = -\frac{1}{4}, \quad (58)$$

$$\beta_{0k} = \alpha_{1k} = \beta_{1k} = 0 \quad \forall k \geq 0, \quad \alpha_{0k} = \alpha_{2k} = \beta_{2k} = 0 \quad \forall k \geq 1, \quad (59)$$

and

$$\alpha_{nk} = \beta_{nk} = 0 \quad \forall n \geq 3, \quad \forall k \geq 0. \quad (60)$$

Thus, with the use of these values in expression (52), the Hamiltonian (47) leads straightforwardly, when we take the limit  $\lambda \rightarrow 0$ , to the well-known QSHJE given by (1).

It is also interesting to remark that by using (58), (59) and (60) in expression (42), we deduce that  $\lim_{\hbar \rightarrow 0} T = \mu \dot{x}^2/2$ . Thus, the classical expression for the kinetic energy is a consequence of the substitution of the general form of the Hamiltonian that we have constructed in the well-known QSHJE when we take the limit  $\hbar \rightarrow 0$ .

Since there is a unique physical solution for the set  $\{\alpha_{nk}, \beta_{nk}\}$  and the fact that  $\beta_{20}$  is different from 0, we conclude that without the linear term in  $\dot{x}$  in Eq. (33), our initial Lagrangian never allows to reach the QSHJE. This constitutes the mathematical reason for which we have imposed the presence of this term.

## 5 The Quantum Law of Motion

With the use of (58), (59 and (60), although we obtain from (52)

$$\frac{dS_0}{dx} = \mu \dot{x} , \quad (61)$$

recalling the famous Bohm's relation postulated a half century ago [7], the principal conjugate momentum can be derived from (44) and takes the form

$$P = \mu \dot{x} - \frac{\hbar^2}{4\mu} \left[ 2 \frac{\ddot{x}^2}{\dot{x}^5} - \frac{\dot{\ddot{x}}}{\dot{x}^4} \right] . \quad (62)$$

From (43), the expression of the Lagrangian ( $\lambda \rightarrow 0$ ) is

$$L = \frac{1}{2} \mu \dot{x}^2 + \frac{\hbar^2}{4\mu} \left[ \frac{5}{2} \frac{\ddot{x}^2}{\dot{x}^4} - \frac{\dot{\ddot{x}}}{\dot{x}^3} \right] - V(x) . \quad (63)$$

It is clear that, thanks to the obtained values for  $\{\alpha_{nk}, \beta_{nk}\}$ , the kinetic term does not depend on  $x$ . This means that in the absence of the external potential,  $V(x) = 0$ , we have  $\partial L / \partial x = 0$  and then we deduce that the space is homogeneous. This property was not imposed in the present formalism but it is a consequence of the QSHJE. We also deduce that the principal conjugate momentum,  $P$ , is a constant of motion in the case where  $V(x) = 0$ .

By applying (9) or (14), the Hamiltonian becomes

$$H = \frac{1}{2} \mu \dot{x}^2 - \frac{\hbar^2}{4\mu} \left[ \frac{5}{2} \frac{\ddot{x}^2}{\dot{x}^4} - \frac{\dot{\ddot{x}}}{\dot{x}^3} \right] + V(x) . \quad (64)$$

First, this expression can also be obtained from the QSHJE, Eq. (1), by using (61) and substituting  $E$  by  $H$ . Second, it is interesting to remark that if we define

$$Q = -\frac{\hbar^2}{4\mu} \left[ \frac{5}{2} \frac{\ddot{x}^2}{\dot{x}^4} - \frac{\dot{\ddot{x}}}{\dot{x}^3} \right] , \quad (65)$$

representing the well-known quantum potential in the QSHJE, Eqs. (63) and (64) can be written as

$$L = \frac{1}{2} \mu \dot{x}^2 - [Q + V(x)] , \quad H = \frac{1}{2} \mu \dot{x}^2 + [Q + V(x)] . \quad (66)$$

These two last relations constitute a proof that  $Q$  really plays a role of an additional potential, as predicted by Bohm [7], despite it is a part of the kinetic term  $T$ . This is possible because of the particular values we have obtained for  $\alpha_{20}$  and  $\beta_{20}$ .

By construction, we have automatically  $dH/dt = 0$  since we are in the stationary case. Thus, by substituting in (64)  $H$  by  $E$ , we obtain a first integral of the analogue of the quantum Newton's law. Calculating the total derivative with respect to  $x$  or  $t$  of the two members of (64), we get

$$\mu\ddot{x} + \frac{\hbar^2}{\mu} \left[ \frac{5}{2} \frac{\dot{x}^3}{x^6} - 2 \frac{\dot{x}\ddot{x}}{x^5} + \frac{1}{4} \frac{\ddot{x}}{x^4} \right] + \frac{dV}{dx} = 0, \quad (67)$$

representing the analogue of the quantum Newton's law. It can also be obtained by applying (8) or (11). As expected, it is a fourth order differential equation. The general solution  $x(t)$  will contain four integration constants which can be determined by the knowledge of the initial conditions  $x(t_0) = x_0$ ,  $\dot{x}(t_0) = \dot{x}_0$ ,  $\ddot{x}(t_0) = \ddot{x}_0$  and  $\ddot{\dot{x}}(t_0) = \ddot{\dot{x}}_0$ . In the case where the energy  $E$  is known, the three first conditions are sufficient. In contrast to the law established in [6], through (64) we see that there is no derivative of the external potential in the first integral of (67).

We remark that in all the above relations, if we put  $\hbar = 0$ , we reproduce the classical formulas.

We observe also that relation (61) can be considered as a law of motion. In fact, by using (2), we have

$$\mu \frac{dx}{dt} = \frac{\hbar a W}{a^2 \phi_1^2 + (1 + b^2) \phi_2^2 + 2ab\phi_1\phi_2}, \quad (68)$$

where  $W = \phi_2 d\phi_1/dx - \phi_1 d\phi_2/dx$  is a constant representing the Wronskian of the two independent solutions  $(\phi_1, \phi_2)$  of the SE. Relation (68) is a first order differential equation in which we see the presence of three integration constants:  $a$ ,  $b$  and  $E$  through  $\phi_1$  and  $\phi_2$ . We stress to indicate that Eq. (68) is independent on the choice of the couple  $(\phi_1, \phi_2)$ . In fact, if we use another couple  $(\theta_1, \theta_2)$  in (2), with the same procedure used in Ref. [12], we can find two other parameters  $(\tilde{a}, \tilde{b})$ , which we must use instead of  $(a, b)$  in expression (2), in such a way as to guarantee that  $dS_0/dx$  remains invariant.

In Section 3, we have indicated that a problem about some divergences may occur in the kinetic term (42). Concerning  $\ddot{x}$ , since  $\beta_{00} = \beta_{01} = \beta_{10} = 0$ ,  $\ddot{x}$  is never present in the denominator. With regard to  $\dot{x}$ , since  $a$  and  $W$  are both different from 0, Eq. (68) indicates that  $\dot{x}$  never reaches a vanishing value.

As an example, let us consider the free particle case for which  $V(x) = 0$ . Choosing for the SE the two following solutions

$$\phi_1 = \sin(kx), \quad \phi_2 = \cos(kx), \quad (69)$$

where  $k = \sqrt{2\mu E}/\hbar$ , Eq. (68) leads to the quantum time equation

$$a\sqrt{\frac{2E}{\mu}}(t - t_0) = \frac{a^2 + b^2 + 1}{2}x + \frac{1 + b^2 - a^2}{4k} \sin(2kx) - \frac{ab}{2k} \cos(2kx). \quad (70)$$

First, if we put  $a = 1$  and  $b = 0$ , we reproduce the classical relation

$$x = \sqrt{\frac{2E}{\mu}}(t - t_0), \quad (71)$$

as it is the case in the earlier formulations [6, 17] of trajectory representation. Second, in contrast to the trajectories obtained in [12], we have no nodal structure. Third, in the classical limit  $\hbar \rightarrow 0$ , as in [17], a residual indeterminacy subsists. However, in this limit, (70) differs from (71) only by the proportionality factor between  $x$  and  $(t - t_0)$ . For a particular condition on the parameters  $a$  and  $b$ , (70) reduces to (71).

In our point of view, in order to obtain a realistic model, it is necessary to generalize the present formulation to the three-dimensional space.

## 6 Conclusion

Before concluding, let us summarize the principal steps of the present approach.

After having proposed a generalization of the classical mechanics by starting from any Lagrangian depending on  $(x, \dot{x}, \ddot{x}, \dot{\ddot{x}}, t)$ , our goal was to reach the SE by reproducing the well-known QSHJE.

Our task consisted in establishing a fourth order differential equation to describe the quantum motion. For this purpose, by appealing to the dimensional analysis, we constructed in the stationary case a Lagrangian depending on  $(x, \dot{x}, \ddot{x}, \dot{\ddot{x}})$  from which we deduced a conjugate momentum which was a constant of motion in the absence of the external potential. Although we observed that a Lagrangian depending only on  $(x, \dot{x}, \ddot{x})$  was sufficient to establish a fourth order law, we indicated that in order to obtain a Lagrangian which described the “mechanical state” of the particle, it had also to depend on  $\dot{\ddot{x}}$ . Furthermore, without this dependence, from a mathematical point of view it was impossible to reach the QSHJE from our initial Lagrangian. Nevertheless, this dependence was linear in order to guarantee that the resulting law of motion should be a fourth order equation. We kept  $\ddot{x}$  in the terms proportional to  $\dot{\ddot{x}}$  because the induced fifth order terms in the equation of motion cancel each other out.

However, we remarked that with such a Lagrangian, it was not possible to obtain a coherent Hamiltonian formulation. This difficulty was surmounted by adding to the Lagrangian a quadratic term in  $\dot{\ddot{x}}$  proportional to one parameter noted  $\lambda$  and which was independent on  $\hbar$  and the mass  $\mu$  of the particle. In order to avoid a sixth order law, we took the limit  $\lambda \rightarrow 0$  after the equation of motion was obtained. We stress that this quadratic term is useless for the Lagrangian formulation. In other words, we have the same result if we do not add this term or if we add it and then take the limit  $\lambda \rightarrow 0$ . However, in the Hamiltonian approach, this term is necessary in order to have a coherent formulation. We showed that the two formulations, the Lagrangian and Hamiltonian ones, are equivalent.

We would like to indicate that it was possible to obtain all the results presented here without appealing to the Hamiltonian formulation, in particular to the canonical equations. It was sufficient to only use expression (9) of the Hamiltonian and to consider expressions (10), (12) and (13) of  $P$ ,  $\Pi$  and  $\Xi$  as mathematical beings but not as conjugate momentums. With this procedure, we insist that the additional quadratic term in  $\dot{\ddot{x}}$  proportional to  $\lambda$  is not required. However, we developed an Hamiltonian formulation in order to prove that such an approach was possible in the context of the generalized classical mechanics, to confirm the result obtained with the Lagrangian formulation and especially to justify that  $P$ ,  $\Pi$  and  $\Xi$  are really conjugate momentums.

The general Lagrangian and Hamiltonian that we constructed depend on

some dimensionless parameters. We substituted in the well-known QSHJE the expression of the Hamiltonian and we determined the values of these parameters in such a way as to guarantee the validity of the obtained relation. We showed that there was a unique physical solution. The obtained values allowed to reproduce the famous Bohm's relation  $dS_0/dx = \mu\dot{x}$  and to show that the quantum potential, although it was a part of the kinetic term, it really played a role of a potential. We also established the fundamental law of the quantum motion and applied our result in the free particle case.

In addition, we would like to mention that we have not imposed in the formalism to the space to be homogeneous in the absence of the external potential since we have not excluded the possibility of the dependence on  $x$  of  $T$ . We showed that this property was a consequence of the substitution in the QSHJE of the general expression of the Hamiltonian that we constructed and, in contrast to the earlier formulations of trajectory representation, we deduced that the principal conjugate momentum was a constant of motion in the absence of the external potential. We have not also imposed to the kinetic term to have as a limit when  $\hbar \rightarrow 0$  the classical expression  $\mu\dot{x}^2/2$ . We showed that this condition was a consequence of the QSHJE. In our point of view, this constitutes a proof that the function  $S_0$  appearing in the QSHJE, Eq. (1), and which is related to the Schrödinger wave function by (5), is really a quantum generalization of the classical reduced action.

To conclude, we would like to emphasize that in the context of the following hypothesis:

- the Hamilton's principal function is represented as an integral of a Lagrangian;
- the Lagrangian is a difference between a kinetic term,  $T$ , and the external potential  $V(x)$ ;
- the kinetic term is a function of  $(\hbar, \mu, x, \dot{x}, \ddot{x}, \dot{\ddot{x}})$  and its form does not depend on  $V(x)$ ;
- the resulting equation of motion is a fourth order one,

we showed that the Lagrangian which leads to the well-known QSHJE, and then to the SE, is unique and is the one given by (63). Of course, in the context of the above hypothesis, the resulting quantum law of motion, Eq. (67), is also unique.

## Appendix I

In this Appendix, we would like to explain in the classical Lagrangian case why linear terms induce ambiguities and how to overcome these difficulties. For this purpose, let us consider the following Lagrangian of the form

$$L_i(x, \dot{x}) = f(x) \dot{x}^i - V(x) , \quad (72)$$

where  $f(x)$  is an arbitrary function and  $i$  an integer number different from 0. The usual Euler-Lagrange equation leads to

$$(i-1) \left[ i f \dot{x}^{i-2} \ddot{x} + \frac{df}{dx} \dot{x}^i \right] + \frac{dV}{dx} = 0 . \quad (73)$$

The conjugate momentum is given by

$$P_i = \frac{\partial L_i}{\partial \dot{x}} = i f \dot{x}^{i-1} . \quad (74)$$

The resulting Hamiltonian is

$$H_i(x, P_i) = P_i \dot{x} - L_i = \frac{i-1}{i} \left( \frac{1}{i f} \right)^{\frac{1}{i-1}} P_i^{\frac{i}{i-1}} + V(x) . \quad (75)$$

The presence of  $(i-1)$  in the denominator indicates clearly that the case  $i=1$  requires a particular treatment. For  $i \neq 1$ , we can check that the canonical equations

$$\dot{x} = \frac{\partial H_i}{\partial P_i} = \left( \frac{P_i}{i f} \right)^{\frac{1}{i-1}} , \quad (76)$$

$$\dot{P}_i = -\frac{\partial H_i}{\partial x} = \left( \frac{P_i}{i f} \right)^{\frac{i}{i-1}} \frac{df}{dx} - \frac{dV}{dx} , \quad (77)$$

are compatible with (73) and (74). For  $i=1$ , if we apply naively the well-known relations

$$P_1 = \frac{\partial L_1}{\partial \dot{x}} = f(x) \quad (78)$$

and

$$H_1(x, P_1) = P_1 \dot{x} - L_1 = V(x) , \quad (79)$$

the canonical equations

$$\dot{x} = \frac{\partial H_1}{\partial P_1} = 0 \quad (80)$$

and

$$\dot{P}_1 = -\frac{\partial H_1}{\partial x} = -\frac{dV}{dx} \quad (81)$$

are not compatible with (73) and (74) for  $i=1$ . In fact, (73) indicates that  $dV/dx = 0$  while (78) and (81) imply that  $dV/dx = -\dot{f}$ . Furthermore, (78) indicates that  $P_1$  and  $x$  are not independent variables. The solution to this problem consists in adding to the Lagrangian (72) a quadratic term,

$$L_1(x, \dot{x}) = \frac{1}{2} \lambda \dot{x}^2 + f(x) \dot{x} - V(x) , \quad (82)$$

$\lambda$  being a constant, and in taking the limit  $\lambda \rightarrow 0$  after having obtained the equation of motion. In the Hamiltonian formulation, it is necessary to keep  $\lambda$  until  $P_1$  and  $\dot{P}_1$  are eliminated from the canonical equations. In this way, we can check that we obtain the same equation of motion with the two formulations.

## Appendix II

Our goal here is to deduce the generalized canonical equations from the Hamiltonian (47) and to show that the Lagrangian formulation and the Hamiltonian one are equivalent.

By applying relations (18)-(23), the canonical equations are

$$\dot{x} \equiv \frac{\partial H}{\partial P} = \dot{x} , \quad (83)$$

$$\ddot{x} \equiv \frac{\partial H}{\partial \Pi} = \ddot{x} , \quad (84)$$

$$\dot{\xi} \equiv \frac{\partial H}{\partial \Xi} = \frac{1}{\lambda} \left[ \Xi - \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n \beta_{nk}}{\mu^{n-1}} \frac{x^k \ddot{x}^{n+k-2}}{\dot{x}^{3n+2k-3}} \right], \quad (85)$$

$$\begin{aligned} \dot{P} \equiv -\frac{\partial H}{\partial x} &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n (k+1) \alpha_{n,k+1}}{\mu^{n-1}} \frac{x^k \ddot{x}^{n+k+1}}{\dot{x}^{3n+2k}} \\ &+ \frac{1}{\lambda} \left[ \Xi - \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n \beta_{nk}}{\mu^{n-1}} \frac{x^k \ddot{x}^{n+k-2}}{\dot{x}^{3n+2k-3}} \right] \\ &\left[ \sum_{p=0}^{\infty} \sum_{j=0}^{\infty} \frac{\hbar^p (j+1) \beta_{p,j+1}}{\mu^{p-1}} \frac{x^j \ddot{x}^{p+j-1}}{\dot{x}^{3p+2j-1}} \right] \\ &- \frac{dV}{dx}, \end{aligned} \quad (86)$$

$$\begin{aligned} \dot{\Pi} \equiv -\frac{\partial H}{\partial \dot{x}} &= -P - \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n (3n+2k-2) \alpha_{nk}}{\mu^{n-1}} \frac{x^k \ddot{x}^{n+k}}{\dot{x}^{3n+2k-1}} \\ &- \frac{1}{\lambda} \left[ \Xi - \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n \beta_{nk}}{\mu^{n-1}} \frac{x^k \ddot{x}^{n+k-2}}{\dot{x}^{3n+2k-3}} \right] \\ &\left[ \sum_{p=0}^{\infty} \sum_{j=0}^{\infty} \frac{\hbar^p (3p+2j-3) \beta_{pj}}{\mu^{p-1}} \frac{x^j \ddot{x}^{p+j-2}}{\dot{x}^{3p+2j-2}} \right], \end{aligned} \quad (87)$$

$$\begin{aligned} \dot{\Xi} \equiv -\frac{\partial H}{\partial \ddot{x}} &= -\Pi + \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n (n+k) \alpha_{nk}}{\mu^{n-1}} \frac{x^k \ddot{x}^{n+k-1}}{\dot{x}^{3n+2k-2}} \\ &+ \frac{1}{\lambda} \left[ \Xi - \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n \beta_{nk}}{\mu^{n-1}} \frac{x^k \ddot{x}^{n+k-2}}{\dot{x}^{3n+2k-3}} \right] \\ &\left[ \sum_{p=0}^{\infty} \sum_{j=0}^{\infty} \frac{\hbar^p (p+j-2) \beta_{pj}}{\mu^{p-1}} \frac{x^j \ddot{x}^{p+j-3}}{\dot{x}^{3p+2j-3}} \right]. \end{aligned} \quad (88)$$

Firstly, as we have indicated at the end of Section 2, Eqs. (18) and (20), now become (83) and (84), represent identities and do not give any information about the motion.

Secondly, remark that (85) is equivalent to (46).

Thirdly, if we calculate the temporal derivative of the two members of (85), we can deduce that

$$\begin{aligned} \dot{\Xi} &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n}{\mu^{n-1}} \left\{ [-(3n+2k-3)\beta_{nk} + (k+1)\beta_{n,k+1}] \frac{x^k \ddot{x}^{n+k-1}}{\dot{x}^{3n+2k-2}} \right. \\ &\quad \left. + (n+k-2)\beta_{nk} \frac{x^k \ddot{x}^{n+k-3} \dot{\ddot{x}}}{\dot{x}^{3n+2k-3}} \right\} + \lambda \dot{\ddot{x}}. \end{aligned}$$

Then, by substituting in this last relation  $\dot{\Xi}$  by its expression (88) and taking into account relation (85), we reproduce for  $\Pi$  the same expression as the one given by (45).

Fourthly, if we calculate the temporal derivative of the obtained result for  $\Pi$



(or of (45)), we get

$$\begin{aligned}
\dot{\Pi} = & \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n}{\mu^{n-1}} \left\{ \right. \\
& \left[ -(3n+2k-2)[(n+k)\alpha_{nk} + (3n+2k-3)\beta_{nk}] + (k+1)[(n \right. \\
& \left. +k+1)\alpha_{n,k+1} + (6n+4k-3)\beta_{n,k+1} - (k+2)\beta_{n,k+2}] \right] \frac{x^k \ddot{x}^{n+k}}{\dot{x}^{3n+2k-1}} \\
& + (n+k-1) \left[ (n+k)\alpha_{nk} + (3n+2k-3)\beta_{nk} \right. \\
& \left. \left. - (k+1)\beta_{n,k+1} \right] \frac{x^k \ddot{x}^{n+k-2} \dot{\ddot{x}}}{\dot{x}^{3n+2k-2}} \right\} \\
& - \lambda \ddot{\ddot{x}} .
\end{aligned}$$

Then, by substituting this expression in (87) and taking into account relation (85), we reproduce for  $P$  the same expression as the one given by (44).

It follows that the canonical equations (85), (87) and (88) lead to the same expressions (44), (45) and (46) for the conjugate momentums  $P$ ,  $\Pi$  and  $\Xi$  obtained from the Lagrangian (43).

Fifthly, from (43) and (47), we can easily check that

$$\frac{\partial L}{\partial x} = -\frac{\partial H}{\partial x} , \quad (89)$$

where we have used (85). Thus, since we have obtained the same expression for  $P$  in the two approaches, by comparing (11) and (19), we deduce that the Hamiltonian formulation is equivalent to the Lagrangian one and both of them lead to the same law of motion for any  $\lambda$ .

### Appendix III

In this Appendix, we will search for the numerical values of  $\alpha_{nk}$  and  $\beta_{nk}$  with which relation (57) is valid. For this purpose, let us calculate the second and the third derivatives of  $S_0$  from (52). We obtain

$$\begin{aligned}
\frac{d^2 S_0}{dx^2} = & \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n}{\mu^{n-1}} \left\{ \right. \\
& [(k+1)A_{n,k+1} - (3n+2k-1)A_{nk}] \frac{x^k \ddot{x}^{n+k+1}}{\dot{x}^{3n+2k+1}} \\
& + [(n+k)A_{nk} - (3n+2k-2)B_{nk} + (k+1)B_{n,k+1}] \frac{x^k \ddot{x}^{n+k-1} \dot{\ddot{x}}}{\dot{x}^{3n+2k}} \\
& \left. + (n+k-2)B_{nk} \frac{x^k \ddot{x}^{n+k-3} \dot{\ddot{x}}^2}{\dot{x}^{3n+2k-1}} + B_{nk} \frac{x^k \ddot{x}^{n+k-2} \dot{\ddot{x}}}{\dot{x}^{3n+2k-1}} \right\} , \quad (90)
\end{aligned}$$

and

$$\frac{d^3 S_0}{dx^3} = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\hbar^n}{\mu^{n-1}} \left\{ \right.$$

$$\begin{aligned}
& [(3n+2k-1)(3n+2k+1)A_{nk} - 2(k+1)(3n+2k+1)A_{n,k+1} \\
& \quad + (k+1)(k+2)A_{n,k+2}] \frac{x^k \ddot{x}^{n+k+2}}{\dot{x}^{3n+2k+3}} \\
& + [-(6n^2+4k^2+10nk+2n+k-1)A_{nk} \\
& \quad + (3n+2k)(3n+2k-2)B_{nk} + 2(k+1)(n+k+1)A_{n,k+1} \\
& \quad - 2(k+1)(3n+2k)B_{n,k+1} + (k+1)(k+2)B_{n,k+2}] \frac{x^k \ddot{x}^{n+k} \dot{\ddot{x}}}{\dot{x}^{3n+2k+2}} \\
& + [(n+k)(n+k-1)A_{nk} \\
& \quad - (6n^2+4k^2+10nk-12n-9k+4)B_{nk} \\
& \quad + 2(k+1)(n+k-1)B_{n,k+1}] \frac{x^k \ddot{x}^{n+k-2} \dot{\ddot{x}}^2}{\dot{x}^{3n+2k+1}} \\
& + (n+k-2)(n+k-3)B_{nk} \frac{x^k \ddot{x}^{n+k-4} \dot{\ddot{x}}^3}{\dot{x}^{3n+2k}} \\
& + [(n+k)A_{nk} - (6n+4k-3)B_{nk} + 2(k+1)B_{n,k+1}] \frac{x^k \ddot{x}^{n+k-1} \dot{\ddot{x}}}{\dot{x}^{3n+2k+1}} \\
& + 3(n+k-2)B_{nk} \frac{x^k \ddot{x}^{n+k-3} \dot{\ddot{x}} \ddot{\ddot{x}}}{\dot{x}^{3n+2k}} + B_{nk} \frac{x^k \ddot{x}^{n+k-2} \dot{\ddot{x}}}{\dot{x}^{3n+2k}} \Big\} . \quad (91)
\end{aligned}$$

The values of  $\alpha_{nk}$  and  $\beta_{nk}$  can be determined by writing the two members of (57) as a power series with respect to  $\hbar$  by using (52), (90) and (91). At the classical level ( $\hbar = 0$ ), Eqs. (52) and (57) can be written as

$$\left( \frac{dS_0}{dx} \right)^{(0)} = \sum_{k=0}^{\infty} \mu \left[ A_{0k} \frac{x^k \ddot{x}^k}{\dot{x}^{2k-1}} + B_{0k} \frac{x^k \ddot{x}^{k-2} \dot{\ddot{x}}}{\dot{x}^{2k-2}} \right] , \quad (92)$$

and

$$\dot{x} \left( \frac{dS_0}{dx} \right)^{(0)} - \frac{1}{2\mu} \left[ \left( \frac{dS_0}{dx} \right)^{(0)} \right]^2 - \sum_{k=0}^{\infty} \mu \left[ \alpha_{0k} \frac{x^k \ddot{x}^k}{\dot{x}^{2k-2}} + \beta_{0k} \frac{x^k \ddot{x}^{k-2} \dot{\ddot{x}}}{\dot{x}^{2k-3}} \right] = 0 . \quad (93)$$

The terms proportional to  $\dot{\ddot{x}}^2$  that we obtain by substituting (92) in (93) are linearly independent. Thus, Eq. (93) can not be satisfied without imposing any condition on the particle motion unless one has

$$B_{0k} = 0, \quad \forall k \geq 0 . \quad (94)$$

By using this result in (92), relation (93) turns out to be

$$\sum_{k=0}^{\infty} A_{0k} \frac{x^k \ddot{x}^k}{\dot{x}^{2k-2}} - \frac{1}{2} \left[ \sum_{k=0}^{\infty} A_{0k} \frac{x^k \ddot{x}^k}{\dot{x}^{2k-1}} \right]^2 - \sum_{k=0}^{\infty} \left[ \alpha_{0k} \frac{x^k \ddot{x}^k}{\dot{x}^{2k-2}} + \beta_{0k} \frac{x^k \ddot{x}^{k-2} \dot{\ddot{x}}}{\dot{x}^{2k-3}} \right] = 0 . \quad (95)$$

The terms proportional to  $\dot{\ddot{x}}$  are linearly independent. Thus, as above, (95) can not be satisfied unless one has

$$\beta_{0k} = 0, \quad \forall k \geq 0 . \quad (96)$$

Using expression (54) for  $n = 0$  and taking into account (94) and (96), we deduce that  $k(k-1)\alpha_{0k} = 0 \quad \forall k \geq 0$ . It follows that  $\alpha_{0k} = 0 \quad \forall k \geq 2$ . Therefore, with

the use of (53) for  $n = 0$  and (96), Eq. (95) becomes

$$\begin{aligned} & \left( -2\alpha_{00}^2 - \frac{1}{2}\alpha_{01}^2 + 2\alpha_{00}\alpha_{01} + \alpha_{00} - \alpha_{01} \right) \dot{x}^2 \\ & - \frac{1}{2}\alpha_{01}^2 \frac{x^2 \ddot{x}^2}{\dot{x}^2} - \alpha_{01}(2\alpha_{00} - \alpha_{01})x\dot{x} = 0. \end{aligned} \quad (97)$$

In this relation, we have three terms linearly independent. It follows that  $\alpha_{01} = 0$  and then  $\alpha_{00}(1 - 2\alpha_{00}) = 0$ . The solution  $\alpha_{00} = 0$  corresponds to a trivial solution and does not allow to obtain the classical limit. We keep the other solution and we write

$$\alpha_{00} = \frac{1}{2}, \quad \alpha_{0k} = 0, \quad \forall k \geq 1. \quad (98)$$

Note that with the above results, we have  $A_{00} = 1$  and  $A_{0k} = 0 \forall k \geq 1$ . Then, by taking into account relation (94), expression (92) becomes

$$\left( \frac{dS_0}{dx} \right)^{(0)} = \mu \dot{x}. \quad (99)$$

At the first level, taking into account relations (96) and (98), Eq. (57) yields

$$\hbar \mu^2 \sum_{k=0}^{\infty} \left[ \alpha_{1k} \frac{x^k \ddot{x}^{k+1}}{\dot{x}^{2k-1}} + \beta_{1k} \frac{x^k \ddot{x}^{k-1} \dot{x}}{\dot{x}^{2k-2}} \right] = 0.$$

In this equality, all the terms are linearly independent. If we do not want to impose any condition on the particle motion, it is necessary that

$$\alpha_{1k} = 0, \quad \beta_{1k} = 0, \quad \forall k \geq 0. \quad (100)$$

These relations mean that  $A_{1k} = 0$  and  $B_{1k} = 0 \forall k \geq 0$  and that the derivatives (52), (90) and (91) of  $S_0$  do not contain terms proportional to  $\hbar$ . Then, at the second level, from (57) we have

$$\hbar^2 \mu \left[ \left( \alpha_{20} - \frac{5}{8} \right) \frac{\ddot{x}^2}{\dot{x}^2} + \left( \beta_{20} + \frac{1}{4} \right) \frac{\dot{x}}{\dot{x}} + \sum_{k=1}^{\infty} \left( \alpha_{2k} \frac{x^k \ddot{x}^{k+2}}{\dot{x}^{2k+2}} + \beta_{2k} \frac{x^k \ddot{x}^k \dot{x}}{\dot{x}^{2k+1}} \right) \right] = 0.$$

For the same reasons as above, we deduce that

$$\alpha_{20} = \frac{5}{8}, \quad \beta_{20} = -\frac{1}{4}, \quad \alpha_{2k} = 0, \quad \beta_{2k} = 0, \quad \forall k \geq 1. \quad (101)$$

In order to determine the other parameters  $\alpha_{nk}$  and  $\beta_{nk}$  for  $n > 2$ , it is essential to remark that with the values given in (101), we can see from (53) and (54) that  $A_{2k} = 0$  and  $B_{2k} = 0$  for every  $k$  even for  $k = 0$ . Thus, as it is the case for the terms proportional to  $\hbar$ , the derivatives (52), (90) and (91) of  $S_0$  do not contain terms proportional to  $\hbar^2$ . For the upper levels, we will use a reasoning by recursion. Let us begin by the third level. Taking into account relations (96), (98), (100) and (101), it is easy to check that the validity of (57) requires that

$$\hbar^3 \sum_{k=0}^{\infty} \left[ \alpha_{3k} \frac{x^k \ddot{x}^{k+3}}{\dot{x}^{2k+5}} + \beta_{3k} \frac{x^k \ddot{x}^{k+1} \dot{x}}{\dot{x}^{2k+4}} \right] = 0,$$

which leads to

$$\alpha_{3k} = 0, \quad \beta_{3k} = 0, \quad \forall k \geq 0. \quad (102)$$

Now, let us suppose that for  $n > 3$ , the validity of (57) at the  $(n-1)^{th}$  requires that  $\alpha_{3k} = 0, \dots, \alpha_{n-1,k} = 0$  and  $\beta_{3k} = 0, \dots, \beta_{n-1,k} = 0$  for every  $k \geq 0$ . We have then  $A_{3k} = 0, \dots, A_{n-1,k} = 0$  and  $B_{3k} = 0, \dots, B_{n-1,k} = 0$  for every  $k \geq 0$ . By taking into account the fact that  $A_{00} = 1, A_{0k} = 0$  for every  $k \geq 1$  and  $B_{0k} = A_{1k} = B_{1k} = A_{2k} = B_{2k} = 0$  for every  $k \geq 0$ , at the  $n^{th}$  level, relation (57) yields

$$\frac{\hbar^n}{\mu^{n-3}} \sum_{k=0}^{\infty} \left[ \alpha_{nk} \frac{x^k \ddot{x}^{n+k}}{\dot{x}^{3n+2k-4}} + \beta_{nk} \frac{x^k \ddot{x}^{n+k-2} \dot{x}}{\dot{x}^{3n+2k-5}} \right] = 0.$$

As above, this relation implies that  $\alpha_{nk} = 0$  and  $\beta_{nk} = 0 \forall k \geq 0$ . Now we can assert that, in addition to the conditions (96), (98), (100) and (101), the validity of (57) requires that

$$\alpha_{nk} = 0, \quad \beta_{nk} = 0, \quad \forall n \geq 3, \quad \forall k \geq 0. \quad (103)$$

## References

1. A. Messiah, *Quantum Mechanics*, Vol. 1, (North Holland, New York, 1961).
2. E. R. Floyd, *Phys. Rev. D* 34 (1986) 3246; *Found. Phys. Lett.* 9 (1996) 489.
3. A. E. Faraggi and M. Matone, *Phys. Lett. B* 450 (1999) 34; *Phys. Lett. B* 437 (1998) 369.
4. A. E. Faraggi and M. Matone, *Int. J. Mod. Phys. A* 15 (2000) 1869.
5. A. Bouda, *Found. Phys. Lett.* 14 (2001) 17.
6. A. Bouda and T. Djama, *Phys. Lett. A* 285 (2001) 27.
7. D. Bohm, *Phys. Rev.* 85 (1952) 166; 85 (1952) 180; D. Bohm and J. P. Vigier, *Phys. Rev.* 96 (1954) 208.
8. E. R. Floyd, *Phys. Rev. D* 25 (1982) 1547.
9. E. R. Floyd, *Phys. Rev. D* 26 (1982) 1339.
10. E. R. Floyd, quant-ph/0009070.
11. A. Bouda and T. Djama, *Phys. Lett. A* 296 (2002) 312.
12. A. Bouda and T. Djama, *Physica Scripta* 66 (2002) 97.
13. A. E. Faraggi and M. Matone, *Phys. Lett. A* 249 (1998) 180.
14. A. Bouda and F. Hammad, *Acta Physica Slovaca* 52 (2002) 101.
15. E. R. Floyd, *Phys. Lett. A* 296 (2002) 307.
16. G. Salesi, *Int. J. Mod. Phys. A* 17 (2002) 347.
17. E. R. Floyd, *Int. J. Mod. Phys. A* 15 (2000) 1363.
18. L. Landau et E. Lifchitz, *Mécanique*, Tome I (Editions Mir, Moscou, 1981).