# Semiclassical quantisation rules for the Dirac and Pauli equations

Stefan Keppeler

Abteilung Theoretische Physik Universität Ulm Albert-Einstein-Allee 11 D-89069 Ulm, Germany Matematisk Fysik<sup>1</sup> Lunds Tekniska Högskola Lunds Universitet, Box 118 SE-22100 Lund, Sweden

#### Abstract

We derive explicit semiclassical quantisation conditions for the Dirac and Pauli equations. We show that the spin degree of freedom yields a contribution which is of the same order of magnitude as the Maslov correction in Einstein-Brillouin-Keller quantisation. In order to obtain this result a generalisation of the notion of integrability for a certain skew product flow of classical translational dynamics and classical spin precession has to be derived. Among the examples discussed is the relativistic Kepler problem with Thomas precession, whose treatment sheds some light on the amazing success of Sommerfeld's theory of fine structure [Ann. Phys. (Leipzig) **51** (1916) 1–91].

<sup>&</sup>lt;sup>1</sup>present address, email: stefan.keppeler@matfys.lth.se

### 1 Introduction

Semiclassical quantisation conditions provide the most direct link between the old quantum theory of Bohr and Sommerfeld on the one hand and wave mechanics on the other hand. Unlike other semiclassical tools, like trace formulae or the Van Vleck-Gutzwiller propagator [1], they do not only express quantum mechanical objects in terms of classical properties but also employ exactly the same formulation as was used by the old quantum theory, namely action quantisation.

Before the advent of quantum mechanics quantisation of a system was done by determining action integrals within the classical theory and setting these equal to an integer multiple of Planck's constant  $h = 2\pi\hbar$ , i.e. one required

$$\oint p \, \mathrm{d}x = 2\pi\hbar\,n\tag{1.1}$$

with integer n. This condition, originally put forward by Bohr [2] in order to understand the hydrogen spectrum, was first understood as the quantisation rule for one degree of freedom.

Around 1915 there was an ongoing discussion how this condition should be translated to more than one degree of freedom, see e.g. [3, 4, 5] and the introduction of [6]. Epstein [4]proposed to use that set of coordinates in which the problem separates if such coordinates exist. For each degree of freedom there would then be a condition of the form (1.1), i.e.

$$\oint p_j \,\mathrm{d}x_j = 2\pi\hbar\,n_j \tag{1.2}$$

with integers  $n_j$ , j numbering the degrees of freedom and  $(p_j, x_j)$  being a pair of canonically conjugate variables in that particular set of coordinates. Epstein's point of view was assumed by Sommerfeld who successfully applied this prescription treating many problems in spectroscopy [6].

Shortly after, Einstein [7] pointed out that separability of the equations of motion is not a necessary condition for action quantisation but that merely integrability (in the sense of Liouville and Arnold [8, 9]) is required: If there are sufficiently many integrals of motion with pairwise commuting Poisson brackets then the phase space foliates into invariant tori on which the line integral

$$\int_{\boldsymbol{x}_{i}}^{\boldsymbol{x}_{f}} \boldsymbol{p} \, \mathrm{d}\boldsymbol{x} \tag{1.3}$$

is locally path-independent. The quantisation conditions can then be written in the form

$$\oint_{\mathcal{C}_j} \boldsymbol{p} \, \mathrm{d}\boldsymbol{x} = 2\pi\hbar \, n_j \tag{1.4}$$

where  $\{C_1, \ldots, C_d\}$  denotes a basis of non-contractible loops on a given torus. This formulation has the advantage over Epstein's version of being independent of the coordinate system. Soon after the old quantum theory had been replaced by matrix and wave mechanics the old quantisation conditions were rederived and modified by Wentzel [10], Kramers [11], Brillouin [12] and Jeffreys [13] in a short-wavelength approximation, the so-called WKB or JWKB method. It was shown that depending on the character of the motion the quantum numbers may have to be shifted by a small number, leading e.g. to half-integer quantum numbers for oscillations but integer quantum numbers for rotations. Again the treatment was first for one-dimensional and then for separable systems. A complete derivation of the quantisation conditions from the Schrödinger equation that takes into account both the abstract integrability condition used by Einstein and the small shift of the quantum numbers is due to Keller [14]. He proved the semiclassical quantisation conditions

$$\oint_{\mathcal{C}_j} \boldsymbol{p} \, \mathrm{d}\boldsymbol{x} = 2\pi\hbar \left( n_j + \frac{\mu_j}{4} \right) \tag{1.5}$$

which are now known as Einstein-Brillouin-Keller (EBK) or torus quantisation. The number  $\mu_j \in \mathbb{Z}_4$  denotes the Maslov index, see e.g. [15, 16], a topological invariant of the cycle  $C_j$ . In one dimension it counts the number of turning points encountered along the loop. A good overview on these topics is given in [17].

The discussion so far was only for non-relativistic quantum mechanics, i.e the semiclassical approximation for the Schrödinger equation. In 1916 Sommerfeld applied the quantisation conditions (1.2) also in a relativistic context [6]. His aim was to find small corrections to the hydrogen spectrum which he expected to be due to relativistic effects. The success was overwhelming, the so-called Sommerfeld fine structure formula agreed excellently with the experimental data. More than ten years later [18, 19] it was found that the energy levels of the hydrogen atom, when calculated using the Dirac equation, the correct relativistic wave equation for the electron, are identical to the levels determined by Sommerfeld. The Dirac equation, however, does not only take into account relativistic effects but also the half-integer spin of the electron. The fine structure accounted for by the Sommerfeld formula is to a large extent due to spin-orbit coupling, an effect that was unknown at the time Sommerfeld did his calculations. In fact, even the property of spin itself was yet to be discovered. This seeming paradox has to be explained by a semiclassical analysis of the Dirac equation.

Early semiclassical approaches to the Dirac equation are due to Pauli [20] and Rubinow and Keller [21]. These will be discussed in section 3. The subtleties connected with such an approach are related to the fact that the Dirac equation is a partial differential equation for a spinor and not just for a scalar wave function. Therefore these problems should be discussed in the more general context of semiclassical (or short wavelength) approximations to multicomponent wave equations. It was observed by Yabana and Horiuchi [22] that the occurrence of geometrical or Berry phases [23, 24] plays an important rôle in this context. Kuratsuji and Iida [25, 26], using path integral methods, suggested that the symplectic structure of phase space should be deformed such that it includes the contribution of geometric phases. A review of these results is given in [27]. A general method for the semiclassical quantisation of multi-component wave equation was derived by Littlejohn and Flynn [28, 27]. Their results hold whenever the principal Weyl symbol of the Hamiltonian, which is a matrix valued function on classical phase, has non-degenerate eigenvalues. Thus it does not apply to the Dirac equation, in which case these eigenvalues have a multiplicity of two as will be shown in section 3. In such a situation the geometrical phases that appear in the semiclassical expressions are not simple U(1)-phases any longer but are themselves matrix-valued and thus in general non-Abelian. It was shown by Emmrich and Weinstein [29] that in this case integrability of the dynamics generated by the classical ray Hamiltonians (the eigenvalues of the principal Weyl symbol) is no longer sufficient to guarantee the existence of global semiclassical wave functions and thus of semiclassical quantisation conditions.

In this article we derive semiclassical quantisation conditions from the Dirac equation, the relativistic wave equation for particles with spin  $\frac{1}{2}$ , and from the Pauli equation, describing particles with arbitrary spin in a non-relativistic context. We show how to resolve the problems mentioned in the context with the occurrence of non-Abelian Berry phases by developing a generalisation of the notion of integrability that not only includes the dynamics of the ray Hamiltonians but imposes an additional condition. In this way we effectively reduce the non-Abelian phases to U(1)-phases. The latter enter the semiclassical quantisation conditions by a correction which is of the same order as the Maslov contribution in (1.5). This correction represents the influence of the spin degree of freedom and can be given a clear physical interpretation in terms of classical spin precession. By applying the method to the relativistic Kepler problem we shed some light on the success of Sommerfeld's fine structure formula. A brief account of some of these results was given in [30].

The paper is organised as follows. In section 2 we outline the derivation of EBKquantisation for later reference, thereby emphasising the rôle of integrability. Section 3 deals with the determination of semiclassical wave functions for the Dirac equation. In section 4 we generalise the concept of integrability to the case of group extensions and the skew product of classical translational dynamics and classical spin precession. Based on this characterisation we then derive explicit semiclassical quantisation conditions for the Dirac equation in section 5. In section 6 we show how these results translate to the Pauli equation with arbitrary spin. Before we treat some special examples in sections 8 and 9 (among which is Sommerfeld's theory of fine structure) we derive general formulae which facilitate the semiclassical quantisation of spherically symmetric systems in section 7. We conclude with a summary in section 10. Some important formulae for Weyl quantisation are listed in appendix A.

## 2 EBK quantisation

In this section we briefly summarise the main steps in the derivation of the EBK quantisation rules. We do so in order to introduce some notation and for later reference such that we can explicitly compare to this basic situation when treating systems with spin. This is, however, not intended to be a complete review of EBK quantisation and we thus refer the reader seeking a comprehensive introduction to EBK quantisation to the literature, e.g. [14, 17].

We want to find an asymptotic solution of the stationary Schrödinger equation,

$$\hat{H}\psi(\boldsymbol{x}) = E\psi(\boldsymbol{x}),$$
(2.1)

where  $\psi \in L^2(\mathbb{R}^d)$  and  $\hat{H}$  shall be a Weyl operator (some facts on Weyl quantisation are summarised in appendix A) with symbol

$$H(\boldsymbol{p},\boldsymbol{x}) = H_0(\boldsymbol{p},\boldsymbol{x}) + \hbar H_1(\boldsymbol{p},\boldsymbol{x}) + (\hbar^2), \quad \hbar \to 0.$$
(2.2)

The leading order term  $H_0$  in the semiclassical limit  $\hbar \to 0$  is known as the principal symbol of  $\hat{H}$  and  $H_1$  is called the sub-principal symbol. For simplicity we will choose  $H_1 \equiv 0$  for the rest of this section. For the familiar Hamiltonian

$$\hat{H} = -\frac{\hbar^2}{2m}\Delta + V(\boldsymbol{x}) \tag{2.3}$$

describing a particle of mass m moving under the influence of the external potential  $V(\boldsymbol{x})$  the Weyl symbol reads

$$H(\boldsymbol{p}, \boldsymbol{x}) \equiv H_0(\boldsymbol{p}, \boldsymbol{x}) = \frac{\boldsymbol{p}^2}{2m} + V(\boldsymbol{x})$$
(2.4)

and we can write  $\hat{H} = H(\frac{\hbar}{i}\nabla, \boldsymbol{x}).$ 

For the wave function  $\psi$  one tries the WKB ansatz

$$\psi_{\text{WKB}}(\boldsymbol{x}) = \sum_{k \ge 0} \left(\frac{\hbar}{i}\right)^k a_k(\boldsymbol{x}) e^{\frac{i}{\hbar}S(\boldsymbol{x})}.$$
(2.5)

Inserting into (2.1) yields in leading orders in  $\hbar$ , cf. appendix A,

$$[H(\nabla_{\boldsymbol{x}}S,\boldsymbol{x}) - E] a_0 + \frac{\hbar}{i} \Big\{ [H(\nabla_{\boldsymbol{x}}S,\boldsymbol{x}) - E] a_1 + [(\nabla_{\boldsymbol{p}}H)(\nabla_{\boldsymbol{x}}S,\boldsymbol{x})] (\nabla_{\boldsymbol{x}}a_0) + \frac{1}{2} [\nabla_{\boldsymbol{x}}(\nabla_{\boldsymbol{p}}H)(\nabla_{\boldsymbol{x}}S,\boldsymbol{x})] a_0 \Big\} + (\hbar^2) = 0.$$
(2.6)

This equation is easily confirmed for the particular Hamiltonian (2.3) by direct computation, but it also holds for arbitrary (semiclassical) Weyl operators, see appendix A. The strategy of the WKB method is now to satisfy eq. (2.6) separately order by order in  $\hbar$ . In leading order one finds the Hamilton-Jacobi equation

$$H(\nabla_{\boldsymbol{x}}S, \boldsymbol{x}) = E \tag{2.7}$$

of classical mechanics with the (principal) symbol  $H(\mathbf{p}, \mathbf{x})$  acting as the classical Hamiltonian. From standard Hamilton-Jacobi theory, see e.g. [31, 9], one thus concludes that the phase  $S(\mathbf{x})$  of the WKB ansatz becomes the classical action generating the dynamics

with Hamiltonian  $H(\boldsymbol{p}, \boldsymbol{x})$ : If  $(\boldsymbol{P}(t), \boldsymbol{X}(t))$  is a solution of Hamilton's equations of motion then  $\nabla_{\boldsymbol{x}} S(\boldsymbol{X}(t)) = \boldsymbol{P}(t)$ . A solution of the Hamilton-Jacobi equation (2.7) can thus be obtained by integration along solutions of Hamilton's equations of motion in the following way. Denote by  $\boldsymbol{y}$  an arbitrary point in configuration space and by  $\boldsymbol{\xi}$  a momentum satisfying  $H(\boldsymbol{\xi}, \boldsymbol{y}) = E$ . Let  $\phi_H^t$  be the (Hamiltonian) flow generated by the classical Hamiltonian  $H(\boldsymbol{p}, \boldsymbol{x})$ , i.e.  $\phi_H^t(\boldsymbol{\xi}, \boldsymbol{y}) =: (\boldsymbol{P}(t), \boldsymbol{X}(t))$  denotes the point reached at time t on the trajectory starting at  $(\boldsymbol{\xi}, \boldsymbol{y})$ . Then we have

$$S(\boldsymbol{x}) - S(\boldsymbol{y}) = \int_0^t \left[ \frac{\mathrm{d}}{\mathrm{d}t'} S(\boldsymbol{X}(t')) \right] \mathrm{d}t' = \int_0^t \nabla_{\!\!\boldsymbol{x}} S(\boldsymbol{X}(t')) \, \dot{\boldsymbol{X}}(t') \, \mathrm{d}t' = \int_{\boldsymbol{y}}^{\boldsymbol{x}} \boldsymbol{P} \, \mathrm{d}\boldsymbol{X} \qquad (2.8)$$

where in the last expression integration is along the trajectory  $\phi_H^t(\boldsymbol{\xi}, \boldsymbol{y})$ . Finally the action reads

$$S(\boldsymbol{x}) = S(\boldsymbol{y}) + \int_{\boldsymbol{y}}^{\boldsymbol{x}} \boldsymbol{P} \, \mathrm{d}\boldsymbol{X} \,, \qquad (2.9)$$

where  $S(\boldsymbol{y})$  is the arbitrarily chosen value of S at the point  $\boldsymbol{y}$ . Given a solution (2.9) of the Hamilton-Jacobi equation the next-to-leading order equation deriving from (2.6) reduces to

$$[(\nabla_{\boldsymbol{p}} H)(\nabla_{\boldsymbol{x}} S, \boldsymbol{x})](\nabla_{\boldsymbol{x}} a_0) + \frac{1}{2} [\nabla_{\boldsymbol{x}} (\nabla_{\boldsymbol{p}} H)(\nabla_{\boldsymbol{x}} S, \boldsymbol{x})] a_0 = 0.$$
(2.10)

This is known as the transport equation for the leading order amplitude  $a_0(\boldsymbol{x})$ . Due to Hamilton's equations of motion the first term can now be interpreted as a time derivative along the the trajectory  $\phi_H^t(\boldsymbol{\xi}, \boldsymbol{y})$  which we shall denote by  $\frac{d}{dt}$  or simply by a dot,

$$\dot{a} \equiv \frac{\mathrm{d}a}{\mathrm{d}t} := \left[ (\nabla_{\boldsymbol{p}} H) (\nabla_{\boldsymbol{x}} S, \boldsymbol{x}) \right] (\nabla_{\boldsymbol{x}} a) \,. \tag{2.11}$$

The solution of (2.10) is locally given by

$$a_0(\boldsymbol{x}) = \sqrt{\det \frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}}},$$
 (2.12)

see e.g. [14, 17]. Together with (2.9) one has thus found an approximate solution

$$\psi_{\text{WKB}}(\boldsymbol{x}) \sim \sqrt{\det \frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}}} \exp\left(\frac{\mathrm{i}}{\hbar}S(\boldsymbol{y}) + \frac{\mathrm{i}}{\hbar}\int_{\boldsymbol{y}}^{\boldsymbol{x}} \boldsymbol{P} \,\mathrm{d}\boldsymbol{X}\right)$$
 (2.13)

of the Schrödinger equation for points  $\boldsymbol{x}$  in a neighbourhood of  $\boldsymbol{y}$  that are visited along a a solution of Hamilton's equations of motion starting at  $\boldsymbol{y}$ .

However, one still needs to find a way to integrate the Hamilton-Jacobi equation along paths transversal to  $\phi_H^t$ . Furthermore, the approximation (2.13) breaks down at points where  $\phi_H^t(\boldsymbol{\xi}, \boldsymbol{y})$  touches a caustic and thus  $\frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}}$  becomes singular. For one degree of freedom these points are given by the turning points. In order to address both these problems one has to know more about the classical phase space structure. For arbitrary Hamiltonians one can in general not proceed far beyond this point as was already pointed out by Einstein [7] in the context of the old quantum theory. Instead one has to invoke the concept of integrability.

Following Liouville [8] we say that a Hamiltonian is integrable if there are d constants of motion  $A_1 := H, A_2, \ldots, A_d$  which are in involution, i.e. whose Poisson brackets vanish pairwise. A complete proof of the consequences of this definition is due to Arnold whose version [9, chapter 10] we shall quote here.

**Theorem 1.** (Liouville-Arnold) Suppose that we are given d functions in involution on a d-dimensional symplectic manifold

$$A_1, \dots, A_d, \quad \{A_j, A_k\} \equiv 0, \quad j, k = 1, \dots, d.$$
 (2.14)

Consider a level set of the functions  $A_i$ ,

$$M_{a} = \{ (\boldsymbol{p}, \boldsymbol{x}) \mid A_{j}(\boldsymbol{p}, \boldsymbol{x}) = a_{j}, \ j = 1, \dots, d \}.$$
(2.15)

Assume that the d functions  $A_j$  are independent on  $M_a$  (i.e. the d 1-forms  $dA_j$  are linearly independent at each point of  $M_a$ ). Then

- 1.  $M_{a}$  is a smooth manifold, invariant under the phase flow with Hamiltonian function  $H = A_{1}$ .
- 2. If the manifold  $M_{\mathbf{a}}$  is compact and connected, then it is diffeomorphic to the ddimensional torus

$$\mathbb{T}^d = \{ (\vartheta_j, \dots, \vartheta_d) \mod 2\pi \}.$$
(2.16)

3. The phase flow with Hamiltonian function H determines a conditionally periodic motion on  $M_{\mathbf{a}}$ , i.e. in angular coordinates  $\boldsymbol{\vartheta} = (\vartheta_j, \ldots, \vartheta_d)$  we have

$$\frac{\mathrm{d}\boldsymbol{\vartheta}}{\mathrm{d}t} = \boldsymbol{\omega}, \quad \boldsymbol{\omega} = \boldsymbol{\omega}(\boldsymbol{a}).$$
 (2.17)

4. The canonical equations with Hamiltonian function H can be integrated by quadratures.

We refrain from providing a proof of the theorem here but refer the reader to Arnold's book [9]. Instead we remark on some aspects which are relevant for the following sections.

In order to prove property 2 one first shows that the conditions (2.14) imply that the flows  $\phi_j^t$  generated by the observables  $A_j$ ,  $j = 1, \ldots, d$ , commute on  $M_a$ , i.e.

$$\phi_j^t \circ \phi_k^{t'} = \phi_k^{t'} \circ \phi_j^t \quad \forall \ j, k = 1, \dots, d.$$

$$(2.18)$$

this yields a transitive action of  $\mathbb{R}^d$  on  $M_a$ . In the following we will refer to  $M_a$  as a Liouville-Arnold torus.

In general the coordinates  $\vartheta$  and the observables A are not canonically conjugate. However, (locally) there exists a mapping  $A \mapsto I$  such that  $(I, \vartheta)$  form a set of canonically conjugate variables. The new constants of motion I(A) are called action variables and the explicit construction of action and angle variables  $(I, \vartheta)$  is the desired integration of of Hamilton's equations of motion by quadratures. The Hamiltonian expressed in the new variables becomes a function  $\overline{H}(I)$  of the action variables only and thus the frequencies in (2.17) are given by  $\boldsymbol{\omega} = \nabla_{I} \overline{H}(I)$ .

Theorem 1 can be used in order to derive the EBK quantisation conditions as follows. Since the flows  $\phi_j^t$  commute we can define the action  $S(\boldsymbol{x})$  analogously to (2.9) by integration along the flow lines of  $\phi_2^t, \ldots, \phi_d^t$  instead of  $\phi_1^t \equiv \phi_H^t$ . To this end define the multi-time flow

$$\Phi^{\boldsymbol{t}} := \phi_d^{\boldsymbol{t}_d} \circ \dots \circ \phi_1^{\boldsymbol{t}_1} \,, \tag{2.19}$$

where due to (2.18) the ordering is unimportant. Since  $\Phi^t$  is a transitive action of  $\mathbb{R}^d$  on a Liouville-Arnold torus, for any  $\boldsymbol{x}$  in a small neighbourhood of  $\boldsymbol{y}$  there is a unique  $\boldsymbol{t} \in \mathbb{R}^d$  such that

$$\Phi^t(\boldsymbol{\xi}, \boldsymbol{y}) = (\boldsymbol{p}, \boldsymbol{x}) \tag{2.20}$$

with some momentum p. The rapidly oscillating phase of the WKB wave function is then given by

$$S(\boldsymbol{x}) = S(\boldsymbol{y}) + \int_{\boldsymbol{y}}^{\boldsymbol{x}} \boldsymbol{P} \, \mathrm{d}\boldsymbol{X}$$
(2.21)

where integration is along  $\Phi^t(\boldsymbol{\xi}, \boldsymbol{y})$ . Since the flows  $\phi_j^t$ ,  $j = 1, \ldots, d$ , commute this is not in conflict with the requirement of S solving the Hamilton-Jacobi equation (2.7). This answers the question of how to integrate transversal to  $\phi_H^t$ .

One still has to solve the problem of  $a_0 = \sqrt{\det \frac{\partial y}{\partial x}}$  becoming singular for certain values of t. To overcome this difficulty one has to glue together various local solutions of the form (2.13). Here the crucial observation is that whenever an eigenvalue of  $\frac{\partial x}{\partial y}$  changes sign (causing  $\frac{\partial y}{\partial x}$  to become singular) this results in a phase jump of the wave function by  $-\frac{\pi}{2}$ . For a closed curve C the number of times this happens along C is a topological invariant of C, its Maslov index  $\mu$ , see [15, 16]. On a d-torus  $\mathbb{T}^d$  let us choose a set of basis cycles  $\{C_j\}$ ,  $j = 1, \ldots, d$ , such that along  $C_j$  the angle  $\vartheta_j$  increases by  $2\pi$ . and all other angles  $\vartheta_{k\neq j}$ remain constant. See figure 1 for an illustration of this basis for a 2-torus. Then every closed curve on  $\mathbb{T}^d$  is a linear combination of the basis cycles  $C_j$ .

All we have to do now in order to get a globally well-defined WKB wave function is to make sure that  $\psi_{\text{WKB}}(\boldsymbol{x})$  returns to its initial value when we follow its value along  $C_j$ . In other words the phase change along a loop has to be an integer multiple of  $2\pi$ , i.e.

$$\frac{1}{\hbar} \oint_{\mathcal{C}_j} \boldsymbol{P} \,\mathrm{d}\boldsymbol{X} - \mu_j \frac{\pi}{2} = 2\pi \, n_j \,, \quad n_j \in \mathbb{Z} \,.$$
(2.22)

These are the quantisation conditions (1.5). Since the action variables I are given by

$$I_j = \frac{1}{2\pi} \oint_{\mathcal{C}_j} \boldsymbol{P} \,\mathrm{d}\boldsymbol{X}$$
(2.23)



Figure 1: Sketch of a basis  $\{C_1, C_2\}$  of loops on a two-torus  $\mathbb{T}^2$ .

the EBK energies read

$$E_{\boldsymbol{n}} = \overline{H} \left( \hbar \left( \boldsymbol{n} + \frac{\boldsymbol{\mu}}{4} \right) \right) \tag{2.24}$$

where  $\overline{H}$  is the Hamiltonian transformed to action and angle variables and  $\mu_j$  denotes the Maslov index of the basis cycle  $C_j$ . Further restrictions on the values which the integers  $\boldsymbol{n}$  may assume usually arise from the possible values the action variables  $\boldsymbol{I}$  can take in the particular problem of classical mechanics, cf. the examples given below.

## 3 The semiclassical wave function for the Dirac equation

In this section we review the determination of semiclassical wave functions for the Dirac equation and introduce the relevant notation for the following sections.

The first steps towards a semiclassical wave function for the Dirac equation were undertaken by Pauli [20]. He inserted an ansatz similar to (2.5) into the Dirac equation and found that the phase function S has to solve a relativistic Hamilton-Jacobi equation. He then solved the analogue of the transport equation (2.10) for a particular case but did not derive a general expression like (2.12). The problem was taken up again many years later by Rubinow and Keller [21] who proceeded one step further. They showed that the solution of the transport equation can be related to the Thomas precession [32, 33], see also [34], of a classical spin vector. However, also in this work no general quantisation conditions similar to (1.5) were given. As we will see in the following sections their construction is complicated by the occurrence of non-Abelian Berry phases for which additional integrability conditions are needed.

We also list some related literature which we will, however, not directly refer to in the following: Semiclassical approximations to the radial Dirac equation were studied in [35, 36, 37]. The semiclassical time evolution of the Dirac equation was examined in [38]. Semiclassical quantisation of subspectra of the Dirac Hamiltonian based on the complex germ method was discussed in [39, 40]. The time evolution of semiclassical Wigner functions for the Dirac equation is addressed in [41].

We briefly repeat the basic steps in the derivation of semiclassical wave functions for the Dirac equation. For details we refer the reader to [21] or to [42] where the notation is similar to that used here.

The aim is to find asymptotic solutions to the stationary Dirac equation

$$\hat{H}_{\rm D}\Psi(\boldsymbol{x}) = E\Psi(\boldsymbol{x}) \tag{3.1}$$

with Dirac Hamiltonian

$$\hat{H}_{\rm D} = c\boldsymbol{\alpha} \left(\frac{\hbar}{\rm i} \nabla - \frac{e}{c} \boldsymbol{A}(\boldsymbol{x})\right) + \beta m c^2 + e\phi(\boldsymbol{x})$$
(3.2)

in the semiclassical limit  $\hbar \to 0$ . The wave function is now a four-spinor,  $\Psi \in L^2(\mathbb{R}^3) \otimes \mathbb{C}^4$ , and the 4 × 4 matrices  $\alpha$  and  $\beta$  are given by

$$\boldsymbol{\alpha} = \begin{pmatrix} 0 & \boldsymbol{\sigma} \\ \boldsymbol{\sigma} & 0 \end{pmatrix}, \quad \boldsymbol{\beta} = \begin{pmatrix} \mathbb{1}_2 & 0 \\ 0 & -\mathbb{1}_2 \end{pmatrix}, \quad (3.3)$$

where each entry is to be understood as a  $2 \times 2$  matrix. The Pauli matrices  $\sigma$  are given by

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$
(3.4)

and  $\mathbb{1}_n$  denotes the  $n \times n$  unit matrix. The Dirac equation describes a particle of mass m and charge e moving under the influence of the external electro-magnetic potentials  $(\phi(\boldsymbol{x}), \boldsymbol{A}(\boldsymbol{x}))$ , i.e. we have fixed a frame of reference in which the potentials are static.

We now modify the semiclassical ansatz (2.5) such that the amplitudes  $a_k(\boldsymbol{x})$  take values in  $\mathbb{C}^4$  but the phase  $S(\boldsymbol{x})$  is kept scalar. Inserting this ansatz into the Dirac equation (3.1) yields

$$\left[H_{\mathrm{D}}(\nabla_{\boldsymbol{x}}S,\boldsymbol{x}) - E\right]a_0 + \frac{\hbar}{\mathrm{i}}\left\{\left[H_{\mathrm{D}}(\nabla_{\boldsymbol{x}}S,\boldsymbol{x}) - E\right]a_1 + c\boldsymbol{\alpha}(\nabla_{\boldsymbol{x}}a_0)\right\} + (\hbar^2) = 0, \quad (3.5)$$

where  $H_{\rm D}(\boldsymbol{p}, \boldsymbol{x})$  denotes the Weyl symbol of the Dirac Hamiltonian (3.2),

$$H_{\rm D}(\boldsymbol{p}, \boldsymbol{x}) = \begin{pmatrix} mc^2 + e\phi(\boldsymbol{x}) & \boldsymbol{\sigma} \left(c\boldsymbol{p} - e\boldsymbol{A}(\boldsymbol{x})\right) \\ \boldsymbol{\sigma} \left(c\boldsymbol{p} - e\boldsymbol{A}(\boldsymbol{x})\right) & mc^2 - e\phi(\boldsymbol{x}) \end{pmatrix}.$$
(3.6)

When comparing (3.5) with (2.6) notice that  $c\boldsymbol{\alpha} = \nabla_{\boldsymbol{p}} H_{\rm D}$ . Since the leading order equation

$$[H_{\rm D}(\nabla_{\!\boldsymbol{x}} S, \boldsymbol{x}) - E] a_0 = 0 \tag{3.7}$$

is a matrix equation it implies the necessary condition that the expression in square brackets has an eigenvalue which vanishes identically, i.e.

$$H^{\pm}(\nabla_{\!\boldsymbol{x}}S,\boldsymbol{x}) = E, \qquad (3.8)$$

where the eigenvalues

$$H^{\pm}(\boldsymbol{p}, \boldsymbol{x}) = e\phi(\boldsymbol{x}) \pm \sqrt{(c\boldsymbol{p} - e\boldsymbol{A}(\boldsymbol{x}))^2 + m^2 c^4}$$
(3.9)

of  $H_{\rm D}$  act as classical Hamiltonians for our problem. In the following we will use the notation  $S^{\pm}$  indicating which of the two Hamilton-Jacobi equations (3.8) is solved by the respective phase function; the solution itself is again given by (2.9) where now the integration is along the flow lines of  $\phi_{H^{\pm}}^t$ . Each of the eigenvalues (3.9) has a multiplicity of two and the corresponding eigenvectors can be chosen as the columns of the  $4 \times 2$  matrices  $V_{\pm}$ ,

$$V_{+}(\boldsymbol{p}, \boldsymbol{x}) = \frac{1}{\sqrt{2\varepsilon(\varepsilon + mc^{2})}} \begin{pmatrix} \varepsilon + mc^{2} \\ \boldsymbol{\sigma}(c\boldsymbol{p} - e\boldsymbol{A}(\boldsymbol{x})) \end{pmatrix},$$
  
$$V_{-}(\boldsymbol{p}, \boldsymbol{x}) = \frac{1}{\sqrt{2\varepsilon(\varepsilon + mc^{2})}} \begin{pmatrix} \boldsymbol{\sigma}(c\boldsymbol{p} - e\boldsymbol{A}(\boldsymbol{x})) \\ -(\varepsilon + mc^{2}) \end{pmatrix},$$
(3.10)

where we have introduced the abbreviation

$$\varepsilon(\boldsymbol{p}, \boldsymbol{x}) := \sqrt{(c\boldsymbol{p} - e\boldsymbol{A}(\boldsymbol{x}))^2 + m^2 c^4}.$$
(3.11)

With this choice the eigenvectors are orthonormal and complete, i.e.

$$V_{\pm}^{\dagger}(\boldsymbol{p}, \boldsymbol{x}) V_{\pm}(\boldsymbol{p}, \boldsymbol{x}) = \mathbb{1}_{2}, \quad V_{\mp}^{\dagger}(\boldsymbol{p}, \boldsymbol{x}) V_{\pm}(\boldsymbol{p}, \boldsymbol{x}) = 0,$$
  
and  $V_{+}(\boldsymbol{p}, \boldsymbol{x}) V_{+}^{\dagger}(\boldsymbol{p}, \boldsymbol{x}) + V_{-}(\boldsymbol{p}, \boldsymbol{x}) V_{-}^{\dagger}(\boldsymbol{p}, \boldsymbol{x}) = \mathbb{1}_{4}.$  (3.12)

Now  $S^{\pm}(\boldsymbol{x})$  solving one of the Hamilton-Jacobi equations (3.8) is not sufficient in order to satisfy eq. (3.7). Instead we also need  $a_0^{\pm}(\boldsymbol{x})$  to be of the form

$$a_0^{\pm}(\boldsymbol{x}) = V_{\pm}(\nabla_{\boldsymbol{x}}S, \boldsymbol{x}) b^{\pm}(\boldsymbol{x})$$
(3.13)

with the still unknown function  $b^{\pm}(\boldsymbol{x})$  taking values in  $\mathbb{C}^2$ .

An equation for  $b^{\pm}$  can be obtained from the next-to-leading order equation deriving from (3.5) by inserting (3.13) and multiplying with  $V_{\pm}^{\dagger}(\nabla_{\!\!\boldsymbol{x}} S, \boldsymbol{x})$  from the left yielding

$$cV_{\pm}^{\dagger}(\nabla_{\boldsymbol{x}}S,\boldsymbol{x})\boldsymbol{\alpha}V(\nabla_{\boldsymbol{x}}S,\boldsymbol{x})(\nabla_{\boldsymbol{x}}b^{\pm}) + cV_{\pm}^{\dagger}(\nabla_{\boldsymbol{x}}S,\boldsymbol{x})\boldsymbol{\alpha}[(\nabla_{\boldsymbol{x}}V)(\nabla_{\boldsymbol{x}}S,\boldsymbol{x})]b^{\pm} = 0.$$
(3.14)

After some algebra one arrives at

$$\left[ (\nabla_{\boldsymbol{p}} H^{\pm}) (\nabla_{\boldsymbol{x}} S, \boldsymbol{x}) \right] (\nabla_{\boldsymbol{x}} b^{\pm}) + \frac{1}{2} \left[ \nabla_{\boldsymbol{x}} (\nabla_{\boldsymbol{p}} H^{\pm}) (\nabla_{\boldsymbol{x}} S^{\pm}, \boldsymbol{x}) \right] b^{\pm} + \frac{\mathrm{i}}{2} \boldsymbol{\sigma} \boldsymbol{\mathcal{B}}^{\pm} (\nabla_{\boldsymbol{x}} S^{\pm}, \boldsymbol{x}) b^{\pm} = 0$$

$$\text{with} \quad \boldsymbol{\mathcal{B}}^{\pm} (\boldsymbol{p}, \boldsymbol{x}) = \mp \frac{ec}{\varepsilon} \boldsymbol{B}(\boldsymbol{x}) + \frac{ec}{\varepsilon(\varepsilon + mc^2)} \left[ c\boldsymbol{p} - e\boldsymbol{A}(\boldsymbol{x}) \right] \times \boldsymbol{E}(\boldsymbol{x}) .$$

$$(3.15)$$

Here we have introduced the electric and magnetic fields

$$\boldsymbol{E}(\boldsymbol{x}) = -\nabla \phi(\boldsymbol{x}) \text{ and } \boldsymbol{B}(\boldsymbol{x}) = \nabla \times \boldsymbol{A}(\boldsymbol{x}).$$
 (3.16)

Equation (3.15) is of the same form as the scalar transport equation (2.10) except for the extra term  $\frac{i}{2}\sigma \mathcal{B}^{\pm} b^{\pm}$ . Thus we can exploit our knowledge of how to solve (2.10) by making the ansatz

$$b^{\pm}(\boldsymbol{x}) = \sqrt{\det \frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}}} u^{\pm}(\boldsymbol{x})$$
 (3.17)

leaving us with the spin transport equation

$$\dot{u}^{\pm} + \frac{\mathrm{i}}{2} \boldsymbol{\sigma} \boldsymbol{\mathcal{B}}^{\pm} (\nabla_{\!\!\boldsymbol{x}} S^{\pm}, \boldsymbol{x}) \, u^{\pm} = 0 \tag{3.18}$$

for the  $\mathbb{C}^2$ -valued function  $u^{\pm}(\boldsymbol{x})$ , where the dot denotes a derivative along the Hamiltonian flow  $\phi_{H^{\pm}}^t$ , cf. eq. (2.11).

Given an initial value  $u^{\pm}(\boldsymbol{y})$  at a point  $\boldsymbol{y}$  its value at a point  $\boldsymbol{x}$ , which is connected to  $\boldsymbol{y}$  by the trajectory  $\phi_{H^{\pm}}^{t}(\boldsymbol{\xi}, \boldsymbol{y})$ , is given by

$$u^{\pm}(\boldsymbol{x}) = d_{\pm}(\boldsymbol{\xi}, \boldsymbol{y}, t) \, u^{\pm}(\boldsymbol{y}) \,, \qquad (3.19)$$

where  $d_{\pm}(\boldsymbol{\xi}, \boldsymbol{y}, t)$  is a 2 × 2 matrix. We have explicitly indicated the dependence of  $d_{\pm}$  on the initial point in phase space  $(\boldsymbol{\xi}, \boldsymbol{y})$  where we start the integration and the time t until which we proceed. Clearly,  $d_{\pm}$  also has to solve a spin transport equation,

$$\dot{d}_{\pm}(\boldsymbol{\xi},\boldsymbol{y},t) + \frac{\mathrm{i}}{2}\boldsymbol{\sigma}\boldsymbol{\mathcal{B}}^{\pm}(\phi_{H^{\pm}}^{t}(\boldsymbol{\xi},\boldsymbol{y})) d_{\pm}(\boldsymbol{\xi},\boldsymbol{y},t) = 0, \quad d_{\pm}(\boldsymbol{\xi},\boldsymbol{y},0) = \mathbb{1}_{2}.$$
(3.20)

Since the coefficient  $\frac{i}{2} \boldsymbol{\sigma} \boldsymbol{\mathcal{B}}^{\pm}$  takes values in the Lie algebra  $\mathfrak{su}(2)$  it follows that  $d_{\pm}(\boldsymbol{\xi}, \boldsymbol{y}, 0) \in$  SU(2). More precisely, the matrix valued function  $d_{\pm}$  is an SU(2)-valued cocycle of the flow  $\phi_H^t$  as one easily verifies the composition law

$$d_{\pm}(\boldsymbol{\xi}, \boldsymbol{y}, t+t') = d_{\pm}(\phi_{H^{\pm}}^{t}(\boldsymbol{\xi}, \boldsymbol{y}), t') d_{\pm}(\boldsymbol{\xi}, \boldsymbol{y}, t) .$$
(3.21)

Accordingly we can define the skew product flow [43]

$$Y_{\pm}^{t}: \mathbb{R}^{d} \times \mathbb{R}^{d} \times \mathrm{SU}(2) \to \mathbb{R}^{d} \times \mathbb{R}^{d} \times \mathrm{SU}(2)$$
  
( $\boldsymbol{p}, \boldsymbol{x}, g$ )  $\mapsto \left(\phi_{H^{\pm}}^{t}(\boldsymbol{p}, \boldsymbol{x}), d_{\pm}(\boldsymbol{p}, \boldsymbol{x}, t)g\right)$  (3.22)

which preserves the product of Liouville measure on phase space  $\mathbb{R}^d \times \mathbb{R}^d$  and Haar measure on SU(2), see [43]. Since the cocycle  $d_{\pm}$  takes values in the group G = SU(2) this construction is also known as a group extension or, more precisely as an SU(2)-extension.

At this point the spin degree of freedom is still described on a quantum mechanical level in the sense that it is represented by elements of  $\mathbb{C}^2$  which are evolved in time by an SU(2)-valued propagator. It is, however, possible to switch to a purely classical description. To this end consider the adjoint representation of SU(2) defined by

$$\begin{aligned} \operatorname{Ad}_{g} : \mathfrak{su}(2) &\to \mathfrak{su}(2) \\ Z &\to gZg^{-1} \,. \end{aligned} \tag{3.23}$$

Since an element of the Lie algebra  $\mathfrak{su}(2)$  can be written as a linear combination of the Pauli matrices,  $Z = \mathbf{z}\boldsymbol{\sigma}, \mathbf{z} \in \mathbb{R}^3$ , expanding also the right hand side of (3.23) in this basis,  $gZg^{-1} = (\varphi(g)\mathbf{z})\boldsymbol{\sigma}$ , provides us with a rotation matrix  $\varphi(g)$ . The map  $\varphi : \mathrm{SU}(2) \to \mathrm{SO}(3)$  is two-to-one and known as the covering map. Anticipating classical spin as a vector of constant length let us consider  $\mathbf{s}_0 \in S^2 \hookrightarrow \mathbb{R}^3$ . On easily verifies that  $\mathbf{s} = \varphi(d(\boldsymbol{\xi}, \boldsymbol{y}, t)) \mathbf{s}_0$  solves

$$\dot{\boldsymbol{s}} = \boldsymbol{\mathcal{B}}^{\pm}(\phi_{H^{\pm}}^{t}(\boldsymbol{\xi}, \boldsymbol{y})) \times \boldsymbol{s}$$
(3.24)

i.e. the equation of Thomas precession [32, 33], which has thus been derived from the Dirac equation [21, 44, 42, 41]. Classical spin precession (3.24) and the Hamiltonian flow  $\phi_{H^{\pm}}^{t}$  can now be combined into the classical skew product [45]

$$Y_{cl\pm}^{t}: \mathbb{R}^{d} \times \mathbb{R}^{d} \times S^{2} \to \mathbb{R}^{d} \times \mathbb{R}^{d} \times S^{2}$$

$$(\boldsymbol{p}, \boldsymbol{x}, \boldsymbol{s}) \mapsto \left(\phi_{H^{\pm}}^{t}(\boldsymbol{p}, \boldsymbol{x}), \varphi(d_{\pm}(\boldsymbol{p}, \boldsymbol{x}, t))\boldsymbol{s}\right).$$

$$(3.25)$$

As opposed to  $Y_{\pm}^t$  this is a symplectic flow conserving the product of Liouville measure on the phase space  $\mathbb{R}^d \times \mathbb{R}^d$  of the translational degrees of freedom and the surface element on the sphere  $S^2$  which acts as the phase space for the spin degree of freedom. As we will see below the properties of  $Y_{cl\pm}^t$  determine the semiclassical solutions of the Dirac equation as the properties of  $\phi_{H^{\pm}}^t$  determine that of the Schrödinger equation.

Collecting the results of the previous paragraphs, so far we have obtained the following expression for the semiclassical wave function for the Dirac equation,

$$\Psi_{\rm sc}^{\pm}(\boldsymbol{x}) \sim \sqrt{\det \frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}}} e^{\frac{i}{\hbar}S^{\pm}(\boldsymbol{x})} V_{\pm}(\nabla_{\!\!\boldsymbol{x}}S^{\pm}, \boldsymbol{x}) d_{\pm}(\boldsymbol{\xi}, \boldsymbol{y}, t) u^{\pm}(\boldsymbol{y}).$$
(3.26)

The new ingredients as compared to (2.13) are the projection matrices  $V_{\pm}$  selecting either positive or negative kinetic energies and the spinor part  $d_{\pm}(\boldsymbol{\xi}, \boldsymbol{y}, t) u^{\pm}(\boldsymbol{y})$ . We can now explain why (3.26) does not immediately lead to a generalisation of the EBK quantisation conditions (2.22). Assume that the Hamiltonian flow generated by either  $H^+(\boldsymbol{p}, \boldsymbol{x})$  or  $H^-(\boldsymbol{p}, \boldsymbol{x})$  is integrable in the sense of Theorem 1. Then we know that the factors  $\sqrt{\det \frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}}}$ and  $e^{\frac{i}{\hbar}S^{\pm}(\boldsymbol{x})}$  can be defined globally by the construction described in the preceeding section. This also applies to the projection matrix  $V_{\pm}$  which simply has to be evaluated along the respective path of integration; in particular, for a closed path also the value of  $V_{\pm}$  returns to its initial value thus not contributing to the quantisation conditions.

In contrast, integrating the spin transport equation (3.20) along a closed path  $C_j$  will yield an SU(2) matrix  $d_j$ . Thus the initial and final values of the semiclassical wave function (3.26) would not just differ by a phase factor but by an SU(2) transformation which would have to be compensated for in order to turn  $\Psi_{sc}(\boldsymbol{x})$  into a globally single-valued object. Even worse, integration along a different loop  $\mathcal{C}_{k\neq j}$  (which we still have to define, so far we are only able to integrate (3.20) along the flow lines of  $\phi_{H^{\pm}}^{t}$ ) will in general yield a different SU(2) matrix  $d_{k\neq j}$  which need not commute with the first one. This can make it impossible to find a globally well-defined semiclassical solution as was pointed out by Emmrich and Weinstein in a general setting [29]. From the point of view of physics this is not surprising. We know that in order to be able to use EBK quantisation the corresponding classical system has to be integrable. Now we are dealing with a system that has an additional degree of freedom, namely spin. So far, however, we have not imposed any condition on the spin dynamics but we have only required the Hamiltonian flow  $\phi_{H^{\pm}}^{t}$  to be integrable. In the following section we will show how the notion of integrability can be extended to skew products of the form (3.22) or (3.25) and prove a generalisation of Theorem 1.

## 4 Integrability of skew products

From the point of view of semiclassics the crucial aspect in the characterisation of integrable systems by Theorem 1 is the geometrical description of the invariant manifolds which is a consequence of the existence of commuting flows  $\phi_1^t, \ldots, \phi_d^t$ . Therefore, one has to find a generalisation of the condition (2.14). We will show that in this sense the following definition provides a good generalisation of the notion of integrability.

**Definition 2.** The skew product  $Y_{cl\pm}^t$  is called integrable, if

(i) the underlying Hamiltonian flow φ<sup>t</sup><sub>H<sup>±</sup></sub> is integrable in the sense of Liouville and Arnold (Theorem 1), i.e. besides the Hamiltonian H<sup>±</sup>(**p**, **x**) =: A<sub>1</sub>(**p**, **x**) there are d−1 more independent integrals of motion, A<sub>2</sub>(**p**, **x**),..., A<sub>d</sub>(**p**, **x**) with

$$\{A_j, A_k\} = 0 \quad \forall \ j, k = 1, \dots, d$$
(4.1)

and

(ii) the flows  $\phi_2^t, \ldots, \phi_d^t$  can also be extended to skew products  $Y_{cl_j}^t$  on  $\mathbb{R}^d \times \mathbb{R}^d \times S^2$  $(Y_{cl\pm}^t \equiv Y_{cl_1}^t)$  with fields  $\mathcal{B}_j(\boldsymbol{p}, \boldsymbol{x})$ , i.e

$$Y_{\text{cl}j}^{t}(\boldsymbol{p}, \boldsymbol{x}, \boldsymbol{s}) = \left(\phi_{j}^{t}(\boldsymbol{p}, \boldsymbol{x}), \varphi(d_{j}(\boldsymbol{p}, \boldsymbol{x}, t))\boldsymbol{s}\right)$$
(4.2)

$$\dot{d}_j(\boldsymbol{p},\boldsymbol{x},t) + \frac{\mathrm{i}}{2}\boldsymbol{\sigma}\boldsymbol{\mathcal{B}}_j(\phi_j^t(\boldsymbol{p},\boldsymbol{x})) \, d_j(\boldsymbol{p},\boldsymbol{x},t) = 0 \,, \quad d_j(\boldsymbol{p},\boldsymbol{x},0) = \mathbb{1}_2 \,, \tag{4.3}$$

fulfilling

$$\{A_j, \mathcal{B}_k\} + \{\mathcal{B}_j, A_k\} - \mathcal{B}_j \times \mathcal{B}_k = 0 \quad \forall \ j, k = 1, \dots, d.$$

$$(4.4)$$

In view of what was said above let us first show that this is indeed a good definition by the following lemma. **Lemma 3.** Two skew products  $Y_{cl_j}^t$  and  $Y_{cl_k}^t$  of the type (4.2) commute if and only if the corresponding base flows  $\phi_j^t$  and  $\phi_k^t$  commute and if the fields  $\mathcal{B}_j$  and  $\mathcal{B}_k$  fulfil

$$\{A_j, \mathcal{B}_k\} + \{\mathcal{B}_j, A_k\} - \mathcal{B}_j \times \mathcal{B}_k = 0.$$
(4.5)

PROOF: Since the two skew products can only commute if the corresponding base flows commute it remains to show that

$$\varphi\left(d_j(\phi_k^t(\boldsymbol{p},\boldsymbol{x}),t')\right)\varphi(d_k(\boldsymbol{p},\boldsymbol{x},t))\,\boldsymbol{s} = \varphi\left(d_k(\phi_j^{t'}(\boldsymbol{p},\boldsymbol{x}),t)\right)\varphi(d_j(\boldsymbol{p},\boldsymbol{x},t'))\,\boldsymbol{s} \tag{4.6}$$

for all  $(\boldsymbol{p}, \boldsymbol{x}, \boldsymbol{s}) \in \mathbb{R}^d \times \mathbb{R}^d \times S^2$  and all  $t, t' \in \mathbb{R}$ , or equivalently

$$\varphi\left(d_j(\phi_k^t(\boldsymbol{p},\boldsymbol{x}),t')\right)\varphi(d_k(\boldsymbol{p},\boldsymbol{x},t)) = \varphi\left(d_k(\phi_j^{t'}(\boldsymbol{p},\boldsymbol{x}),t)\right)\varphi(d_j(\boldsymbol{p},\boldsymbol{x},t'))$$
(4.7)

for all  $(\boldsymbol{p}, \boldsymbol{x}) \in \mathbb{R}^d \times \mathbb{R}^d$  and all  $t, t' \in \mathbb{R}$ . Moreover, since  $\varphi$  is a double covering, (4.5) is also equivalent to

$$d_j(\phi_k^t(\boldsymbol{p},\boldsymbol{x}),t') \, d_k(\boldsymbol{p},\boldsymbol{x},t) \, \left[ d_k(\phi_j^{t'}(\boldsymbol{p},\boldsymbol{x}),t) \, d_j(\boldsymbol{p},\boldsymbol{x},t') \right]^{-1} = \pm \mathbb{1}_2 \,. \tag{4.8}$$

However, due to

$$d_{j}(\phi_{k}^{t}(\boldsymbol{p},\boldsymbol{x}),t') d_{k}(\boldsymbol{p},\boldsymbol{x},t) \left[ d_{k}(\phi_{j}^{t'}(\boldsymbol{p},\boldsymbol{x}),t) d_{j}(\boldsymbol{p},\boldsymbol{x},t') \right]^{-1} \Big|_{t'=0} = +\mathbb{1}_{2}.$$
(4.9)

we conclude that

(4.5) 
$$\Leftrightarrow \Delta(t,t') = 0,$$
 (4.10)

where the difference  $\Delta(t, t')$  is defined by

$$\Delta(t,t') := d_j(\phi_k^t(\boldsymbol{p},\boldsymbol{x}),t') \, d_k(\boldsymbol{p},\boldsymbol{x},t) - d_k(\phi_j^{t'}(\boldsymbol{p},\boldsymbol{x}),t) \, d_j(\boldsymbol{p},\boldsymbol{x},t') \,. \tag{4.11}$$

It is easy to see that  $\Delta(t,0)=\Delta(0,t')=0 ~\forall~t,t'$  and thus

$$\Delta(t,t') = \frac{\partial^2 \Delta}{\partial t \partial t'}(0,0) tt' + (t^2 + t'^2), \quad t,t' \to 0.$$
(4.12)

The relevant second derivative is given by

$$\frac{\partial^{2}\Delta}{\partial t \partial t'}(0,0) = \frac{\partial}{\partial t} \left[ -\frac{i}{2} \boldsymbol{\sigma} \boldsymbol{\mathcal{B}}_{j}(\phi_{k}^{t}(\boldsymbol{p},\boldsymbol{x})) d_{k}(\boldsymbol{p},\boldsymbol{x},t) \right]_{t=0} \\
- \frac{\partial}{\partial t'} \left[ -\frac{i}{2} \boldsymbol{\sigma} \boldsymbol{\mathcal{B}}_{k}(\phi_{j}^{t'}(\boldsymbol{p},\boldsymbol{x})) d_{j}(\boldsymbol{p},\boldsymbol{x},t') \right]_{t'=0} \\
= \left[ -\frac{i}{2} \boldsymbol{\sigma} \{A_{k}, \boldsymbol{\mathcal{B}}_{j}\}(\boldsymbol{p},\boldsymbol{x}) + \frac{i}{2} (\boldsymbol{\sigma} \boldsymbol{\mathcal{B}}_{j}(\boldsymbol{p},\boldsymbol{x})) \frac{i}{2} (\boldsymbol{\sigma} \boldsymbol{\mathcal{B}}_{k}(\boldsymbol{p},\boldsymbol{x})) \right] \\
- \left[ -\frac{i}{2} \boldsymbol{\sigma} \{A_{j}, \boldsymbol{\mathcal{B}}_{k}\}(\boldsymbol{p},\boldsymbol{x}) + \frac{i}{2} (\boldsymbol{\sigma} \boldsymbol{\mathcal{B}}_{k}(\boldsymbol{p},\boldsymbol{x})) \frac{i}{2} (\boldsymbol{\sigma} \boldsymbol{\mathcal{B}}_{j}(\boldsymbol{p},\boldsymbol{x})) \right] \\
= \frac{i}{2} \boldsymbol{\sigma} \left[ \{ \boldsymbol{\mathcal{B}}_{j}, A_{k} \} + \{A_{j}, \boldsymbol{\mathcal{B}}_{k} \} - \boldsymbol{\mathcal{B}}_{j} \times \boldsymbol{\mathcal{B}}_{k} \right] (\boldsymbol{p},\boldsymbol{x}),$$
(4.13)

which already proves one half of Lemma 3: If the skew products  $Y_{cl_j}^t$  and  $Y_{cl_k}^t$  commute then (4.5) holds. For the reverse direction notice that (4.5) now implies

$$\Delta(t,t') = (t^2 + t'^2), \quad t,t' \to 0.$$
(4.14)

Dividing the time intervals t and t' into N subintervals of length  $\varepsilon = t/N$  and  $\varepsilon' = t'/N$ , respectively, we can rewrite the first term in (4.11) as follows,

$$d_{j}(\phi_{k}^{t}(\boldsymbol{p},\boldsymbol{x}),t') d_{k}(\boldsymbol{p},\boldsymbol{x},t) = d_{j}(\phi_{k}^{t}\circ\phi_{j}^{\varepsilon'}(\boldsymbol{p},\boldsymbol{x}),t'-\varepsilon') \underbrace{d_{j}(\phi_{k}^{t}(\boldsymbol{p},\boldsymbol{x}),\varepsilon') d_{k}(\phi_{k}^{t-\varepsilon}(\boldsymbol{p},\boldsymbol{x}),\varepsilon)}_{=:(*)} d_{k}(\boldsymbol{p},\boldsymbol{x},t-\varepsilon) \qquad (4.15)$$

$$(*) = d_{k}(\phi_{k}^{t-\varepsilon}\circ\phi_{j}^{\varepsilon'}(\boldsymbol{p},\boldsymbol{x}),\varepsilon) d_{j}(\phi_{k}^{t-\varepsilon}(\boldsymbol{p},\boldsymbol{x}),\varepsilon') + (\varepsilon^{2}+\varepsilon'^{2}).$$

Repeating this procedure  $N^2$  times yields

$$\Delta(t,t') = N^2 \left(\varepsilon^2 + \varepsilon'^2\right) = N^2 \left(1/N^2\right) = (1), \quad N \to \infty,$$
(4.16)

i.e.  $\Delta(t, t')$  vanishes which proves Lemma 3.

Having thus established a reasonable generalisation of the notion of integrability we can now state the main result of this section.

**Theorem 4.** If the skew product flow  $Y_{cl\pm}^t$  is integrable, the combined phase space  $\mathbb{R}^d \times \mathbb{R}^d \times S^2$  can be decomposed into invariant bundles  $\mathcal{T}_{\theta} \xrightarrow{\pi} \mathbb{T}^d$  over Liouville-Arnold tori  $\mathbb{T}^d$  with typical fibre  $S^1$ . The bundles can be embedded in  $\mathbb{T}^d \times S^2$  such that the fibres are characterised by the latitude with respect to a local direction  $\boldsymbol{n}(\boldsymbol{p}, \boldsymbol{x})$ , i.e.

$$\mathcal{T}_{\theta} = \{ (\boldsymbol{p}, \boldsymbol{x}, \boldsymbol{s}) \in \mathbb{T}^d \times S^2 \, | \, \sphericalangle(\boldsymbol{s}, \boldsymbol{n}(\boldsymbol{p}, \boldsymbol{x})) = \theta \} \,.$$
(4.17)

As we have seen above, integrability of  $Y_{cl}^t$  also implies similar properties of  $Y^t$ . Let us therefore state the following proposition for group extensions, the proof of which will facilitate the proof of Theorem 4.

**Proposition 5.** Let  $\phi_1^t$  be an integrable Hamiltonian flow in the sense of Theorem 1. A group extension  $Y_1^t$  of  $\phi_1^t$  with group G = U(n) or a subgroup thereof,

$$Y_1^t: \mathbb{R}^d \times \mathbb{R}^d \times G \to \mathbb{R}^d \times \mathbb{R}^d \times G (\boldsymbol{p}, \boldsymbol{x}, g) \mapsto (\phi_1^t(\boldsymbol{p}, \boldsymbol{x}), d(\boldsymbol{p}, \boldsymbol{x}, t) g) ,$$

$$(4.18)$$

$$\dot{d}(\boldsymbol{p},\boldsymbol{x},t) + M(\phi_1^t(\boldsymbol{p},\boldsymbol{x})) \ d(\boldsymbol{p},\boldsymbol{x},t) = 0, \quad d(\boldsymbol{p},\boldsymbol{x},0) = \mathbb{1}_n, \quad M: \ \mathbb{R}^d \times \mathbb{R}^d \to \mathfrak{g}, \quad (4.19)$$

where  $\mathfrak{g}$  is the Lie-albegra of G, is called integrable if the flows  $\phi_1^t, \ldots, \phi_d^t$  can also be extended to G-extensions  $Y_2^t, \ldots, Y_d^t$  with

$$\{A_j, M_k\} + \{M_j, A_k\} + [M_j, M_k] = 0 \quad \forall \ j, k = 1, \dots, d.$$
(4.20)

Then (4.19) defines a connection in the trivial principal bundle  $\mathbb{T}^d \times G$  whose holonomy group is an Abelian subgroup of G.

We remark that in the case of G = SU(2) the matrices  $M_j$  take the form

$$M_j(\boldsymbol{p}, \boldsymbol{x}) = \frac{\mathrm{i}}{2} \boldsymbol{\sigma} \boldsymbol{\mathcal{B}}_j(\boldsymbol{p}, \boldsymbol{x})$$
(4.21)

thus reducing condition (4.20) to (4.4).

PROOF OF PROPOSITION 5: First notice that condition (4.20) ensures the commutativity of the *G*-extensions  $Y_1^t, \ldots, Y_d^t$ . This can be seen by repeating the proof of Lemma 3 where in the definition (4.11) of the difference  $\Delta(t, t')$  the matrices  $d_j$  and  $d_k$  take values in *G*, see (4.19). Then equation (4.12) is still valid and (4.13) reads

$$\frac{\partial^2 \Delta}{\partial t \partial t'}(0,0) = \left( \{M_j, A_k\} + \{A_j, M_k\} + [M_j, M_k] \right) (\boldsymbol{p}, \boldsymbol{x}) \,. \tag{4.22}$$

The following steps up to (4.16) can be adopted identically.

Now define the multi-time flow

$$\mathbb{Y}^{\boldsymbol{t}} := Y_d^{t_d} \circ \dots \circ Y_1^{t_1} \,, \tag{4.23}$$

where due to commutativity of the flows  $Y_1^{t_1}, \ldots, Y_d^{t_d}$  ordering is irrelevant. Explicitly we have

$$\mathbb{Y}^{\boldsymbol{t}}(\boldsymbol{p},\boldsymbol{x},g) = \left( (\phi_d^{t_d} \circ \cdots \circ \phi_1^{t_1})(\boldsymbol{p},\boldsymbol{x}), \, \mathfrak{d}(\boldsymbol{p},\boldsymbol{x},\boldsymbol{t}) \, g \right) \quad \text{with}$$
(4.24)

$$d(\boldsymbol{p}, \boldsymbol{x}, \boldsymbol{t}) := d_d \left( (\phi_{d-1}^{t_{d-1}} \circ \cdots \circ \phi_1^{t_1})(\boldsymbol{p}, \boldsymbol{x}), t_d \right) \cdots d_1(\boldsymbol{p}, \boldsymbol{x}, t_1) g \,. \tag{4.25}$$

Consider a Liouville-Arnold torus  $\mathbb{T}^d$  which is invariant under the restriction of  $\mathbb{Y}^t$  to  $\mathbb{R}^d \times \mathbb{R}^d$ . The multi-time cocycle (4.25) then defines a connection in  $\mathbb{T}^d \times G$ . In order to determine the holonomy of this connection choose a basis  $\{\mathcal{C}_j\}$  of closed loops on  $\mathbb{T}^d$  as described in section 2. With each loop  $\mathcal{C}_j$  we can associate a unique (minimal) tuple  $t_j$  such that

$$\Phi^{\boldsymbol{t}}(\boldsymbol{p},\boldsymbol{x}), \quad t_k \in [0,(t_j)_k], \quad k = 1,\dots,d$$
(4.26)

topologically describes  $C_j$  for any  $(\boldsymbol{p}, \boldsymbol{x}) \in \mathbb{T}^d$ . We denote the cocycle  $d(\boldsymbol{p}, \boldsymbol{x}, \boldsymbol{t}_j)$  associated with  $C_j$  by  $d_j(\boldsymbol{p}, \boldsymbol{x})$ , i.e.

$$\mathbb{Y}^{\boldsymbol{t}_j}(\boldsymbol{p}, \boldsymbol{x}, g) = (\boldsymbol{p}, \boldsymbol{x}, d_j(\boldsymbol{p}, \boldsymbol{x}) g).$$
(4.27)

We immediately see that due to the commutativity of the skew products  $Y_j^t$  the cocycles  $d_j(\boldsymbol{p}, \boldsymbol{x})$  commute thus generating the Abelian subgroup

$$H_{(\boldsymbol{p},\boldsymbol{x})} = \left\{ g \in G \; \middle| \; g = \prod_{j=1}^{d} [d_j(\boldsymbol{p},\boldsymbol{x})]^{n_j} , \boldsymbol{n} \in \mathbb{Z}^d \right\}$$
(4.28)

of G. In order to see how two such groups, say  $H_{(\boldsymbol{p},\boldsymbol{x})}$  and  $H_{(\boldsymbol{p}',\boldsymbol{x}')}$ , are related, recall that for any two points  $(\boldsymbol{p},\boldsymbol{x})$  and  $(\boldsymbol{p}',\boldsymbol{x}')$  on a Liouville-Arnold torus there exists a d-tuple  $\boldsymbol{t}$ such that

$$\Phi^{\boldsymbol{t}}(\boldsymbol{p},\boldsymbol{x}) = (\boldsymbol{p}',\boldsymbol{x}'). \tag{4.29}$$

Again due to commutativity we have the equality

$$\mathbb{Y}^{t_j} = \mathbb{Y}^{-t} \circ \mathbb{Y}^{t_j} \circ \mathbb{Y}^t \tag{4.30}$$

implying

$$d_j(\boldsymbol{p}, \boldsymbol{x}) = \mathfrak{d}(\boldsymbol{p}', \boldsymbol{x}', -\boldsymbol{t}) \, d_j(\boldsymbol{p}', \boldsymbol{x}') \, \mathfrak{d}(\boldsymbol{p}', \boldsymbol{x}', \boldsymbol{t}) \,.$$
(4.31)

Moreover,

$$\mathbb{Y}^{-t} \circ \mathbb{Y}^{t} = \mathrm{id} \quad \Rightarrow \quad \mathrm{d}(\boldsymbol{p}', \boldsymbol{x}', -\boldsymbol{t}) = [\mathrm{d}(\boldsymbol{p}', \boldsymbol{x}', \boldsymbol{t})]^{-1}$$

$$(4.32)$$

and thus

$$H_{(\boldsymbol{p}',\boldsymbol{x}')} = g H_{(\boldsymbol{p},\boldsymbol{x})} g^{-1}, \quad g := d(\boldsymbol{p}',\boldsymbol{x}',\boldsymbol{t}), \qquad (4.33)$$

i.e. the subgroups  $H_{(\mathbf{p}',\mathbf{x}')}$  at different points are obtained by conjugation with a group element  $g \in G$ . Thus  $H_{(\mathbf{p},\mathbf{x})}$  and  $H_{(\mathbf{p}',\mathbf{x}')}$  are isomorphic and we have identified the Abelian group (4.28) as the holonomy group of the connection defined by (4.19).  $\Box$ PROOF OF THEOREM 4: Applying Proposition 5 to the SU(2)-extension (3.22), with each point  $(\mathbf{p}, \mathbf{x})$  on a Liouville-Arnold torus is associated an Abelian subgroup  $H_{(\mathbf{p},\mathbf{x})}$  of SU(2). Abelian subgroups of SU(2) are either one-parameter subgroups or discrete subgroups thereof, i.e. we can associate with each point  $(\mathbf{p}, \mathbf{x}) \in \mathbb{T}^d$  a one-parameter subgroup of SU(2). The latter can be parametrised as

$$H_{\boldsymbol{n}} := \left\{ g \in \mathrm{SU}(2) \mid g = \mathrm{e}^{-\mathrm{i}\frac{\alpha}{2}\boldsymbol{\sigma}\boldsymbol{n}}, \alpha \in [0, 4\pi) \right\}$$
(4.34)

with a direction characterised by the unit vector  $\mathbf{n} \in S^2 \hookrightarrow \mathbb{R}^3$ . By means of the covering map  $\varphi$ , see definition below (3.23), this construction uniquely determines a one-parameter subgroup of SO(3),  $\varphi(H_{\mathbf{n}(\mathbf{p},\mathbf{x})})$ , at each point  $(\mathbf{p},\mathbf{x})$  of a Liouville-Arnold torus. This fact in turn allows for a construction of invariant manifolds of  $Y_{cl}^t$ . Consider a point  $(\mathbf{p},\mathbf{x}) \in \mathbb{T}^d$ and a spin vector  $\mathbf{s} \in S^2 \hookrightarrow \mathbb{R}^3$ . Transporting  $\mathbf{s}$  along a path on  $\mathbb{T}^d$  by means of the multi-time flow

$$\mathbb{Y}_{cl}^{\boldsymbol{t}} := Y_{cl_d}^{t_d} \circ \cdots \circ Y_{cl_1}^{t_1} : (\boldsymbol{p}, \boldsymbol{x}, \boldsymbol{s}) \mapsto \left( (\phi_d^{t_d} \circ \cdots \circ \phi_1^{t_1}) (\boldsymbol{p}, \boldsymbol{x}), \varphi(\mathfrak{d}(\boldsymbol{p}, \boldsymbol{x}, \boldsymbol{t})) \, \boldsymbol{s} \right)$$
(4.35)

gives rise to rotations  $\varphi(d(\boldsymbol{p}, \boldsymbol{x}, \boldsymbol{t}))$  of  $\boldsymbol{s}$ . The rotation associated with a closed path  $C_j$  is given by the rotation matrix  $\varphi(d_j(\boldsymbol{p}, \boldsymbol{x})) \in \varphi(H_{\boldsymbol{n}(\boldsymbol{p}, \boldsymbol{x})}) \subset SO(3)$ . Thus whenever the path on the Liouville-Arnold torus is closed the spin vector  $\boldsymbol{s}$  returns to a point on the circle  $\varphi(H_{\boldsymbol{n}(\boldsymbol{p}, \boldsymbol{x})})\boldsymbol{s}$ , which is a parallel of latitude with respect to the axis  $\boldsymbol{n}(\boldsymbol{p}, \boldsymbol{x})$ . Corresponding circles at different points  $(\boldsymbol{p}', \boldsymbol{x}')$  are obtained as follows, see figure 2. The one-parameter subgroups at different points of the torus are related by conjugation with  $d(\boldsymbol{p}, \boldsymbol{x}, \boldsymbol{t})$ , see (4.31). However, from the definition of the covering map  $\varphi$ , see below (3.23), it follows that

$$H_{\boldsymbol{n}(\boldsymbol{p}',\boldsymbol{x}')} = \mathbb{d}(\boldsymbol{p},\boldsymbol{x},\boldsymbol{t}) H_{\boldsymbol{n}(\boldsymbol{p},\boldsymbol{x})} [\mathbb{d}(\boldsymbol{p},\boldsymbol{x},\boldsymbol{t})]^{-1} = H_{\varphi(\mathbb{d}(\boldsymbol{p},\boldsymbol{x},\boldsymbol{t}))\boldsymbol{n}}.$$
(4.36)

On the other hand from

$$\mathbb{Y}_{cl}^{t}(\boldsymbol{p}, \boldsymbol{x}, \boldsymbol{t}) = (\boldsymbol{p}', \boldsymbol{x}', \, \varphi(\mathfrak{d}(\boldsymbol{p}, \boldsymbol{x}, \boldsymbol{t}))\boldsymbol{s}) \tag{4.37}$$



Figure 2: Illustration of the invariant manifolds  $\mathcal{T}_{\theta}$ , see (4.17). At different points of the Liouville-Arnold torus  $\mathbb{T}^d$  spin vectors are restricted to parallels of latitude with different axes  $\boldsymbol{n}$ ; the angle  $\sphericalangle(\boldsymbol{s}, \boldsymbol{n}(\boldsymbol{p}, \boldsymbol{x})) = \sphericalangle(\boldsymbol{s}', \boldsymbol{n}(\boldsymbol{p}', \boldsymbol{x}'))$  is conserved

we see that by moving from  $(\boldsymbol{p}, \boldsymbol{x})$  to  $(\boldsymbol{p}', \boldsymbol{x}')$  the spin vector  $\boldsymbol{s}$  is rotated in the same way as is the axis  $\boldsymbol{n}$ . Therefore the angle

$$\theta := \sphericalangle(\boldsymbol{s}, \boldsymbol{n}(\boldsymbol{p}, \boldsymbol{x})) \tag{4.38}$$

is conserved by  $\mathbb{Y}_{cl}^t$  and thus by  $Y_{cl}^t$ , and the bundles  $\mathcal{T}_{\theta}$ , see (4.17), are invariant, concluding the proof of Theorem 4.

## 5 Quantisation and spin rotation angles

With the novel notion of integrability for skew products at hand we can now return to the semiclassical wave function (3.26) of the Dirac equation. We have already seen that except for the spin transporter  $d_{\pm}(\boldsymbol{\xi}, \boldsymbol{y}, t)$  all terms in the local expression (3.26) can be given a global meaning provided that the classical translational dynamics is integrable. In the case of the Schödinger equation, this observation led to the EBK quantisation conditions in a straightforward way, see section 2. We will now show that this is also the case for the Dirac equation if the skew product flow  $Y_{cl\pm}^t$  is integrable in the sense of Definition 2.

Theorem 4 provides us with a method for integrating the spin transport equation not only along the flow lines of  $\phi_{H^{\pm}}^t$  but also along those of the flows  $\phi_2^t, \ldots, \phi_d^t$ , which are

defined in the Theorem 1. The resulting spin transporter is defined on the whole Liouville-Arnold torus and given by the multi-time cocycle  $d(\boldsymbol{p}, \boldsymbol{x}, \boldsymbol{t})$ , see (4.25), for a point  $(\boldsymbol{p}, \boldsymbol{x})$ which is reached from  $(\boldsymbol{\xi}, \boldsymbol{y})$  by  $\Phi^{\boldsymbol{t}}(\boldsymbol{\xi}, \boldsymbol{y}) = (\boldsymbol{p}, \boldsymbol{x})$ . For a closed path  $C_j$  the initial and final values of the spin part  $u^{\pm}$  of the semiclassical wave function are still related by the SU(2) transformation  $d_j(\boldsymbol{\xi}, \boldsymbol{y})$ . However, provided that the skew product  $Y_{cl\pm}^t$  is integrable we know that the matrices  $d_j(\boldsymbol{\xi}, \boldsymbol{y})$  for different loops  $C_j$  commute. Therefore we can choose  $u^{\pm}$  to be a simultaneous eigenvector of these SU(2)-matrices which effectively reduces the SU(2)-holonomy to a simple phase, i.e. a U(1)-holonomy.

Introducing the local axis  $\boldsymbol{n}(\boldsymbol{\xi}, \boldsymbol{y})$ , see Theorem 4, we have

$$d_j(\boldsymbol{\xi}, \boldsymbol{y}) = e^{-\frac{i}{2}\alpha_j \boldsymbol{\sigma} \boldsymbol{n}(\boldsymbol{\xi}, \boldsymbol{y})}, \qquad (5.1)$$

where  $\alpha_j$  is the angle by which a classical spin vector is rotated about the axis  $\boldsymbol{n}(\boldsymbol{\xi}, \boldsymbol{y})$  when transported along  $\mathcal{C}_j$ ; note that  $\alpha_j$  has to measured modulo  $4\pi$ . The spin part  $u^{\pm}(\boldsymbol{y})$  of the wave function (3.26) is chosen to be an eigenvector of  $\boldsymbol{\sigma}\boldsymbol{n}(\boldsymbol{\xi}, \boldsymbol{y})$  with eigenvalue either +1 or -1. Thus the phase shift resulting from spin transport is given by  $e^{\mp i\alpha_j/2}$ . Together with the contribution from the classical action and the Maslov phase, cf. (2.22), the total phase shift suffered by the wave function when going through the cycle  $\mathcal{C}_j$  is

$$\frac{1}{\hbar} \oint_{\mathcal{C}_j} \boldsymbol{P} \,\mathrm{d}\boldsymbol{X} - \frac{\pi}{2} \mu_j \mp \frac{\alpha_j}{2} \,. \tag{5.2}$$

Requiring this to be an integer multiple of  $2\pi$  in order to obtain a single valued wave function yields the novel quantisation conditions

$$\oint_{\mathcal{C}_j} \boldsymbol{P} \,\mathrm{d}\boldsymbol{X} = 2\pi\hbar \left( n_j + \frac{\mu_j}{4} + m_s \frac{\alpha_j}{2\pi} \right)$$
(5.3)

where we have introduced the spin quantum number  $m_s = \pm \frac{1}{2}$ . Equation (5.3) is our central result replacing the EBK quantisation condition (1.5) in the case of relativistic particles with spin  $\frac{1}{2}$ . If the classical Hamiltonian  $H^{\pm}(\boldsymbol{p}, \boldsymbol{x})$  expressed in action and angle variables  $(\boldsymbol{I}, \boldsymbol{\vartheta})$ , see (2.23), is given by  $\overline{H}(\boldsymbol{I})$  then the semiclassical eigenvalues resulting from (5.3) read

$$E_{\boldsymbol{n},m_s}^{\pm} = \overline{H} \left( \hbar \left( \boldsymbol{n} + \frac{\boldsymbol{\mu}}{4} + m_s \frac{\boldsymbol{\alpha}}{2\pi} \right) \right) \,. \tag{5.4}$$

So far we have only employed the flows  $Y_{cl_j}^t$  resulting from the "Hamiltonians"  $A_1 = H^{\pm}, A_2, \ldots, A_d$  complemented with fields  $\mathcal{B}_j(\boldsymbol{p}, \boldsymbol{x})$ . For practical purposes it is helpful to directly make use of the flows  $\phi_{I_j}^t$  generated by the action variables  $\boldsymbol{I}$  in the semiclassical quantisation process. To this end these flows have to be extended to skew products on  $\mathbb{R}^d \times \mathbb{R}^d \times S^2$  by some suitable fields  $\mathcal{B}_{I_j}$ . Transport along a basis cycle  $\mathcal{C}_j$  is then given by  $Y_{I_j}^{2\pi}$  and the relevant SU(2) transformation  $d_j = \exp(-i\alpha_j \boldsymbol{\sigma} \boldsymbol{n}/2)$  is given by the cocycle of just one flow instead of the a linear combination of the d flows. Since

$$\phi_{H^{\pm}}^t = \phi_{I_d}^{\omega_d t} \circ \dots \circ \phi_{I_1}^{\omega_1 t}, \qquad (5.5)$$

where  $\boldsymbol{\omega}$  are the fundamental frequencies, see (2.17), we have to require that

$$Y_{\rm cl\pm}^t \stackrel{!}{=} Y_{I_d}^{\omega_d t} \circ \dots \circ Y_{I_1}^{\omega_d 1}, \qquad (5.6)$$

yielding the consistency condition

$$\boldsymbol{\mathcal{B}}^{\pm} = \sum_{j=1}^{d} \omega_j \boldsymbol{\mathcal{B}}_{I_j} \,. \tag{5.7}$$

We will illustrate these remarks when applying the method to explicit examples in sections 7-9.

## 6 Non-relativistic limit: The Pauli equation

In this section we show that the semiclassical quantisation scheme developed in the preceding sections for the Dirac equation carries over to the Pauli equation. The latter arises as a non-relativistic approximation to the Dirac equation but can also be generalised to describe particles with spin  $s \in \mathbb{N}_0/2$  other than  $s = \frac{1}{2}$ . We show that also in the case with  $s \neq \frac{1}{2}$  the semiclassical analysis of the Pauli equation gives rise to a skew product on  $\mathbb{R}^d \times \mathbb{R}^d \times S^2$  which can then be quantised along the same lines as in the case of the Dirac equation. We keep the presentation short relying heavily on our treatment of the Dirac equation in the preceding sections; details on the semiclassical analysis of the Pauli equation with arbitrary spin can be found in [46].

The Pauli equation for a spin- $\frac{1}{2}$  particle can be obtained from the Dirac equation (3.1) in the non-relativistic limit  $c \to \infty$ , see e.g. [47, 48]. With the representation (3.3) of the matrices  $\boldsymbol{\alpha}$  and  $\boldsymbol{\beta}$  and writing the Dirac spinor as  $\Psi^T = (\psi^T, \chi^T)$  with  $\psi, \chi \in L^2(\mathbb{R}^3) \otimes \mathbb{C}^2$ one finds the following equation for the upper two components,

$$\hat{H}_{\rm P}\psi(\boldsymbol{x}) = E\psi(\boldsymbol{x})$$
(6.1)

with Pauli Hamiltonian

$$\hat{H}_{\rm P} = -\frac{\hbar^2}{2m} \left( \frac{\hbar}{\rm i} \nabla - \frac{e}{c} \boldsymbol{A}(\boldsymbol{x}) \right)^2 + e\phi(\boldsymbol{x}) - \frac{e}{mc} \boldsymbol{B}(\boldsymbol{x}) \frac{\hbar}{2} \boldsymbol{\sigma} \,. \tag{6.2}$$

Let us generalise our discussion to Pauli Hamiltonians

$$\hat{H}_{\rm P} = \hat{H}_{\rm S} + \frac{\hbar}{2} \mathrm{d}\pi_s(\boldsymbol{\sigma})\hat{\boldsymbol{\mathcal{B}}}, \qquad (6.3)$$

where  $\hat{H}_{s}$  is a Schrödinger Hamiltonian with Weyl symbol  $H(\boldsymbol{p}, \boldsymbol{x})$ ,  $d\pi_{s}(\boldsymbol{\sigma})$  denotes the 2s+1 dimensional (derived) irreducible representation of  $\mathfrak{su}(2)$  and  $\hat{\boldsymbol{\beta}}$  is the Weyl quantisation of the classical vector valued function  $\boldsymbol{\mathcal{B}}(\boldsymbol{p}, \boldsymbol{x})$  on phase space. The wave function  $\psi$  has now 2s + 1 components, i.e  $\psi \in L^{2}(\mathbb{R}^{d}) \otimes \mathbb{C}^{2s+1}$ . The special case (6.2) is recovered by

the choices d = 3,  $s = \frac{1}{2}$ ,  $H(\boldsymbol{p}, \boldsymbol{x}) = \frac{1}{2m} \left( \boldsymbol{p} - \frac{e}{c} \boldsymbol{A}(\boldsymbol{x}) \right)^2 + e\phi(\boldsymbol{x})$  and  $\boldsymbol{\mathcal{B}}(\boldsymbol{p}, \boldsymbol{x}) = -\frac{e}{mc} \boldsymbol{B}(\boldsymbol{x})$ . Spin-orbit coupling can, e.g., be described by

$$\boldsymbol{\mathcal{B}}_{so}(\boldsymbol{p}, \boldsymbol{x}) = f(r)\boldsymbol{L}, \qquad (6.4)$$

where  $\boldsymbol{L} = \boldsymbol{x} \times \boldsymbol{p}$  is orbital angular momentum and f is an arbitrary function of the radial coordinate  $r = |\boldsymbol{x}|$ .

As in (2.5) we make the ansatz

$$\psi(\boldsymbol{x}) = \sum_{k \ge 0} \left(\frac{\hbar}{i}\right)^k a_k(\boldsymbol{x}) e^{\frac{i}{\hbar}S(\boldsymbol{x})}$$
(6.5)

with scalar S, and  $a_k$  taking values in  $\mathbb{C}^{2s+1}$ . Upon inserting (6.5) into the Pauli equation (6.1) with Hamiltonian (6.3), in leading order we find the Hamilton-Jacobi equation

$$H(\nabla_{\!\boldsymbol{x}}S,\boldsymbol{x}) = E \tag{6.6}$$

and in next-to-leading order we obtain the transport equation

$$[(\nabla_{\boldsymbol{p}}H)(\nabla_{\boldsymbol{x}}S,\boldsymbol{x})](\nabla_{\boldsymbol{x}}a_0) + \frac{1}{2}[\nabla_{\boldsymbol{x}}(\nabla_{\boldsymbol{p}}H)(\nabla_{\boldsymbol{x}}S,\boldsymbol{x})]a_0 + \frac{\mathrm{i}}{2}\mathrm{d}\pi_s(\boldsymbol{\sigma})\boldsymbol{\mathcal{B}}(\nabla_{\boldsymbol{x}}S,\boldsymbol{x})a_0 = 0.$$
(6.7)

As below (3.15) the ansatz

$$a_0(\boldsymbol{x}) = \sqrt{\det \frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}}} u(\boldsymbol{x})$$
(6.8)

leaves us with the spin transport equation

$$\dot{u}(\boldsymbol{x}) + \frac{\mathrm{i}}{2} \mathrm{d}\pi_s(\boldsymbol{\sigma}) \boldsymbol{\mathcal{B}}(\nabla_{\!\!\boldsymbol{x}} S, \boldsymbol{x}) \, u(\boldsymbol{x}) = 0 \,.$$
(6.9)

Integration along a trajectory with  $\phi_H^t(\boldsymbol{\xi}, \boldsymbol{y}) = (\boldsymbol{p}, \boldsymbol{x})$  yields

$$u(\boldsymbol{x}) = \pi_s(d(\boldsymbol{\xi}, \boldsymbol{y}, t)) \, u(\boldsymbol{y}) \,, \tag{6.10}$$

where  $\pi_s$  denotes the 2s + 1 dimensional unitary irreducible representation of SU(2) and d solves the spin transport equation

$$\dot{d}(\boldsymbol{\xi}, \boldsymbol{y}, t) + \frac{\mathrm{i}}{2} \boldsymbol{\sigma} \boldsymbol{\mathcal{B}} \left( \phi_H^t(\boldsymbol{\xi}, \boldsymbol{y}) \right) d(\boldsymbol{\xi}, \boldsymbol{y}, t) = 0, \quad d(\boldsymbol{\xi}, \boldsymbol{y}, 0) = \mathbb{1}_2, \quad (6.11)$$

in SU(2). Thus the covering map  $\varphi : SU(2) \to SO(3)$  still relates spin transport to classical spin precession,

$$\dot{\boldsymbol{s}} = \boldsymbol{\mathcal{B}}(\phi_H^t(\boldsymbol{\xi}, \boldsymbol{y})) \times \boldsymbol{s}, \quad \boldsymbol{s} \in S^2 \hookrightarrow \mathbb{R}^3,$$
(6.12)

and we identify the skew product flow

$$Y_{\rm cl}^t(\boldsymbol{p}, \boldsymbol{x}, \boldsymbol{s}) = \left(\phi_H^t(\boldsymbol{p}, \boldsymbol{x}), \varphi(d(\boldsymbol{p}, \boldsymbol{x}, t))\boldsymbol{s}\right)$$
(6.13)

as the classical system corresponding to the Pauli equation with Hamiltonian (6.3). If  $Y_{cl}^t$  is integrable in the sense of Definition 2 we can use the same construction as in section 5 to define a semiclassical wave function associated with a Liouville-Arnold torus  $\mathbb{T}^d$ . Locally we have

$$\psi_{\rm sc}(\boldsymbol{x}) \sim \sqrt{\det \frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}}} e^{\frac{i}{\hbar}S(\boldsymbol{x})} \pi_s(d(\boldsymbol{\xi}, \boldsymbol{y}, t)) u(\boldsymbol{y}).$$
 (6.14)

and the initial and final values of u after transport along along a basis cycle  $C_j$  are related by  $\pi_s(d_j(\boldsymbol{\xi}, \boldsymbol{y}))$ . As in (5.1) we obtain

$$d_j(\boldsymbol{\xi}, \boldsymbol{y}) = e^{-\frac{i}{2}\alpha_j \boldsymbol{\sigma} \boldsymbol{n}(\boldsymbol{\xi}, \boldsymbol{y})}, \qquad (6.15)$$

where the axis  $\boldsymbol{n}$  is specified by Theorem 4 and the angle  $\alpha_j$  is the rotation angle for a classical spin vector. The representation matrix  $\pi_s(d_j(\boldsymbol{\xi}, \boldsymbol{y}))$  has eigenvalues  $\exp(-im_s\alpha_j)$ ,  $m_s = -s, -s + 1, \ldots, s$ , see e.g. [49]. Upon choosing  $n(\boldsymbol{\xi}, \boldsymbol{y})$  to be an eigenvector of  $\pi_s(d_j(\boldsymbol{\xi}, \boldsymbol{y}))$  the total phase change of experienced by  $\psi_{sc}$  along a cycle  $C_j$  is given by

$$\frac{1}{\hbar} \oint_{\mathcal{C}_j} \boldsymbol{P} \,\mathrm{d}\boldsymbol{X} - \mathrm{i}\frac{\pi}{2}\mu_j - m_s \alpha_j \tag{6.16}$$

thus resulting in the semiclassical quantisation conditions

$$\oint_{\mathcal{C}_j} \boldsymbol{P} \,\mathrm{d}\boldsymbol{X} = 2\pi\hbar \left( n_j + \frac{\mu_j}{4} + m_s \frac{\alpha_j}{2\pi} \right),\tag{6.17}$$

with  $\boldsymbol{n} \in \mathbb{Z}^d$  and  $m_s = -s, -s+1, \ldots, s$ .

## 7 Spherically symmetric systems

Before we treat some explicit examples in order to illustrate the novel quantisation conditions (5.3) and (6.17) we show how to apply them to an important class of integrable systems, namely spherically symmetric systems.

A spherically symmetric Dirac Hamiltonian has the structure

$$\hat{H}_{\rm D} = c \alpha \, \frac{\hbar}{\rm i} \nabla + \beta \, mc^2 + e \phi(r) \,, \tag{7.1}$$

where the electrostatic potential  $\phi$  depends only on the radial variable  $r := |\mathbf{x}|$ . One easily verifies that the Hamiltonian (7.1) commutes with all components of total angular momentum,

$$\hat{\boldsymbol{J}}_{\mathrm{D}} := \hat{\boldsymbol{L}} + \frac{\hbar}{2} \begin{pmatrix} \boldsymbol{\sigma} & 0\\ 0 & \boldsymbol{\sigma} \end{pmatrix}, \quad \hat{\boldsymbol{L}} := \boldsymbol{x} \times \frac{\hbar}{\mathrm{i}} \nabla.$$
 (7.2)

Since one also has

$$[\hat{\boldsymbol{J}}_{D}^{2}, \hat{J}_{Dz}] = 0$$
 (7.3)

one can choose the eigenfunctions of  $\hat{H}_{\rm D}$  to be also simultaneous eigenfunctions of  $\hat{J}_{\rm D}^2$  and  $\hat{J}_{{\rm D}z}$ .

Analogously a spherically symmetric Pauli Hamiltonian is of the form

$$\hat{H}_{\rm P} = -\frac{\hbar^2}{2m} \Delta + e\phi(\boldsymbol{x}) + \frac{\hbar}{2} \mathrm{d}\pi_s(\boldsymbol{\sigma}) f(r) \hat{\boldsymbol{L}}, \qquad (7.4)$$

cf. (6.4), and commutes with all components of total angular momentum

$$\hat{J}_{\mathrm{P}} := \hat{\boldsymbol{L}} + \frac{\hbar}{2} \mathrm{d}\pi_s(\boldsymbol{\sigma}) \,. \tag{7.5}$$

Again due to

$$[\hat{\boldsymbol{J}}_{\mathrm{P}}^2, \hat{J}_{\mathrm{P}z}] = 0 \tag{7.6}$$

the eigenfunctions of  $\hat{H}_{\rm P}$  can be chosen such that they are simultaneous eigenfunctions to the modulus squared  $\hat{J}_{\rm P}^2$  and the z-component  $\hat{J}_{\rm Pz}$  of total angular momentum. In the semiclassical limit the Hamiltonians (7.1) and (7.4) both give rise to skew prod-

In the semiclassical limit the Hamiltonians (7.1) and (7.4) both give rise to skew products  $Y_{cl}^t$  on  $\mathbb{R}^d \times \mathbb{R}^d \times S^2$  with a spherically symmetric classical Hamiltonian  $H(\mathbf{p}, r)$  and spin precession with fields of the form

$$\mathcal{B}(\boldsymbol{p}, \boldsymbol{x}) = f(r) \boldsymbol{L}.$$
(7.7)

It is well-known that  $\phi_H^t$  is integrable in the sense of Liouville and Arnold since

$$\{H, L\} = \{H, M\} = \{L, M\} = 0$$
(7.8)

where L and M are the modulus and z-component of classical orbital angular momentum  $L = \mathbf{x} \times \mathbf{p}$ , respectively. We remark that  $L(\mathbf{p}, \mathbf{x})$  not being smooth at points where  $\mathbf{p} \| \mathbf{x}$  will not play a rôle in the following since all relevant constructions will stay away from these points. Thus (7.8) holds wherever needed.

In order to show that  $Y_{cl}^t$  with field (7.7) is also integrable we have to extend the flows  $\phi_L^t$  and  $\phi_M^t$  to skew products with fields  $\mathcal{B}_L$  and  $\mathcal{B}_M$  fulfilling (4.4). To this end consider the Weyl symbol of the operator  $\hat{J} := \hat{L} + \frac{\hbar}{2}\sigma$ ,

$$\boldsymbol{J} = \boldsymbol{p} \times \boldsymbol{x} + \frac{\hbar}{2} \boldsymbol{\sigma} \,. \tag{7.9}$$

By straightforward calculation one finds

$$J := |\boldsymbol{J}| = L + \frac{\hbar}{2}\boldsymbol{\sigma} \, \frac{\boldsymbol{L}}{L} + (\hbar^2) \quad \text{and} \quad J_z = M + \frac{\hbar}{2}\sigma_z \,. \tag{7.10}$$

Comparing these to the Pauli Hamiltonian (7.4) suggests that

$$\boldsymbol{\mathcal{B}}_L := \frac{\boldsymbol{L}}{L} \quad \text{and} \quad \boldsymbol{\mathcal{B}}_M := \boldsymbol{e}_z \,,$$
 (7.11)

where  $e_z$  is the unit vector in z-direction, might be a good choice. As one easily checks the skew products  $Y_{cl}^t$ ,  $Y_L^t$  and  $Y_M^t$  obtained in this way indeed form a set of commuting flows on  $\mathbb{R}^d \times \mathbb{R}^d \times S^2$  and thus  $Y_{cl}^t$  is integrable, cf. Theorem 4.

Spherically symmetric Hamiltonians  $H(\mathbf{p}, r)$  can be separated in spherical coordinates  $(r, \theta, \phi)$ , see e.g. [31]. Introducing the action variables

$$I_r = \oint p_r \,\mathrm{d}r \,, \quad I_\theta = \oint p_\theta \,\mathrm{d}\theta \quad \text{and} \quad I_\phi = \oint p_\phi \,\mathrm{d}\phi \,, \tag{7.12}$$

where  $p_r$ ,  $p_{\theta}$  and  $p_{\phi}$  denote the canonical momenta conjugate to r,  $\theta$  and  $\phi$ , respectively, one always finds

$$I_{\phi} = M$$
 and  $I_{\theta} = L - M$ . (7.13)

The radial action  $I_r$  depends on the particular system under investigation and solving  $I_r = \oint p_r(E, I_{\theta}, I_{\phi}) dr$  for E yields the Hamiltonian  $\overline{H}$  in action and angle variables. It is a general feature of spherically symmetric systems that  $\overline{H}$  does not depend on  $I_{\theta}$  and  $I_{\phi}$  independently but only on the sum  $L = I_{\theta} + I_{\phi}$ , i.e.  $\overline{H}$  is a function of  $I_r$  and L.

The motion generated by  $M = I_{\phi}$  is a rotation about the z-axis and thus there is no turning point in the time evolution of the coordinate  $\phi$  yielding the Maslov index

$$\mu_M = \mu_\phi = 0. \tag{7.14}$$

On the other hand the motion generated by  $I_{\theta}$  takes place between two turning points for the coordinate  $\theta$  and thus

$$\mu_{\theta} = 2$$
 and accordingly  $\mu_L = 2$ . (7.15)

In order to find the spin rotation angle  $\alpha_M$  one has to integrate the spin precession equation

$$\dot{\boldsymbol{s}} = \boldsymbol{e}_z \times \boldsymbol{s} \tag{7.16}$$

over one cycle of  $\phi_M^t$ . The latter simply changes the angle variable  $\vartheta_M$  conjugate to M with unit speed and we have

$$\alpha_M = 2\pi |\boldsymbol{e}_z| = 2\pi \,. \tag{7.17}$$

Similarly  $\phi_L^t$  changes  $\vartheta_L$  with unit speed and since  $\boldsymbol{\mathcal{B}}_L = \boldsymbol{L}/L$  is constant along such a cycle we also find

$$\alpha_L = 2\pi |\mathbf{L}/L| = 2\pi \,. \tag{7.18}$$

Therefore, for any spherically symmetric Dirac or Pauli equation we have semiclassical quantisation conditions

$$L = \hbar \left( l + \frac{1}{2} + m_s \right) \quad \text{and} \quad M = \hbar \left( m_l + m_s \right)$$
(7.19)

with integers l and  $m_l$ .

Classical mechanics imposes the restrictions

$$L \ge 0 \quad \text{and} \quad M \le L \,.$$
 (7.20)

Defining the new quantum numbers  $j := l + m_s$  and  $m_j := m_l + m_s$  after some trivial algebra one finds that (7.20) translates to

$$j \ge 0$$
 and  $m_j = -j, -j+1, \dots, j$ . (7.21)

Notice that if s is integer or half-integer, respectively, then so are j and  $m_j$ . Finally the semiclassical quantisation conditions for angular momentum read

$$L = \hbar \left( j + \frac{1}{2} \right)$$
 and  $M = \hbar m_j$ . (7.22)

As mentioned at the end of section 5 it will now be useful if we also find a field  $\mathcal{B}_r$  which turns the flow  $\phi_r^t$  generated by the action variable  $I_r$  into a skew product that commutes with  $Y_L^t$  and  $Y_M^t$ . To this end we may exploit the relation (5.7). Keeping in mind that  $\overline{H}$ is a function of  $I_r$  and  $L = I_{\theta} + I_{\phi}$  only, i.e.  $\omega_{\theta} = \omega_{\phi} = \omega_L = \partial \overline{H} / \partial L$ , this yields

$$\boldsymbol{\mathcal{B}}_{r} = \frac{\boldsymbol{\mathcal{B}} - \omega_{L} \boldsymbol{\mathcal{B}}_{L}}{\omega_{r}} \,. \tag{7.23}$$

Since  $\mathcal{B} = f(r)L$  is parallel to  $\mathcal{B}_L = L/L$  and since L is a constant of motion  $\mathcal{B}_r$  does not change its direction along the flow line of either  $\phi_H^t$  or  $\phi_r^t$ . Therefore

$$\alpha_r = \left| \int_0^{2\pi} \frac{\mathcal{B} - \omega_L \frac{\mathbf{L}}{L}}{\omega_r} \mathrm{d}\vartheta_r \right|,\tag{7.24}$$

where  $\vartheta_r$  is the angle variable conjugate to  $I_r$ . Splitting the integrand into two terms,  $\mathcal{B}/\omega_r$  and  $-\omega_L \mathbf{L}/(\omega_r L)$ , the second integration is again trivial. In the first integral we can make use of  $\mathrm{d}\vartheta_r/\omega_r = \mathrm{d}\vartheta_r/\dot{\vartheta}_r = \mathrm{d}t$ , where t is the physical time along a flow line of  $\phi_H^t$ . Therefore we have obtained the handy expression

$$\alpha_r = \left| \oint_{\text{radial}} \mathcal{B} \left( \phi_H^t(\boldsymbol{p}, \boldsymbol{x}) \right) dt - 2\pi \frac{\omega_L}{\omega_r} \frac{\boldsymbol{L}}{L} \right|$$
(7.25)

where the remaining integral extends over one cycle of the radial motion, e.g. from perihelion to aphelion and back. The missing quantisation condition then reads

$$I_r = 2\pi\hbar \left( n_r + \frac{1}{2} + m_s \frac{\alpha_r}{2\pi} \right) , \qquad (7.26)$$

where we have inserted  $\mu_r = 2$  for a typical radial motion between perihelion and aphelion.

## 8 Example 1: Harmonic oscillator with spin-orbit coupling

As a first example we discuss the three dimensional isotropic harmonic oscillator with spin-orbit coupling in a non-relativistic context with arbitrary spin.

The Pauli Hamiltonian

$$\hat{H}_{\rm P} = -\frac{\hbar^2}{2m}\Delta + \frac{m}{2}\omega^2 r^2 + \frac{\hbar}{2}\mathrm{d}\pi_s(\boldsymbol{\sigma})\,\kappa\hat{\boldsymbol{L}}$$
(8.1)

describes an oscillator with frequency  $\omega$  and for the spin-orbit coupling we have chosen a Thomas term, i.e. the leading order expression of  $\mathcal{B}^+$ , see (3.15), in the non-relativistic limit  $c \to \infty$  with

$$e\boldsymbol{E} = -\nabla\left(\frac{m}{2}\omega^2 r^2\right) = -m\omega^2 \boldsymbol{x}$$
(8.2)

i.e.

$$B^{+} = \frac{\omega^{2}}{2mc^{2}}L + (c^{-4}).$$
(8.3)

By comparison with (8.1) we have  $\kappa = \omega^2/(2mc^2)$  but we may keep  $\kappa$  arbitrary for the rest of the section.

One usually determines the eigenvalues of (8.1) by first observing that besides  $\hat{\boldsymbol{J}}_{\mathrm{P}}^2$  and  $\hat{\boldsymbol{J}}_{\mathrm{P}z}$ , see (7.5), also  $\hat{\boldsymbol{L}}^2$  and  $\hat{\boldsymbol{s}}^2 = [\hbar/2\mathrm{d}\pi_s(\boldsymbol{\sigma})]^2$  commute with  $\hat{H}_{\mathrm{P}}$  and that the spin-orbit term can be expressed in terms of these conserved quantities, see e.g. [50].

Obviously, the Hamiltonian (8.1) defines a spherically symmetric system as discussed in the preceeding section. Therefore, we immediately have the quantisation conditions (7.22) for total angular momentum. The semiclassical eigenvalues, however, can even be found in a more straightforward way than to draw on (7.25). To this end notice that the classical Hamiltonian  $H(\mathbf{p}, r) = \frac{\mathbf{p}^2}{2m} + \frac{m}{2}\omega^2 r^2$  transformed to action and angle variables reads

$$\overline{H}(I_r, L) = \omega(2I_r + L).$$
(8.4)

Semiclassical quantisation with s = 0 yields

$$E_{n_r l} = \hbar \omega \left( 2n + l + \frac{3}{2} \right), \tag{8.5}$$

exhibiting the zero point energy of  $3\hbar\omega/2$  of the three independent oscillators in x-, yand z-direction. Since  $\overline{H}$  only depends on a linear combination of  $I_r$  and L the problem has an even higher degeneracy than general spherically symmetric systems. Thus we may introduce the new action variable

$$I_1 := 2I_r + L \tag{8.6}$$

with corresponding frequency  $\omega_1 = \partial \overline{H} / \partial I_1 = \omega$ . From (8.5) we see that without spin  $I_1$  has to be quantised as  $I_1 = \hbar (2n + l + \frac{3}{2})$ . In order to find the correction from the spin

contribution we have to extend  $\phi_{I_1}^t$  to a skew product  $Y_{I_1}^t$  that commutes with  $Y_{cl}^t$ . The consistency condition (5.7) uniquely determines the relevant field  $\mathcal{B}_1$ , since it reduces to

$$\mathcal{B} = \omega_1 \mathcal{B}_1 \quad \Rightarrow \quad \mathcal{B}_1 = \frac{\mathcal{B}}{\omega_1} = \frac{\kappa}{\omega} L.$$
 (8.7)

Integration of the spin precession equation  $\dot{s} = \mathcal{B}_1 \times s$  is once more trivial since L is a constant of motion and we get

$$\alpha_1 = \frac{2\pi\kappa}{\omega} |\mathbf{L}| = \frac{2\pi\hbar\kappa}{\omega} \left( l + \frac{1}{2} + m_s \right) \,. \tag{8.8}$$

Finally, the semiclassical energies read

$$E_{n_r l m_s} = \hbar \omega \left(2n + l + \frac{3}{2}\right) + m_s \hbar^2 \kappa \left(l + \frac{1}{2} + m_s\right).$$
(8.9)

For spin  $s = \frac{1}{2}$  these are the exact eigenvalues of the Hamiltonian (8.1) whereas for  $s \neq \frac{1}{2}$  they are good approximations to the exact eigenvalues if l is large, i.e. if the action variable L is large compared to  $\hbar$  as required for semiclassical approximations.

At this point a short remark concerning the practical application of conditions (5.3) or (6.17) is in order: In general the spin rotation angels  $\boldsymbol{\alpha}$  can depend on the action variables  $\boldsymbol{I}$ . If the dependence is simple as in this example ( $\alpha_1$  depends on L but not on  $I_1$ ,  $\alpha_L$  and  $\alpha_M$  are constant) one can recursively apply (6.17) as was done above: First we quantised L and then used the result when quantising  $I_1$ . In general, however, one may first have to solve the quantisation conditions (6.17) for the action variables  $\boldsymbol{I}$  before inserting the result into the Hamiltonian  $\overline{H}(\boldsymbol{I})$ .

## 9 Example 2: Sommerfeld's theory of fine structure revisited

In this section we address the classic problem of Sommerfeld's theory of fine structure which was already mentioned in the introduction as a motivating example.

To this end we have to consider the relativistic Kepler problem with classical Hamiltonian

$$H(\mathbf{p}, r) = -\frac{e^2}{r} + \sqrt{c^2 \mathbf{p}^2 + m^2 c^4}.$$
(9.1)

Solutions of Hamilton's equations of motion corresponding to bound states are given by "Rosettenbahnen", ellipses with moving perihelia. Since (9.1) is spherically symmetric angular momentum is conserved and the motion takes place in a plane. Introducing polar coordinates in this plane the orbits can be expressed as

$$\frac{1}{r(\phi)} = \underbrace{\frac{e^2 E}{c^2 L^2 - e^4}}_{:=C} + \underbrace{\frac{\sqrt{c^2 L^2 E^2 + (c^2 L^2 - e^4)m^2 c^4}}_{:=A}}_{:=A} \cos\left(\underbrace{\frac{\sqrt{c^2 L^2 - e^4}}{cL}}_{:=\gamma}\phi\right), \tag{9.2}$$

where for later reference we have defined the constants C, A and  $\gamma$ . Detailed information on the classical mechanics of the relativistic Kepler problem can, e.g., be found in Sommerfeld's original article [6] or in his book [51]. Transforming the Hamiltonian to action and angle variables yields

$$\overline{H}(I_r, L) = mc^2 \left[ 1 + \frac{e^4/c^2}{\left(I_r + \sqrt{L^2 - e^4/c^2}\right)^2} \right]^{-1/2}.$$
(9.3)

Sommerfeld quantised the system using (1.2) by demanding

$$I_r = \hbar n_r \quad \text{and} \quad L = \hbar l \tag{9.4}$$

with integers  $n_r$  and l. Classical mechanics imposes the restrictions  $I_r \geq 0$  and  $L \geq 0$  yielding  $n_r, l \geq 0$ . Moreover, Sommerfeld excluded l = 0 because in this case the electron would collide with the nucleus. Thus he found the energy levels

$$E_{n_r l}^{\text{Sommerfeld}} = mc^2 \left[ 1 + \frac{\alpha_{\text{S}}^2}{\left(n_r + \sqrt{l^2 - \alpha_{\text{S}}^2}\right)^2} \right]^{-1/2}, \quad n_r \in \mathbb{N}_0, \quad l \in \mathbb{N}, \qquad (9.5)$$

where  $\alpha_{\rm S} = e^2/(\hbar c)$  denotes Sommerfeld's fine structure constant.

When now quantising the relativistic Kepler problem with spin  $\frac{1}{2}$  using conditions (5.3) first notice that due to spherical symmetry we have, cf. (7.19),

$$L = \hbar \left( l + \frac{1}{2} + m_s \right), \quad m_s = \pm \frac{1}{2}.$$
 (9.6)

Since the additional conditions (7.20) do not allow the combination l = 0,  $m_s = -\frac{1}{2}$  (which would lead to a negative quantum number j) we see that  $L/\hbar \in \mathbb{N}$ , i.e. it assumes the same values as in Sommerfeld's prescription (9.4).

In order to quantise the radial action variable  $I_r$  we also have to calculate the spin rotation angle  $\alpha_r$  to which end we can use the general formula (7.25). The field **B** occurring in the equation of spin precession can be obtained from **B**<sup>+</sup>, see (3.15), by inserting  $\mathbf{E} = -\nabla(-e^2/r)$  yielding

$$\boldsymbol{\mathcal{B}} = \frac{e^2 c^2}{\varepsilon(\varepsilon + mc^2)} \frac{1}{r^3} \boldsymbol{L} \,. \tag{9.7}$$

where  $\varepsilon$  was defined in equation (3.11). Without restriction we may choose  $L \| e_z$  for our calculation leading to

$$\boldsymbol{L} = \frac{\varepsilon}{c^2} r^2 \, \dot{\phi} \, \boldsymbol{e}_z \quad \Rightarrow \quad \boldsymbol{\mathcal{B}} = \frac{e^2}{\varepsilon + mc^2} \frac{1}{r} \, \dot{\phi} \, \boldsymbol{e}_z \,. \tag{9.8}$$

Using  $E = H(\mathbf{p}, r) = -e^2/r + \varepsilon$ , see (9.1), this expression further simplifies to

$$\boldsymbol{\mathcal{B}} = \frac{e^2}{(E+mc^2)r+e^2} \,\dot{\phi} \,\boldsymbol{e}_z \,. \tag{9.9}$$

which is nicely adapted for use in (7.25),

$$\oint_{\text{radial}} \mathcal{B} \, \mathrm{d}t = \mathbf{e}_z \oint_{\text{radial}} \frac{e^2}{(E+mc^2)r(\phi) + e^2} \, \mathrm{d}\phi = \mathbf{e}_z \oint_{\text{radial}} \left[ 1 - \frac{E+mc^2}{E+mc^2 + e^2/r(\phi)} \right] \mathrm{d}\phi \,.$$
(9.10)

Changing variables to  $\eta = \gamma \phi$  and inserting (9.2) we get

$$= \frac{\boldsymbol{e}_z}{\gamma} \oint_{\text{radial}} \left[ 1 - \frac{E + mc^2}{E + mc^2 + e^2 C + e^2 A \cos(\eta)} \right] \mathrm{d}\phi = \frac{\boldsymbol{e}_z}{\gamma} (1 - 2\pi\gamma) \,, \tag{9.11}$$

where the calculation of the last integral is tedious but elementary. From (9.3) we easily determine the frequency ratio in the second term of (7.25) to  $\omega_L/\omega_r = \gamma^{-1}$  and since we have chosen  $\boldsymbol{L} \| \boldsymbol{e}_z$  we obtain

$$\alpha_r = \left| \boldsymbol{e}_z \left( \frac{1}{\gamma} - 2\pi \right) - \frac{\boldsymbol{e}_z}{\gamma} \right| = 2\pi.$$
(9.12)

This remarkable result turns (7.26) into the rather trivial condition

$$I_r = \hbar \left( n_r + \frac{1}{2} + m_s \right) \tag{9.13}$$

with integer  $n_r$  and  $m_s = \pm \frac{1}{2}$ . Moreover, classical mechanics demands  $I_r \ge 0$  and thus  $I_r/\hbar \in \mathbb{N}_0$ , i.e. the values assumed are again identical to those predicted by Sommerfeld's condition (9.4).

Finally we have found the semiclassical energies,

$$E_{n_r l m_s} = mc^2 \left[ 1 + \frac{\alpha_{\rm S}^2}{\left(n_r + \frac{1}{2} + m_s + \sqrt{\left(l + \frac{1}{2} + m_s\right)^2 - \alpha_{\rm S}^2}\right)^2} \right]^{-1/2}, \qquad (9.14)$$

which are identical to those obtained from the exact solution of the Dirac equation (3.1) with potential  $-e^2/r$ , see e.g. [47, 50], as can be most easily checked by introducing the quantum numbers j and  $m_j$  of total angular momentum, see below (7.20), and the principal quantum number n, associated with  $I := I_r + L$ , by  $I = \hbar n$ . Although Sommerfeld's formula (9.5) also yields the correct energy levels of the hydrogen atom it predicts the wrong multiplicities. This problem is rectified by the present treatment as can be checked straightforwardly. Consider, e.g., the ground state which in (9.5) is obtained by  $n_r = 0$ , l = 1, and thus is non-degenerate. On the other hand in (9.14) the same energy is obtained by choosing either  $n_r = -1$ , l = 0,  $m_s = \frac{1}{2}$ , or  $n_r = 0$ , l = 1,  $m_s = -\frac{1}{2}$ . Alternatively these states can be characterised by n = 1,  $j = \frac{1}{2}$ ,  $m_j = \pm \frac{1}{2}$  yielding a multiplicity of 2 as in the exact quantum spectrum.

#### 10 Conclusions

In this article we have derived semiclassical quantisation conditions for the Dirac and Pauli equations. We have shown that spin yields a contribution of the same order of magnitude as the Maslov correction. The spin contribution is determined by rotation angles for a classical spin vector that is transported along orbits of the translational dynamics.

The crucial step in the derivation of the semiclassical quantisation conditions was the generalisation of the notion of integrability to certain skew products and group extensions. The relevant integrability conditions enabled us to effectively reduce the non-Abelian Berry phases, appearing in the analysis of multi-component wave equations, to a U(1)-holonomy, i.e. to an ordinary phase factor. The latter can then be incorporated into the quantisation conditions.

We remark that our treatment generalises to arbitrary multi-component wave equations for which the principal symbol of the Hamiltonian has eigenvalues with arbitrary but constant multiplicity. One then has to deal with G-extensions, where G = U(n) or a subgroup thereof, of the ray dynamics (generated by the eigenvalues of the principal symbol). Proposition 5 provides us with the relevant integrability condition which allows for an effective reduction of the holonomy group G to an Abelian subgroup H. The spin rotation angles are then replaced by the eigenphases of some unitary representation matrices of H.

In section 9 we have applied the novel quantisation conditions to the relativistic Kepler problem. We have seen that by a freak of nature all relevant spin rotation angles are given by  $2\pi$  and thus cancel (or add up to an integer) with the Maslov term. It is this coincidence due to which Sommerfeld was able to calculate the energy levels of the relativistic hydrogen atom including spin-orbit coupling 10 years before the Dirac equation was developed.

#### Acknowledgement

I would like to thank Jens Bolte for helpful discussions. This work was supported by the Deutsche Forschungsgemeinschaft under contract no. Ste 241/10-2.

## A Wigner-Weyl calculus

With a differential operator  $\hat{A}$  one can associate an object on classical phase space, its Weyl symbol  $A(\boldsymbol{p}, \boldsymbol{x})$ , by

$$(\hat{A}\Psi)(\boldsymbol{x}) = \frac{1}{(2\pi\hbar)^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} A\left(\boldsymbol{p}, \frac{\boldsymbol{x}+\boldsymbol{z}}{2}\right) e^{\frac{\mathrm{i}}{\hbar}\boldsymbol{p}(\boldsymbol{x}-\boldsymbol{z})} \Psi(\boldsymbol{z}) \,\mathrm{d}^d \boldsymbol{z} \,\mathrm{d}^d \boldsymbol{p} \,. \tag{A.1}$$

If  $\Psi$  is a multi-component object, e.g.  $\Psi \in L^2(\mathbb{R}^d) \otimes \mathbb{C}^{2s+1}$ , then  $A(\boldsymbol{p}, \boldsymbol{x})$  is matrix valued. Reverting this reasoning, one can also associate an operator  $\hat{A}$  with a more general symbol  $A(\boldsymbol{p}, \boldsymbol{x})$ , which does not necessarily correspond to a differential operator, via (A.1). This procedure is known as Weyl quantisation and certain properties of symbols translate to properties of the operators, thus leading to so-called pseudo-differential operators, see e.g. [52] for an introduction.

If an operator  $\hat{A}$  can be represented by an integral kernel  $K_A(\boldsymbol{x}, \boldsymbol{y})$ , i.e.

$$(\hat{A}\Psi)(\boldsymbol{x}) = \int_{\mathbb{R}^d} K_A(\boldsymbol{x}, \boldsymbol{y}) \,\Psi(\boldsymbol{y}) \,\mathrm{d}^d y \,, \tag{A.2}$$

one obtains its Weyl symbol from

$$A(\boldsymbol{p},\boldsymbol{x}) = \int_{\mathbb{R}^d} K_A\left(\boldsymbol{x} + \frac{\boldsymbol{z}}{2}, \boldsymbol{x} - \frac{\boldsymbol{z}}{2}\right) e^{-\frac{\mathrm{i}}{\hbar}\boldsymbol{z}\boldsymbol{p}} \mathrm{d}^d z \,. \tag{A.3}$$

Inverting this transformation yields

$$K_A(\boldsymbol{x}, \boldsymbol{y}) = \frac{1}{(2\pi\hbar)^d} \int_{\mathbb{R}^d} A\left(\boldsymbol{p}, \frac{\boldsymbol{x} + \boldsymbol{y}}{2}\right) e^{\frac{i}{\hbar}\boldsymbol{p}(\boldsymbol{x} - \boldsymbol{y})} d^d p.$$
(A.4)

If the symbol  $A(\boldsymbol{p}, \boldsymbol{x})$  has an expansion in powers of  $\hbar$ ,

$$A(\boldsymbol{p}, \boldsymbol{x}) = \sum_{k \ge 0} \hbar^k A_k(\boldsymbol{p}, \boldsymbol{x}), \qquad (A.5)$$

the corresponding operator  $\hat{A}$  is called a semiclassical Weyl operator. The leading and subleading terms  $A_0$  and  $A_1$  in the expansion (A.5) are known as the principal symbol and the subprincipal symbol, respectively.

Application of a semiclassical Weyl operator to a rapidly oscillating function

$$\Psi_{\rm sc}(\boldsymbol{x}) = a_{\hbar}(\boldsymbol{x}) e^{\frac{i}{\hbar}S(\boldsymbol{x})} , \quad a_{\hbar}(\boldsymbol{x}) = \sum_{k=0}^{\infty} \left(\frac{\hbar}{i}\right)^k a_k(\boldsymbol{x}) .$$
(A.6)

is governed by the following theorem; the corresponding statement in a slightly different setting can, e.g., be found in [53, chapter 4.3].

**Theorem 6.** Applying a semiclassical Weyl operator  $\hat{A}$  with a symbol  $A(\boldsymbol{p}, \boldsymbol{x})$  of the form (A.5) to a wave function of type (A.6) yields in leading orders as  $\hbar \to 0$ ,

$$(\hat{A}\Psi_{sc})(\boldsymbol{x}) = \left\{ A_0(\nabla_{\boldsymbol{x}}S(\boldsymbol{x}), \boldsymbol{x}) a_0(\boldsymbol{x}) + \frac{\hbar}{i} \left[ A_0(\nabla_{\boldsymbol{x}}S(\boldsymbol{x}), \boldsymbol{x}) a_1(\boldsymbol{x}) + (\nabla_{\boldsymbol{p}}A_0)(\nabla_{\boldsymbol{x}}S(\boldsymbol{x}), \boldsymbol{x}) \nabla_{\boldsymbol{x}}a_0(\boldsymbol{x}) + \frac{1}{2}a_0(\boldsymbol{x})[\nabla_{\boldsymbol{x}}(\nabla_{\boldsymbol{p}}A_0)(\nabla_{\boldsymbol{x}}S(\boldsymbol{x}), \boldsymbol{x})] + A_1(\nabla_{\boldsymbol{x}}S(\boldsymbol{x}), \boldsymbol{x}) a_0(\boldsymbol{x}) \right] + \mathcal{O}(\hbar^2) \right\} e^{\frac{i}{\hbar}S(\boldsymbol{x})}.$$
(A.7)

Notice that the nesting of brackets in the fourth term on the r.h.s. indicates that the gradient with respect to p is taken before we set  $p = \nabla_x S$ , whereas the gradient with respect to x is only taken after doing so, i.e.

$$\left[\nabla_{\boldsymbol{x}}(\nabla_{\boldsymbol{p}}A_0)(\nabla_{\boldsymbol{x}}S(\boldsymbol{x}),\boldsymbol{x})\right] = \sum_{j=1}^d \frac{\partial^2 A_0}{\partial p_j \partial x_j} (\nabla_{\boldsymbol{x}}S(\boldsymbol{x}),\boldsymbol{x}) + \sum_{j=1}^d \sum_{j=1}^d \frac{\partial^2 A_0}{\partial p_j \partial p_k} (\nabla_{\boldsymbol{x}}S(\boldsymbol{x}),\boldsymbol{x}) \frac{\partial^2 S}{\partial x_j \partial x_k} (\boldsymbol{x}) .$$
(A.8)

For the proof we refer the reader to [53, chapter 4.3].

## References

- M. C. Gutzwiller: Chaos in Classical and Quantum Mechanics, Springer-Verlag, New York, (1990).
- [2] N. Bohr: On the Constitution of Atoms and Molecules. I., Phil. Mag. 26 (1913) 1–24.
- [3] M. Planck: Die physikalische Struktur des Phasenraumes, Ann. Phys. (Leipzig) 50 (1916) 385-418.
- 4 P. S. Epstein: Zur Theorie des Starkeffektes, Ann. Phys. (Leipzig) 50 (1916) 489–520.
- [5] K. Schwarzschild: Zur Quantenhypothese, Sitzungsb. Königl. Preuss. Akad. Wiss. (1916) 548-569.
- [6] A. Sommerfeld: Zur Quantentheorie der Spektrallinien, Ann. Phys. (Leipzig) 51 (1916) 1–94, 125–167.
- [7] A. Einstein: Zum Quantensatz von Sommerfeld und Epstein, Verh. Dtsch. Phys. Ges. 19 (1917) 82–92.
- [8] J. Liouville: Sur l'intégration des équations différentielles de la Dynamique, J. Math. Pures Appl. 20 (1855) 137-138.
- [9] V. I. Arnold: Mathematical Methods of Classical Mechanics, Springer-Verlag, New York, (1978).
- [10] G. Wentzel: Eine Verallgemeinerung der Quantenbedingungen für die Zwecke der Wellenmechanik, Z. Physik 38 (1926) 519–529.
- [11] H. A. Kramers: Wellenmechanik und halbzahlige Quantisierung, Z. Physik 39 (1926) 828-840.
- [12] L. Brillouin: La mécanique ondulatoire de Schrödinger; une méthode générale par approximations successives, Compt. Rend. 183 (1926) 24-26.
- [13] H. Jeffreys: On certain approximate solutions of linear differential equations of the second order, Proc. London Math. Soc. 23 (1925) 428–436.
- [14] J. B. Keller: Corrected Bohr-Sommerfeld Quantum Conditions for Nonseparable Systems, Ann. Phys. (NY) 4 (1958) 180–185.
- [15] V. P. Maslov: Théorie des perturbations et méthodes asymptotiques, Dunod, Paris, (1972).
- [16] V. P. Maslov and M. V. Fedoriuk: Semi-Classical Approximation in Quantum Mechanics, D. Reidel, Dodrecht, (1981).

- [17] J. B. Keller: Semiclassical mechanics, SIAM Rev. 27 (1985) 485–504.
- [18] W. Gordon: Die Energieniveaus des Wasserstoffatoms nach der Diracschen Quantentheorie des Elektrons, Z. Phys. 48 (1928) 11–14.
- [19] C. G. Darwin: The Wave Equations of the Electron, Proc. R. Soc. London Ser. A 118 (1928) 654-680.
- [20] W. Pauli: Diracs Wellengleichung des Elektrons und geometrische Optik, Helv. Phys. Acta 5 (1932) 179–199.
- [21] S. I. Rubinow and J. B. Keller: Asymptotic Solution of the Dirac Equation, Phys. Rev. 131 (1963) 2789-2796.
- [22] K. Yabana and H. Horiuchi: Adiabatic Viewpoint for the WKB Treatment of Coupled Channel System, Prog. Theor. Phys. 75 (1986) 592–618.
- [23] M. V. Berry: Quantal phase factors accompanying adiabatic changes, Proc. R. Soc. London Ser. A 392 (1984) 45–57.
- [24] A. Shapere and F. Wilczek: Geometric phases in physics, World Scientific Publishing, Singapore, (1989).
- [25] H. Kuratsuji and S. Iida: Effective Action for Adiabatic Process, Prog. Theor. Phys. 74 (1985) 439-445.
- [26] H. Kuratsuji and S. Iida: Deformation of symplectic structure and anomalous commutators in field theory, Phys. Rev. D 37 (1988) 441-447.
- [27] R. G. Littlejohn and W. G. Flynn: Geometric phases in the asymptotic theory of coupled wave equations, Phys. Rev. A 44 (1991) 5239-5256.
- [28] R. G. Littlejohn and W. G. Flynn: Geometric Phases in the Bohr-Sommerfeld Quantization of Multicomponent Wave Fields, Phys. Rev. Lett. 66 (1991) 2839-2842.
- [29] C. Emmrich and A. Weinstein: Geometry of the transport equation in multicomponent WKB approximations, Commun. Math. Phys. 176 (1996) 701-711.
- [30] S. Keppeler: Torus Quantization for Spinning Particles, Phys. Rev. Lett. 89 (2002) 210405.
- [31] H. Goldstein: Classical Mechanics, Addison-Wesley, Reading, Massachusetts, 2nd edn., (1980).
- [32] L. H. Thomas: The Motion of the Spinning Electron, Nature **117** (1926) 514.
- [33] L. H. Thomas: The Kinematics of an Electron with an Axis, Philos. Mag. 3 (1927) 1-22.

- [34] V. Bargman, L. Michel and V. L. Telegdi: Precession of the polarization of particles moving in a homogeneous electromagnetic field, Phys. Rev. Lett. 2 (1959) 435-436.
- [35] R. H. Good: The Generalization of the WKB Method to Radial Wave Equations, Phys. Rev. 90 (1953) 131–137.
- [36] M. Rosen and D. R. Yennie: A Modified WKB Approximation for Phase Shifts, J. Math. Phys. 5 (1964) 1505-1515.
- [37] P. Lu: Derivation of Sommerfeld-Dirac Fine-Structure Formula by WKB Method, Phys. Rev. A 1 (1970) 1283-1285.
- [38] K. Yajima: The quasiclassical approximation to Dirac equation. I, J. Fac. Sci. Univ. Tokyo Sect. IA Math. 29 (1982) 161–194.
- [39] V. G. Bagrov, V. V. Belov, A. Y. Trivonov and A. A. Yevseyevich: Quantization of closed orbits in Dirac Theory by Maslov's complex germ method, J. Phys. A 27 (1994) 1021–1043.
- [40] V. G. Bagrov, V. V. Belov, A. Y. Trivonov and A. A. Yevseyevich: Quasiclassical spectral series of the Dirac operators corresponding to quantized two-dimensional Lagrangian tori, J. Phys. A 27 (1994) 5273-5306.
- [41] H. Spohn: Semiclassical limit of the Dirac equation and spin precession, Ann. Phys. (NY) 282 (2000) 420-431.
- [42] J. Bolte and S. Keppeler: A semiclassical approach to the Dirac equation, Ann. Phys. (NY) 274 (1999) 125–162.
- [43] J. Bolte and S. Keppeler: Semiclassical form factor for chaotic systems with spin 1/2, J. Phys. A 32 (1999) 8863–8880.
- [44] J. Bolte and S. Keppeler: Semiclassical Time Evolution and Trace Formula for Relativistic Spin-1/2 Particles, Phys. Rev. Lett. 81 (1998) 1987–1991.
- [45] J. Bolte, R. Glaser and S. Keppeler: Quantum and classical ergodicity of spinning particles, Ann. Phys. (NY) 293 (2001) 1–14.
- [46] S. Keppeler: Spinning Particles: Semiclassical quantisation and spectral statistics, PhD thesis, Abteilung Theoretische Physik, Universität Ulm, (2002).
- [47] J. D. Bjorken and S. D. Drell: *Relativistic Quantum Mechanics*, McGraw-Hill, New York, St. Louis, San Francisco, (1964).
- [48] B. Thaller: *The Dirac Equation*, Springer-Verlag, Berlin, Heidelberg, (1992).
- [49] A. O. Barut and R. Rączka: Theory of group representations and applications, PWN-Polish Scientific Publishers, Warsaw, (1977).

- [50] P. Strange: Relativistic Quantum Mechanics with applications in condensed matter and atomic physics, Cambridge University Press, (1998).
- [51] A. Sommerfeld: Atombau und Spektrallinien, vol. I, Friedr. Viehweg & Sohn GmbH, Braunschweig, 8th edn., (1969).
- [52] G. B. Folland: *Harmonic Analysis in Phase Space*, no. 122 in Annals of Mathematics Studies, Princeton University Press, Princeton, (1989).
- [53] J. J. Duistermaat: Fourier Integral Operators, Birkhäuser, Boston, (1996).