We show that the geometric phase of Lévy-Leblond arises from a law of parallel transport for wave functions and point out that this phase belongs to a new class of geometric phases due to the presence of a quantum potential.

03.65.-w; 03.65.Vf

Geometric Phase and Quantum Potential

R. Dandoloff^{\dagger}

Laboratoire de Physique Théorique et Modélisation, Université de Cergy-Pontoise, 5 Mail Gay-Lussac, 95031 Cergy-Pontoise Cedex, France

 $\mathbf{2}$

Since M. Berry [1] has shown that the phase of the wave function may play an important role, there have been many papers investigating this effect, which now is widely known as Berry's phase. J.-M. Lévy-Leblond [2] has discovered yet another geometric phase. He has shown that a quanton that propagates through a tube, within which it is confined by impenetrable walls, acquires a phase when it comes out of the tube. For simplicity he has considered a tube with square section of side a. The tube has a length L. Before entering the tube the quanton's wave function is: $\phi = e^{\frac{i}{\hbar}pz}$, where zis the axis of the tube and p is its initial momentum. In the tube the wave function has the following form:

$$\psi = \sin(n_x \pi \frac{x}{a}) \sin(n_y \pi \frac{y}{a}) e^{\frac{i}{\hbar} p' z},\tag{1}$$

with appropriate transverse boundary conditions. After entering the tube the energy E of the quanton is unchanged but satisfies:

$$E = \frac{p^{\prime 2}}{2m} + (n_x^2 + n_y^2) \frac{\pi^2}{2ma}.$$
 (2)

For the simplest case: $n_x = n_y = 1$, Lévy-Leblond obtained the following additional phase for the wave function, after the quantum has left the tube:

$$\Delta \Phi = \frac{\pi^2 \hbar^2}{p a^2} L. \tag{3}$$

Subsequently R.E. Kastner [3] has related this to the quantum potential that arises in the tube. Now let us write the wave function in the tube in polar form: $\psi_1 = Re^{\frac{i}{\hbar}S + \frac{i}{\hbar}pz}$ (the eventual changes in the phase of the wave function, due to the presence of the tube, are now concentrated in S). In order to single out the influence of the tube on the wave function, we will write the wave function in the following form: $\psi_1 = \psi e^{\frac{i}{\hbar}pz}$. The quantum potential which corresponds to ψ is given by the following expression:

$$Q = -\frac{\hbar^2}{2m} \frac{\Delta R}{R} = \frac{\pi^2 \hbar^2}{ma^2}.$$
 (4)

Now, let us turn our attention to the laws of parallel transport for the wave function, related to the appearance of the different geometric phases. For the Berry's phase the law of parallel transport for the wave function has been established by B. Simon [4]:

$$Im < \psi | \dot{\psi} >= 0, \tag{5}$$

where (') stands for a time derivative. For a spin- $\frac{1}{2}$ in magnetic field this law of parallel transport is the analog of the Fermi-Walker parallel transport [5].

For the Lévy-Leblond's phase the law of parallel transport for the wave function ψ is given by the following expression:

$$Im < \psi |\dot{\psi}\rangle = -\frac{1}{\hbar}Q|\psi|^2.$$
(6)

In Lévy-Leblond's *gedanken experiment* the wave function acquires an additional phase after the quanton has left the tube:

$$\psi(t + \Delta t) = e^{-\frac{i}{\hbar}Q\Delta t}\psi(t),\tag{7}$$

which after expansion in Δt leads to the law of parallel transport: $Im < \psi |\dot{\psi} > = -\frac{1}{\hbar}Q|\psi|^2$ eqn.(6). Indeed:

$$Q\Delta t = \frac{\pi^2 \hbar^2}{ma^2} \Delta t = \frac{\pi^2 \hbar^2}{ma^2} \frac{mL}{p} = \frac{\pi^2 \hbar^2}{pa^2} L = \Delta \Phi \qquad (8).$$

If we use the polar form for the wave function, eq.(6) gives: $\frac{\partial S}{\partial t} = -Q$. This means that this new law of parallel transport eliminates the quantum potential from the "quantum" Hamilton-Jacobi equation and we are left with the classical Hamilton-Jacobi equation for a free particle. The whole "quantum information" is now carried by the phase of the wave function.

Now, after we have established the law of parallel transport for ψ in the case of the geometric phase of Lévy-Leblond, we see that the nature of this phase is quite different from the Berry's phase. The appearance of the geometric phase of Lévy-Leblond is related to the presence or not of constraints in the system. In this case the quanton is constrained to propagate in a tube.

Let us now turn our attention to quite a different type of constrained system and see how a new geometric phase, based on the new law of parallel transport may appear there. We will consider a quantum particle constrained to be on a circle [6–8]. In this case the wave function has the form: $\psi \sim sin(\frac{ns}{\rho_0})$, where ρ_0 is the radius of the circle and s is the arc-length with origin at the tangent point. In this way the wave function on the circle has a node at the tangent point. In the case of a particle constrained to move on a circle the value $n = \frac{1}{2}$ is also allowed [7,8]. The corresponding quantum potential for $n = \frac{1}{2}$ now is:

$$Q = \frac{\hbar^2}{8m\rho_0^2}.$$
(9)

The quantum potential Q is exactly equal to the constant E_0 which appears in the Hamiltonian for a particle on a circle with radius ρ_0 following the Dirac quantization procedure for constrained systems [6]. The phase which would acquire a quanton travelling along the circle is then:

$$Q\frac{2\pi\rho_0 m}{p} = \frac{\pi\hbar^2}{4\rho_0 p}.$$
(10)

Note that if the circle becomes very small, then the geometric phase can not get bigger than $\sim \frac{\hbar}{m}$. This limit is imposed by the Heisenberg's uncertainty relation $\rho_0 p \sim \hbar$. This is not the case for the Lévy-Leblond's phase which can get very big provided $L \gg a$. One can imagine the following gedanken experiment in order to test the appearance of this new phase: a flow of quantons propagating on a straight line with a circle of radius ρ_0 tangent to it. The scattering at the point where the circle touches the straight line will split this flow into two: one following the straight line and the other following the circle and then the straight line (we are not interested in the reflected flow which goes back). At the exit these two flows will produce an interference picture which may then be measured.



FIG. 1. A flow of quantons propagating along a straight line and a tangent circle.

As a conclusion, we have established the law of parallel transport for the wave function in the case of Lévy-Leblond's geometric phase and have shown that this phase is not another manifestation of Berry's phase. This phase is directly related to the constraint imposed on the system. We have also shown that another constrained system, a quantum particle on a circle, may exhibit the same kind of geometric phase. The quantum potential gives a direct insight into the problem why there is an additional constant appearing in the Hamiltonian of this system. Acknowledgments:The author thanks T.T.Truong for discussions.

- [†] rossen.dandoloff@ptm.u-cergy.fr
- [1] M.V. Berry, Proc.R.Soc.London, A392, 45, (1984).
- [2] J.-M. Lévy-Leblond, Phys.Lett. A 125, 44, (1987).
- [3] R.E. Kastner, AM.J.Phys; **61**, 852, (1993).
- [4] B. Simon, *Phys. Rev. Lett.*, **51**, 2167, (1983).
- [5] R. Dandoloff, *Phys.Lett.* A, 139, 19, (1989).
- [6] A. Scardicchio, *Phys.Lett.* A, 300, 7, (2002).
- [7] V.B.Ho and M.Morgan, J.Phys.A:Math.Gen.,29, 1497, (1996).
- [8] K.Kowalski, J. Rembielinski and L.C.Papalucas, *J.Phys.* A: Math.Gen., 29, 4149, (1996).