

Analytical formula connecting entangled state and the closest disentangled state

Satoshi Ishizaka

Fundamental Research Laboratories, NEC Corporation,
34 Miyukigaoka, Tsukuba, Ibaraki, 305-8501 Japan

(Dated: November 20, 2018)

The separable state closest to a given entangled state in the relative entropy measure is called the closest disentangled state. We provide an analytical formula connecting the entangled state and the closest disentangled state in two qubits. Using this formula, when any disentangled state (σ) located at the entangle-disentangle boundary is given, entangled states to which σ is closest can be obtained analytically. Further, this formula naturally defines the direction normal to the boundary surface. The direction is uniquely determined by σ in almost all cases.

PACS numbers: 03.67.-a, 03.65.Ud

Quantum entanglement is the most striking feature of quantum mechanics. Several measures have been proposed to quantify the amount of entanglement [1]. One of the important measures is relative entropy of entanglement [2, 3], which is defined as

$$E_R(\varrho) = \min_{\sigma \in \mathcal{D}} S(\varrho||\sigma) = \min_{\sigma \in \mathcal{D}} [\text{Tr} \varrho \ln \varrho - \text{Tr} \varrho \ln \sigma], \quad (1)$$

where the minimization is performed over all density matrices in the set of disentangled states (ln denotes the natural logarithm throughout this paper). The states achieving the minimum are called the closest disentangled states. The efficient algorithm for the numerical minimization has been proposed in [4]. However, the analytical minimization is difficult in general, and the analytical solutions only for several classes of states have been shown so far.

In this paper, we provide an analytical formula connecting entangled states and the closest disentangled state in two qubits. Using this formula, when any disentangled state (σ) located at the entangle-disentangle boundary is given, entangled states ϱ to which σ is closest can be obtained analytically. Further, this formula naturally defines the direction normal to the boundary surface when relative entropy is regarded as the distant measure. The direction is uniquely determined by σ in almost all cases. The discussions in this paper are limited to the case of two qubits.

The strategy of the minimization is based on the following fact: *Any full rank density matrix can be transformed into a unique Bell diagonal state by applying a suitable local filtering operation* [5, 6]. The explicit formulation using the Lorentz matrices has been shown in [5], and that using the Wootters basis [7] has been shown in [6]. The entanglement manipulation by local filtering on a single copy was originally considered in [8, 9]. Further, the connection between the Wootters basis and Lewenstein-Sanpera decomposition [10] has been shown in [11]. According to these results, any full rank density matrix can be written as $\varrho = \frac{1}{N} (F_A \otimes F_B) \varrho_{\text{BD}} (F_A^\dagger \otimes F_B^\dagger)$, where $N = \text{Tr}[(F_A \otimes F_B) \varrho_{\text{BD}} (F_A^\dagger \otimes F_B^\dagger)]$ and $\varrho_{\text{BD}} = \sum_{i=0}^3 p_i |e_i\rangle\langle e_i|$ is the Bell diagonal state. ϱ_{BD} can be always chosen in a canonical form, in which p_0 is maximum ($p_0 \geq p_1, p_2, p_3$) and $|e_i\rangle$ is a fixed set of Bell basis, since suitable local unitary operations transform

any Bell diagonal state into the canonical form [12, 13]. On the other hand, ϱ can be always expressed using the Wootters basis as $\varrho = \sum_{i=0}^3 \lambda_i |\phi_i\rangle\langle \phi_i|$ [7], where $\langle \phi_i | \tilde{\phi}_j \rangle = \langle \tilde{\phi}_i | \phi_j \rangle = \delta_{ij}$ defining the tilde operation as $|\tilde{\psi}\rangle = (\sigma_2 \otimes \sigma_2) |\psi^*\rangle$. The Bell basis is one of the Wootters basis by itself (adding a suitable global phase if necessary), and hence $|\tilde{e}_i\rangle = |e_i\rangle$. The concurrence of ϱ is $C = \lambda_0 - \lambda_1 - \lambda_2 - \lambda_3$. Although the actual definition of the concurrence is $\max(\lambda_0 - \lambda_1 - \lambda_2 - \lambda_3, 0)$ [7], we define it here such that C can take negative when ϱ is disentangled. Further, we can put $\det F_A = \det F_B = 1$ without loss of generality. In this choice, $|\phi_i\rangle = (F_A \otimes F_B) |e_i\rangle$ (using $A^T \sigma_2 A = (\det A) \sigma_2$, it can be checked that $|\phi_i\rangle = (F_A \otimes F_B) |e_i\rangle$ satisfy $\langle \phi_i | \tilde{\phi}_j \rangle = \delta_{ij}$) and $\lambda_i = p_i/N$ [5, 6].

Considering these facts carefully, the following structure of the Hilbert space is revealed: The total 15-dimensional space is divided into subspaces in such a way that only the states in the same subspace can be transformed to each other by local filtering (we only consider the full rank density matrices for the sake of simplicity [5, 6]). Each subspace contains the canonical Bell diagonal states having the same p_i only (we do not care about the exchange among p_1, p_2 , and p_3). Therefore, the subspaces can be specified by three parameters (p_i with $\sum p_i = 1$), and each subspace constitutes a 12-dimensional manifold. Further, since $C > 0$ for entangled states and $C \leq 0$ for disentangled states, and the local filtering does not change the sign of C (see Theorem 1 in [5]), we can define *boundary subspaces*, in which all the states have just zero of C . The boundary subspaces are specified by two parameters, and the assemble of the boundary subspaces constitutes the 14-dimensional boundary surface separating the region of entangled states and disentangled states.

Then, we perform the minimization of $S(\varrho||\sigma)$ separately: First we obtain the extremal conditions by minimizing within a fixed subspace, and second we obtain the conditions to determine the boundary subspace containing the closest disentangled state. Finally we solve the equations obtained. Hereafter, the entangled state and the corresponding closest disentangled state is denoted by ϱ and σ , respectively. Further, we first assume that σ is full rank. The case of σ with lower rank will be discussed in the last part of this paper.

We have already performed the first step and obtained the set of equations σ must satisfy [14]. We briefly repeat the results here. The states obtained from σ by local filtering belong to the same subspace as σ by its definition. Let us consider Bob's local filtering of $\sigma' = \frac{1}{N}(I \otimes e^{it\vec{n}\cdot\vec{\sigma}/2})\sigma(I \otimes e^{it\vec{n}\cdot\vec{\sigma}/2})$, where t is a real parameter and $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ is the vector of Pauli matrices. In order that $S(\varrho|\sigma)$ is minimum, the linear coefficient of t in the expansion of $\text{Tr}\varrho \ln \sigma'$ must be zero. As the result, using $\ln(X + tY) = \ln X + t \int_0^\infty \frac{1}{X+z} Y \frac{1}{X+z} dz + \dots$ we obtained

$$\text{Tr}(I \otimes \sigma_i)Z\sigma + \text{Tr}Z(I \otimes \sigma_i)\sigma = 0 \quad \text{for } i = 1, 2, 3 \quad (2)$$

where

$$Z \equiv \int_0^\infty \frac{1}{\sigma+z} \varrho \frac{1}{\sigma+z} dz - I \quad (3)$$

(Eq. (2) holds for $i = 0$ also, but we omitted it since $\text{Tr}Z\sigma = 0$ is obvious from Eq. (3)). In the same manner, considering Bob's local unitary transformation of $\sigma' = (I \otimes e^{it\vec{n}\cdot\vec{\sigma}/2})\sigma(I \otimes e^{-it\vec{n}\cdot\vec{\sigma}/2})$, we obtained

$$\text{Tr}(I \otimes \sigma_i)Z\sigma - \text{Tr}Z(I \otimes \sigma_i)\sigma = 0 \quad \text{for } i = 1, 2, 3. \quad (4)$$

From Eqs. (2) and (4), we have

$$\text{Tr}(I \otimes \sigma_i)Z\sigma = 0, \quad (5)$$

which is sufficient for $\text{Tr}Z(I \otimes \sigma_i)\sigma = 0$ since Z is Hermitian. Therefore, σ must satisfy Eqs. (5) and Alice's counterparts

$$\text{Tr}(\sigma_i \otimes I)Z\sigma = 0. \quad (6)$$

These are fundamental equations for the closest disentangled state (and even for the closest positive partial transposed states), because these equations must hold in any multi-party systems and any higher dimensional systems [14].

The second step is to obtain the conditions to determine the boundary subspace containing σ . The canonical Bell diagonal state in the boundary subspace must have zero of $C = p_0 - p_1 - p_2 - p_3$, but $\sum p_i = 1$ due to the normalization. As the result, p_0 must be $1/2$ (this is also obvious from the fact that the concurrence of the Bell diagonal state is $2p_0 - 1$ [7, 15]). Therefore, σ , which belongs to the boundary subspace, must be obtained by local filtering from $\sigma_{\text{BD}} = \frac{1}{2}|e_0\rangle\langle e_0| + \sum_{i=1}^3 p_i|e_i\rangle\langle e_i|$, and hence σ must be written as

$$\begin{aligned} \sigma &= \frac{1}{N}(F_A \otimes F_B)\sigma_{\text{BD}}(F_A^\dagger \otimes F_B^\dagger) \\ &= \frac{1}{2N}|\phi_0\rangle\langle\phi_0| + \sum_{i=1}^3 \frac{p_i}{N}|\phi_i\rangle\langle\phi_i|, \end{aligned} \quad (7)$$

where N is again normalization, and the set of $|\phi_i\rangle$'s is the Wootters basis of σ .

Adopting this parameterization, to determine the subspace containing σ is equivalent to determine p_1 , p_2 , and p_3 under the constraint of $p_1 + p_2 + p_3 = 1/2$ and

$0 \leq p_i \leq 1/2$. However, we assumed that σ is full rank, which ensures that the minimum of $S(\varrho|\sigma)$ is achieved at $0 < p_i < 1/2$. Therefore, we can simply minimize

$$f \equiv -\text{Tr}\rho \ln \sigma + l(p_1 + p_2 + p_3 - 1/2), \quad (8)$$

with l being a Lagrange multiplier. Considering the small change of p_i (note that N also contains p_i), we obtain

$$\frac{1}{N}\text{Tr}Z|\phi_i\rangle\langle\phi_i| = l \quad \text{for } i = 1, 2, 3. \quad (9)$$

Further, multiplying p_i on both side of the above equations and adding them, we find

$$\sum_{i=1}^3 \frac{p_i}{N}\text{Tr}Z|\phi_i\rangle\langle\phi_i| = \text{Tr}Z\sigma - \frac{1}{2N}\text{Tr}Z|\phi_0\rangle\langle\phi_0| = \frac{l}{2}, \quad (10)$$

and we finally obtain the extremal conditions as

$$\langle\phi_i|Z|\phi_i\rangle + \langle\phi_0|Z|\phi_0\rangle = 0 \quad \text{for } i = 1, 2, 3. \quad (11)$$

There are three equations in the above, in spite that the boundary subspaces are specified by two parameters as mentioned before. This is because Eqs. (11) includes the trace condition of $\text{Tr}\varrho = \text{Tr}\sigma$.

Then, all equations of the extremal conditions are Eqs. (5), (6), and (11) with $p_1 + p_2 + p_3 = 1/2$ (and thus $\lambda_0 - \lambda_1 - \lambda_2 - \lambda_3 = 0$). The total number of these equations is 16 including the normalization condition. These must be solved for any given ϱ in order to obtain a true closed formula for the closest disentangled state (and for $E_R(\varrho)$). However, since it is a hard problem unfortunately, we follow the strategy in [16]. Namely, for any given σ , we obtain the set of ϱ for which $S(\varrho|\sigma)$ is minimum. We can obtain it in an analytical form as shown below.

Let us assume that σ with $C=0$ is given. Here, we still assume that σ is full rank. It is convenient to express σ in two representations:

$$\sigma = \sum_{ij} \Lambda_{ij}|\phi_i\rangle\langle\phi_j| = \sum_{ij} \Gamma_{ij}|i\rangle\langle j|, \quad (12)$$

where $|\phi_i\rangle$'s and $|i\rangle$'s are the Wootters basis and eigenstates of σ , respectively, and hence $\Lambda_{ij} = \lambda_i\delta_{ij}$ and $\Gamma_{ij} = \gamma_i\delta_{ij}$ with γ_i 's being eigenvalues of σ . These two basis sets are connected through a unitary matrix as $\sqrt{\lambda_i}|\phi_i\rangle = \sum_j V_{ij}^* \sqrt{\gamma_j}|j\rangle$ [7]. Since $|\phi_i\rangle$'s are non-orthogonal in general, it is also convenient to introduce the matrices describing the non-orthogonality as follows:

$$Q_{ij} = \langle\phi_i|\phi_j\rangle, \quad \Pi_{ij} = \langle\tilde{\phi}_i|\tilde{\phi}_j\rangle. \quad (13)$$

Using the completeness of $I = \sum_i |\phi_i\rangle\langle\tilde{\phi}_i| = \sum_i |\tilde{\phi}_i\rangle\langle\phi_i|$ [11, 17] and $I = \sum_{ij} Q_{ij}|\tilde{\phi}_i\rangle\langle\tilde{\phi}_j| = \sum_{ij} \Pi_{ij}|\phi_i\rangle\langle\phi_j|$, it is easy to check that Hermitian Q and Π satisfy $QQ^T = I$, $\Pi = Q^*$, and $Q\Pi = \Pi Q = I$. Further,

$$\Gamma = V^\dagger \sqrt{\Lambda} Q \sqrt{\Lambda} V = U \sqrt{Q} \Lambda \sqrt{Q} U^\dagger, \quad (14)$$

where we introduced the unitary U by the singular value decomposition of $U\sqrt{Q}\sqrt{\Lambda}V = \sqrt{\Gamma}$. Then, we shall obtain ϱ in the same representation as σ . Namely, using $|\phi_i\rangle$ and $|i\rangle$ of σ ,

$$\varrho = \sum_{ij} R_{ij}^\Lambda |\phi_i\rangle\langle\phi_j| = \sum_{ij} R_{ij}^E |i\rangle\langle j|. \quad (15)$$

Hereafter, the coefficient matrix in the Wootters basis and eigenstates is denoted by suffix Λ and E , respectively. Two representations are connected through $R^E = \sqrt{\Gamma}V^\dagger\sqrt{\Lambda}R^\Lambda\sqrt{\Lambda}V\sqrt{\Gamma} = U\sqrt{Q}R^\Lambda\sqrt{Q}U^\dagger$.

Viewing the form of Eqs. (5), (6), and (11), it is found that, when ϱ is a solution, $\varrho' = x\varrho + (1-x)\sigma$ is also a solution, since $Z \rightarrow xZ$ for $\varrho \rightarrow x\varrho + (1-x)\sigma$. This implies that the normalization of Z can be taken as a free parameter, and we express Z as

$$\langle i|Z|j\rangle = \int_0^\infty \frac{1}{\gamma_i + z} R_{ij}^E \frac{1}{\gamma_j + z} dz - \delta_{ij} \equiv xW_{ij}^E. \quad (16)$$

Then, we have $xW^E = [R^E \circ \hat{G}] - I$, where $A \circ B$ is the Hadamard product defined as $[A \circ B]_{ij} = A_{ij}B_{ij}$, $\hat{G}_{ij} = 1/G_{ij}$, and

$$G_{ij} \equiv \begin{cases} \gamma_i & \text{for } \gamma_i = \gamma_j \\ \frac{\gamma_i - \gamma_j}{\ln \gamma_i - \ln \gamma_j} & \text{for } \gamma_i \neq \gamma_j. \end{cases} \quad (17)$$

By inverting the above (we can safely invert since $G_{ij} > 0$ by $\gamma_i > 0$), we have

$$R^E = \Gamma + x[W^E \circ G], \quad (18)$$

and therefore the problem is reduced to find W^E satisfying the set of all extremal conditions.

For this purpose, it is convenient to express Z in the $|\tilde{\phi}_i\rangle$ basis as $Z = \sum xW_{ij}|\tilde{\phi}_i\rangle\langle\tilde{\phi}_j|$. Then, the extremal conditions of Eqs. (11) become quite simple:

$$W_{ii} = -W_{00} \quad \text{for } i = 1, 2, 3. \quad (19)$$

Since the normalization of W was absorbed in x , we adopt $W_{00} = 1$ and the diagonal elements of W are determined to be $\{1, -1, -1, -1\}$. The remaining extremal conditions are Eqs. (5) and (6), those are

$$\begin{cases} \sum_{lm} \langle \phi_l | I \otimes \sigma_i | \tilde{\phi}_m \rangle W_{ml} \lambda_l = 0 \\ \sum_{lm} \langle \phi_l | \sigma_i \otimes I | \tilde{\phi}_m \rangle W_{ml} \lambda_l = 0. \end{cases} \quad (20)$$

On the other hand, by virtue of $F_A^\dagger \tilde{F}_A = F_A^\dagger \sigma_2 F_A^* \sigma_2 = I$ [5, 6, 17] (we put $\det F_A = \det F_B = 1$), we find

$$\begin{aligned} \langle \phi_l | I \otimes \sigma_i | \tilde{\phi}_l \rangle &= \langle e_l | F_A^\dagger \tilde{F}_A \otimes F_B^\dagger \sigma_i \tilde{F}_B | \tilde{e}_l \rangle \\ &= \text{Tr}(I \otimes F_B^\dagger \sigma_i \tilde{F}_B) |e_l\rangle\langle e_l| \\ &= \frac{1}{2} \text{Tr} \sigma_i \sigma_2 F_B^* \sigma_2 F_B^\dagger = \frac{1}{2} \text{Tr} \sigma_i = 0, \end{aligned}$$

and therefore, Eqs. (20) are the set of linear equations only for the off-diagonal elements of W . As the result, $W_{m \neq l} = 0$ obviously satisfy Eqs. (20). After all,

$$W = \text{diag}\{1, -1, -1, -1\} \quad (21)$$

is a solution of all extremal conditions. Since $W^E = U\sqrt{\Pi}W\sqrt{\Pi}U^\dagger$, we finally obtain a relation between ϱ and σ in an analytical form as

$$\begin{aligned} R^E &= \Gamma + x\Delta^E \\ \Delta^E &= [(U\sqrt{\Pi}W\sqrt{\Pi}U^\dagger) \circ G] \end{aligned} \quad (22)$$

with G by Eq. (17), W by Eq. (21), and Π by Eq. (13). It should be noted that the phases in U , as well as V , should be chosen so that $\langle \phi_i | \tilde{\phi}_j \rangle = \delta_{ij}$. In order to avoid the complication, it is convenient to express it in the Wootters basis, that is

$$\begin{aligned} R^\Lambda &= \Lambda + x\Delta^\Lambda \\ \Delta^\Lambda &= \sqrt{\Pi}U^\dagger [(U\sqrt{\Pi}W\sqrt{\Pi}U^\dagger) \circ G] U\sqrt{\Pi}. \end{aligned} \quad (23)$$

In this expression, any U satisfying $U\sqrt{Q}\Lambda\sqrt{Q}U^\dagger = \Gamma$ can be used safely. From the relation, it can be seen that the non-orthogonality of the Wootters basis plays an important role.

The trace of the matrix $\delta = \sum_{ij} \Delta_{ij}^E |i\rangle\langle j|$ is confirmed to be zero as

$$\begin{aligned} \text{Tr} \Delta^E &= \sum_i [U\sqrt{\Pi}W\sqrt{\Pi}U^\dagger]_{ii} \gamma_i = \text{Tr} U\sqrt{\Pi}W\sqrt{\Pi}U^\dagger \Gamma \\ &= \text{Tr} W\Lambda = \lambda_0 - \lambda_1 - \lambda_2 - \lambda_3 = 0, \end{aligned} \quad (24)$$

and δ obviously plays the role of the direction *normal* to the boundary surface at the position of σ , when $S(\varrho||\sigma)$ is regarded as the distant measure between states. Which side does the entangled region spread? For an infinitesimally small $|x|$, by treating Δ^Λ as the perturbation, the diagonal elements of Δ^Λ contribute to the concurrence of ϱ in the lowest order of $|x|$, and then

$$\begin{aligned} C(\varrho) &\sim x\Delta C \\ \Delta C &= \text{Tr} \Delta^\Lambda W = \sum_{ij} |[U\sqrt{\Pi}W\sqrt{\Pi}U^\dagger]_{ij}|^2 G_{ij} > 0, \end{aligned}$$

since $G_{ij} > 0$ and $U\sqrt{\Pi}W\sqrt{\Pi}U^\dagger$ is not zero operator (it is invertible). Therefore, the entangled region always spreads in $x > 0$. When σ is full rank, we can always find the positive ϱ for some infinitesimally small $|x|$. This implies that there are both entangled states and disentangled states in the vicinity of σ . Therefore, the full rank σ with $C=0$ is the true boundary state separating the entangled and disentangled region. Further, $E_R(\varrho)$ for $|x| \ll 1$ is also calculated as

$$S(\varrho||\sigma) \sim x \text{Tr}(Z + I)\delta = x^2 \text{Tr} W\Delta^\Lambda = x^2 \Delta C. \quad (25)$$

It is important to note that there are entangled states satisfying Eqs. (5), (6), and (11) for which σ is not optimal. However, the solution we obtained are not the case: σ is indeed optimal for $\varrho = \sigma + x\delta$ with any $x \geq 0$. This can be proven as follows. The equations we solved take into account any small deviation from σ along the boundary surface, and the solutions are ensured to be extremal along any path on the boundary. This extremeness does not immediately imply the global minimum of $S(\varrho||\sigma)$ among the all disentangled states by itself. However, for

our solution, $S(\varrho||\sigma + x\delta) > S(\varrho||\sigma)$ holds for $x < 0$ (in the vicinity of σ), and hence the relative entropy inside the boundary is larger than that on the boundary. In this situation, considering the semi-continuity and convexity of the relative entropy [18], it can be shown that the extremeness along the boundary must be local minimum among the disentangled states around σ , and it must be global minimum [3]. Further, the assumption we first made that ϱ has σ of full rank is also justified in the same reason.

It should be noted further about the uniqueness of W . (if it is not unique, there might exist entangled states other than those we obtained, for which σ is optimal). Since W is Hermitian, the number of real parameters of the off-diagonal elements is 12, and the number of linear Eqs. (20) is also 12. Therefore, Eq. (21) is a unique solution unless some of Eqs. (20) are dependent to each other. We have numerically generated random σ 's and confirmed that Eqs. (20) are always independent. However, such Monte Carlo type calculations cannot pick up the states of measure zero, such as those having some special symmetry, and the other solutions ϱ have not been completely excluded. Further, it is an open question whether the solution we obtained is enough to express whole entangled states (does $\sigma + |x|\delta$ span whole entangled states, when σ moves on the boundary?)

In so far, we solely discussed the case that σ is full rank. In the case of σ with a lower rank (denoted by σ_L), let us consider a sequence constituted by all full rank σ_n 's converging as $\sigma_n \rightarrow \sigma_L$. We can always find such sequences because there always exist full rank boundary states in the vicinity of σ_L (if λ_k of σ_L is zero, adding infinitesimally small portion of $|\phi_k\rangle\langle\phi_k|$ to σ_L). In this sequence, ϱ_n obtained from Eq. (22) also converge as $\varrho_n \rightarrow \varrho_\infty$ (since G_{ij} is non-singular). Then, the continuity of the relative entropy of entanglement shown in [19] ensures that $E_R(\varrho_\infty) = S(\varrho_\infty||\sigma_L)$, and σ_L is one of the closest disentangled states of ϱ_∞ . After all, Eq. (22) must hold

under such limiting sequences even for low rank σ_L .

Finally, we show some simple examples of Eq. (23). The Wootters basis can be easily constructed by $|\phi_i\rangle = (F_A \otimes F_B)|e_i\rangle$. For $F_A = F_B = I$, σ becomes a Bell diagonal state, and $|\phi_i\rangle = |e_i\rangle$ constitute the orthogonal set. Then, $Q = \Pi = U = I$ and $\Gamma = \Lambda$, and we obtain

$$\begin{aligned} \Delta^\Lambda &= \text{diag}\{1/2, -\lambda_1, -\lambda_2, -\lambda_3\} \\ R^\Lambda &= \text{diag}\left\{\frac{1+x}{2}, (1-x)\lambda_1, (1-x)\lambda_2, (1-x)\lambda_3\right\}. \end{aligned}$$

For the choice of $F_A = F_B = \text{diag}\{t, 1/t\}$, $|e_0\rangle = i|\phi^+\rangle$, and $|e_1\rangle = (I \otimes \sigma_t)|\phi^+\rangle$, where $|\phi^+\rangle = (00 + |11\rangle)/\sqrt{2}$, we have

$$Q = \Pi^* = \frac{1}{2} \begin{pmatrix} t^4 + 1/t^4 & & -it^4 + i/t^4 \\ & 1 & \\ it^4 - i/t^4 & & t^4 + 1/t^4 \end{pmatrix}, \quad (26)$$

and the limiting sequence of $\lambda_1 = \lambda_2 \rightarrow 0$ and $\lambda_3 \rightarrow \lambda_0$ gives $\Gamma = \lambda_0 \text{diag}\{t^4, 0, 0, 1/t^4\}$, and

$$\begin{aligned} \Delta^\Lambda &\propto \text{diag}\{1, 0, 0, -1\} \\ R^\Lambda &= \text{diag}\{1 + x', 0, 0, 1 - x'\}. \end{aligned} \quad (27)$$

These examples reproduce the correct relation between Bell diagonal ϱ and σ [2], and between pure state and its closest disentangled state of rank 2 [3], respectively.

In summary, we provided an analytical formula connecting σ and ϱ in two qubits [Eq. (22)]. Using this formula, when σ with $C=0$ is given, ϱ for which $S(\varrho||\sigma)$ is minimum can be obtained analytically. Further, this formula naturally defines the vector (δ) normal to the entangle-disentangle boundary surface. The normal vector is uniquely determined by σ in almost all cases.

This work was supported by CREST, Japan Science and Technology Corporation (JST).

-
- [1] M. Horodecki, *Quant. Inf. Comp.* **1**, 3 (2001).
[2] V. Vedral, M. B. Plenio, M. A. Rippin, and P. L. Knight, *Phys. Rev. Lett.* **78**, 2275 (1997).
[3] V. Vedral and M. B. Plenio, *Phys. Rev. A* **57**, 1619 (1998).
[4] J. Řeháček and Z. Hradil, *Phys. Rev. Lett.* **90**, 127904-1 (2003).
[5] F. Verstraete, J. Dehaene, and B. DeMoor, *Phys. Rev. A* **64**, 010101 (2001).
[6] L.-X. Cen, N.-J. Wu, F.-H. Yang, and J.-H. An, *Phys. Rev. A* **65**, 52318 (2002).
[7] W. K. Wootters, *Phys. Rev. Lett.* **80**, 2245 (1998).
[8] N. Linden, S. Massar, and S. Popescu, *Phys. Rev. Lett.* **81**, 3279 (1998).
[9] A. Kent, N. Linden, and S. Massar, *Phys. Rev. Lett.* **83**, 2656 (1999).
[10] M. Lewenstein and A. Sanpera, *Phys. Rev. Lett.* **80**, 2261 (1998).
[11] S. Akhtarshenas and M. Jafarizadeh, *quant-ph/0211051*.
[12] Any Bell diagonal states can be transformed to the canonical Hilbert-Schmidt form where $|e_0\rangle = i|\phi^+\rangle$, and $|e_1\rangle = (I \otimes \sigma_t)|\phi^+\rangle$ [13]. Then, $I \otimes \sigma_i$ can transform $|e_i\rangle$ to $|e_0\rangle$ (the other bases are simply exchanged).
[13] R. Horodecki and M. Horodecki, *Phys. Rev. A* **54**, 1838 (1996).
[14] S. Ishizaka, *J. Phys. A: Math. Gen.* **35**, 8075 (2002).
[15] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, *Phys. Rev. A* **54**, 3824 (1996).
[16] E. M. Rains, *Phys. Rev. A* **60**, 179 (1999).
[17] L.-X. Cen and X.-Q. Li, and Y. Yan, *quant-ph/0212152*.
[18] M. Ohya and D. Petz, *Quantum Entropy and Its Use* (Springer-Verlag, 1993).
[19] M. J. Donald and M. Horodecki, *Phys. Lett. A* **264**, 257 (1999).