

Klein Paradox in Bosons

Partha Ghose and Manoj K. Samal

S. N. Bose National Centre for Basic Sciences, Block JD, Sector III, Salt Lake, Kolkata 700 098, India

Animesh Datta *

Department of Electrical Engineering, Indian Institute of Technology, Kanpur, 208 016, India

November 5, 2018

Abstract

We show analytically that the *Zitterbewegung* and Klein Paradox, such well known aspects of the Dirac Equation are not found in the case of Bosons. We use the Kemmer-Duffin-Harish Chandra formalism with β matrices to arrive at our results.

1 Introduction

In discussing problems and interactions in which the electron (or any fermion) is "spread out" over distances large compared to its Compton wavelength, we may simply ignore the existence of the uninterpreted negative-energy solutions of the one-particle relativistic Dirac Equation [1] and hope of obtaining physically sensible and accurate results. This will not work, however, in situations which find electrons (or fermions) localized to distances comparable with $\frac{\hbar}{mc}$. The negative frequency amplitudes will then be appreciable, *Zitterbewegung* [2] terms in the current important and indeed we shall find ourselves beset by paradoxes and dilemmas which defy interpretation within the framework of the one particle Dirac equation. A celebrated example of these difficulties is the **Klein Paradox** [3] for electrons, and in general for fermions. It is discussed in detail in numerous texts [4],[5] and resolved using the hole-theory which provides the treatment of the negative energy states. The actual paradox is that in the situations under question, the coefficient of reflection for an electron beam or any current from a barrier is more than one. It is to be noted that all the energies under consideration are below the threshold of pair creation.

In this Letter we address the same problem for bosons. We explicitly show that the reflection coefficient is less than one and hence the scope of any paradox is nonexistent.

The plan of the Letter is as follows. In Section 2 we explain the formalism in some detail for the massive and massless bosons. Section 3 contains our results about the absence of Klein Paradox like phenomenon in the case of massive bosons. The corresponding result for massless bosons is presented in Section 4. We make some concluding remarks in Section 5.

2 Formalism

In order to address the problem of Klein Paradox, a single-particle relativistic quantum mechanics is necessary. For fermions, such a framework is provided by the Dirac Equation, though it eventually turns out to be a multiparticle equation. Such a consistent relativistic quantum mechanics for spin 0 and spin 1 bosons was provided by one of us [6]. It has been shown in Ref [6] that a conserved four-current with a positive definite time component does exist for relativistic bosons, and is associated not with charge current but the flow of energy.

*Corresponding author: animesh.datta@iitk.ac.in (Animesh Datta)

2.1 The Kemmer Formalism for massive bosons

The Kemmer equations for a massive spin 0 or spin 1 boson are

$$(i\beta^\mu \partial_\mu + m)\psi = 0 \quad (1)$$

$$i\partial^\mu \bar{\psi} \beta_\mu - m\bar{\psi} = 0 \quad (2)$$

where the Kemmer-Duffin-Petiau matrices satisfy the algebra

$$\beta_\mu \beta_\nu \beta_\lambda + \beta_\lambda \beta_\nu \beta_\mu = \beta_\mu g_{\nu\lambda} + \beta_\lambda g_{\nu\mu} \quad (3)$$

and $\bar{\psi} = \psi^\dagger \eta_0$ with $\eta_0 = 2\beta_0^2 - 1$. As a consequence of Eqn (3) one has

$$\eta_0 \beta_0 = \beta_0 \eta_0 = \beta_0, \quad \eta_0^2 = 1 \quad (4)$$

$$\eta_0 \beta_i + \beta_i \eta_0 = 0, \quad i = 1, 2, 3. \quad (5)$$

We choose a representation in which $\beta_i^\dagger = -\beta_i (i = 1, 2, 3)$. The β_i in Ref [7] correspond to what we call $i\beta_i$ and $\beta_0^\dagger = \beta_0$. There are two nontrivial representations of these β 's: one is 5×5 , which describe spin 0 bosons, the other is 10×10 , which describe spin 1 bosons. For their explicit forms the reader is referred to [8].

Multiplying (1) by $\partial^\rho \beta_\rho \beta_\nu$, one can deduce,

$$\partial_\nu \psi = \partial^\rho \beta_\rho \beta_\nu \psi \quad (6)$$

and similarly

$$\partial_\nu \bar{\psi} = \partial^\rho \bar{\psi} \beta_\nu \beta_\rho \quad (7)$$

The second order wave equation

$$(\square + m^2)\psi = 0 \quad (8)$$

$$(\square + m^2)\bar{\psi} = 0 \quad (9)$$

follow from (1) and (6) and (2) and (7) respectively. A further consequence of (1) and (2) is the conservation equation

$$\partial_\mu j^\mu = 0 \quad (10)$$

$$j^\mu = \bar{\psi} \beta^\mu \psi \quad (11)$$

Unfortunately, although j_μ is a conserved 4-vector, j_0 is not positive definite and cannot be interpreted as probability density, a feature common with the Klein- Gordon and Proca equations.

2.2 Massless Bosons

Harish Chandra [9] showed how the Kemmer formalism could be applied for the case of massless bosons *without taking the limit* $m \rightarrow 0$. We restrict considerations to neutral particles only. Instead of (1), one writes

$$i\beta_\mu \partial^\mu \psi + m\gamma\psi = 0 \quad (12)$$

where γ is a matrix satisfying

$$\gamma^2 = \gamma \quad (13)$$

$$\gamma\beta_\mu + \beta_\mu\gamma = \beta_\mu \quad (14)$$

Multiplying (12) by $(1 - \gamma)$ on the left, one obtains

$$i\beta_\mu \partial^\mu (\gamma\psi) = 0 \quad (15)$$

Multiplying (12) by $\partial_\lambda \beta^\lambda \beta^\nu$ on the left, one also gets

$$\partial^\lambda \beta_\lambda \beta_\nu (\gamma\psi) = \partial_\nu (\gamma\psi) \quad (16)$$

The second order wave equation

$$\square(\gamma\psi) = 0 \quad (17)$$

follows from (15) and (16) which show that $\gamma\psi$ is a massless boson.

2.3 Probability Interpretation

Although j_0 is not positive definite, the fact that a conserved four- vector current with a positive definite time component can be defined using this formalism as follows. Multiplying (1) by β_0 , one obtains the Schrödinger form of the equation,

$$i\frac{\partial\psi}{\partial t} = [-i\tilde{\beta}_i\partial_i - m\beta_0]\psi \quad (18)$$

where $\tilde{\beta}_i = \beta_0\beta_i - \beta_i\beta_0$. Multiplying (1) by $(1 - \beta_0^2)$ one obtains

$$i\beta_i\beta_0^2\partial_i\psi = -m(1 - \beta_0^2)\psi. \quad (19)$$

If one multiplies Eqn (18) by ψ^\dagger from the left, its Hermitian conjugate by ψ from the right and adds the equations, one obtains the continuity equation

$$\frac{\partial\psi^\dagger\psi}{\partial t} + \partial_i\psi^\dagger\tilde{\beta}_i\psi = 0 \quad (20)$$

whence $S_i = \psi^\dagger\tilde{\beta}_i\psi$ is the conserved four-vector current density in the direction i with $S_0 = \psi^\dagger\psi > 0$. For details the reader is referred to Ref [10], Sec 1.

3 Massive Bosons

In this section we present our results. We always consider a potential step at $x = 0$ with the incident wave travelling along the positive x axis.

3.1 Spin 1 Bosons

The Kemmer-Duffin-HarishChandra wave function in this case is given by

$$\Psi = \frac{A}{\sqrt{m}} \begin{pmatrix} \psi_1 \\ \psi_2 \\ \vdots \\ \psi_9 \\ \psi_{10} \end{pmatrix} \quad (21)$$

We consider the situation of a wave propagating along the positive x-axis with a potential step at $x=0$,

$$\Psi_{inc} = \frac{A}{\sqrt{m}}(0, 0, E_0, 0, -E_0, 0, 0, 0, 0, 0)^T e^{i(kx - \omega t)} \quad (22)$$

$$\Psi_{ref} = \frac{B}{\sqrt{m}}(0, 0, E_0, 0, E_0, 0, 0, 0, 0, 0)^T e^{-i(kx + \omega t)} \quad (23)$$

$$\Psi_{trans} = \frac{C}{\sqrt{m}}(0, 0, E_0, 0, -\sqrt{\epsilon}E_0, 0, 0, 0, 0, 0)^T e^{i(k'x - \omega t)}. \quad (24)$$

where ϵ contains the information about the height of the barrier. The exact relation is not essential to the context of the problem.

Then $\frac{S_{inc}}{S_{ref}} = \frac{\Psi_{inc}^\dagger\tilde{\beta}_i\Psi_{inc}}{\Psi_{ref}^\dagger\tilde{\beta}_i\Psi_{ref}}$ for $i = 1, 2, 3$. In particular case that we treat we have $i = 1$. Then some algebra gives $\frac{S_{inc}}{S_{ref}} = \frac{|B|^2}{|A|^2}$.

Matching $\Psi_{inc} + \Psi_{ref} = \Psi_{trans}$ at $x = 0, t = 0$, readily gives

$$A + B = C \quad (25)$$

$$A - B = \sqrt{\epsilon}C \quad (26)$$

whence $\frac{|B|^2}{|A|^2} = \left(\frac{1 - \sqrt{\epsilon}}{1 + \sqrt{\epsilon}}\right)^2$. Thus the ratio $\frac{S_{ref}}{S_{inc}} < 1$ which means that no effect analogous to Klein paradox occurs in the case of Spin 1 massive bosons.

3.2 Spin 0 Bosons

For a wave propagating along $i = 1$, with $x_0 = t$, we have

$$\begin{aligned}\Psi_{inc} &= A(k, 0, 0, -ik_0, 1)e^{i(k_0x_0 - kx)} \\ \Psi_{ref} &= B(-k, 0, 0, -ik_0, 1)e^{-i(k_0x_0 + kx)} \\ \Psi_{trans} &= C(k', 0, 0, -ik'_0, 1)e^{i(k'_0x_0 - k'x)}\end{aligned}\tag{27}$$

which are special cases of

$$\Psi = \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \\ \phi_0 \\ \phi \end{pmatrix}\tag{28}$$

where $\phi_i = \partial_i \phi$ and $\phi = e^{ik_\mu x_\mu} = e^{i(k_0x_0 - k \cdot x)}$. The barrier is as usual at $x = 0$ and the information about $x > 0$ is incorporated in k' and k'_0 . Since $S_i = \Psi^\dagger \beta_i \Psi$ for all i and

$$\beta_1 = \begin{pmatrix} 0 & 0 & 0 & -i & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}\tag{29}$$

$$\frac{S_{ref}}{S_{inc}} = \frac{|B|^2}{|A|^2} = \left| \frac{B}{A} \right|^2.$$

The set of Eqns (27) evaluated at $x = 0, x_0 = 0$ give

$$\begin{aligned}A - B &= C(k'/k) \\ A + B &= C(k'_0/k_0) \\ A + B &= C\end{aligned}\tag{30}$$

Now $E^2 = \hbar^2 c^2 (k_0^2 - k^2) + m_0^2 c^4$ for $x < 0$ and $E^2 = \hbar^2 c^2 (k_0'^2 - k'^2) + m_0^2 c^4$ for $x > 0$. If the frequency of the wave does not change on moving into the potential *i.e.*, changing the media, which is indeed the case, then $k'_0 = k_0$ and consequently $k'^2 > k^2$.

Finally, Eqn (30) gives $\left| \frac{B}{A} \right| = \left(\frac{1 - k'/k}{1 + k'/k} \right)^2$.

Hence there is no evidence of Klein Paradox in the system under question as the reflection coefficient is less than 1.

4 Massless Bosons

In this case we will only be considering neutral bosons of spin 1, *i.e.*, photons. Using the Harish Chandra formalism [9], we have for a beam of photons propagating along the positive x axis,

$$\gamma \Psi = \begin{pmatrix} 0 \\ 0 \\ E_z \\ 0 \\ H_y \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}\tag{31}$$

The barrier is again placed at $x = 0$. Let

$$\begin{aligned} E_z^i &= E_0 e^{i(kx - \omega t - \phi)} \\ E_z^r &= E_0 \sqrt{R} e^{-i(kx - \omega t + \phi)} \\ E_z^t &= E_0 \sqrt{T} e^{i(k'x - \omega t + \chi)}. \end{aligned} \quad (32)$$

Maxwell's Equations then at once give (using $\partial_x E_z = \partial_t H_y$)

$$\begin{aligned} H_y^i &= -\frac{k}{\omega} E_0 e^{i(kx - \omega t - \phi)} \\ H_y^r &= -\frac{k}{\omega} E_0 \sqrt{R} e^{-i(kx + \omega t + \phi)} \\ H_y^t &= -\frac{k}{\omega} E_0 \sqrt{T} e^{i(k'x - \omega t - \chi)}. \end{aligned} \quad (33)$$

Now, $S_i = [E \times H]_i$ is the (Poynting vector) current density as given by [8]. Thus from Eqns (32) and (33)

$$\frac{S_x^{ref}}{S_x^{inc}} = R. \quad (34)$$

Matching $E_z^i - E_z^r = E_z^t$ at $x = 0, t = 0$ gives at once

$$R = \frac{S_x^{ref}}{S_x^{inc}} = \left(\frac{k'/k - 1}{k'/k + 1} \right)^2 < 1. \quad (35)$$

Thus there is no signature of Klein Paradox in the case of photons.

5 Conclusion

The Klein Paradox was a generic problem for the Dirac equation that was resolved very soon after its inception. The Klein-Gordon equation also shows the paradox [12] where it is an artefact of its not being a proper quantum mechanical equation with a positive definite probability density. However, to the best of our knowledge the question has never been asked, lest answered for bosons. This Letter asks that natural question and conclusively answers it by proving that the Klein paradox does not exist in the case of bosons. The paradox for fermions was resolved using the Hole theory by Dirac. Here too, the relativistic equations for spin 0 and spin 1 bosons are used. As has been shown in Section 2, the Klein-Gordon equation which is generally used to describe spin 0 particles can be deduced from the basic equation of Kemmer. For a similar deduction of Maxwell's equations the reader is referred to Ref [6], Sec 5. Our results, based on sound mathematical footing answer a basic question. However, there have been recent results based on Bohmian trajectories [11] showing the absence of Klein paradox in the case of outgoing scattering asymptotics.

Acknowledgements

P. G. acknowledges financial support from DST, Government of India. A. D. thanks the S. N. Bose National Centre for Basic Sciences for hospitality that enabled this work to be undertaken.

The authors also thank the reviewers whose comments helped improve the clarity of presentation.

References

- [1] P. A. M. Dirac, Proc. Roy. Soc. (London), A 117, 610 (1928), *ibid* A 118, 351 (1928). "Principles of Quantum Mechanics", *op. cit.*
- [2] E. Schrodinger, Sitzber. Preuss. Akad. Wiss. Physik-Math, 24, 418 (1930).

- [3] O. Klein, Z. Physik, 53, 157 (1929).
- [4] J. D. Bjorken, S. D. Drell, Relativistic Quantum Mechanics, McGraw Hill, New York, 1965.
- [5] W. Greiner, Relativistic Quantum Mechanics, Springer- Verlag, Berlin, 1997.
- [6] P. Ghose, Found. of Physics, 26 (11), 1441, 1996.
- [7] N. Kemmer, Proc. Roy. Soc A, 173, 91, 1939.
- [8] P. Ghose, M. K. Samal, Phys. Rev. E, 036620, 2001.
- [9] Harish Chandra, Proc. Roy. Soc(London), 186, 502, 1946.
- [10] P. Ghose, A. S. Majumdar, S. Guha, J. Sau, Phys. Lett. A, 290 (2001), 205.
- [11] G. Grubl *et al*, J. Phys. A: Math, Gen, 34, 2753- 2764 (2001).
- [12] G. Baym, Lectures of Quantum Mechanics, W. A. Benjamin. Inc., New York, 1969.