Dobinski-type relations and the Log-normal distribution

P Blasiak†‡, K A Penson† and A I Solomon†

[†]Université Pierre et Marie Curie, Laboratoire de Physique Théorique des Liquides, CNRS UMR 7600

Tour 16, $5^{i\hat{e}me}$ étage, 4, place Jussieu, F 75252 Paris Cedex 05, France

[‡]H. Niewodniczański Institute of Nuclear Physics, ul.Eliasza Radzikowskiego 152, PL 31-342 Kraków, Poland

E-mail:

blasiak@lptl.jussieu.fr, penson@lptl.jussieu.fr, a.i.solomon@open.ac.uk

Abstract.

We consider sequences of generalized Bell numbers $B(n)$, $n = 0, 1, \ldots$ which can be represented by Dobinski-type summation formulas, i.e. $B(n) = \frac{1}{C} \sum_{k=0}^{\infty} \frac{[P(k)]^n}{D(k)}$ $\frac{P(k)}{D(k)},$ with $P(k)$ a polynomial, $D(k)$ a function of k and $C = const.$ They include the standard Bell numbers $(P(k) = k, D(k) = k!, C = e)$, their generalizations $B_{r,r}(n)$, $r = 2, 3, \ldots$ appearing in the normal ordering of powers of boson monomials $(P(k) = \frac{(k+r)!}{k!}, D(k) = k!, C = e)$, variants of "ordered" Bell numbers $B_o^{(p)}(n)$ $(P(k) = k, D(k) = (\frac{p+1}{p})^k, C = 1+p, p=1,2...$, etc. We demonstrate that for α, β, γ, t positive integers $(\alpha, t \neq 0)$, $[B(\alpha n^2 + \beta n + \gamma)]^t$ is the *n*-th moment of a positive function on $(0, \infty)$ which is a weighted infinite sum of log-normal distributions.

$Dobi\acute{n}ski-type~relations~and~the~Log-normal~distribution$ 2

In a recent investigation [1] we analysed sequences of integers which appear in the process of normal ordering of powers of monomials of boson creation a^{\dagger} and annihilation *a* operators, satisfying the commutation rule $[a, a^{\dagger}] = 1$. For *r*, *s* integers such that $r \geq s$, we define the generalized Stirling numbers of the second kind $S_{r,s}(n, k)$ as

$$
[(a^{\dagger})^r a^s]^n = (a^{\dagger})^{n(r-s)} \sum_{k=s}^{ns} S_{r,s}(n,k) (a^{\dagger})^k a^k
$$
 (1)

and the corresponding Bell numbers $B_{r,s}(n)$ as

$$
B_{r,s}(n) = \sum_{k=s}^{ns} S_{r,s}(n,k).
$$
 (2)

In [1] explicit and exact expressions for $S_{r,s}(n, k)$ and $B_{r,s}(n)$ were found. In a parallel study [2] it was demonstrated that $B_{r,s}(n)$ can be considered as the *n*-th moment of a probability distribution on the positive half-axis. In addition, for every pair (r, s) the corresponding distribution can be explicitly written down. These distributions constitute the solutions of a family of Stieltjes moment problems, with $B_{r,s}(n)$ as moments. Of particular interest to us are the sequences with $r = s$, for which the following representation as an infinite series has been obtained:

$$
B_{r,r}(n) = \frac{1}{e} \sum_{k=0}^{\infty} \frac{1}{k!} \left[\frac{(k+r)!}{k!} \right]^{n-1}
$$
 (3)

$$
= \frac{1}{e} \sum_{k=0}^{\infty} \frac{\left[k(k+1)\dots(k+r-1)\right]^n}{(k+r-1)!}, \qquad n > 0. \tag{4}
$$

Eqs.(3) and (4) are generalizations of the celebrated Dobinski formula $(r = 1)$ [3]:

$$
B_{1,1}(n) = \frac{1}{e} \sum_{k=0}^{\infty} \frac{k^n}{k!}, \quad n \ge 0,
$$
\n(5)

which expresses the conventional Bell numbers $B_{1,1}(n)$ as a rapidly convergent series. Its simplicity has inspired combinatorialists such as G.-C. Rota [4] and H.S. Wilf [5]. Eq.(5) has far-reaching implications in the theory of stochastic processes $[6]$, $[7]$, $[8]$.

The probability distribution whose *n*-th moment is $B_{r,r}(n)$ is an infinite ensemble of weighted Dirac delta functions located at a specific set of integers (a so-called Dirac comb):

$$
B_{r,r}(n) = \int_0^\infty x^n \left\{ \frac{1}{e} \sum_{k=0}^\infty \frac{\delta(x - k(k+1) \dots (k+r-1))}{(k+r-1)!} \right\} dx, \quad n \ge 0. \tag{6}
$$

For $r = 1$ the discrete distribution of Eq.(6) is the weight function for the orthogonality relation for Charlier polynomials [9]. In contrast we emphasize that for $r \neq s$ the $B_{r,s}(n)$ are moments of continuous distributions [2].

In this note we wish to point out an intimate relation between the formulas of Eqs. (3) , (4) , (5) and the log-normal distribution [10], [11]:

$$
P_{\sigma,\mu}(x) = \frac{1}{\sqrt{2\pi}\sigma x} e^{-\frac{(\ln(x) - \mu)^2}{2\sigma^2}}, \qquad x \ge 0, \quad \sigma, \mu > 0.
$$
 (7)

Figure 1. Weight function $W_{1,1}(1,1,1;x)$, see Eq.(13).

First we quote the standard expression for its n -th moment:

$$
M_n = \int_0^\infty x^n P_{\sigma,\mu}(x) dx = e^{n\left(\mu + n\frac{\sigma^2}{2}\right)}, \quad n \ge 0,
$$
\n(8)

which can be reparametrized for $k > 1$ as

$$
M_n = k^{\alpha n^2 + \beta n},\tag{9}
$$

with

$$
\mu = \beta \ln(k),\tag{10}
$$

$$
\sigma = \sqrt{2\alpha \ln(k)} > 0. \tag{11}
$$

Given three integers α, β, γ (where $\alpha > 0$), we wish to find a weight function $W_{1,1}(\alpha, \beta, \gamma; x) > 0$ such that

$$
B_{1,1}(\alpha n^2 + \beta n + \gamma) = \int_0^\infty x^n W_{1,1}(\alpha, \beta, \gamma; x) dx.
$$
 (12)

Eqs. (5) , (7) and (9) provide an immediate solution:

$$
W_{1,1}(\alpha,\beta,\gamma;x) = \frac{1}{e} \left[\delta(x-1) + \sum_{k=2}^{\infty} \frac{k^{\gamma} \exp\left(-\frac{(\ln(x)-\beta\ln(k))^2}{4\alpha\ln(k)}\right)}{2xk!\sqrt{\pi\alpha\ln(k)}}\right],
$$
(13)

which is an *infinite* sum of weighted log-normal distributions supplemented by a single Dirac peak of weight e^{-1} located at $x = 1$. Thus it is a *superposition* of discrete and continuous distributions. Virtually the same approach can be adopted for the sequences $B_{r,r}(n), r > 1$. In this case the $k = 1$ term in the numerator of Eq.(3) is larger than one

Figure 2. Weight functions $W_{r,r}(1,1,1;x)$ for $r = 2,3,4$.

and so there will be no Dirac peak in the formula. Then the function $W_{r,r}(\alpha,\beta,\gamma;x) > 0$ defined by $(\alpha, \beta, \gamma \text{ integers}, \alpha, \gamma > 0)$:

$$
B_{r,r}(\alpha n^2 + \beta n + \gamma) = \int_0^\infty x^n W_{r,r}(\alpha, \beta, \gamma; x) dx,
$$
\n(14)

is a purely continuous probability distribution given again by an infinite sum of weighted log-normal distributions:

$$
W_{r,r}(\alpha,\beta,\gamma;x) = \frac{1}{e} \sum_{k=0}^{\infty} \frac{\left[\frac{(k+r)!}{k!}\right]^{\gamma-1} \exp\left(-\frac{\left[\ln(x)-\beta\ln\left(\frac{(k+r)!}{k!}\right)\right]^2}{4\alpha\ln\left[\frac{(k+r)!}{k!}\right]}\right)}{2xk!\sqrt{\pi\alpha\ln\left[\frac{(k+r)!}{k!}\right]}}.
$$
(15)

The solutions of the moment problems of Eqs.(9), (12) and (14) are not unique. More general solutions may be obtained by the method of the inverse Mellin transform, see [12].

Several other types of combinatorial sequences have properties exemplified by Eqs.(12) and (14). We quote for example the so-called "ordered" Bell numbers $B_o(n)$ defined as [5]:

$$
B_o(n) = \sum_{k=1}^{n} S(n,k)k!,
$$
\n(16)

where $S(n, k)$ are the Stirling numbers of the second kind, $S_{1,1}(n, k)$ in our notation.

These ordered Bell numbers satisfy the following Dobinski-type relation:

$$
B_o(n) = \frac{1}{2} \sum_{k=0}^{\infty} \frac{k^n}{2^k},\tag{17}
$$

from which a formula analogous to Eq.(13) readily follows. A more general identity of type (17) is $|13|$:

$$
B_o^{(p)}(n) = \frac{1}{p+1} \sum_{k=1}^{\infty} k^n \left(\frac{p}{p+1}\right)^k = \sum_{k=0}^n S(n,k)k! \ p^k, \qquad p = 2, 3, \dots (18)
$$

We will not discuss other types of sequences but rather observe that the relations of Eqs. (3) , (4) , (5) , (17) , (18) naturally imply that any power of these numbers also satisfies a Dobinski-type relation. As an example we give explicitly the simplest case of Eq.(5). For integer $t > 0$:

$$
[B_{1,1}(n)]^t = \frac{1}{e^t} \sum_{k_1, k_2, \dots k_t = 0}^{\infty} \frac{(k_1 k_2 \dots k_t)^n}{k_1! k_2! \dots k_t!},
$$
\n(19)

with correspondingly more complicated formulas of a similar nature for powers of $B_{r,r}(n)$, $B_o(n)$ and $B_o^{(p)}(n)$. For combinatorial applications of Eq.(19) see [14], [15] and [16].

We conclude that for any sequences of the type $B(n)$ specified above $\left[B(\alpha n^2 + \beta n + \gamma)\right]^t$ is always given as an *n*-th moment of a positive function on $(0, \infty)$ expressible by sums of weighted log-normal distributions. We illustrate such a function for $B_{1,1}(n)$ in Fig.(1). The application to $B_{r,r}(n)$ for $r = 2, 3, 4$ is presented in Fig.(2). The area under every curve is equal to 1 on extrapolating to large x (not displayed). In both examples we have chosen $\alpha = \beta = \gamma = 1$. Observe the exceedingly slow decrease of these probabilities for $x \to \infty$. This is confirmed by the fact that the moment sequences $[B(\alpha n^2 + \beta n + \gamma)]^t$ are extremely rapidly increasing. In the simplest case $\alpha = \beta = \gamma = t = 1$ we find $B_{1,1}(n^2 + n + 1) = 1, 5, 877, 27644437, 474869816156751$ for $n = 0 \dots 4$.

The circumstance that we can determine the positive solutions of the Stieltjes moment problem with both $B(n)$ (discrete distribution) and $[B(\alpha n^2 + \beta n + \gamma)]^t$ (continuous distribution) is a very specific consequence of the existence of Dobinskitype expansions. To our knowledge it has no equivalent in standard solutions of the moment problem. For instance, if the moments are $n!$ the solution e^{-x} does not give any indication as how one might obtain the solution for the moments equal to $(n^2)!$.

The strict positivity of $W_{r,r}(\alpha,\beta,\gamma;x)$, for $r=1,2...$, suggests their use in the construction of coherent states, which satisfy the resolution of identity property [17], [18], [19], [20]. This can be done by the substitution $n! \rightarrow B_{r,r}(\alpha n^2 + \beta n + \gamma)$ in the definition of standard coherent states. More precisely for a complete and orthonormal set of wave functions $|n\rangle$ such that $\langle n|n'\rangle = \delta_{n,n'}$ and complex z we define the normalized coherent state as

$$
|z; \alpha, \beta, \gamma\rangle = \frac{1}{\mathcal{N}^{1/2}(\alpha, \beta, \gamma; |z|^2)} \sum_{n=0}^{\infty} \frac{z^n}{\sqrt{B_{r,r}(\alpha n^2 + \beta n + \gamma)}} |n\rangle, \tag{20}
$$

with the normalization

$$
\mathcal{N}(\alpha, \beta, \gamma; x) = \sum_{n=0}^{\infty} \frac{x^n}{B_{r,r}(\alpha n^2 + \beta n + \gamma)},
$$
\n(21)

which is a rapidly converging function of x for $0 \leq x < \infty$, $x = |z|^2$. Then, using the procedure of [18] we can demonstrate that the states of $Eq.(20)$ along with $Eq.(14)$ automatically satisfy the resolution of unity

$$
\int \int_{\mathbb{C}} d^2 z |z; \alpha, \beta, \gamma \rangle \tilde{W}_{r,r}(\alpha, \beta, \gamma; |z|^2) \langle z; \alpha, \beta, \gamma \rangle = I = \sum_{n=0}^{\infty} |n\rangle \langle n|, \qquad (22)
$$

with

$$
W_{r,r}(\alpha,\beta,\gamma;|z|^2) = \pi \frac{\tilde{W}_{r,r}(\alpha,\beta,\gamma;|z|^2)}{\mathcal{N}(\alpha,\beta,\gamma;|z|^2)}.
$$
\n(23)

We are currently investigating the quantum-optical properties of states defined in $Eq.(20)$.

We close by quoting from [7] that, "the idea of representing the combinatorially defined numbers by an infinite sum or an integral, typically with a probabilistic interpretation, has proved to be a very fruitful one". In our particular case it has allowed us to reveal quite an unexpected relation between the Dobinski-type summation relations, which by themselves are reflections of boson statistics, and the log-normal distribution.

Acknowledgments

We thank D. Barsky, G. Duchamp, L. Haddad, A. Horzela and M.Yor for numerous fruitful discussions. N.J.A. Sloane's Encyclopedia of Integer Sequences (http://www.research.att.com/˜njas/sequences) was an essential help in analyzing the properties of sequences we have encountered in this work.

References

- [1] Blasiak P, Penson K A and Solomon A I 2003 The general boson normal ordering problem Phys. Lett. A 309 198
- [2] Penson K A and Solomon A I 2002 Coherent state measures and the extended Dobinski relations Preprint quant-ph/0211061
- [3] Comtet L 1974 Advanced Combinatorics (Dordrecht: Reidel)
- [4] Rota G-C 1964 Amer. Math. Monthly **71** 498
- [5] Wilf H S 1994 Generatingfunctionology (New York: Academic Press)
- [6] Constantine G M and Savits T H 1994 A stochastic process interpretation of partition identities SIAM J. Discrete Math. 7 194
- [7] Pitman J 1997 Some probabilistic aspects of set partitions Amer. Math. Monthly 104 201
- [8] Constantine G M 1999 Identities over set partitions Discrete Math. 204 155
- [9] Koekoek R and Swarttouv R F 1998 The Askey scheme of hypergeometric polynomials and its q-analogue, Dept. of Technical Mathematics and Informatics Report No 98-17 (Delft University of Technology)
- [10] Crow E L and Shimizu K (Eds.) 1988 Log-Normal Distributions, Theory and Applications (New York: Dekker)
- [11] Bertoin J, Biane P and Yor M 2003 Poissonian exponential functionals, q-series, q- integrals and the moment problem for log-normal distributions Proceedings of Rencontre d'Ascona, Mai 2002 Russo F and Dozzi M (Eds.) (Basel: Birkhäuser)
- [12] Sixdeniers J-M 2001 Constructions de nouveaux états cohérents à l'aide de solutions des problèmes des moments Ph.D. Thesis (Paris: Univ. Pierre et Marie Curie); Penson K A and Sixdeniers J-M (unpublished)
- [13] Weisstein E W World of Mathematics, entry: Stirling Numbers of the Second Kind (http://mathworld.wolfram.com/)
- [14] Pittel B 2000 Where the typical set partitions meet and join Electron. J. of Combin. 7 R5
- [15] Canfield E R 2001 Meet and join within the lattice of set partitions Electron. J. of Combin. 8 R15
- [16] Bender C M, Brody D C and Meister B K 1999 Quantum field theory of partitions J. Math. Phys. 40 3239
- [17] Sixdeniers J-M, Penson K A and Solomon A I 1999 Mittag-Leffler coherent states J. Phys. A 32 7543
- [18] Klauder J R, Penson K A and Sixdeniers J-M 2001 Constructing coherent states through solutions of Stieltjes and Hausdorff moment problems Phys. Rev. A 64 013817
- [19] Quesne C 2001 Generalized coherent states associated with the C_{λ} -extended oscilator Ann. Phys., NY 293 147
- [20] Quesne C 2002 New q-deformed coherent states with an explicitly known resolution of unity J. Phys A 35 9213