## Quantum Information as Reduced Classical Information

B.A.Nikolov

Department of Theoretical and Applied Physics, Shoumen University, 9712 Shoumen, Bulgaria E-mail: b.nikolov@shu-bg.net

## Abstract

It is shown on a simple classical model of a quantum particle at rest that information contained into the quantum state (quantum information) can be obtained by integrating the corresponding probability distribution on phase space, i.e. by reduction of the information contained into the classical state.

1.A physical theory must give specification of three items: a) state space, b) observables, and c)the value of an observable in a given state.For example in the classical model state space is the phase space of the system, observables are smooth real functions on phase space, and the value of the observable A in a state (q, p) is A(q, p).In general however the phase space point is not known and the information about the real state (classical information) is given by a phase space distribution  $\rho(q, p)$ .In this case one can predict only the mean value (expectation) of the observable

$$\langle A \rangle = \int A(q,p)\rho(q,p)dqdp$$
 (1)

Thus it is reasonable under a *(classical)* state to understand not a point in phase space but rather a probability distribution on it. Then instead about the value of an observable in some state one should speak about the *mean* value of the observable in that state.

In the quantum model the state space is a Hilbert space, the information about the state (quantum state) is represented by a density operator, i.e. a positive operator  $\rho$  satisfying  $tr\rho = 1$ , observables are Hermitian operators, and the mean value of the observable A in state  $\rho$  is given by the Born rule

$$\langle A \rangle = tr A \rho \tag{2}$$

Our aim in the present note is ,using the simple model of a quantum particle given in a previous work [1], to show that ,in fact, the quantum state can be obtained by integrating of the classical one.

2.As a model of a quantum particle at rest we consider a d = 2s + 1 dimensional oscillator ,where s is the spin of the particle. The mass and the frequency of the oscillator are taken equal to unit. The coordinates  $q_n$ , (n = 1, 2, ..., d) of the oscillator describe the microscopic (unobservable) deviation from the rest position. If  $p_n$  are the corresponding momenta, and energy is assumed to be  $E = \hbar \omega/2 = \hbar/2$  then the phase space trajectory is given by the equation

$$\sum_{n=1}^{d} (p_n^2 + q_n^2) = \hbar$$
 (3)

Introducing new coordinates and momenta , viz.  $x_n = \hbar^{-1/2}, y_n = \hbar^{-1/2}, \psi_n = x_n + iy_n$  we can rewrite Eq.(1) in the form

$$\sum_{n} \psi_n^* \psi_n = 1 \tag{4}$$

Now using the same techniques as in Ref.1 we can show that arbitrary observable generating canonical transformation that preserve Eq.(4) is a hermitian form:

$$A(\psi^*, \psi) = \hbar \sum_n A_{nm} \psi_n^* \psi_n \tag{5}$$

Here  $A_{nm}$  is a Hermitian matrix completely determining the observable.

3. Let  $\rho(\psi^*, \psi)$  is the probability distribution of phase space coordinates, and

$$d\Omega_{\psi} = (d-1)!\delta(1-\psi^*\psi)d\psi^*d\psi \tag{6}$$

is the normalized measure over the unit sphere (4) [2]. Then, applying Eq.(1) taken in the form

$$\langle A \rangle = \int A(\psi^*, \psi) \rho(\psi^*, \psi) d\Omega_{\psi}$$
(7)

we obtain

$$\langle A \rangle = \sum_{n,m} A_{nm} \rho_{mn} \tag{8}$$

where

$$\rho_{mn} = \hbar \int \psi_n^* \psi_m \rho(\psi^*, \psi) d\Omega_\psi \tag{9}$$

In fact we obtained the Born rule in its most general form.

## References

- [1] B.A.Nikolov, "A Thermodynamic Approach to Quantum Measurement and Quantum Probability", quant-ph 0303121.
- [2] K.R.W.Jones, Ann. Physics, v. 207, p. 140.