# ALGORITHM OF REDUCTION

N. K. Solovarov

Zavoisky Physical-Technical Institute of the Russian Academy of Sciences, Sibirskii trakt 10/7, 420029 Kazan, Russia Kazan State Pedagogical University, ul. Mezhlauk 1, 420021 Kazan, Russia\*

, **,** ,

(Dated: July 29, 2018)

# Abstract

We show that the von Neumann's algorithm of reduction (i.e. the algorithm of calculating the density matrix of the observable subsystem from the density matrix of the closed quantum system) corresponds to the special approximation at which the unobservable subsystem is supposed to be in the steady state of minimum information (infinite temperature). We formulate the generalized algorithm of reduction that includes as limiting cases the von Neumann's reduction and the self-congruent correlated reduction most corresponding to the quantum nondemolition measurement. We demonstrate the correlation in dynamics of subsystems with exactly soluble models of quantum optics: 1) about the dynamics of a pair of interacting two-level atoms, and 2) about the dynamics of a two-level atom interacting with a single-mode resonant field.

PACS numbers: 03.65.Ta, 03.65.Ud, 42.50.Ct, 42.50.Pq

Keywords: reduction, reduced density matrix, quantum nondemolition measurements, EPR-pair, Janes-Cummings model

<sup>\*</sup>Electronic address: solovar@kfti.knc.ru

#### I. INTRODUCTION

The old question of the quantum theory on the reduction of a quantum system and on the decoherence of an observable subsystem (transition from a pure state to a mixed state) due to its interaction with an unobservable subsystem and the measurement is formulated anew in quantum optics [1, 2]. This process is initiated by problems of quantum computing [3], by hope to clarify the quantum nature of the irreversibility [2, 4] and, mainly, by the new experimental potentials to check consequences of axioms of quantum mechanics accepted previously. The qualitatively new experiments allow one to observe dynamics of subsystems of simple quantum systems (e.g., the dynamics of a pair of correlated photons emitted in a nonlinear process, or the dynamics of a single two-level atom in the course of its interaction with the resonant field of extremely small intensity [1, 2, 3, 5]). The mathematical operation of reduction defines the rule of calculating the density operator of an observable subsystem from the known density operator of a closed quantum system. By the von Neumann's postulate, this rule consists in calculating the partial trace of the density matrix of a closed quantum system over the unobservable subsystem [6, 7, 8, 9]. Though this nonunitary operation has been universally accepted to account for the influence of the classical measuring device on the result of measurement, to realize the transition from the closed system to the open one, and to introduce the irreversibility into the dynamics of an observable subsystem [2, 4, 6, 7, 9], the algorithm of accounting for the interaction of the closed quantum system with a measuring device remains in so doing reduction implicit. Moreover, when calculating the mean values of observables of subsystems, the realization of reduction is implicit ("disguised") by itself.

The initial point of the our treatment lies in the identity of the mathematical formulation of the reduction (other than the interpretation!) with the formulation of the quantum relaxation theory problem [8, 9]. In both cases the closed composite quantum system is considered. Consequently the unitary time evolution of the system is described by the density operator  $\rho(t)$ , which may be approximately calculated (basically in general case). In both cases the dynamics, i.e. the density operator, of one subsystem  $\hat{\rho}_{\alpha}(t)$  of the system should be calculated. This extraction of the one subsystem implies implicitly the separation of the system into two parts: the observable subsystem  $\alpha$  and the unobservable subsystem  $\beta$ . In general case the subsystems of the arbitrary closed system are equal in rights. Let us suppose the dynamics of the subsystem  $\beta$  has described by some density operator  $\hat{\rho}_{\beta}(t)$ . The exact representation of the density operator of the closed system through the density operators of subsystems is known for the noninteracting subsystems only:  $\hat{\rho}(t) = \hat{\rho}_{\alpha}(t) \otimes \hat{\rho}_{\beta}(t)$ [8, 9]. The considered problem consist in calculating of the density operator of one subsystem in the general case of interacting or interacted previously (entangled) subsystems.

In the quantum relaxation theory the "unobservable" subsystem is believed to be the thermostat at a temperature of T from physical reasons. Consequently its density operator supposed to be known and stationary:  $\hat{\rho}_{\beta}(T)$ . The desired density operator  $\hat{\rho}_{\alpha}(t)$  next is calculated to some approximation from the approximate equation  $\hat{\rho}(t) \approx \hat{\rho}_{\alpha}(t) \otimes \hat{\rho}_{\beta}(T)$ , where two operators  $\hat{\rho}(t)$  and  $\hat{\rho}_{\beta}(T)$  are known. In the reduction problem the state of the unobservable subsystem is believed unknown. The algorithm was instead postulated to calculate  $\hat{\rho}_{\alpha}(t)$  from  $\hat{\rho}(t)$ . The question is arises then: is this postulation equivalent to some approach on the state of unobservable subsystem? As a matter of fact this question is identical with the mathematical problem of the quantum relaxation theory: how the density operator of subsystem  $\hat{\rho}_{\beta}(t)$  should be approximately calculated from the known density operators  $\hat{\rho}(t), \hat{\rho}_{\alpha}(t)$ ?

We apply the method similar to used by von Neumann for demonstrating the correlation (the one-to-one correspondence) between the physically observables of the different interacting subsystems of the closed quantum system [6]. In this way the approximate equation system demonstrating the mutual relations between the density operators of subsystems and the density operator of the closed system was deduced. This equation system reveals that given in some way the density operator of one subsystem, the density operator of the other subsystem can be calculated approximately. We prove that the von Neumann's algorithm of reduction is identical to the approximation when the subsystem  $\beta$  is constantly in the state of minimum information  $\hat{\rho}_{\beta min}$  (identical with the state of infinite temperature [6, 8]). It should emphasize this approximation is solely employing when the reduction carring out, other than when the  $\hat{\rho}(t)$  calculating!

By this means the von Neumann's reduction is mathematically equivalent to the replacement  $\hat{\rho}(t) \rightarrow \hat{\rho}_{\alpha}(t) \otimes \hat{\rho}_{\beta min}$  with exactly defined operators  $\hat{\rho}_{\alpha}(t), \hat{\rho}_{\beta min}$ . This replacement corresponds on other hand to the changing of the considered quantum system and to the traditional interpretation of reduction as the transition from the closed to an open system or the accounting for the measurement. The state of minimum information (of infinite temperature) corresponds to the complete absence of the quantum coherence in the subsystem  $\beta$ . This implicit approach leads to some decoherence of the observable subsystem although all the calculations are done in framework of the closed quantum system.

We get the conclusion that any one of the possible algorithms of reduction is mathematically equivalent to the replacement  $\hat{\rho}(t) \rightarrow \hat{\rho}_{\alpha}(t) \otimes \hat{\rho}_{\beta}(t)$ , i.e. to the approximate representation of the closed quantum system as two quasy-independent subsystems with the definite dynamics. This raises the question of whether the accepted algorithm of reduction (i.e. the corresponding replacement of the density operator) corresponds to the initial assumption of the closed quantum system. Or mathematically: how much would this replacement be in error with the believed to be true  $\hat{\rho}(t)$ ?

As it is evident from ours consideration, the generally accepted von Neumann's algorithm of reduction is self-contradictory: on the one hand the closed quantum system is assumed, and on the other the infinite temperature of the subsystem is taken implisitly. It is possible to perfect the algorithm of reduction: for example, to accept some more reasonable state of the unobservable subsystem (the thermostat, as in the quantum relaxation theory, or one of eigenstates, as in the theory of quantum nondemolition measurements of photons [12, 13], or any intermediate variant). But in any case this step corresponds to the changing of the considered system and to the coming to the new system with the new dynamics, what is customary interpreted as the coming to the open system.

If the both subsystems of the considered system are the essentially quantum systems and the approximation of the closed system is believed true, it seems reasonable to use the principle of the interchangeability of subsystems [6] to make the reduction. We use the mutual relations between the density operators  $\hat{\rho}(t)$ ,  $\hat{\rho}_{\alpha}(t)$ ,  $\hat{\rho}_{\beta}(t)$  to calculate the correlated density operators  $\hat{\rho}_{\alpha C}(t)$ ,  $\hat{\rho}_{\beta C}(t)$  simultaneously by the self-congruent procedure of the successive approximations. In this way the reduction making the minimum (the vanishing in the limit) changing of  $\hat{\rho}(t)$  is obtainable. In other words, the correlated reduction procedure is most consistent with the initial approximation of the closed quantum system or to the quantum nondemolition measurement.

The structure of the paper is as follows: in Sec. II we demonstrate that the von Neumann's reduction corresponds to the approximation of the steady state of minimum information of the unobservable subsystem and formulate the generalized reduction algorithm as an approximation of a given state of the unobservable subsystem of a closed quantum system. Sec. III contains the algorithm of correlated reduction that defines the procedure of calculating self-congruent density operators of subsystems of a closed quantum system in the limit of the quantum nondemolition measurement. In Sec. IV we use the correlated reduction technique to consider three known problems of quantum optics: 1) the separable representation of the entangled EPR(Einshtein-Podolsky-Rosen) state, 2) the dynamics of a pair of interacting two-level atoms, 3) the dynamics of a two-level atom interacting with the resonant single-mode electromagnetic field (the model of cavity quantum electrodynamics).

# **Designations and definitions** [6, 7, 8, 9]:

1.  $\hat{H} = \hat{H}_{\alpha} + \hat{H}_{\beta} + \hat{H}_{\alpha\beta}$  is a hamiltonian of a closed quantum system,  $\hat{H}_{\alpha}$  and  $\hat{H}_{\beta}$  are hamiltonians of uncoupled subsystems  $\alpha$  and  $\beta$ , respectively,  $\hat{H}_{\alpha\beta}$  is the interaction hamiltonian of subsystems;  $\hat{A}$  and  $\hat{B}$  are operators of observables of subsystems  $\alpha$  and  $\beta$ , respectively.

2.  $\hat{\rho}(t) = \hat{U}(t)\hat{\rho}(0)\hat{U}^{\dagger}(t)$  is the solution of the Schrödinger equation determining the dynamics of the system  $\hat{H}, \hat{U}(t) = \exp[-(it/\hbar)\hat{H}]$  is the operator of evolution.

3. Operators  $\hat{H}_{\alpha}$ ,  $\hat{A}$  and  $\hat{H}_{\beta}$ ,  $\hat{B}$  are determined in spaces of subsystems  $\alpha$  and  $\beta$ , respectively; operators  $\hat{H}$ ,  $\hat{H}_{\alpha\beta}$ ,  $\hat{\rho}(t)$ ,  $\hat{U}(t)$  are determined in the complete space that is the direct (exterior  $\leftrightarrow \otimes$ ) product of spaces  $\alpha, \beta$ ;  $\hat{A}' = \hat{A} \otimes \hat{1}_{\beta}$ ,  $\hat{B}' = \hat{1}_{\alpha} \otimes \hat{B}$  are extended operators of subsystems in the complete space;  $\hat{1}_{\alpha(\beta)}$  are unity operators in spaces of subsystems,  $\hat{1}_{\alpha} \otimes \hat{1}_{\beta} = \hat{1}$  is the unity operator of the complete space;  $a_{ij}, b_{i'j'}$  are matrix elements of the operators  $\hat{A}, \hat{B}; o_{\{ii'\}\{jj'\}}$  are matrix elements of operators ( $\hat{O}$ ) of the closed system, a pair of subscripts  $\{ii'\}$  designates basis state of the closed system.

4. The von Neumann's algorithm of calculating the reduced density operator of the observable subsystem is determined by:

$$\hat{\rho}_{\alpha N}(t) = S p_{\beta} \hat{\rho}(t), \quad \text{or} \quad \hat{\rho}_{\beta N}(t) = S p_{\alpha} \hat{\rho}(t), \quad (1.1)$$

where  $Sp_{\alpha(\beta)}$  means the partial trace over the subsystem  $\alpha(\beta)$ , the subscript N hereinafter designates quantities calculated in accordance with the von Neumann's algorithm of reduction.

## II. THE THEOREM ON THE CORRELATION OF SUBSYSTEMS

## Theorem

The reduction (1.1) of the closed quantum system  $\hat{H}$  for the observable subsystem  $\alpha$  corresponds to the approximation at which: i) the closed system is represented as two uncoupled correlated subsystems, and ii) the unobservable subsystem  $\beta$  is supposed to be in the steady state of minimum information.

#### Proof.

i). Let us suppose that the dynamics of the closed system  $\hat{H}$  is approximately represented as the dynamics of two uncoupled subsystems. It means [8, 9] that there are such density operators  $\hat{\rho}_{\alpha C}(t)$  and  $\hat{\rho}_{\beta C}(t)$   $(Sp_{\alpha}\hat{\rho}_{\alpha C}(t) = Sp_{\beta}\hat{\rho}_{\beta C}(t) = 1)$  of subsystems for which

$$\hat{\rho}_{\alpha C}(t) \otimes \hat{\rho}_{\beta C}(t) \approx \hat{\rho}(t).$$
(2.1)

We transform (2.1) in much the same way as von Neumann did to prove the mutual correlation of mean values  $\langle \hat{A}(t) \rangle$ ,  $\langle \hat{B}(t) \rangle$  (see [6, Ch.6.2]). If we multiply (2.1) on both sides by  $\hat{\rho}'_{\beta C}(t) = \hat{1}_{\alpha} \otimes \hat{\rho}_{\beta C}(t)$  or  $\hat{\rho}'_{\alpha C}(t) = \hat{\rho}_{\alpha C} \otimes \hat{1}_{\beta}$  and take partial traces over the subsystem  $\beta$  or  $\alpha$  we obtain the system of two connected equations [10, 11]:

$$\hat{\rho}_{\alpha C}(t) \approx \frac{S p_{\beta} \hat{\rho}(t) \hat{\rho}'_{\beta C}(t)}{S p_{\alpha \beta} \hat{\rho}(t) \hat{\rho}'_{\beta C}(t)},\tag{2.2}$$

$$\hat{\rho}_{\beta C}(t) \approx \frac{S p_{\alpha} \hat{\rho}(t) \hat{\rho}_{\alpha C}'(t)}{S p_{\alpha \beta} \hat{\rho}(t) \hat{\rho}_{\alpha C}'(t)}.$$
(2.3)

The right-hand sides of (2.2), (2.3) represent the normalized quantum averaging of the density operator of the closed system over one of the subsystems. Each equation determines the density operator of one subsystem through the density operator of the closed system and density operator of another subsystem. From (2.1) and (2.2), (2.3), it is evident that density operators  $\hat{\rho}_{\alpha C}(t), \hat{\rho}_{\beta C}(t)$  are correlated, that is why we may call **i**) the approximation of correlated subsystems (is marked by a subscript C).

ii). Let us suppose that the unobservable subsystem  $\beta$  is in the steady state of minimum information, i.e. its density operator equals [6, 8]:

$$\hat{\rho}_{\beta C}(t) = \hat{\rho}_{\beta \min} = (1/N_{\beta}) \hat{1}_{\beta}, \qquad (2.4)$$

where  $N_{\beta}$  is the dimension (number of eigenstates) of the subsystem  $\beta$ . The theorem then follows if we insert (2.4) into the right-hand side of (2.2):

$$\hat{\rho}_{\alpha}(t) \approx \frac{Sp_{\beta}\hat{\rho}(t)(1/N_{\beta})\left(\hat{1}_{\alpha}\otimes\hat{1}_{\beta}\right)}{Sp_{\alpha\beta}\hat{\rho}(t)(1/N_{\beta})\left(\hat{1}_{\alpha}\otimes\hat{1}_{\beta}\right)} = Sp_{\beta}\hat{\rho}(t) = \hat{\rho}_{\alpha N}(t).$$
(2.5)

#### Consequence 1

By (2.1)-(2.4), the von Neumann's algorithm of reduction (1.1) is equivalent to the replacement of the density operator  $\hat{\rho}(t)$  with the reduced density operator  $\hat{\rho}_{RN}(t)$  according to the rule:

$$\hat{\rho}(t) \to \hat{\rho}_{RN}(t) = \hat{\rho}_{\alpha N}(t) \otimes \hat{\rho}_{\beta min} = [Sp_{\beta}\hat{\rho}(t)] \otimes (1/N_{\beta}) \hat{1}_{\beta}.$$
(2.6)

The relation (2.6) agrees in form with " the basic equation of irreversibility " or "the first assumption" of the quantum relaxation theory ( see, e.g. [9, Ch.7]):

$$\hat{\rho}(t) \to [Sp_{\beta}\hat{\rho}(t)] \otimes \frac{\exp\left[-\left(1/k_{B}T\right)\hat{H}_{\beta}\right]}{Sp_{\beta}\exp\left[-\left(1/k_{B}T\right)\hat{H}_{\beta}\right]},\tag{2.7}$$

that redefines  $\hat{\rho}(t)$  in a case, when the unobservable subsystem is taken to be the thermostat at a temperature of T ( $k_B$  is the Boltzmann constant). It follows from the comparison of (2.6) with (2.7) that the approximation **ii**) within the quantum relaxation theory corresponds to the representation of the unobservable subsystem by the thermostat at the infinite temperature, no matter of actual physical properties of the unobservable subsystem.

#### Consequence 2

i), and ii) represent qualitatively distinct approximations. The first approximation (i) represents a closed quantum system as two of such quasi-uncoupled correlated subsystems with mean values of all observables of the closed system being approximately correct. The second approximation (ii) describes the action of the measurement on the state of the observable subsystem and implicitly carries out the transition from a closed system to an open one. By Consequence 1, the model of measurement  $(1.1)\equiv(2.6)$  used in the von Neumann's reduction may be considered as the identification of the classical measuring device with a thermostat of the infinite temperature and the measurement with the ideal thermal contact of the unobservable subsystem with this device. The limited applicability of such model and hence the von Neumann's reduction seems physically evident.

In the form similar to **ii**), the model of measurement was included into the algorithm of reduction in the theory of quantum nondemolition measurements of quantum optics [12, 13]. The "probe" (unobservable) subsystem  $\beta$  was postulated to be determined with certainty in its eigenstate  $|j'\rangle$  at the measurement point of time t, i.e. the probe subsystem density operator is to be presented by a projector  $\hat{\rho}_{\beta j'}(t) = \hat{P}_{\beta j'} \equiv |j'\rangle\langle j'|$ . The equation of such "projective" reduction (is marked by the subscript P) was determined by [12, 13]:

$$\hat{\rho}(t) \to \hat{\rho}_{RPj'}(t) = \hat{\rho}_{\alpha j'}(t) \otimes \hat{P}_{\beta j'} = \frac{Sp_{\beta}\hat{\rho}(t)\hat{P}'_{\beta j'}}{Sp_{\alpha\beta}\hat{\rho}(t)\hat{P}'_{\beta j'}} \otimes \hat{P}_{\beta j'}, \qquad (2.8)$$

being the special case of the equation (3a).

Uniting  $(2.6) \equiv (1.1)$  and (2.8), it is possible to formulate the generalized algorithm of reduction as the procedure of calculating the density operator of an observable subsystem in the approximation of a given state of the unobservable subsystem in the form of two consecutive operations:

- 1. A model of interaction of a closed quantum system with a measuring device is postulated from physical reasons and is expressed mathematically in the form of the approximation of a given state of an unobservable subsystem.
- 2. The density operator of the observable subsystem is calculated from equation (2.2) of correlated quasi-uncoupled subsystems of a closed quantum system.

In (2.6) and (2.8) two special approximations of the given state of the subsystem  $\beta$  are realized. In (2.6) the subsystem  $\beta$  is permanently in each of eigenstates with equal probability and in (2.8) it is with certainty in one of them at the time t.

## Consequence 3

It is known [8, 9] that calculating the mean value of the observable of a closed system described by the operator  $\hat{A}'$ , the von Neumann's reduction of a closed system is carried out implicitly:

$$\langle \hat{A}'(t) \rangle = Sp_{\alpha\beta}\hat{\rho}(t)\hat{A}' = Sp\hat{\rho}_{\alpha N}(t)\hat{A} = \langle \hat{A}(t) \rangle_N.$$
(2.9)

In doing so not the state of the subsystem  $\beta$  is postulated but the operator of the closed system that corresponds to the observable  $\hat{A}$  of the independent subsystem  $\alpha$  ( $\hat{A}' \leftrightarrow \hat{A}$ ) (see [6, Ch. 6.2]). Hence, by Theorem and Consequence 2, the choice of the required operator as  $\hat{A}'$  is equivalent to realization of both approximations **i**) and **ii**).

Such equivalence becomes evident by considering the correlator of observables  $\hat{A}$  and  $\hat{B}$  of subsystems  $\langle (\hat{A} \otimes \hat{B})(t) \rangle = Sp_{\alpha\beta}\hat{\rho}(t) (\hat{A} \otimes \hat{B})$ . If we use the obvious identity  $\hat{A} \otimes \hat{B} = (\hat{A} \otimes \hat{1}_{\beta}) (\hat{1}_{\alpha} \otimes \hat{B})$ , the commutativity  $[\hat{A}', \hat{B}'] = 0$  and the interchangeability of partial

traces  $Sp_{\alpha\beta} = Sp_{\beta\alpha}$ , the correlator can be presented in two forms symmetric relative to subsystems:

$$\left\langle \left( \hat{A} \otimes \hat{B} \right)(t) \right\rangle = Sp_{\alpha\beta}\rho(t) \left( \hat{A} \otimes \hat{B} \right) = \begin{cases} \left\langle \hat{A}(t) \right\rangle_B \left\langle \hat{B}(t) \right\rangle_N, \\ \left\langle \hat{A}(t) \right\rangle_N \left\langle \hat{B}(t) \right\rangle_A, \end{cases}$$
(2.10)

where

$$\left\langle \hat{A}(t) \right\rangle_{B} = Sp_{\alpha}\hat{\rho}_{\alpha B}(t)\hat{A}, \quad \left\langle \hat{B}(t) \right\rangle_{A} = Sp_{\beta}\hat{\rho}_{\beta A}(t)\hat{B}, \tag{2.11}$$

$$\hat{\rho}_{\alpha B}(t) = \frac{Sp_{\beta}\hat{\rho}(t)\hat{\rho}_{\beta B}'}{Sp_{\alpha\beta}\hat{\rho}(t)\hat{\rho}_{\beta B}'}, \quad \hat{\rho}_{\beta A}(t) = \frac{Sp_{\alpha}\hat{\rho}(t)\hat{\rho}_{\alpha A}'}{Sp_{\alpha\beta}\hat{\rho}(t)\hat{\rho}_{\alpha A}'}.$$
(2.12)

$$\hat{\rho}_{\alpha A} = \hat{A}/Sp_{\alpha}\hat{A}, \quad \hat{\rho}_{\beta B} = \hat{B}/Sp_{\beta}\hat{B}, \qquad (2.13)$$

It is supposed here for simplicity that  $\langle \hat{A}(t) \rangle_N \neq 0$ ,  $\langle \hat{B}(t) \rangle_N \neq 0$ , and  $\hat{A}, \hat{B}$  are the nonnegative operators.

The expressions (2.10)-(2.13) are the exact equations for arbitrary Hermitian  $\hat{A}, \hat{B}$ . The meaning of values  $\langle \hat{A}(t) \rangle_B$  and  $\langle \hat{B}(t) \rangle_A$  is clear from the comparison of (2.12) with (2.2),(2.3). For example, the operator  $\hat{\rho}_{\alpha B}(t)$  is the reduced density operator of the subsystem  $\alpha$  in the approximation of the given state  $\hat{\rho}_{\beta B}$  of the subsystem  $\beta$  in accordance with Consequence 2. Accordingly,  $\langle \hat{A}(t) \rangle_B$  is the mean value of the observable  $\hat{A}$  of the subsystem  $\alpha$  if the subsystem  $\beta$  is in the state  $\hat{\rho}_{\beta B}$ . Thus, (2.10) shows that the correlator  $\langle (\hat{A} \otimes \hat{B})(t) \rangle$  of the closed system  $\hat{H}$  is equal to the product of mean values of observables of subsystems calculated for the system  $\hat{H}$  reduced in the different ways. Namely: 1) the reduced state of a subsystem  $\alpha$  is defined at the given  $\hat{\rho}_{\beta} = \hat{\rho}_{\beta B}$ , but the reduced state of the subsystem  $\beta$  is defined simultaneously at the given  $\hat{\rho}_{\alpha} = \hat{\rho}_{\alpha min}$ ; or, leading to the same result, 2) the reduced state of the subsystem  $\alpha$  is defined at the given  $\hat{\rho}_{\beta} = \hat{\rho}_{\beta min}$  but that of the subsystem  $\beta$  at  $\hat{\rho}_{\alpha} = \hat{\rho}_{\alpha A}$ .

In the specific case  $\hat{B} = \hat{1}_{\beta}$  we have  $\hat{\rho}_{\beta B} = \hat{\rho}_{\beta min}$ ,  $\left\langle \hat{B}(t) \right\rangle_{N} = \left\langle \hat{B}(t) \right\rangle_{A} = 1$ ,  $\left\langle \hat{A}(t) \right\rangle_{B} = \left\langle \hat{A}(t) \right\rangle_{N}$  and (2.10) = (2.9).

For this axiomatic choice of the operator of an observable of the closed system  $\hat{A}' \leftrightarrow \hat{A}$ , commonly accepted in the quantum mechanics, in the limits of uncoupled subsystems  $(\hat{H}_{\alpha\beta} = 0)$  the expression of the observable  $\langle \hat{A}'(t) \rangle = \langle \hat{A}(t) \rangle$  is exact. But at  $\hat{H}_{\alpha\beta} \neq 0$  the mathematically exact equality (2.9) is the physical approximation as it contains implicitly approximations **i**) and **ii**). One can see from (2.6), (2.8) - (2.13) and Consequence 2 that depending on a choice of a model of measurements (or of an algorithm of reduction) the

calculated mean value of the observable described by the same operator  $\hat{A}$  and the same density operator of the closed system  $\hat{\rho}(t)$  is different:  $\langle \hat{A}(t) \rangle_N \neq \langle \hat{A}(t) \rangle_{Pj'} \neq \langle \hat{A}(t) \rangle_B$ . The physical reason of this difference is obvious — a choice of a model of measurement (of a algorithm of reduction) corresponds to a choice of the back effect of the measuring device on the state of the observable subsystem and, hence, on the result of measurement. Or, alternatively, it is possible to say following (2.10) that each choice of the model of measurement corresponds to the calculation of different observables of the closed quantum system.

# **III. ALGORITHM OF THE CORRELATED REDUCTION**

If the approximation of the closed system is true, it seems reasonable to seek the algorithm of reduction such that minimize the changing of  $\hat{\rho}(t)$ .

Let us consider expressions:

$$\left\langle \hat{A}(t) \right\rangle_{C} = Sp\hat{\rho}_{\alpha C}(t)\hat{A}, \qquad \left\langle \hat{B}(t) \right\rangle_{C} = Sp\hat{\rho}_{\beta C}(t)\hat{B};$$
(3.1)

$$\left\langle \left(\hat{A} \otimes \hat{B}\right)(t) \right\rangle_{C} = \left[ Sp\hat{\rho}_{\alpha C}(t)\hat{A} \right] \left[ Sp\hat{\rho}_{\beta C}(t)\hat{B} \right].$$
 (3.2)

The equalities (3.1) define mean values of observables  $\hat{A}$  and  $\hat{B}$  of independent subsystems with the correlated density operators  $\hat{\rho}_{\alpha C}(t)$  and  $\hat{\rho}_{\beta C}(t)$ , respectively. But these correlated density operators of subsystems can be determined only approximately for the lack of the exact expression of  $\hat{\rho}(t)$  in the form of (2.1) at  $\hat{H}_{\alpha\beta} \neq 0$ . Let us calculate  $\hat{\rho}_{\alpha C}(t), \hat{\rho}_{\beta C}(t)$  from the system of equations (2.2),(2.3) by the successive approximation method [10, 11]:

$$\hat{\rho}_{\alpha C}(t) = \lim_{n \to \infty} \hat{\rho}_{\alpha}^{(n+1)}(t) = \lim_{n \to \infty} \frac{Sp_{\beta}\hat{\rho}(t)\hat{\rho}_{\beta}^{(n)}(t)}{Sp_{\alpha\beta}\hat{\rho}(t)\hat{\rho}_{\beta}^{(n)}(t)},\tag{3.3}$$

$$\hat{\rho}_{\beta C}(t) = \lim_{n \to \infty} \hat{\rho}_{\beta}^{(n+1)}(t) = \lim_{n \to \infty} \frac{Sp_{\alpha}\hat{\rho}(t)\hat{\rho}_{\alpha}^{(n)}(t)}{Sp_{\alpha\beta}\hat{\rho}(t)\hat{\rho}_{\alpha}^{(n)}(t)},\tag{3.4}$$

where, for an example, the von Neumann's reduced density operators  $\hat{\rho}_{\alpha N}(t)$ ,  $\hat{\rho}_{\beta N}(t)$  may be used as a zeroth-order (n=0) iteration. For the convergent sequences (3.3),(3.4), the calculated self-congruent density operators of subsystems most closely correspond to (2.1). It means that the mathematical operation of reduction (3.3),(3.4) does not change approximately the state of the closed quantum system and hence may be considered to correspond to the quantum nondemolition measurement. A matter of principle is if (2.9) or (3.1) describes observables of subsystems of a closed system more properly.

The expression (2.9) is the mathematical equality but the physical approximation by Consequences 1-3. Each of (3.1) is the physical equality but mathematical approximation by (2.1) and (3.3), (3.4). The accuracy of (3.1) or (3.3),(3.4) can be estimated by a comparison of matrix elements at the left-hand and right-hand sides of (2.1) or by a comparison of mean values at the left-hand and right-hand sides of (3.2). It should be particularly emphasized that the correlated reduction (3.3),(3.4) seems to be more appropriate to the equivalency of subsystems of the closed system than the reduction (1.1) including the physically impossible property ( $T = \infty$ ) of the unobservable subsystem. Therefore we expect that the correlated self-congruent density operators (3.3), (3.4) would more properly describe the dynamics of subsystems in experimental cases when the approximation of the closed system holds.

Let us consider the previous procedure of a correlated reduction and its physical consequences on model problems of quantum optics.

#### IV. DYNAMICS OF A PAIR OF CORRELATED TWO-LEVEL SUBSYSTEMS

Two coupled two-level subsystems constitute the compound closed quantum system now in use to describe many basic problems of quantum optics and quantum informatics [1, 2, 3]. Let us look after the dynamics of such simplest correlated subsystems. To visualize the distinction in the dynamics of subsystems described by the traditional reduction (1.1),(2.6), and by the correlated reduction (2.2),(2.3),(3.3),(3.4) we present at first the consideration in general.

Let us designate eigenstates of uncoupled subsystems  $\alpha$  and  $\beta$  by  $|2\rangle_{\alpha}$ ,  $|1\rangle_{\alpha}$  and  $|2\rangle_{\beta}$ ,  $|1\rangle_{\beta}$ , respectively. Operators of the closed system are represented by matrices of the fourth order, each element of which is designated by two pairs of subscripts (see, e.g., [9, Appendix A]). The first subscript of each pair is attributed to a subsystem  $\alpha$  and second subscript to a subsystem  $\beta$ . The general expression of the density matrix of the considered closed system is then:

$$(\hat{\rho}) = \begin{pmatrix} \rho_{22,22} & \rho_{22,21} & \rho_{22,12} & \rho_{22,11} \\ \rho_{21,22} & \rho_{21,21} & \rho_{21,12} & \rho_{21,11} \\ \rho_{12,22} & \rho_{12,21} & \rho_{12,12} & \rho_{12,11} \\ \rho_{11,22} & \rho_{11,21} & \rho_{11,12} & \rho_{11,11} \end{pmatrix}.$$

$$(4.1)$$

For an observable subsystem  $\alpha$  the reduced von Neumann's density matrix in accordance with (1.1) is defined by expression:

$$(\hat{\rho}_{\alpha N}) = \begin{pmatrix} \rho_{22,22} + \rho_{21,21} & \rho_{22,12} + \rho_{21,11} \\ \rho_{12,22} + \rho_{11,21} & \rho_{12,12} + \rho_{11,11} \end{pmatrix}.$$
(4.2)

Accordingly, for an observable subsystem  $\beta$ :

$$(\hat{\rho}_{\beta N}) = \begin{pmatrix} \rho_{22,22} + \rho_{12,12} & \rho_{22,21} + \rho_{12,11} \\ \rho_{21,22} + \rho_{11,12} & \rho_{21,21} + \rho_{11,11} \end{pmatrix}.$$
(4.3)

In accordance with (2.6), such definition of the operation of reduction is equivalent to the replacement of (4.1) by expressions:

$$\begin{aligned} (\hat{\rho}) \to & (\hat{\rho}_{R\alpha N}) = (\hat{\rho}_{\alpha N}) \otimes (\hat{\rho}_{\beta \min}) \\ & = \frac{1}{2} \begin{pmatrix} \rho_{22,22} + \rho_{21,21} & 0 & \rho_{22,12} + \rho_{21,11} & 0 \\ 0 & \rho_{22,22} + \rho_{21,21} & 0 & \rho_{22,12} + \rho_{21,11} \\ \rho_{12,22} + \rho_{11,21} & 0 & \rho_{12,12} + \rho_{11,11} & 0 \\ 0 & \rho_{12,22} + \rho_{11,21} & 0 & \rho_{12,12} + \rho_{11,11} \end{pmatrix}, \end{aligned}$$
(4.4)

or

$$(\hat{\rho}) \rightarrow (\hat{\rho}_{R\beta N}) = (\hat{\rho}_{\alpha \min}) \otimes (\hat{\rho}_{\beta N})$$

$$= \frac{1}{2} \begin{pmatrix} \rho_{22,22} + \rho_{12,12} & \rho_{22,21} + \rho_{12,11} & 0 & 0 \\ \rho_{21,22} + \rho_{11,12} & \rho_{21,21} + \rho_{11,11} & 0 & 0 \\ 0 & 0 & \rho_{22,22} + \rho_{12,12} & \rho_{22,21} + \rho_{12,11} \\ 0 & 0 & \rho_{21,22} + \rho_{11,12} & \rho_{21,21} + \rho_{11,11} \end{pmatrix}.$$

$$(4.5)$$

For comparison, we write the system of equations (2.2), (2.3) determining correlated reduced density matrices of subsystems of the closed system (4.1):

$$(\hat{\rho}_{\alpha C}) = \begin{pmatrix} \alpha_{22} & \alpha_{21} \\ \alpha_{12} & \alpha_{11} \end{pmatrix} \simeq \begin{pmatrix} (\rho_{22,22}\beta_{22} + \rho_{21,21}\beta_{11} + & (\rho_{22,12}\beta_{22} + \rho_{21,11}\beta_{11} + \\ +\rho_{22,21}\beta_{12} + \rho_{21,22}\beta_{21}) & +\rho_{22,11}\beta_{12} + \rho_{21,12}\beta_{21}) \\ (\rho_{12,22}\beta_{22} + \rho_{11,21}\beta_{11} + & (\rho_{12,12}\beta_{22} + \rho_{11,11}\beta_{11} + \\ +\rho_{12,21}\beta_{12} + \rho_{11,22}\beta_{21}) & +\rho_{12,11}\beta_{12} + \rho_{11,12}\beta_{21}) \end{pmatrix}$$

$$\times \left(\rho_{22,22}\beta_{22} + \rho_{21,21}\beta_{11} + \rho_{22,21}\beta_{12} + \rho_{21,22}\beta_{21} + \rho_{12,12}\beta_{22} + \rho_{11,11}\beta_{11} + \rho_{12,11}\beta_{12} + \rho_{11,12}\beta_{21}\right)^{-1}, \quad (4.6)$$

$$(\hat{\rho}_{\beta C}) = \begin{pmatrix} \beta_{22} & \beta_{21} \\ \beta_{12} & \beta_{11} \end{pmatrix} \simeq \begin{pmatrix} (\rho_{22,22}\alpha_{22} + \rho_{12,12}\alpha_{11} + & (\rho_{22,21}\alpha_{22} + \rho_{12,11}\alpha_{11} + \\ +\rho_{22,12}\alpha_{12} + \rho_{12,22}\alpha_{21}) & +\rho_{22,11}\alpha_{12} + \rho_{12,21}\alpha_{21}) \\ (\rho_{21,22}\alpha_{22} + \rho_{11,12}\alpha_{11} + & (\rho_{21,22}\alpha_{22} + \rho_{11,11}\alpha_{11} + \\ +\rho_{21,12}\alpha_{12} + \rho_{11,22}\alpha_{21}) & +\rho_{21,11}\alpha_{12} + \rho_{11,21}\alpha_{21}) \end{pmatrix} \times (\rho_{22,22}\alpha_{22} + \rho_{12,12}\alpha_{11} + \rho_{22,12}\alpha_{12} + \rho_{12,22}\alpha_{21} \\ & + \rho_{21,21}\alpha_{22} + \rho_{11,11}\alpha_{11} + \rho_{21,11}\alpha_{12} + \rho_{11,21}\alpha_{21})^{-1}. \quad (4.7)$$

It is seen that in the general case the mutual correlation of subsystems (4.6) and (4.7) as distinct from (4.2) and (4.3) implies the following.

1) The possibility of the simultaneous existence of quantum coherence, i.e. nonzero nondiagonal elements of reduced density matrices of both subsystems ( $\alpha_{12} \neq 0, \beta_{12} \neq 0$ ).

2) The inequality of probabilities to find the subsystems in their eigenstates ( $\alpha_{11} \neq \alpha_{22}, \beta_{11} \neq \beta_{22}$ ).

Let us use (4.1)-(4.7) in considering two known physical problems.

# A. Representation of an entangled EPR (Einstein-Podolsky-Rosen) pair of photons as the superposition of separated correlated photons

The model of the entangled EPR state is of considerable current use in the description of experiments on the correlation of photons [1, 2, 3]. The density matrix of such closed quantum system of two noninteracting but correlated photons in designations (4.1) is of the form [1, 2, 3]:

$$(\hat{\rho}_S) = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$
(4.8)

States of subsystems (i.e. individual photons) by the von Neumann's reduction (1.1), (4.2), (4.3) are:

$$(\hat{\rho}_{\alpha N}) = (\hat{\rho}_{\alpha \min}) = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \text{ or } (\hat{\rho}_{\beta N}) = (\hat{\rho}_{\beta \min}) = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$
 (4.9)

By (2.6), (4.4), (4.5), for both subsystems it is equivalent to the replacement of the density matrix (4.8) by the reduced density matrix

$$(\hat{\rho}_S) \to (\hat{\rho}_{SRN}) = \hat{\rho}_{min} = \frac{1}{4}\hat{1}.$$
 (4.10)

The expressions (4.9) and (4.10) show that each photon can be registered in any eigenstate with equal probability (1/2) but they do not determine the mutual correlation of results of simultaneous or consecutive measurements of both photons. The approximate nature of the von Neumann's reduction makes itself evident in the nonequality (4.8) $\neq$ (4.10). On the other hand, as photons in the pure state (4.8) are noninteracting, one might expect the exact expression of the system (4.8) through correlated subsystems in terms similar to (3.3),(3.4) to exist. With (4.6) and (4.7) for (4.8) the iterative procedure (3.3),(3.4) converges already in the first iteration for the arbitrary zeroth-order iteration and results in the following relations between the elements of correlated reduced density matrices of subsystems:  $\alpha_{22} = \beta_{11}$ ,  $\alpha_{21} = -\beta_{21}$ , in addition to usual relations of normalizing and of hermiticity:  $\alpha_{11} = 1 - \alpha_{22}$ ,  $\alpha_{12} = \alpha_{21}^*$ . With these correlation conditions, let us write down the system of equations for calculating the elements  $\alpha_{22}$ ,  $\alpha_{21}$  from (2.1). This system is separated into three groups of equations: for diagonal elements, nondiagonal elements and equations of their relation:

$$\begin{aligned} \alpha_{22}\beta_{22} \leftrightarrow 0 &=: \alpha_{22}(1-\alpha_{22}), \ \alpha_{21}\beta_{12} \leftrightarrow -1/2 =: -|\alpha_{21}|^2, \ \alpha_{22}\beta_{21} \leftrightarrow 0 =: \alpha_{22}(-\alpha_{21}), \\ \alpha_{22}\beta_{11} \leftrightarrow 1/2 &=: \alpha_{22}^2, \qquad \alpha_{21}\beta_{21} \leftrightarrow 0 =: -(\alpha_{21})^2, \qquad \alpha_{21}\beta_{22} \leftrightarrow 0 =: \alpha_{21}(1-\alpha_{22}), \\ \alpha_{11}\beta_{22} \leftrightarrow 1/2 &=: (1-\alpha_{22})^2, \qquad \alpha_{21}\beta_{11} \leftrightarrow 0 =: \alpha_{21}\alpha_{22}, \\ \alpha_{11}\beta_{11} \leftrightarrow 0 &=: (1-\alpha_{22})\alpha_{22}, \qquad \alpha_{11}\beta_{21} \leftrightarrow 0 =: -(1-\alpha_{22})\alpha_{21}. \end{aligned}$$

$$(4.11)$$

The system (4.11) is obviously incompatible as an algebraic system of equation, but it is a compatible system of statistical equations that we have designated by "=:" instead of "=". For example, the system of four equations from the first column of (4.11) is compatible, if it is granted that  $\alpha_{22}$  takes two values:  $\alpha_{22}^{(1)} = 1$  and  $\alpha_{22}^{(2)} = 0$  with equal probability (1/2), and each equation is considered as a result of statistical averaging. In this way, for the first equation of the first column from (4.11) we have:  $0 = (1/2) \left[ \alpha_{22}^{(1)} (1 - \alpha_{22}^{(1)}) + \alpha_{22}^{(2)} (1 - \alpha_{22}^{(2)}) \right]$ . For the second equation:  $1/2 = (1/2) \left[ \left( \alpha^{(1)} \right)^2 + \left( \alpha^{(2)} \right)^2 \right]$ . In solving the system (4.11) as a system of the statistical equations, we obtain within the arbitrary phase factor  $(\exp(i\theta))$ :

$$(\hat{\rho}_S) = \sum_{i=1}^{4} p_i \left\{ \left( \hat{\rho}_{\alpha C}^{(i)} \right) \otimes \left( \hat{\rho}_{\beta C}^{(i)} \right) \right\}$$

$$= \frac{1}{4} \left\{ \begin{pmatrix} 1 & \frac{1}{\sqrt{2}} e^{i\theta} \\ \frac{1}{\sqrt{2}} e^{-i\theta} & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & \frac{1}{\sqrt{2}} e^{i(\theta+\pi)} \\ \frac{1}{\sqrt{2}} e^{-i(\theta+\pi)} & 1 \end{pmatrix} \\ + \begin{pmatrix} 1 & \frac{1}{\sqrt{2}} e^{i(\theta+\pi)} \\ \frac{1}{\sqrt{2}} e^{-i(\theta+\pi)} & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & \frac{1}{\sqrt{2}} e^{i\theta} \\ \frac{1}{\sqrt{2}} e^{-i\theta} & 1 \end{pmatrix} \\ + \begin{pmatrix} 0 & \frac{1}{\sqrt{2}} e^{i(\theta+\pi/2)} \\ \frac{1}{\sqrt{2}} e^{-i(\theta+\pi/2)} & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & \frac{1}{\sqrt{2}} e^{i(\theta+3\pi/2)} \\ \frac{1}{\sqrt{2}} e^{-i(\theta+3\pi/2)} & 0 \end{pmatrix} \\ + \begin{pmatrix} 0 & \frac{1}{\sqrt{2}} e^{i(\theta+3\pi/2)} \\ \frac{1}{\sqrt{2}} e^{-i(\theta+3\pi/2)} & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & \frac{1}{\sqrt{2}} e^{i(\theta+\pi/2)} \\ \frac{1}{\sqrt{2}} e^{-i(\theta+\pi/2)} & 0 \end{pmatrix} \right\}.$$
(4.12)

Consequently, the entangled pure EPR state is exactly represented as a sum of four hidden equiprobable ( $p_i = const = 1/4$ ) states of separated but correlated ( $\hat{\rho}_{\alpha C}^{(i)}, \hat{\rho}_{\beta C}^{(i)}$ ) subsystems. Each of summands presents by itself the distinct state of EPR pare. Each of  $\hat{\rho}_{\alpha C}^{(i)}, \hat{\rho}_{\beta C}^{(i)}$  corresponds to the existence of the subsystem in one of eigenstates: the diagonal elements of density matrices are 1 and 0 only. Note simultaneously the unusual nonzero values of nondiagonal elements. Nondiagonal elements of this type reflect not the usual superposition of eigenstates of one subsystem but the correlation in phase of the eigenstates of the correlated subsystems.

The considered model of a pair of noninteracting two-level subsystems is the elementary example of the compound closed quantum system. The obvious next step of the generalization of the model is taking into account the interaction between subsystems.

#### B. Dynamics of a pair of interacting two-level atoms

The interaction of a pair of two-level subsystems was in detail considered in the theory of magnetic resonance, therefore it is convenient to take advantage of designations accepted there. The hamiltonian of a pair of identical spins-1/2 in an external d-c magnetic field with a simple form of the spin-spin interaction is [14]:

$$(\hat{H}) = \hbar \begin{pmatrix} \omega + J & 0 & 0 & d \\ 0 & -J & c & 0 \\ 0 & c & -J & 0 \\ d & 0 & 0 & -\omega + J \end{pmatrix},$$
(4.13)

where  $\hbar\omega$  is the Zeeman energy of a single spin-1/2 in an external d-c magnetic field, parameters J, c, d characterize the energy of the spin-spin interaction, which for definiteness is believed to be rather weak ( $\omega > J, c, d$ ). Designations of basic states accepted in (4.1) are used, i.e.  $|2\rangle, |1\rangle$  are eigenstates of a single spin-1/2 in the d-c magnetic field with higher and lower energies, respectively.

The operator of evolution of quantum system (4.13) results in:

$$\exp\left(\mp \frac{it}{\hbar}\hat{H}\right) = \begin{cases} \hat{U}(t)\\ \hat{U}^{\dagger}(t) \end{cases} = \\ \begin{pmatrix} e^{\mp iJt} \left[C(\Omega t) \mp \frac{i\omega}{\Omega}S(\Omega t)\right] & 0 & 0 & \mp (id/\Omega)e^{\mp iJt}S(\Omega t) \\ 0 & e^{\pm iJt}\cos(ct) & \mp ie^{\pm iJt}\sin(ct) & 0 & (4.14) \\ 0 & \mp ie^{\pm iJt}\sin(ct) & e^{\pm iJt}\cos(ct) & 0 & (4.14) \\ \mp (id/\Omega)e^{\mp iJt}S(\Omega t) & 0 & 0 & e^{\mp iJt} \left[C(\Omega t) \pm \frac{i\omega}{\Omega}S(\Omega t)\right] \end{cases}$$

where  $C(\Omega t) = \cos(\Omega t)$ ,  $S(\Omega t) = \sin(\Omega t)$ . Thus, the density matrix  $(\hat{\rho}(t))$ , i.e. the dynamics of the system (4.13) at the given initial condition  $(\hat{\rho}(0))$  is known exactly. To describe the dynamics of each spin one should calculate correlated reduced density matrices of spin-1/2 subsystems according to Sec. III. Let the initial state of the system (4.13) be:

$$(\hat{\rho}(0)) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \cos^2 \varphi & -\sin \varphi \cos \varphi & 0 \\ 0 & -\sin \varphi \cos \varphi & \sin^2 \varphi & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$
 (4.15)

For example, the initial condition of a radical pair born during photosynthesis is described in this form [15]. Similarly to the consideration of Subsec. IV A, this initial density matrix is presented as:

$$\hat{\rho}(0) \simeq \hat{\rho}_{RC}(0) = \begin{cases}
\cos^2 \varphi \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} + \sin^2 \varphi \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, |\tan \varphi| > 1, \\
\cos^2 \varphi \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \sin^2 \varphi \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, |\tan \varphi| < 1, , (4.16) \\
\hat{\rho}_{S}, & \tan \varphi = 1, \\
\hat{\rho}_{T}, & \tan \varphi = -1,
\end{cases}$$

where

$$\begin{aligned} (\hat{\rho}_{\mathrm{T}}) &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} = \frac{1}{4} \left[ \begin{pmatrix} 1 & \frac{1}{\sqrt{2}}e^{i\theta} \\ \frac{1}{\sqrt{2}}e^{-i\theta} & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & \frac{1}{\sqrt{2}}e^{i\theta} \\ \frac{1}{\sqrt{2}}e^{-i\theta} & 1 \end{pmatrix} \right. \\ &+ \begin{pmatrix} 0 & \frac{1}{\sqrt{2}}e^{i(\theta+\pi/2)} \\ \frac{1}{\sqrt{2}}e^{-i(\theta+\pi/2)} & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & \frac{1}{\sqrt{2}}e^{i(\theta+\pi/2)} \\ \frac{1}{\sqrt{2}}e^{-i(\theta+\pi/2)} & 0 \end{pmatrix} \\ &+ \begin{pmatrix} 1 & \frac{1}{\sqrt{2}}e^{i(\theta+\pi)} \\ \frac{1}{\sqrt{2}}e^{-i(\theta+\pi)} & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & \frac{1}{\sqrt{2}}e^{i(\theta+\pi)} \\ \frac{1}{\sqrt{2}}e^{-i(\theta+\pi)} & 1 \end{pmatrix} \\ &+ \begin{pmatrix} 0 & \frac{1}{\sqrt{2}}e^{i(\theta+3\pi/2)} \\ \frac{1}{\sqrt{2}}e^{-i(\theta+3\pi/2)} & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & \frac{1}{\sqrt{2}}e^{i(\theta+3\pi/2)} \\ \frac{1}{\sqrt{2}}e^{-i(\theta+3\pi/2)} & 0 \end{pmatrix} \end{bmatrix}. \end{aligned}$$
(4.17)

Unlike (4.12) expression (4.16) is exact only for  $|\tan \varphi| = 1$  (i.e. for the last two rows in (4.16)).

As a result of correlated reduction (3.3),(3.4) the density matrix  $\hat{\rho}(t)$  is represented at any time as follows:

$$\begin{split} (\hat{\rho}(t)) &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & P(\varphi, t) & i\cos(2\varphi)\sin(2ct) - \sin(2\varphi) & 0 \\ 0 & -i\cos(2\varphi)\sin(2ct) - \sin(2\varphi) & M(\varphi, t) & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \to (\hat{\rho}_{RC}(t) \\ &= \begin{cases} \frac{1}{2} \begin{bmatrix} P(\varphi, t) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} + M(\varphi, t) \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \\ \frac{1}{2} \begin{bmatrix} P(\varphi, t) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + M(\varphi, t) \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \\ \frac{1}{2} \begin{bmatrix} P(\varphi, t) \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + M(\varphi, t) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \end{bmatrix}, \quad C(\varphi, t) > 0, \\ \frac{\hat{\rho}_{S}}{\hat{\rho}_{T}}, & \tan(\varphi) = 1, \\ \hat{\rho}_{T}, & \tan(\varphi) = -1, \end{split}$$

where  $C(\varphi, t) = \cos(2\varphi)\cos(2ct)$ ,  $P(\varphi, t) = 1 + C(\varphi, t)$ ,  $M(\varphi, t) = 1 - C(\varphi, t)$ .

Thus, the dynamics of the closed system of two interacting spins-1/2 (4.13), according to the initial state (the value of  $\varphi$ ), may be of two types. For  $|\tan \varphi| = 1$  the system is in the stable statistically defined state S $\leftrightarrow$ (4.12) or T $\leftrightarrow$ (4.17). For  $|\tan \varphi| \neq 1$  there are the harmonic oscillations of probabilities to detect spins in basic states opposite in phase. At  $t_n = (2n+1)\pi/8c$ , (n = 0, 1, 2, ...) for an arbitrary initial state these probabilities are the same and equal 1/2.

# V. DYNAMICS OF A TWO-LEVEL ATOM IN A RESONANT FIELD

To demonstrate the decoherence phenomenon, the model problem of quantum optics on the dynamics of the inversion of a two-level atom interacting with a resonant electromagnetic field of a single-mode cavity (see, e.g., [2, 5]) is used traditionally. It is attractive due to the existence of the experimental setup of the cavity quantum electrodynamics entangling machine [5] and the existence of the exact solution of the theoretical Janes-Cummings model (JCM) [16, 17].

Let us trace basic distinctions in the dynamics of atom and field following from definitions  $(1.1)\leftrightarrow(2.6)$  and  $(2.2),(2.3)\leftrightarrow(3.3),(3.4)$ .

The hamiltonian of the resonant JCM looks like [16, 17, 18, 19, 20, 21]:

$$\hat{H} = \hat{H}_a + \hat{H}_f + \hat{H}_{af} = \frac{\hbar\omega}{2} \left( \hat{P}_{22} - \hat{P}_{11} \right) + \frac{\hbar\omega}{2} \left( \hat{a}^+ \hat{a} + \hat{a}\hat{a}^+ \right) + \frac{i\hbar\Omega}{2} \left( \hat{P}_{21}\hat{a} - \hat{P}_{12}\hat{a}^+ \right), \quad (5.1)$$

where  $\hat{H}_a, \hat{H}_f, \hat{H}_{af}$  are hamiltonians of the atom, field and atom-field interactions,  $\omega$  is the resonant frequency of the field,  $\Omega$  is the constant of the atom-field interaction,  $\hat{P}_{ij}$ are projection atomic operators representing square matrices of the second order, the ijth element of which is equal to 1, others are equal to 0, i, j = 1, 2 and the subscript 2 corresponds to the level with the higher energy.

It is convenient to write down the solution of the JCM (obtained in a general form in the Heisenberg picture in Ref. [19]) in the Schrödinger picture in the form of expression for the operator of evolution [20, 21]:

$$\begin{cases} \hat{U}(t) \\ \hat{U}^{\dagger}(t) \end{cases} = \exp\left(\mp \frac{it}{\hbar}\hat{H}\right) = \hat{P}_{22}\exp\left(\mp i\omega t\hat{a}\hat{a}^{+}\right)\cos\left[\frac{\Omega t}{2}\left(\hat{a}\hat{a}^{+}\right)^{1/2}\right] \\ + \hat{P}_{11}\exp\left(\mp i\omega t\hat{a}^{+}\hat{a}\right)\cos\left[\frac{\Omega t}{2}\left(\hat{a}^{+}\hat{a}\right)^{1/2}\right] \pm \hat{P}_{21}\exp\left(\mp i\omega t\hat{a}\hat{a}^{+}\right)\exp(i\hat{\varphi})\sin\left[\frac{\Omega t}{2}\left(\hat{a}^{+}\hat{a}\right)^{1/2}\right] \\ \mp \hat{P}_{12}\exp\left(\mp i\omega t\hat{a}^{+}\hat{a}\right)\exp(-i\hat{\varphi})\sin\left[\frac{\Omega t}{2}\left(\hat{a}\hat{a}^{+}\right)^{1/2}\right], \qquad (5.2)$$

where  $\exp(\pm i\hat{\varphi}) = \begin{cases} (\hat{a}^+ \hat{a} + 1)^{-1/2} \hat{a} \\ \hat{a}^+ (\hat{a}^+ \hat{a} + 1)^{-1/2} \end{cases}$  is the operator of a phase of a field [22]. Use of the operators  $\exp(\pm i\hat{\varphi})$  and  $\hat{P}_{ij}$  (instead of more widespread Pauli operators) allows us to

write down the operator of evolution in the symmetric compact form (5.2), convenient for the further analysis.

The elementary example (traditionally used in educational purposes [16, 17]) is the consideration of the dynamics of the inversion of the atom, if the initial condition was  $\hat{\rho}(0) = \hat{\rho}_a(0)\hat{\rho}_f(0) = \hat{P}_{22}|0\rangle\langle 0|$ , i.e. the atom was in the excited state  $|2\rangle$  and the field was in the vacuum state. The known result, at once following from (5.2) and (1.1), consists in oscillations with the frequency  $\Omega$  of the inversion of the atom or the probability of the existence of the photon in the cavity (vacuum Rubi oscillation [5]):

$$\hat{\rho}_{aN}(t) = \hat{P}_{22} \cos^2(\Omega t/2) + \hat{P}_{11} \sin^2(\Omega t/2), \qquad (5.3)$$

or

$$\hat{\rho}_{fN}(t) = \sin^2(\Omega t/2)|1\rangle\langle 1| + \cos^2(\Omega t/2)|0\rangle\langle 0|.$$
(5.4)

At  $t \neq 2\pi l/\Omega$ , (l = 0, 1, 2, ...) the atom and the field are in a mixture state corresponding to decoherence of the observed subsystem.

It is easy to notice that expressions (5.3) and (5.4) are incompatible for the system (2.2),(2.3). We search for the correlated reduced density operators of the atom and the field with (3.3),(3.4), using (5.3) and (5.4) as the zeroth-order iteration. As a result, we obtain:

$$\hat{\rho}_a(t) = C(t)\hat{P}_{22} + S(t)\hat{P}_{11}, \quad \hat{\rho}_f(t) = S(t)|0\rangle\langle 0| + C(t)|1\rangle\langle 1|, \tag{5.5}$$

where S(t) = 1 - C(t),  $s = \sin(\Omega t/2), c = \cos(\Omega t/2)$ ,

$$C(t) = \lim_{n \to \infty} \frac{c^{2n}}{c^{2n} + s^{2n}} = \begin{cases} 1, & [(4l-1)\pi]/2\Omega < t < [(4l+1)\pi]/2\Omega, \\ 1/2, & t = [(2l+1)\pi]/2\Omega, \\ 0, & [(4l+1)\pi]/2\Omega < t < [(4l+3)\pi]/2\Omega. \end{cases}$$
(5.6)

According to (5.5) and (5.6), probabilities to find the atom and the field in their eigenstates are stepwise periodic functions of time with the period  $\pi/\Omega$ . Practically all time the atom and the field are in pure states and only at the discrete moments of time they are in a mixture state equiprobable for two probable pure states.

The expression (5.3) is employed at once to describe the experimentally observed vacuum Rubi oscillations [5, 23]. We call attention to the experimental "dead time" prior to the first-time photon emission in these experiments. The authors of [5, 23] included the effective interaction time taking into account the spatial variation of the atom-field coupling when atom moves across the cavity. It seems us unreasonable for the initially vacuum field up to the first photon emission. All the space: the interior-space and the outer-space of the cavity, is uniform for the vacuum field. Thus all the atom way after the excitation must be included into consideration and not just the interior of the cavity. Then the experimental results [5, 23] contain the dead time when atom is in the excited state and there is not spontaneous emission that corresponds qualitatively to the time interval  $(0 \rightarrow \pi/2\Omega)$  of (5.5), (5.6).

The other experimental result corresponding qualitatively to (5.5), (5.6) consist in the observing of the omnidirectional superfluorescense [26, 27]. There is the well-known experimental situation, when each atom of the system of many uncoupled two-level atoms for a short (in comparison with  $\pi/\Omega$ ) time interval is transferred into the excited state and the spontaneous emission of the inverted system of atoms is investigated without cavity. The pulse of radiation experimentally observed in this case after a time interval  $\tau$  is traditionally interpreted, after the authors of first observation [24], as a pulse of superradiation or superfluorescence [25]. And  $\tau$  is considered to be defined by the time of self-organization in the system of atoms at the expense of their interaction through a common radiation field [24].

According to (5.6), each atom emits a photon at the same time  $\tau = \pi/2\Omega$  if at t = 0 each atom was in the excited state, i.e. the pulsed emission of spontaneous radiation by the system of noninteracting atoms may be due to the correlation of each atom with the initially vacuum field. And for the gaseous spherical sample [26, 27] the directions of the spontaneous emission of photons are equiprobable.

#### VI. CONCLUSION

Results of the present work may be formulated as the following assertions:

- 1. The identification of an operator of an observable of a subsystem of a closed quantum system with the appropriate extended operator of the independent observable subsystem  $(\hat{A}' \leftrightarrow \hat{A})$  corresponds to the physical approximation, at which this obserable is calculated provided that the unobservable subsystem is in the steady state of minimum information (infinite temperature).
- 2. The known algorithms of reduction correspond to the approximation of a given state

of the unobservable subsystem.

- 3. The algorithm of correlated reduction that most corresponds to the ideal quantum measurement of observables of subsystems of the closed quantum sistem may be presented as the self-congruent calculation of mutually correlated reduced density operators of subsystems  $\hat{\rho}_{\alpha C}(t)$ ,  $\hat{\rho}_{\beta C}(t)$  by the successive approximation method (3.3),(3.4). The mean values of observables of subsystems at the ideal quantum measurements are defined by expressions  $\langle \hat{A}(t) \rangle_C = Sp\hat{\rho}_{\alpha C}(t)\hat{A}, \langle \hat{B}(t) \rangle_C = Sp\hat{\rho}_{\beta C}(t)\hat{B}.$
- The correlations in the dynamics of subsystems are demonstrated with examples of simple compound quantum systems: 1) the pair of the coupled two-level atoms (spins-1/2), and 2) the two-level atom in a single-mode resonant field.

## Acknowledgments

The author acknoledge numerous discussions with V. N. Lisin.

- [1] D. N. Klyshko, Uspechi **168**, 975 (1998).
- [2] M. B. Mensky, Uspechi **170**, 631 (2000).
- [3] S. Ya. Kilin, Uspechi **169**, 507 (1999).
- [4] I. Prigogine, *The End of Certainty* (The Free Press, NY-London-Toronto-Sidney-Singapore, 1997).
- [5] J. M. Raimond, M. Brune and S. Haroche, Rev. Mod. Phys. 73, 565 (2001).
- [6] J. von Neumann, Mathematical Foundations of Quantum Mechanics (Princeton University Press, Princeton, 1955).
- [7] P. A. M. Dirac, The Principles of Quantum Mechanics (Clarendon Press, Oxford, 1958).
- [8] U. Fano, Rev. Mod. Phys. **29**, 74 (1957).
- [9] K. Blum, Density Matrix. Theory and Applications (Plenum Press, New York and London, 1981).
- [10] N. K. Solovarov, in Proceedings 2 of Intern. Conf. Geometrization of Physics V, edited by V. I. Bashkov, (Kazan State University, Kazan, Russia, 2001), p. 112.

- [11] N. K. Solovarov, in Zavoisky Physical-Technical Institute, Annual 2001, edited by K. M. Salikhov, (Phystech Press, Kazan, 2002), p. 104.
- [12] N. Imoto, M. Ueda and T. Ogawa, Phys. Rev. A41, 4127 (1990).
- [13] T. Fukuo, T. Ogawa and K. Nakamura, Phys. Rev. A58, 3293 (1998).
- [14] A. Abragam and B. Bleaney, Electron Paramagnetic Resonance of Transition Ions (Clarendon Press, Oxford, 1970), v. 1, ch. 9.5.
- [15] Yu. E. Kandrashkin, K. M. Salikhov and D. Shtelik, Appl. Magn. Reson. 12, 141 (1997).
- [16] W. H. Louisell, Quantum statistical properties of radiation (John Wiley & Sons, New York, 1973).
- [17] L. Allen, J. H. Eberly, Optical Resonance and Two-level Atoms (John Wiley and Sons, NY-London-Sydney-Toronto, 1975).
- [18] H. I. Yoo and J. H. Eberly, Phys. Reports **118**, 239 (1985).
- [19] J. C. Ackerhalt and K. Rzążewski, Phys. Rev. A12, 2549 (1975).
- [20] N. K. Solovarov, Optica Acta 27, 393 (1980).
- [21] D. A. Demidov and N. K. Solovarov, Laser Physics 5, 997 (1995).
- [22] R. Loudon, The Quantum Theory of Light (Clarendon Press, Oxford, 1973).
- [23] M. Brune, E. Hagley, J. Dreyer et al., Phys. Rev. Lett. 77, 4887 (1996).
- [24] N. Skribanovitz, I. P. Herman, J. C. MacGillivray et al., Phys. Rev. Lett. 30, 309 (1973).
- [25] R. H. Dicke, Phys. Rev. **93**, 99 (1954).
- [26] A. I. Lvovsky, S. R. Hartmann and F. Moshary, Phys. Rev. Lett. 82, 4420 (1999).
- [27] A. I. Lvovsky, S. R. Hartmann and F. Moshary, Phys. Rev. Lett. 89, 263602 (2002).