

Special functions, raising and lowering operators

N Cotfas

Faculty of Physics, University of Bucharest, PO Box 76-54, Postal Office 76,
Bucharest, Romania, E-mail address: ncotfas@yahoo.com

Abstract. The Schrödinger equations which are exactly solvable in terms of associated special functions are directly related to some self-adjoint operators defined in the theory of hypergeometric type equations. The fundamental formulae occurring in a supersymmetric approach to these Hamiltonians are consequences of some formulae concerning the general theory of associated special functions. We use this connection in order to obtain a *general theory of Schrödinger equations exactly solvable in terms of associated special functions*, and to extend certain results known in the case of some particular potentials.

1. Introduction

It is well-known [2, 4] that, in the case of certain potentials, the Schrödinger equation is exactly solvable and its solutions can be expressed in terms of the so-called *associated special functions*. Our purpose is to present a general theory of these quantum systems. Our systematic study recovers a number of earlier results in a natural unified way and also leads to new findings.

The number of articles concerning exactly solvable quantum systems and related subjects is very large (see [2, 4, 5] and references therein). Our approach is based on the raising/lowering operators presented in general form (for the first time to our knowledge) by Jafarizadeh and Fakhri [5]. We reobtain these operators in a much simpler way, and use them in a rather different way. More details can be found in [3].

2. Orthogonal polynomials and associated special functions

Many problems in quantum mechanics and mathematical physics lead to equations of hypergeometric type

$$\sigma(s)y''(s) + \tau(s)y'(s) + \lambda y(s) = 0 \quad (1)$$

where $\sigma(s)$ and $\tau(s)$ are polynomials of at most second and first degree, respectively, and λ is a constant. This equation can be reduced to the self-adjoint form

$$[\sigma(s)\varrho(s)y'(s)]' + \lambda\varrho(s)y(s) = 0 \quad (2)$$

by choosing a function ϱ such that $[\sigma(s)\varrho(s)]' = \tau(s)\varrho(s)$. For $\lambda = \lambda_l = -\frac{1}{2}l(l-1)\sigma'' - l\tau'$ with $l \in \mathbb{N}$ there exists a polynomial Φ_l of degree l satisfying (1), that is,

$$\sigma(s)\Phi_l''(s) + \tau(s)\Phi_l'(s) + \lambda_l\Phi_l(s) = 0. \quad (3)$$

If there exists a finite or infinite interval (a, b) such that

$$\sigma(s)\varrho(s)s^k|_{s=a} = 0 \quad \sigma(s)\varrho(s)s^k|_{s=b} = 0 \quad \text{for all } k \in \mathbb{N} \quad (4)$$

and if $\sigma(s) > 0$, $\varrho(s) > 0$ for all $s \in (a, b)$, then the polynomials Φ_l are orthogonal with weight function $\varrho(s)$ in the interval (a, b) . In this case Φ_l are known as *classical orthogonal polynomials* [7].

Let $\kappa(s) = \sqrt{\sigma(s)}$. By differentiating the equation (3) m times and multiplying it by $\kappa^m(s)$, we get for each $m \in \{0, 1, 2, \dots, l\}$ the *associated differential equation* which can be written as $H_m \Phi_{l,m} = \lambda_l \Phi_{l,m}$, where

$$H_m = -\sigma(s) \frac{d^2}{ds^2} - \tau(s) \frac{d}{ds} + \frac{m(m-2)}{4} \frac{\sigma'^2(s)}{\sigma(s)} + \frac{m\tau(s)}{2} \frac{\sigma'(s)}{\sigma(s)} - \frac{1}{2} m(m-2) \sigma''(s) - m\tau'(s)$$

and $\Phi_{l,m}(s) = \kappa^m(s) \Phi_l^{(m)}(s)$ are known as the *associated special functions*. The set $\{\Phi_{m,m}, \Phi_{m+1,m}, \Phi_{m+2,m}, \dots\}$ is an orthogonal sequence ([7], pag. 8) in the Hilbert space

$$\mathcal{H} = \left\{ \varphi : (a, b) \longrightarrow \mathbb{R} \mid \int_a^b |\varphi(s)|^2 \varrho(s) ds < \infty \right\} \quad \text{with} \quad \langle \varphi, \psi \rangle = \int_a^b \varphi(s) \psi(s) \varrho(s) ds.$$

For each $m \in \mathbb{N}$, let \mathcal{H}_m be the linear span of $\{\Phi_{m,m}, \Phi_{m+1,m}, \Phi_{m+2,m}, \dots\}$. In the sequel we shall restrict us to the case when \mathcal{H}_m is dense in \mathcal{H} for all $m \in \mathbb{N}$. For this it is sufficient the interval (a, b) to be finite, but not necessary.

3. Raising and lowering operators. Factorizations for H_m

Lorente has shown recently [6] that a factorization of H_0 can be obtained by using the well-known three term recurrence relation satisfied by Φ_l and a consequence of Rodrigues formula. Following Lorente's idea we obtain a factorization of H_m by using the definition $\Phi_{l,m}(s) = \kappa^m(s) \Phi_l^{(m)}(s)$ and a three term recurrence relation.

Differentiating $\Phi_{l,m}(s) = \kappa^m(s) \Phi_l^{(m)}(s)$ we get the relation

$$\Phi_{l,m+1}(s) = \left(\kappa(s) \frac{d}{ds} - m\kappa'(s) \right) \Phi_{l,m}(s) \quad \text{for all } m \in \{0, 1, \dots, l-1\}. \quad (5)$$

If we differentiate (3) $m-1$ times and multiply the obtained relation by κ^{m-1} then we get for each $m \in \{1, 2, \dots, l-1\}$ the three term recurrence relation

$$\Phi_{l,m+1}(s) + \left(\frac{\tau(s)}{\kappa(s)} + 2(m-1)\kappa'(s) \right) \Phi_{l,m}(s) + (\lambda_l - \lambda_{m-1}) \Phi_{l,m-1}(s) = 0 \quad (6)$$

and $\left(\frac{\tau(s)}{\kappa(s)} + 2(l-1)\kappa'(s) \right) \Phi_{l,l}(s) + (\lambda_l - \lambda_{l-1}) \Phi_{l,l-1}(s) = 0$. A direct consequence of these formulae is the relation

$$(\lambda_l - \lambda_m) \Phi_{l,m}(s) = \left(-\kappa(s) \frac{d}{ds} - \frac{\tau(s)}{\kappa(s)} - (m-1)\kappa'(s) \right) \Phi_{l,m+1}(s) \quad (7)$$

satisfied for all $m \in \{0, 1, \dots, l-1\}$.

The operators $A_m : \mathcal{H}_m \longrightarrow \mathcal{H}_{m+1}$ and $A_m^+ : \mathcal{H}_{m+1} \longrightarrow \mathcal{H}_m$ defined by

$$A_m = \kappa(s) \frac{d}{ds} - m\kappa'(s) \quad A_m^+ = -\kappa(s) \frac{d}{ds} - \frac{\tau(s)}{\kappa(s)} - (m-1)\kappa'(s) \quad (8)$$

satisfy the relations $A_m \Phi_{l,m} = \Phi_{l,m+1}$ and $A_m^+ \Phi_{l,m+1} = (\lambda_l - \lambda_m) \Phi_{l,m}$ (see figure 1).




Figure 1. The functions $\Phi_{l,m}$ satisfy the relation $H_m \Phi_{l,m} = \lambda_l \Phi_{l,m}$, and are related (up to some multiplicative constants) through the operators A_m , A_m^+ , a_m , a_m^+ , U_m and $U_m^{-1} = U_m^+$.

Since $\sigma^m(s)\Phi_l^{(m)}(s)\Phi_k^{(m+1)}(s)$ is a polynomial and the function $\sigma(s)\varrho(s)$ satisfies (4), integrating by parts one obtains $\langle A_m \Phi_{l,m}, \Phi_{k,m+1} \rangle = \langle \Phi_{l,m}, A_m^+ \Phi_{k,m+1} \rangle$, that is, the operators A_m and A_m^+ are mutually adjoint. From the relation

$$\|\Phi_{l,m+1}\|^2 = \langle \Phi_{l,m+1}, \Phi_{l,m+1} \rangle = \langle A_m \Phi_{l,m}, \Phi_{l,m+1} \rangle = \langle \Phi_{l,m}, A_m^+ \Phi_{l,m+1} \rangle = (\lambda_l - \lambda_m) \|\Phi_{l,m}\|^2$$

it follows that $\lambda_l > \lambda_m$ for all $l > m$, and $\|\Phi_{l,m+1}\| = \sqrt{\lambda_l - \lambda_m} \|\Phi_{l,m}\|$. This is possible only if $\sigma''(s) \leq 0$ and $\tau'(s) < 0$. Particularly, we have $\lambda_l \neq \lambda_k$ if and only if $l \neq k$.

The operators $H_m : \mathcal{H}_m \rightarrow \mathcal{H}_m$ are self-adjoint, admit the factorizations

$$H_m - \lambda_m = A_m^+ A_m \quad H_{m+1} - \lambda_m = A_m A_m^+ \quad (9)$$

and satisfy the intertwining relations $H_m A_m^+ = A_m^+ H_{m+1}$, $A_m H_m = H_{m+1} A_m$.

4. Creation and annihilation operators. Coherent states

For each $m \in \mathbb{N}$, the sequence $\{|m, m \rangle, |m+1, m \rangle, |m+2, m \rangle, \dots\}$, where $|l, m \rangle = \Phi_{l,m} / \|\Phi_{l,m}\|$ is an orthonormal basis of \mathcal{H} , and $U_m : \mathcal{H} \rightarrow \mathcal{H}$, $U_m |l, m \rangle = |l+1, m+1 \rangle$ is a unitary operator. The mutually adjoint operators (see figure 1) $a_m, a_m^+ : \mathcal{H}_m \rightarrow \mathcal{H}_m$, $a_m = U_m^+ A_m$, $a_m^+ = A_m^+ U_m$ satisfy the relations

$$a_m |l, m \rangle = \sqrt{\lambda_l - \lambda_m} |l-1, m \rangle \quad a_m^+ |l, m \rangle = \sqrt{\lambda_{l+1} - \lambda_m} |l+1, m \rangle \quad (10)$$

and allow us to factorize H_m as $H_m - \lambda_m = a_m^+ a_m$.

The Lie algebra \mathcal{L}_m generated by $\{a_m^+, a_m\}$ is isomorphic to $su(1, 1)$ if $\sigma'' < 0$, and it is isomorphic to the Heisenberg-Weyl algebra $h(2)$ if $\sigma'' = 0$.

Let $m \in \mathbb{N}$ be a fixed natural number, and let $|n \rangle = |m+n, m \rangle$, $e_n = \lambda_{m+n} - \lambda_m$, $\varepsilon_0 = 1$, $\varepsilon_n = e_1 e_2 \dots e_n$. Since $0 = e_0 < e_1 < e_2 < \dots < e_n < \dots$ and

$$a_m |n \rangle = \sqrt{e_n} |n-1 \rangle \quad a_m^+ |n \rangle = \sqrt{e_{n+1}} |n+1 \rangle \quad (H_m - \lambda_m) |n \rangle = e_n |n \rangle \quad (11)$$

we can define a system of coherent states by using the general setting presented in [1].

If $R = \limsup_{n \rightarrow \infty} \sqrt[n]{\varepsilon_n} \neq 0$ then we can define

$$|z \rangle = \frac{1}{N(|z|^2)} \sum_{n \geq 0} \frac{z^n}{\sqrt{\varepsilon_n}} |n \rangle \quad \text{where} \quad (N(|z|^2))^2 = \sum_{n=0}^{\infty} \frac{|z|^{2n}}{\varepsilon_n} \quad (12)$$

for any z in the open disk $C(0, R)$ of center 0 and radius R . We get in this way a continuous family $\{|z \rangle | z \in C(0, R)\}$ of normalized coherent states such that $a_m |z \rangle = z |z \rangle$.

5. Application to Schrödinger type operators

The problem of factorization of operators H_m is a very important one since it is directly related to the factorization of some Schrödinger type operators [2, 4]. If we use a change of variable $s = s(x)$ such that $ds/dx = \kappa(s(x))$ or $ds/dx = -\kappa(s(x))$ and define the new functions $\Psi_{l,m}(x) = \sqrt{\kappa(s(x)) \varrho(s(x))} \Phi_{l,m}(s(x))$ then the relation $H_m \Phi_{l,m} = \lambda_l \Phi_{l,m}$ becomes an equation of Schrödinger type

$$-\frac{d^2}{dx^2} \Psi_{l,m}(x) + V_m(x) \Psi_{l,m}(x) = \lambda_l \Psi_{l,m}(x). \quad (13)$$

For example, by starting from the equation of Jacobi polynomials with $\alpha = \mu - 1/2$, $\beta = \eta - 1/2$, and using the change of variable $s(x) = \cos x$ we obtain the Schrödinger equation corresponding to the Pöschl-Teller potential [1]

$$V_0(x) = \frac{1}{4} \left[\frac{\mu(\mu-1)}{\cos^2(x/2)} + \frac{\eta(\eta-1)}{\sin^2(x/2)} \right] - \frac{(\mu+\eta)^2}{4}. \quad (14)$$

If we choose the change of variable $s = s(x)$ such that $ds/dx = \kappa(s(x))$, then the operators corresponding to A_m and A_m^+ are the adjoint conjugate operators

$$\begin{aligned} \mathcal{A}_m &= [\kappa(s)\varrho(s)]^{1/2} A_m [\kappa(s)\varrho(s)]^{-1/2} \Big|_{s=s(x)} = \frac{d}{dx} + W_m(x) \\ \mathcal{A}_m^+ &= [\kappa(s)\varrho(s)]^{1/2} A_m^+ [\kappa(s)\varrho(s)]^{-1/2} \Big|_{s=s(x)} = -\frac{d}{dx} + W_m(x) \end{aligned} \quad (15)$$

where the *superpotential* $W_m(x)$ is given by the formula

$$W_m(x) = -\frac{\tau(s(x))}{2\kappa(s(x))} - \frac{2m-1}{2\kappa(s(x))} \frac{d}{dx} \kappa(s(x)). \quad (16)$$

From the relations satisfied by A_m and A_m^+ we get the formulae

$$-\frac{d^2}{dx^2} + V_m(x) - \lambda_m = \mathcal{A}_m^+ \mathcal{A}_m \quad -\frac{d^2}{dx^2} + V_{m+1}(x) - \lambda_m = \mathcal{A}_m \mathcal{A}_m^+ \quad (17)$$

$$V_m(x) - \lambda_m = W_m^2(x) - \dot{W}_m(x) \quad V_{m+1}(x) - \lambda_m = W_m^2(x) + \dot{W}_m(x) \quad (18)$$

$$W_m(x) = -\frac{\dot{\Psi}_{m,m}(x)}{\Psi_{m,m}(x)} \quad V_m(x) = \frac{\ddot{\Psi}_{m,m}(x)}{\Psi_{m,m}(x)} + \lambda_m \quad (19)$$

where the dot sign means derivative with respect to x .

If we choose the change of variable $s = s(x)$ such that $ds/dx = -\kappa(s(x))$ then the formulae are very similar (only some signs are changed). In the case of Pöschl-Teller potential (14) we get

$$W_0(x) = \frac{1}{2} \left[\mu \cot \frac{x}{2} - \eta \tan \frac{x}{2} \right].$$

References

- [1] Antoine J-P, Gazeau J-P, Monceau P, Klauder J R and Penson K A 2001 *J. Math. Phys.* **42** 2349-87
- [2] Cooper F, Khare A and Sukhatme U 1995 *Phys. Rep.* **251** 267-385
- [3] Cotfas N 2002 *J. Phys. A: Math. Gen.* **35** 9355-9365 (quant-ph/0206129)
- [4] Infeld L and Hull T E 1951 *Rev. Mod. Phys.* **23** 21-68
- [5] Jafarizadeh M A and Fakhri H 1998 *Ann. Phys., NY* **262** 260-76
- [6] Lorente M 2001 *J. Phys. A: Math. Gen.* **34** 569-88
- [7] Nikiforov A F, Suslov S K and Uvarov V B 1991 *Classical Orthogonal Polynomials of a Discrete Variable* (Berlin: Springer) pp 2-17