

Remarks on the Bose description of the Pauli spin operators*

^{1,2}Hong-yi Fan and ¹Hai-liang Lu

¹Department of Physics, Shanghai Jiao Tong University, Shanghai, 200030, China

²Department of Material Science and Engineering, University of Science and Technology of China, Hefei, Anhui 230026, China

October 26, 2018

Abstract

Using both the fermionic-like and the bosonic-like properties of the Pauli spin operators we discuss the Bose description of the Pauli spin operators firstly proposed by Shigefumi Naka, and derive another new bosonic representation of Pauli operators. The eigenvector of σ_- in the bosonic representation is a nonlinear coherent state with the eigenvalues being the Grassmann numbers.

1 Introduction

The purpose of this paper is to make a supplement to the Bose description of the Pauli spin operators proposed by Shigefumi Naka [1]. This Bose operator description can also be applied to the Fermi operators, since the quantization of the anti-commuting variables by using anti-commutators generates the elements of Clifford algebra such as Pauli matrices. To see the close relationship between Fermi operators and the Pauli spin operators more clearly, we use the 2×2 matrix representation of the Fermi operators. Let f and f^\dagger be fermion annihilation and creation operators respectively, satisfying anti-commutative relations

$$\{f, f^\dagger\}_+ = 1, f^2 = 0, f^{\dagger 2} = 0, \quad (1)$$

*Work supported by the National Natural Science Foundation of China under grant 10175057 and The President Foundation of Chinese Academy of Science

and let $|0\rangle$ be the fermion vacuum state, then $f^\dagger |0\rangle = |1\rangle$, $f |0\rangle = 0$, $f^\dagger |1\rangle = |0\rangle$. Using the matrix representation of $|1\rangle$ and $|0\rangle$,

$$|0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, |1\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad (2)$$

one can see

$$f = |0\rangle \langle 1| = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \sigma_-, \quad f^\dagger = |1\rangle \langle 0| = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} = \sigma_+, \quad (3)$$

so the 2×2 representations of f and f^\dagger are just the Pauli matrices. Therefore, the Bose description of the Pauli spin operators is equivalent to the Bose description of Fermi operators.

In Ref. [1] Shigefumi firstly proposed the representation (he called the Bose discription of fermion) of the anti-commuting operators (or of the pseudospin operator) in the Hilbert space from the simple observation such that dividing the Hilbert space into two subspaces, the anti-commuting operators can be realized as the operators having matrix elements between these subspaces. By setting

$$\begin{aligned} |n, 1\rangle &= \frac{1}{\sqrt{(2n+1)!}} (a^\dagger)^{2n+1} |0\rangle = |2n+1\rangle, \\ |n, 2\rangle &= \frac{1}{\sqrt{(2n)!}} (a^\dagger)^{2n} |0\rangle = |2n\rangle, \end{aligned} \quad (4)$$

in Fock space, here $|n\rangle = a^{\dagger n} / \sqrt{n!} |0\rangle$ is the Fock state, and noting that

$$\cos^2 \frac{\pi N}{2} = \sum_{n=0}^{\infty} |2n\rangle \langle 2n|, \quad (5)$$

Shigefumi identified the Bose description of Fermi operators, though without giving a manifest derivation, as

$$f = \frac{\cos^2 \frac{\pi N}{2}}{\sqrt{N+1}} a = \sum_{n=0}^{\infty} |2n\rangle \langle 2n+1|, \quad f^\dagger = a^\dagger \frac{\cos^2 \frac{\pi N}{2}}{\sqrt{N+1}} = \sum_{n=0}^{\infty} |2n+1\rangle \langle 2n|. \quad (6)$$

Later the bosonization structure of fermions are also discussed in Ref. [2].

In the following we shall show how to directly employ both the bosonic-like property and fermionic-like property of Pauli spin operators σ_i to derive the Shigefumi's description. To go a step further, we make a supplement to this description, which means that another Bose representation of Pauli spin operators can be found. We then note that a nonlinear coherent state formalism can be introduced for the Bose description of the eigenvectors of Pauli spin operator σ_- . However, its eigenvalue is a Grassmann number, which Ohnuki et al used to construct their fermionic coherent states [3].

2 Derivation of Shigefumi's description

Pauli spin operators σ_i obeys the anticommutative relation

$$\{\sigma_i, \sigma_j\}_+ = 2\delta_{ij}. \quad (7)$$

Introducing

$$\sigma_{\pm} = \frac{1}{2}(\sigma_1 \pm i\sigma_2), \quad \sigma_+ = \sigma_-^{\dagger}, \quad (8)$$

Louisell [4] summarizes the following commutative relations (bosonic-like property)

$$\begin{aligned} [\sigma_{\pm}, \sigma_1] &= \pm\sigma_3, & [\sigma_{\pm}, \sigma_2] &= i\sigma_3, \\ [\sigma_{\pm}, \sigma_3] &= \mp 2\sigma_{\pm}, & [\sigma_+, \sigma_-] &= \sigma_3, \end{aligned} \quad (9)$$

as well as the anti-commutative relations (fermionic-like property)

$$\begin{aligned} \{\sigma_{\pm}, \sigma_1\}_+ &= I, & \{\sigma_{\pm}, \sigma_2\}_+ &= \pm i, \\ \{\sigma_{\pm}, \sigma_3\}_+ &= 0, & \{\sigma_+, \sigma_-\}_+ &= I. \end{aligned} \quad (10)$$

We can deduce the representation (6) by the following method. Without loss of generality, we assume that the Pauli spin operators can be expressed as

$$\sigma_- = f(N)a, \quad \sigma_+ = a^{\dagger}f(N), \quad (11)$$

where $f(N)$ is an operator-valued function of number operator $N = a^{\dagger}a$. It follows from $[a, a^{\dagger}] = 1$, and

$$f(N)a = af(N-1) \quad (12)$$

that

$$\sigma_- \sigma_+ = (N+1)f^2(N), \quad \sigma_+ \sigma_- = a^{\dagger}f^2(N)a = Nf^2(N-1). \quad (13)$$

Substituting (13) into the anti-commutative relation (10) we see

$$(N+1)f^2(N) + Nf^2(N-1) = I. \quad (14)$$

Due to the eigenvalues of N are 0, 1, 2, ..., the solution to (14) is

$$(N+1)f^2(N) = \cos^{2l} \frac{\pi N}{2}, \quad l = 1, 2, \dots, \quad (15)$$

since

$$\cos^{2l} \frac{\pi N}{2} + \cos^{2l} \frac{\pi(N-1)}{2} = I. \quad (16)$$

It then follows from (11) that

$$f(N) = \frac{\cos^l \frac{\pi N}{2}}{\sqrt{N+1}}, \quad \sigma_- = \frac{\cos^l \frac{\pi N}{2}}{\sqrt{N+1}}a. \quad (17)$$

When $l = 2$, with the use of the completeness relation of Fock state $|n\rangle$

$$\sum_{n=0}^{\infty} |n\rangle \langle n| = 1, \quad (18)$$

we see

$$\cos^2 \frac{\pi N}{2} = \frac{1}{2} (1 + \cos \pi N) = \frac{1}{2} \sum_{n=0}^{\infty} (1 + \cos \pi n) |n\rangle \langle n| = \sum_{n=0}^{\infty} |2n\rangle \langle 2n|, \quad (19)$$

and using $\langle n| a = \sqrt{n+1} \langle n+1|$,

$$\sigma_- = \sum_{n=0}^{\infty} |2n\rangle \langle 2n+1|, \quad (20)$$

which is just the Shigefumi's representation. For $l = 2k$, $k = 1, 2, 3 \dots$, we have

$$\cos^{2k} \frac{\pi N}{2} = \sum_{n=0}^{\infty} |2n\rangle \langle 2n|, \quad \sin^{2k} \frac{\pi N}{2} = \sum_{n=0}^{\infty} |2n+1\rangle \langle 2n+1|, \quad (21)$$

so the representation of σ_- is the same.

Let us recall the even- and odd- coherent state representations [5]

$$\begin{aligned} |z\rangle_e &= e^{-|z|^2/2} \sum_{n=0}^{\infty} \frac{z^{2n}}{\sqrt{2n!}} |2n\rangle, \\ |z\rangle_o &= e^{-|z|^2/2} \sum_{n=0}^{\infty} \frac{z^{2n+1}}{\sqrt{(2n+1)!}} |2n+1\rangle. \end{aligned} \quad (22)$$

so

$$\begin{aligned} \sum_{n=0}^{\infty} |2n\rangle \langle 2n| &= \cos^{2k} \frac{\pi N}{2} = \int \frac{d^2 z}{\pi} |z\rangle_{ee} \langle z|, \\ \sum_{n=0}^{\infty} |2n+1\rangle \langle 2n+1| &= \sin^{2k} \frac{\pi N}{2} = \int \frac{d^2 z}{\pi} |z\rangle_{oo} \langle z|. \end{aligned} \quad (23)$$

3 Another Bose representation

From (17) we see when $l = 1$,

$$\cos \frac{\pi N}{2} = \sum_{n=0}^{\infty} \cos \frac{\pi n}{2} |n\rangle \langle n| = \sum_{n=0}^{\infty} (-1)^n |2n\rangle \langle 2n|. \quad (24)$$

It then follows from (21) and (24) that when $l = 2k + 1$,

$$\cos^{2k+1} \frac{\pi N}{2} = \sum_{n=0}^{\infty} (-1)^n |2n\rangle \langle 2n| \sum_{n'=0}^{\infty} |2n'\rangle \langle 2n'| = \sum_{n=0}^{\infty} (-1)^n |2n\rangle \langle 2n|, \quad (25)$$

In this case

$$\sigma_- = \sum_{n=0}^{\infty} (-1)^n |2n\rangle \langle 2n+1|, \quad \sigma_+ = \sum_{n=0}^{\infty} (-1)^n |2n+1\rangle \langle 2n|, \quad l \text{ is odd}, \quad (26)$$

this is another Bose realization of Pauli spin operators. Due to

$$\begin{aligned} (-)^{N/2} |z\rangle_e &= e^{-|z|^2/2} \sum_{n=0}^{\infty} \frac{(-)^n z^{2n}}{\sqrt{2n!}} |2n\rangle = |iz\rangle_e, \\ (-)^{(N-1)/2} |z\rangle_o &= e^{-|z|^2/2} \sum_{n=0}^{\infty} \frac{(-)^n z^{2n+1}}{\sqrt{(2n+1)!}} |2n+1\rangle = |iz\rangle_o. \end{aligned} \quad (27)$$

so the corresponding coherent representation of (24) and (25) is

$$\begin{aligned} \cos^{2k+1} \frac{\pi N}{2} &= \int \frac{d^2 z}{\pi} |iz\rangle_{ee} \langle z|, \\ \sin^{2k+1} \frac{\pi N}{2} &= \int \frac{d^2 z}{\pi} |iz\rangle_{oo} \langle z|. \end{aligned} \quad (28)$$

For both l being even and odd,

$$\sigma_+ \sigma_- = \sum_{n=0}^{\infty} |2n+1\rangle \langle 2n+1|, \quad \sigma_- \sigma_+ = \sum_{n=0}^{\infty} |2n\rangle \langle 2n|, \quad (29)$$

from (9) we have

$$\begin{aligned} \sigma_3 &= [\sigma_+, \sigma_-] = \sum_{n=0}^{\infty} (|2n+1\rangle \langle 2n+1| - |2n\rangle \langle 2n|) \\ &= \sin^2 \frac{\pi N}{2} - \cos^2 \frac{\pi N}{2} = -\cos \pi N = -\frac{e^{i\pi N} + e^{-i\pi N}}{2} = (-1)^{N/2+1}. \end{aligned} \quad (30)$$

For the new representation we can check its fermion-like property. Using (28) and (30), we see

$$\{\sigma_-, \sigma_3\}_+ = \sum_{n=0}^{\infty} [(-1)^{3n+2} + (-1)^{3n+1}] |2n\rangle \langle 2n+1| = 0. \quad (31)$$

Its bosonic-like property can also be checked, i.e.,

$$\begin{aligned} [\sigma_-, \sigma_3] &= \sum_{n=0}^{\infty} [(-1)^{3n+2} - (-1)^{3n+1}] |2n\rangle \langle 2n+1| \\ &= 2 \sum_{n=0}^{\infty} (-1)^n |2n\rangle \langle 2n+1| = 2\sigma_-. \end{aligned} \quad (32)$$

4 The eigenvector of σ_- as a nonlinear coherent state

In Ref. [6, 7, 8, 9] the nonlinear coherent state is defined as eigenstate of $f(N)a$,

$$f(N)a|z\rangle_f = z|z\rangle_f. \quad (33)$$

From relation (12),

$$\left[f(N)a, \frac{1}{f(N-1)} a^\dagger \right] = 1, \quad (34)$$

the nonlinear coherent state $|z\rangle_f$ can be constructed directly as,

$$|z\rangle_f = \exp \left[\frac{z}{f(N-1)} a^\dagger \right] |0\rangle. \quad (35)$$

Since σ_- is of the form $f(N)a$ (see eq. (17)), the eigenvector of σ_- can be discussed in the context of nonlinear coherent states and be formally expressed as

$$|\xi\rangle_f = \exp \left[\frac{\sqrt{N}}{\cos^l \frac{\pi(N-1)}{2}} a^\dagger \xi \right] |0\rangle. \quad (36)$$

Thus

$$\sigma_- |\xi\rangle_f = \xi |\xi\rangle_f, \quad (37)$$

and because $\sigma_-^2 = 0$,

$$\sigma_- \left(\sigma_- |\xi\rangle_f \right) = \xi^2 |\xi\rangle_f = 0, \quad (38)$$

therefore $\xi^2 = 0$, so ξ must be a Grassmann number³. Ohnuki and Kashiwa have employed Grassmann number in constructing the fermionic coherent states.

In summary, using both the fermionic-like and the bosonic-like properties of the Pauli spin operators we have reviewed the Bose description of the Pauli spin operators, and derived another new representation. The eigenvector of σ_- in this representation is a nonlinear coherent state with the eigenvalues being the Grassmann numbers.

References

- [1] Shigefumi Naka, Prog. Theor. Phys., 1978, 59,: 2107,
- [2] Ruan Tu-nan, Jing Si-cong and Wang An-ming, Euro. Phys. Lett., 23 (1993): 317
- [3] Ohnuki Y, Kashiwa T., Prog.Theor. Phys., 1978, 60: 548
- [4] Louisell W. H., Quantum Statistical Properties of Radiation, New York: John Wiley, 1973
- [5] Hillery M., Phys. Rev., 1987, A36 (6): 3796T. M. Seligman, K. B. Wolf, Phys. Scr. 55 (1997) 528
- [6] Fan Hong-yi and Cheng Hai-ling, Phys. Lett. A, 2001, 285: 256
- [7] S. Sivakumar, J. Phys. A: Math. Gen. 29 (1996) 5637
- [8] S. Mancini, Phys. Lett. A 233 (1997) 291
- [9] V. I. Manko, G. Marmo, E. C. G. Sudarshan, F. Zaccaria, in : N. M. Atakishiev (Ed.), Proc. IV Wigner Symp. (Guadalajara, Mexico, July 1995), World Scientific, Singapore, 1996, p. 421