Least paradoxical states of the Schrödinger cat

Andrey Pereverzev*

Department of Chemistry, Trinity University, San Antonio, Texas 78212

(Dated: November 10, 2018)

Modeling the Schrödinger cat by a two state system and assuming that the cat is coupled to the environment we look for the least paradoxical states of the Schrödinger cat in the following way. We require the reduced density matrix of the cat for one of the two states in the superposition to be the same as the one for the total state while distinct from the reduced density matrix of the cat for the other state in the superposition. We then look for the reduced density matrices for which the cat is as alive as possible for the first state (and as dead as possible for the second state). The resulting states are those in which the probability for the cat to be alive (or dead) is $1/2 + \sqrt{2}/4 \approx 0.854$.

PACS numbers: 03.65.-w, 03.65.Ud, 03.65.Ta

In the thought experiment introduced by Schrödinger [1] a superposition of microscopically distinct orthogonal states leads to a superposition of macroscopically distinct orthogonal states of the type

$$|\chi\rangle = \frac{1}{\sqrt{2}}|\chi_1\rangle + \frac{1}{\sqrt{2}}|\chi_2\rangle,\tag{1}$$

where $|\chi_1\rangle$ describes the Universe in which the cat is alive, while $|\chi_2\rangle$ describes the Universe in which the cat is dead. Rather then discussing possible resolutions of the paradox (see, e.g. [2]) we would like to consider for what states the paradoxical nature of superposition (1) is least pronounced. Let us denote the reduced density matrices of the cat for states $|\chi\rangle$, $|\chi_1\rangle$, and $|\chi_2\rangle$ by, respectively, ρ , ρ_1 , and ρ_2 . The paradox would not exit if ρ_1 and ρ_2 corresponded, respectively, to pure states for alive and dead cat, and ρ was identical to either ρ_1 or ρ_2 . Clearly, this is not possible. However, since the cat can be entangled with the environment, we can relax the requirement that ρ_1 and ρ_2 correspond to pure states while keeping the requirement that ρ is the same as either ρ_1 or ρ_2 . This would imply that the cat has some probability of being dead for state ρ_1 and some probability of being alive for state ρ_2 . Then we can look for the density matrices ρ_1 and ρ_2 for which the cat is, respectively, as alive and as dead as possible. Our analysis show that the resulting states of the cat are those for which its probability to be alive (or dead) is $1/2 + \sqrt{2}/4 \approx 0.854$.

To formulate the problem more precisely we model the cat by a two state system with states $|1\rangle$ and $|2\rangle$ which correspond, respectively, to alive and dead states of the cat. We impose no restrictions on the form of the environment. Since the cat is modeled by the two state system one can visualize the problem by using the Bloch vector formalism [3]. Since we require that ρ is equal to, say, ρ_1 , the Bloch vectors associated with ρ and ρ_1 have to be the same. We also would like the reduced density matrices ρ_1 and ρ_2 to correspond to distinct states of the cat. However, once ρ_1 is given, there is no unique way to choose a distinct ρ_2 . One possible choice is based on the fact that in the Bloch picture two orthogonal pure states correspond to two vectors of unit length pointing in the opposite directions. Generalizing this to the density matrices we will require the vector for ρ_2 to be of the same length as for ρ_1 but pointing in the opposite direction. Let the Bloch vector for state $|1\rangle$ to lie in the positive z direction. Then our problem is to find vectors for ρ , ρ_1 , and ρ_2 which, in addition to the requirements imposed above, have the largest absolute value of the z component. It is clear from this picture that one necessary condition is that all three vectors are parallel to the z axis. To find the maximum possible length of the vectors and the corresponding probabilities it is convenient to write states $|\chi_1\rangle$ and $|\chi_2\rangle$ which appear in (1) as the Schmidt decompositions, i. e.,

$$\begin{aligned} |\chi_1\rangle &= \lambda |1\rangle |\psi_1\rangle + \sqrt{1 - \lambda^2} |2\rangle |\psi_2\rangle, \\ |\chi_2\rangle &= \sqrt{1 - \lambda^2} |1\rangle |\phi_1\rangle + \lambda |2\rangle |\phi_2\rangle. \end{aligned}$$
(2)

Here λ^2 is the probability that the cat is alive in state $|\chi_1\rangle$ and that it is dead in state $|\chi_2\rangle$. Since Eqs. (2) are the Schmidt decompositions, functions $|\psi_1\rangle$ and $|\psi_2\rangle$ are orthonormal as are functions $|\phi_1\rangle$ and $|\phi_2\rangle$. Our analysis now reduces to the search for states $|\psi_1\rangle$, $|\psi_2\rangle$, $|\phi_1\rangle$, and $|\phi_2\rangle$ which lead to the largest value for λ^2 and ensure that all the requirements imposed above are satisfied.

Since we require states $|\chi_1\rangle$ and $|\chi_2\rangle$ to be orthogonal to each other we must have

$$\langle \phi_1 | \psi_1 \rangle + \langle \phi_2 | \psi_2 \rangle = \langle \phi_1 | \psi_1 \rangle^* + \langle \phi_2 | \psi_2 \rangle^* = 0.$$
 (3)

Substituting Eqs. (2) into Eq. (1) we obtain for $|\chi\rangle$

$$\begin{aligned} |\chi\rangle &= |1\rangle \left(\frac{\lambda}{\sqrt{2}} |\psi_1\rangle + \sqrt{\frac{1-\lambda^2}{2}} |\phi_1\rangle\right) \\ &+ |2\rangle \left(\sqrt{\frac{1-\lambda^2}{2}} |\psi_2\rangle + \frac{\lambda}{\sqrt{2}} |\phi_2\rangle\right). \end{aligned} (4)$$

As we want the reduced density matrix obtained from state $|\chi\rangle$ to be the same as the one obtained from state $|\chi_1\rangle$, state $|\chi\rangle$ should have the Schmidt decomposition

$$|\chi\rangle = \lambda |1\rangle |\xi_1\rangle + \sqrt{1 - \lambda^2} |2\rangle |\xi_2\rangle.$$
 (5)

Comparing Eq. (4) and Eq. (5) we obtain for $|\xi_1\rangle$ and $|\xi_2\rangle$

$$\begin{aligned} |\xi_1\rangle &= \sqrt{\frac{1}{2}}|\psi_1\rangle + \sqrt{\frac{1-\lambda^2}{2\lambda^2}}|\phi_1\rangle, \\ |\xi_2\rangle &= \sqrt{\frac{1}{2}}|\psi_2\rangle + \frac{\lambda}{\sqrt{2(1-\lambda^2)}}|\phi_2\rangle. \end{aligned} \tag{6}$$

For Eq. (5) to be the required Schmidt decomposition functions $|\xi_1\rangle$ and $|\xi_2\rangle$ have to be orthonormal. It is sufficient to require that $|\xi_1\rangle$ and $|\xi_2\rangle$ are orthogonal and that one of them is normalized since conditions (3) will ensure that the other state is also normalized. Requiring that state $|\xi_1\rangle$ is normalized leads to the following equation for λ

$$2\lambda^2 - A\lambda\sqrt{1-\lambda^2} - 1 = 0, \tag{7}$$

where $A = \langle \psi_1 | \phi_1 \rangle + \langle \phi_1 | \psi_1 \rangle$. The requirement of orthogonality of $|\xi_1 \rangle$ and $|\xi_2 \rangle$ leads to

$$(1 - \lambda^2)\langle\phi_1|\psi_2\rangle + \lambda^2\langle\psi_1|\phi_2\rangle = (1 - \lambda^2)\langle\phi_1|\psi_2\rangle^* + \lambda^2\langle\psi_1|\phi_2\rangle^* = 0.$$
(8)

Solving Eq. (7) for λ and keeping only the positive solution (since the Schmidt decompositions allow only for positive expansion coefficients) we obtain

$$\lambda = \sqrt{\frac{1}{2} - \frac{A}{2\sqrt{4 + A^2}}}.$$
(9)

States $|\psi_1\rangle$ and $|\phi_1\rangle$ are normalized, as a result A can take values between -2 and 2. The maximum value of λ , λ_m is reached for A = -2. The corresponding probability λ_m^2 is given by

$$\lambda_m^2 = \frac{1}{2} + \frac{\sqrt{2}}{4} \approx 0.854.$$
 (10)

For A to be equal to -2 we must have $|\phi_1\rangle = -|\psi_1\rangle$. Condition (3) then requires that $|\phi_2\rangle = |\psi_2\rangle$. The last two equalities ensure that condition (8) is also satisfied.

Thus, states $|\chi\rangle$, $|\chi_1\rangle$, and $|\chi_2\rangle$ with the sought after properties will have the form

$$\begin{aligned} |\chi\rangle &= \lambda_m |1\rangle |\psi_1\rangle - \sqrt{1 - \lambda_m^2} |2\rangle |\psi_2\rangle, \\ |\chi_1\rangle &= \lambda_m |1\rangle |\psi_1\rangle + \sqrt{1 - \lambda_m^2} |2\rangle |\psi_2\rangle, \\ |\chi_2\rangle &= \sqrt{1 - \lambda_m^2} |1\rangle |\psi_1\rangle - \lambda_m |2\rangle |\psi_2\rangle, \end{aligned}$$
(11)

where $|\psi_1\rangle$ and $|\psi_2\rangle$ are two arbitrary orthonormal states of the environment and λ_m is obtained from (10). To sum up, the reduced density matrices of the cat for states $|\chi\rangle$ and $|\chi_1\rangle$ are identical and diagonal in the basis of states $|1\rangle$ and $|2\rangle$, the probability for the cat to be alive is about 0.854. The reduced density matrix of the cat for state $|\chi_2\rangle$ is also diagonal in the same basis, the probability for the cat to be dead is about 0.854. It is easy to see that the limit λ_m appears already on the level of an isolated two state system. Consider the following orthonormal states for two state system

$$|I\rangle = \lambda_m |1\rangle + \sqrt{1 - \lambda_m^2} |2\rangle,$$

$$|II\rangle = \sqrt{1 - \lambda_m^2} |1\rangle - \lambda_m |2\rangle,$$
 (12)

and their superposition

$$|III\rangle = \frac{1}{\sqrt{2}}|I\rangle + \frac{1}{\sqrt{2}}|II\rangle = \lambda_m|1\rangle| - \sqrt{1 - \lambda_m^2}|2\rangle.$$
(13)

Note that states $|I\rangle$, $|II\rangle$, and $|III\rangle$ have the same probabilities for the cat to be alive as, respectively, states $|\chi_1\rangle$, $|\chi_2\rangle$, and $|\chi\rangle$. If we use the Bloch vector formalism and chose the vector for state $|1\rangle$ to lie along the positive z axis, then state $|I\rangle$ corresponds to vector \mathbf{P}_{I} at 45 degrees to the z axis. The Bloch vector \mathbf{P}_{II} for state $|II\rangle$ is of the same length as \mathbf{P}_I but pointing in the opposite direction, while vector \mathbf{P}_{III} for state $|III\rangle$ is at 45 degrees to the z axis and perpendicular to both \mathbf{P}_{I} and \mathbf{P}_{II} . It is clear from this picture that vectors \mathbf{P}_{I} and \mathbf{P}_{II} will have z components equal to $\sqrt{2}/2$ with the corresponding probability for the cat to be alive given by λ_m^2 while the z component for vector \mathbf{P}_{III} will be $-\sqrt{2}/2$. By using this geometrical picture one can convince oneself that this arrangement of vectors insures the largest absolute value of the z component for the three vectors. All three vectors, however, will have non zero and different x and y components. Thus, coupling the cat to the environment in states (11) cannot increase the absolute value of the z components and the corresponding probabilities. However, it allows to remove the x and y components of the Bloch vectors for all three states making the reduced density matrices obtained from states $|\chi\rangle$ and $|\chi_1\rangle$ identical. It can be argued that in the resulting states (11)the fate of the cat is less dramatic than in the original formulation of the paradox.

The author would like to thank Dr. Gonzalo Ordóñez for useful discussions. Part of this research was supported by grants from the Robert A. Welch Foundation (Grant No. W-1442) and the Petroleum Research Fund, administered by the American Chemical Society.

- [2] A. J. Leggett, J. Phys. -Condens. Mat. 14, R415 (2002)
- [3] R. P. Feynman, F. L. Vernon Jr., and R. W. Hellwarth, J. Appl. Phys. 28, 49 (1957)

^{*} Electronic address: andrey.pereverzev@trinity.edu

E. Schrödinger, Naturwiss. 23, 807 (1935); English translation in J. A. Wheeler and W. H. Zurek, eds., *Quantum Theory and Measurement* (Princeton University Press, Princeton, N.J., 1983).