Generalized Partitioned Quantum Cellular Automata and Quantization of Classical CA

Shuichi INOKUCHI^{*} Yoshihiro MIZOGUCHI[†]

November 11, 2018

Abstract

In this paper, in order to investigate natural transformations from discrete CA to QCA, we introduce a new formulation of finite cyclic QCA and generalized notion of partitioned QCA. According to the formulations, we demonstrate the condition of local transition functions, which induce a global transition of well-formed QCA. Following the results, extending a natural correspondence of classical cells and quantum cells to the correspondence of classical CA and QCA, we have the condition of classical CA such that CA generated by quantization of its cells is well-formed QCA. Finally we report some results of computer simulations of quantization of classical CA.

1 Introduction

J. Watrous introduced the notion of quantum cellular automata(QCA) and showed that any quantum Turing machine can be efficiently simulated by a QCA with constant slowdown in 1995.

CA with quantum cells is well-formed QCA if and only if its global transition function is unitary. Generally quantization of cells of a classical CA dose not make it become QCA, because usually classical CA dose not have reversibility. Morita and Harao show that we can get reversible CA by partition a cell into three part and partitioned CA(PCA) can simulate non-partitioned

^{*}Faculty of Mathematics, Kyushu University, E-mail:inokuchi@math.kyushu-u.ac.jp [†]Faculty of Mathematics, Kyushu University, E-mail:ym@math.kyushu-u.ac.jp

CA(NPCA)[10]. But there is not a trivial inclusion relation between PCA and NPCA.

In this paper, we introduce a new formulation of finite cyclic QCA and generalized notion of partitioned QCA in order to investigate natural transformations from discrete CA to QCA. According to the formulations, we demonstrate a condition of a local transition function, which induce a wellformed QCA. A natural correspondence of classical cells and quantum cells can be extended to the correspondence of classical CA and QCA. If a classical CA satisfies our conditions then the extended QCA is well-formed. Finally we report some results of computer simulations of quantization of classical CA.

2 Preliminaries

Let Q be a set of states of cells and |Q| = s. We consider Q^n is the set of configurations of CA where n is the size of CA. q_i denotes the *i*th element of the configuration $q \in Q^n$, and assume that $q_0 = q_n$ and $q_{n+1} = q_1$.

Before considering computing process of quantum states, we recall that of deterministic states. We use the set of all subset of Q, that is, the set 2^{Q} of all functions from Q to $2 = \{0, 1\}$ to represent nondeterministic states.

A element q of Q is normally considered as an element $\{q\}$ of 2^Q . Let Σ be the finite set of input characters, then a transition function of input characters of deterministic finite automata is provided as $\delta : Q \times \Sigma \to Q$, and expanded naturally into the transition function $\delta_* : Q \times \Sigma^* \to Q$ of input strings. Let $[\delta] : Q \times \Sigma \to 2^Q$ be a function defined by $[\delta](q, \alpha) = [\delta(q, \alpha)]$.

The function $[\delta]$ is a state transition function of nondeterministic finite automata and we can expand $[\delta]$ into $[\delta]_* : Q \times \Sigma^* \to 2^Q$. Recall $[\delta_*] : Q \times \Sigma^* \to 2^Q$ be a function defined by $[\delta_*](q, \alpha) = [\delta_*(q, \alpha)]$, and we can show that $[\delta_*] = [\delta]_*$ easily. This shows that the set of all deterministic finite automata are included in the set of all nondeterministic finite automata naturally.

We can consider quantum states as generalized states of classical state and extend to a quantum formulation of computer system. But a classical computer system is not always a quantum computer system generally, because a quantum computing process should be a unitary operator and every classical computing process is not so.

A quantum state denotes a function from a finite set Q to a set of complex

numbers \mathbf{C} and \mathbf{C}^Q is denoted by the set of all functions from Q to \mathbf{C} . For $g \in Q$ we define $[q] \in \mathbf{C}^Q$ as follows;

$$[q](x) = \begin{cases} 1 & (x=q) \\ 0 & (x \neq q) \end{cases}$$

 \mathbf{C}^{Q} is a linear space on \mathbf{C} such that its bases is Q. An inner product in \mathbf{C}^{Q} is defined by $\langle p, q \rangle = \sum_{x \in Q} (p(x) \cdot q(x))$ where $p, q \in \mathbf{C}^{Q}$. A linear space $\mathbf{C}^{(Q^{n})}$ on \mathbf{C} is considered as a tensor product $\underbrace{\mathbf{C}^{Q} \otimes \mathbf{C}^{Q} \otimes \cdots \otimes \mathbf{C}^{Q}}_{n}$,

that is, for $p \in Q^n$ $[p] = [p_1] \otimes [p_2] \otimes \cdots \otimes [p_n]$. For $p, q \in Q^n$ the inner product $\langle [p], [q] \rangle$ on $\mathbf{C}^{(Q^n)}$ is as follows;

$$\begin{array}{l} \langle [p], [q] \rangle \\ = & \sum_{x \in Q^n} ([p](x) \cdot [q](x)) \\ = & \sum_{(x_1, x_2, \dots, x_n) \in Q^n} (([p_1](x_1)[p_2](x_2) \cdots [p_n](x_n))) \\ & \cdot ([q_1](x_1)[q_2](x_2) \cdots [q_n](x_n))) \\ = & \sum_{x_1 \in Q} ([p_1](x_1)[q_2](x_1)) \cdot \sum_{x_2 \in Q} ([p_2](x_2)[q_2](x_2)) \\ & \cdots \sum_{x_n \in Q} ([p_n](x_n)[q_n](x_n)) \\ = & \langle [p_1], [q_1] \rangle \langle [p_2], [q_2] \rangle \cdots \langle [p_n], [q_n] \rangle. \end{array}$$

For a function $F: Q^n \to \mathbf{C}^{(Q^n)}$, we define functions $\alpha_F: Q^n \times Q^n \to \mathbf{C}$ and $\overline{F}: \mathbf{C}^{(Q^n)} \to \mathbf{C}^{(Q^n)}$ by $\alpha_F(p,q) = F(p)(q)$ and $\overline{F}(X) = \sum_{q \in Q^n} (X(q)(F(q)))$.

We call \overline{F} is unitary if $||\overline{F}(X)|| = 1$ for any $X \in \mathbf{C}^{(Q^n)}$ such that ||X|| = 1.

Since Q is finite, we can label elements of Q numbers from 1 to s and also elements of Q^n numbers from 1 to s^n by lexicographical ordering. We define a $s^n \times s^n$ matrix (α_{ij}) by $\alpha_{ij} = \alpha_F(p,q)$ where numbers of elements p and q are *i* and j.

Proposition 1 If the matrix $(\alpha_{i,j})$ is unitary then \overline{F} is unitary.

Proof. Assume that (α_{ij}) is a unitary matrix and $\langle X, X \rangle = 1$.

$$\begin{split} &\langle \overline{F}(X), \overline{F}(X) \rangle \\ = &\langle \sum_{p \in Q^n} X(p) \overline{F}(p), \sum_{q \in Q^n} X(q) \overline{F}(q) \rangle \\ = &\sum_{p \in Q^n} \sum_{q \in Q^n} \langle X(p) \overline{F}(p), X(q) \overline{F}(q) \rangle \\ = &\sum_{p \in Q^n} (X(p) \sum_{q \in Q^n} (X(q) \sum_{r \in Q^n} (\overline{F}(p)(r) \cdot \overline{F}(q)(r)))) \\ = &\sum_{p \in Q^n} (X(p) \sum_{q \in Q^n} (X(q) \sum_{r \in Q^n} (\alpha(p, r) \cdot \alpha(q, r)))) \\ = &\sum_{p \in Q^n} (X(p) \sum_{q \in Q^n} (X(q) \sum_{r \in Q^n} (\alpha(p, r) \cdot \overline{\alpha(r, q)}))) \\ = &\sum_{p \in Q^n} (X(p) \cdot X(p)) \\ = &1 \end{split}$$

So \overline{F} is unitary.

Proposition 2 Let $\hat{\sigma} : \mathbb{C}^{Q^n} \to \mathbb{C}^{Q^n}$ be defined by $\hat{\sigma}(X)(p) = X(\sigma(p))$ where $\sigma : Q^n \to Q^n$. Then the followings are equivalent.

- $\hat{\sigma} = \overline{[\sigma]}$ and $\hat{\sigma}$ are unitary.
- σ is a bijection.

A classical CA is a transition system in Q defined by a global transition function $F: Q^n \to Q^n$ where $F(q)_i = f(q_{i-1}, q_i, q_{i+1})$ and $f: Q \times Q \times Q \to Q$ is a local transition function.

When $Q = \{0, 1\}$, a local transition function is defined by the eight values $f(0,0,0) = r_0$, $f(0,0,1) = r_1$, $f(0,1,0) = r_2$, $f(0,1,1) = r_3$, $f(1,0,0) = r_4$, $f(1,0,1) = r_5$, $f(1,1,0) = r_6$ and $f(1,1,1) = r_7$ ($r_i = 0,1$). The rule number R of a local transition function f is defined by

$$R = 2^{7}r_{7} + 2^{6}r_{6} + 2^{5}r_{5} + 2^{4}r_{4} + 2^{3}r_{3} + 2^{2}r_{2} + 2^{1}r_{1} + r_{0}.$$

The local transition function of rule number R is denoted by f_R . The local transition rules with rule number 204,240 and 170 are illustrated as follows;

	111	110	101	100	011	010	001	000
204	1	1	0	0	1	1	0	0
	111	110	101	100	011	010	001	000
240	1	1	1	1	0	0	0	0
	111	110	101	100	011	010	001	000

 f_{204} , f_{240} and f_{170} are identity, shift-right and shift-left functions respectively.

3 Quantum Cellular Automata

The global transition function $F : Q^n \to \mathbf{C}^{(Q^n)}$ is defined by $F(q)(x) = f(q_0, q_1, q_2)x_1 \cdot f(q_1, q_2, q_3)x_2 \cdots f(q_{n-1}, q_n, q_{n+1})x_n$. For a local transition function $f : Q \times Q \times Q \to \mathbf{C}^Q$, A transition system in $\mathbf{C}^{(Q^n)}$ is defined by $\overline{F} : \mathbf{C}^{(Q^n)} \to \mathbf{C}^{(Q^n)}$. The local transition function f is called 'forming a quantum cellular automaton' if \overline{F} is unitary.

Let $F_f : Q^n \to Q^n$ be a transition function of a local transition function $f : Q^3 \to Q$, and $[F_f] : Q^n \to \mathbf{C}^{Q^n}$ a function such that $[F_f](x) = F_f(x)$ for $x \in Q^n$. And we let $[f] : Q^3 \to \mathbf{C}^Q$ be a local transition function of QCA such that [f](x) = f(x) for $x \in Q$, and $F_{[f]} : Q^n \to \mathbf{C}^{(Q^n)}$ its global transition function. Then we can show that $[F_f] = F_{[f]}$ by easy computation. But a local transition functions $f : Q^3 \to Q$ does not always form a QCA, that is, a $\overline{[F_f]}$ is not always unitary.

Proposition 3 [f] is forming a quantum cellular automata if and only if $F_f: Q^n \to Q^n$ is a bijection.

4 Partitioned Quantum Cellular Automata


We define functions $G: Q^n \to \mathbf{C}^{(Q^n)}$ and $\lambda: Q \times Q \to \mathbf{C}$ by $G(q)(x) = g(q_1)x_1 \cdot g(q_2)x_2 \cdots g(q_n)x_n$ and $\lambda(p,q) = g(p)(q)$ for a function $g: Q \to C^Q$.

We label the elements of Q^n numbers from 1 to s^n and define a $s^n \times s^n$ matrix (α_{ij}) from $\alpha_G : Q^n \times Q^n \to \mathbf{C}$. And we label the elements of Qnumbers from 1 to s and define a $s \times s$ matrix (λ_{ij}) from the function λ .

Proposition 4 (λ_{ij}) is a unitary matrix if and only if (α_{ij}) is a unitary matrix.

Proposition 5 If $\sigma : Q^n \to Q^n$ is a bijection, then the followings hold;

- (i) $\overline{G \circ \sigma} = \overline{G} \circ \hat{\sigma}$.
- (ii) $\overline{G \circ \sigma}$ is unitary if and only if \overline{G} is unitary.



Theorem 6 The composition function $f = g \circ e : Q^3 \to \mathbb{C}^Q$ of functions $e : Q^3 \to Q$ and $g : Q \to \mathbb{C}^Q$ is forming a quantum cellular automaton if both of the following two conditions hold:

(i) $F_e: Q^n \to Q^n$ is a bijection.

(ii) The matrix (λ_{ij}) defined from $g: Q \to \mathbf{C}^Q$ is unitary.

Example 7 Let $Q = L \times M \times R$ for finite sets L, M and R. We define $e: Q^3 \to Q$ and $g: Q \to \mathbb{C}^Q$ by $e(((l_1, m_1, r_1), (l_2, m_2, r_2), (l_3, m_3, r_3))) = (l_3, m_2, r_1)$ and g(q) = [q]. Then the composition function $f = g \circ e$ is forming a quantum cellular automaton. Because $F_e: Q^n \to Q^n$ is a bijection, $F_e(((l_i, m_i, r_i)))_j = (l_{j+1}, m_j, r_{j-1})$, and (λ_{ij}) defined from g is an identity matrix.

In example 7, we can replace g to another function $g: Q \to \mathbb{C}^Q$ where the matrix (λ_{ij}) defined from g is unitary. On that occasion $f: (L \times M \times R)^3 \to (L \times M \times R)$ is also forming a quantum cellular automaton. Consequently a partitioned quantum cellular automaton introduced in [13] is demonstrated as a special case of our general formulation.

Example 8 Let $Q = \{0, 1\}$, $e : Q^3 \to Q$ be a function such that $F_e : Q^n \to Q^n$ is a bijection, and $\Lambda = (\lambda_{ij})$ defined from $g : Q \to \mathbb{C}^Q$ be as follows;

$$\Lambda = \left(\begin{array}{cc} \cos\theta & -\sin\theta\\ \sin\theta & \cos\theta \end{array} \right).$$

Then the local transition function defined from $f = g \circ e$ forms a quantum cellular automaton. This shows a quantum cellular automaton formed by a

synthesised function of a local transition function e and its reverse function. That is, if $\theta = 0$ then f = e and if $\theta = \frac{\pi}{2}$ then f is the reversed function of e, so we can consider f for $0 < \theta < \frac{\pi}{2}$ as a synthesised function of two classical local functions.

Example 9 Let $Q = \{0, 1\} \times \{0, 1\}$, $e((a_1, b_1), (a_2, b_2), (a_3, b_3)) = (a_1, b_3)$, and $\Lambda = (\lambda_{ij})$ defined by $g : Q \to \mathbb{C}^Q$ be as follows;

$$\Lambda = \left(\begin{array}{rrrr} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{array}\right)$$

Then $f = g \circ e$ forms a QCA and $f((a_1, b_1), (a_2, b_2), (a_3, b_3)) = (a_1, a_1 \oplus b_3).$

5 Computer Analysis

The following is a table of the size n of CA and rule number R where the local transition function $f_R: Q^3 \to Q$ forms QCA ($Q = \{0, 1\}, 3 \le n \le 22$ and $128 \le R \le 255$). We note that if f_R forms QCA then f_{255-R} also forms QCA.

Size n	Rule number R
3	142, 154, 156, 166, 170, 172, 178, 180, 184
	198,202,204,210,212,216,226,228,240
4, 8, 10, 14	150, 170, 204, 240
16, 20, 22	
5, 7, 11, 13, 19	150, 154, 166, 170, 180, 204, 210, 240
9, 15, 21	154, 166, 170, 180, 204, 210, 240
6, 12, 18	170, 204, 240

In the case of size 6, 12 and 18, trivial transitions, that is, identity, right shift and left shift functions only form QCA. In other case, there is a nontrivial transition function forming QCA.

From the above table, we can guess the following table of sizes of CA and rule numbers of local transition functions forming QCA, but we have not proved it yet.

Size $(k \ge 1)$	Rule number
$6k \pm 2$	150, 170, 204, 240
$6k \pm 1$	150, 154, 166, 170, 180, 204, 210, 240
6k + 3	154, 166, 170, 180, 204, 210, 240
6k	170, 204, 240

Example 10 When the size of CA is 4 or 5, the local transition function $f_{150}(x, y, z) = x + y + z \pmod{2}$ forms a quantum CA. Let F_f be the global transition function, then the following hold;

$$F_{f}(x)_{i} = x_{i-1} + x_{i} + x_{i+1}$$

$$F_{f}^{2}(x)_{i} = x_{i-2} + x_{i-1} + x_{i}$$

$$+ x_{i-1} + x_{i} + x_{i+1}$$

$$+ x_{i} + x_{i+1} + x_{i+2}$$

$$= x_{i-2} + x_{i} + x_{i+2}$$

$$F_{f}^{3}(x)_{i} = x_{i-3} + x_{i-2} + x_{i-1}$$

$$+ x_{i-1} + x_{i} + x_{i+1}$$

$$+ x_{i+1} + x_{i+2} + x_{i+3}$$

$$= x_{i-3} + x_{i-2} + x_{i} + x_{i+2} + x_{i+3}$$

In the case that the size of CA is 4, $F_f^2(x)_i = x_{i+2} + x_i + x_{i+2} = x_i$. So $F_f^2(x) = x$. In the case that the size of CA is 5, $F_f^3(x)_i = x_{i+2} + x_{i+3} + x_i + x_{i+2} + x_{i+3} = x_i$. So $F_f^3(x) = x$. Namely there exists y such that $F_f(y) = x$ for any $x \in Q^n$ (n = 4, 5), and F_f is a bijection. So f forms a QCA.

6 Related Works and Conclusion

Cellular automata dealt in this paper is finite cyclic CA and different from CA without boundary, that is, infinite CA dealt by Watrous[13], Morita and Harao[10]. Because the size of CA is finite it does not have the universal computability[11, 1, 2]. But the conditions of local transition functions forming QCA is formulated clearly in our framework.

Injectivity of global maps of classical CA is an essential property for extending to a QCA. The injectivity are considered in [7, 8, 9, 11] for classical CA without boundary and in [5, 3, 6] for classical finite cyclic CA.

A further direction of this study will be to consider properties on construction and synthesis of general quantum computer system by examining construction and synthesis of local transition function of realizable QCA.

References

- [1] A. Birthiaume and G. Brassard: *The Quantum Challenge to Structural Complexity Theory*, Proc. 7th IEEE Conf. Struct. Comp. Theo., 1992.
- [2] D. Deutsch: Quantum Theory, The Church Turing Principle and The Universal Quantum Computer, Proc. the Royal Society of London A400, pp. 97–117, 1985.
- [3] R.A. Dow: Additive Cellular Automata and Global Injectivity, Physica D, 110, pp. 67–91, 1997.
- [4] J. Gruska: Quantum Computing, McGraw-Hill, 1999.
- [5] M. Harao and S. Noguchi: On the Algebraic Properties of the Iterative Automaton, Trans. IEICE, J55-D, pp.767–774, 1972. [Japanese]
- [6] K. Ichikawa, H. Aso, Y. Inagaki and N. Honda: Dynamical Chracteristics of Linear Circular Cellular Automata, Trans. IEICE, J65-D, pp.1129– 1136, 1982. [Japanese]
- [7] M. Ito, N. Osato and M. Nasu: *Linear Cellular Automata over* Z_m , J. Computer and System Sciences, 27, pp.125–140, 1983.
- [8] A. Maruoka and M. Kimura: Condition for Injectivity of Global Maps for Tesselation Automata, Information and Control, 32, pp.158–162, 1976.
- [9] A. Maruoka and M. Kimura: Injectivity and Surjectivity of Parallel Maps for Cellular Automata, J. Computer and System Sciences, 18, pp.47–64, 1979.
- [10] K. Morita and M. Harao: Computation Universality of One-Dimensional Reversible (Injective) Cellular Automata, Trans. IEICE, E72, pp. 758– 762, 1989.
- [11] T. Toffoli: Computation and Construction Universality of Reversible Cellular Automata, J. Comput. System Sci., 15, pp. 213–231, 1977.
- [12] T. Toffoli and H. Margolus: Invertible Cellular Automata: A Review, Physica D, 45, pp.229–253, 1990.

[13] J. Watrous: On One-Dimensional Quantum Cellular Automata, Proceedings of the 36th Annual Symposium on Foundations of Computer Science, pp. 528–537, 1995.