

Nonlocal Hamiltonians and Information Conservation Law

Jian-Ming Cai, Xiang-Fa Zhou, Zheng-Wei Zhou,* and Guang-Can Guo

*Key Laboratory of Quantum Information,
University of Science and Technology of China (CAS),
Hefei 230026, People's Republic of China.*

Abstract

We use fidelity to characterize the information about a quantum state contained in another quantum state. In a multi-qubit system, each qubit has individual information about a prescribed quantum state. We investigate how the total information changes under the interactions of nonlocal Hamiltonians. We show that for some special physical interactions the total information is conserved. This type of nonlocal Hamiltonians are named as *Conservative Hamiltonians*, similar to the Conservative Forces. We then apply this Information Conservation Law in the quantum spin system, and propose methods for improving the efficiency of quantum communications through an open end spin chain.

PACS numbers: 03.67.Pp, 03.65.Ud

*Electronic address: zwzhou@ustc.edu.cn

Quantum computation (QC) has demonstrated prominent speed-up potential compared to classical computation in several situations [1]. Grover's disordered database search algorithm [2, 3] and Shor's algorithm [4] for prime decomposition of large numbers are the well-known representations. This has stimulated a lot of research aimed at the implementation of a quantum computer in a suitable physical system. One of the most important requirements for such a physical system is scalability. In other words, it should be composed of many qubits and interactions exist between different qubits. In such a scalable multi-qubit system, nonlocal interactions are the central issues not only from the theoretical point of view but also from the experimental one [5].

In recent years, substantial work has shown that nonlocal interactions are necessary and sufficient for the implementation of universal quantum computation. As is well known that an arbitrary unitary transformation can be implemented by a sequence of single-qubit gates and controlled-NOT (CNOT) gates [6, 7], which is a kind of two-qubit gate. In order to perform a CNOT gate, bipartite interactions must be employed. R. Raussendorf et al [8] propose a new kind of quantum computation model, based on Ising-type next-neighbor interactions. Besides, nonlocal interactions can be used for short distance quantum communication, which may be useful in a scalable quantum computer architecture. It is much different from quantum teleportation. S. Bose [9] showed that an unknown quantum state can be transmitted through an open end chain with always-on nearest-neighbor interactions by locating the input and output quantum state at the two opposite ends of the chain. The efficiency exceeds $2/3$, which is the efficiency of classical transmission of a quantum state, even when the length of the chain is as high as 80. Very recently, Christandl et al [10] put forward a modified scheme that can improve the efficiency to 1 in quantum spin networks. Both spin chains and spin networks can be considered as multi-qubit systems with nonlocal interactions.

In this Letter, we study the interactions in a multi-qubit system from a new viewpoint. It is well known that *fidelity* is a fundamental concept in quantum computation and quantum information. It indicates that how much a quantum state is different from the *desired* quantum state, which is named as *standard quantum state*, e.g. the state to be cloned or the state to be transmitted through quantum channels. If the fidelity is larger, the information about the desired quantum state, contained in another quantum state, is more. For a multi-qubit system governed by some nonlocal system Hamiltonian, every individual

qubit has some information about a prescribed standard quantum state respectively. Our interest focus on how the fidelity of each qubit and the total fidelity of all qubits evolve under the interactions of different parts of this system. We find that the total fidelity maybe conserved, while the individual fidelity changes with time. Our work reflects the principle of conservation of quantum information proposed and investigated in [11, 12, 13, 14, 15, 16] from another point of view.

First we consider two qubits with the interaction described by a Hamiltonian in the canonical form [17]:

$$\mathbf{H}_I^{12} = c_1 \sigma_x^1 \otimes \sigma_x^2 + c_2 \sigma_y^1 \otimes \sigma_y^2 + c_3 \sigma_z^1 \otimes \sigma_z^2 \quad (1)$$

where $\pi/4 \geq c_1 \geq c_2 \geq |c_3|$. The prescribed standard quantum state is denoted by the density matrix $(\rho^0)_{2 \times 2} = |\phi\rangle\langle\phi|$, where $|\phi\rangle = \cos(\theta/2)|0\rangle + e^{i\varphi} \sin(\theta/2)|1\rangle$. And the initial state of the two qubits is $[\rho(0)]_{4 \times 4}$. The information about the standard quantum state encoded in the i th qubit ($i = 1, 2$) of this two-qubit system at time t is characterized by the fidelity $F_i(t) = \langle\phi|\rho^i(t)|\phi\rangle$. The reduced density matrix $\rho^i(t) = Tr_{\tilde{i}}\rho(t)$ represents the state of the i th qubit, where \tilde{i} indicates a partial trace over the other qubit except the i th. For the sake of discussion later, we define two quantities:

$$\begin{aligned} CF_i(t) &= \cos^2(\theta/2)\rho_{00}^i(t) + \sin^2(\theta/2)\rho_{11}^i(t) \\ QF_i(t) &= Re[e^{-i\varphi} \sin\theta\rho_{10}^i(t)] \end{aligned} \quad (2)$$

It can be seen that $CF_i(t)$ is related to the probability character of quantum state, and $QF_i(t)$ is determined by the non-diagonal terms of the density matrix which is relevant to quantum coherence. So we can write the fidelity in a simple form as follows:

$$F_i(t) = CF_i(t) + QF_i(t) \quad (3)$$

And the total fidelity is $F(t) = \sum_{i=1}^2 F_i(t) = \sum_{i=1}^2 [CF_i(t) + QF_i(t)]$.

After a specific time interval t , the initial state $\rho(0)$ evolves to a final state $\rho(t) = e^{-i\mathbf{H}_I^{12}t}\rho(0)e^{i\mathbf{H}_I^{12}t}$. We find that the individual fidelity $F_i(t)$ surges with time periodically. The question we are interested in is that dose the total fidelity $F(t)$ changes too? By straightforward calculation, we find that if $c_1 = c_2$, then $\sum_{i=1}^2 CF_i(t)$ is invariable. In this situation $[\mathbf{H}_I^{12}, \sum_{i=1}^2 \sigma_z^i] = 0$. But $\sum_{i=1}^2 QF_i(t)$ changes with time under this condition. So, if we hope that $F(t)$ is invariable too, the Hamiltonian \mathbf{H}_I^{12} must satisfy a more strict

condition. Substitute $\rho(t)$ into the expression of $\sum_{i=1}^2 QF_i(t)$, we can see that if and only if $c_1 = c_2 = c_3$ then $F(t)$ is conserved during the evolution of this two-qubit system.

In the following, we will extend the above conclusions to a multi-qubit system. Suppose that the system consists of n qubits. The nonlocal system Hamiltonian is H_I . The prescribed standard quantum state is still denoted by the density matrix $(\rho^0)_{2 \times 2} = |\phi\rangle\langle\phi|$. And the initial state of the n qubits now is $[\rho(0)]_{2^n \times 2^n}$. Again we will try to find for what kind of nonlocal Hamiltonians $F(t)$ is an invariant. The following theorem give the answer of this problem.

Theorem Information Conservation Law

The total fidelity $F(t)$ of a n -qubit system is invariable for any initial system state if and only if $[H_I, \sum_{i=1}^n C_i] = 0$, where $C_i = \frac{1}{2^{n-1}} I_1 \otimes I_2 \otimes \cdots \otimes \rho_i^0 \otimes I_{i+1} \otimes \cdots \otimes I_n$ denotes that the reduced density matrix of the i th qubit is ρ^0 and the reduced density matrix of other qubits is $\frac{I}{2}$.

Proof The total fidelity $F(t) = \sum_{i=1}^n F_i(t)$. So after the evolution of the multi-qubit system:

$$F(t) = \sum_{i=1}^n Tr \left\{ Tr_{\bar{i}} [\exp(-iH_I t) \rho(0) \exp(iH_I t)] \rho_i^0 \right\} \quad (4)$$

We can write the derivative of $F(t)$ straightforwardly:

$$\frac{\partial F(t)}{\partial t} = i Tr \left\{ \left[H_I, \sum_{i=1}^n C_i \right] \exp(-iH_I t) \rho(0) \exp(iH_I t) \right\} \quad (5)$$

Thus $\frac{\partial F(t)}{\partial t} = 0 \Leftrightarrow [H_I, \sum_{i=1}^n C_i] = 0$

At time $t = 0$, each qubit in the multi-qubit system contain certain amount information about the standard quantum state. Due to the interactions between different parts of this system, the states of these qubits change with time t , and the information is exchanged between them. Thus the fidelity between each individual qubit and the standard quantum state is different at different time t . However, if the nonlocal system Hamiltonian H_I satisfied the condition stated in the above theorem, then the total fidelity of all qubits is an invariant. The interactions only changes the information distributing among the qubits of this multi-qubit system, while the total information is conserved. We name this type of nonlocal Hamiltonians *Conservative Hamiltonians*. This is something like Energy Conservation Law. The interactions between different parts of a closed system make energy exchange between them. But the total energy of this closed system is conserved.

Information Conservation Law can be useful in many situations relevant to fidelity in quantum computation and quantum information. For example, we can understand the fundamental problem: no cloning theorem [18] from a new perspective. The pure state to be cloned $|\phi\rangle$ can be looked on as the standard quantum state. All the qubits involved make up of a multi-qubit system. And the clone procedure can be characterized by some nonlocal Hamiltonian H . So if H belongs to *Conservative Hamiltonians*, then it is easy to obtain the upper bounds of this type cloning from *Information Conservation Law*. Another meaningful scenario is quantum communications in which the source, channel and destination form a closed system governed by an inner Hamiltonian. If this Hamiltonian is conservative, then the total information contained in the source, channel and destination is an constant according to *Information Conservation Law*. So if we want to improve the capacity of this channel, we can try to diminish the information contained in the source and channel. We will discuss a particular example in the following.

Furthermore, *Information Conservation Law* leads to an interesting phenomenon. Both two qubits are not the same with the standard quantum state at time $t = 0$. They contain some but not all the information about the standard quantum state, $F_1(0) = 0.737$ and $F_2(0) = 0.806$. Then information is exchanged between them without changing the total information. It can be seen that at some time t_0 , $F_1(t_0) > \max\{F_1(0), F_2(0)\}$. So we can say that information is partially concentrated spontaneously in the 1st qubit at time t_0 . And how much the information can be concentrated is determined by the initial state. Of course, this phenomenon can also exist in a multi-qubit system. So we can expect to obtain a more perfect state, with a higher fidelity, from some unperfect states in comparison to the standard quantum state. This is somewhat similar to Entanglement Concentration. We name this phenomenon as *Information Concentration*. If we choose the initial state as $|\psi\rangle = |\phi\rangle \otimes |0\rangle$, then an inverse process will happen, that is *Information Dilution*. At time $t = 0$, the 1st qubit has all the information about the standard quantum state, i.e. $F_1(0) = 1$. For time $t > 0$, the information is diluted into these two qubits, which means $F_1(t) \leq F_1(0)$.

The general form of nonlocal Hamiltonian with pairwise interactions can be expressed as

follows:

$$\mathbf{H}_I = \sum_{(i,j)} c_x^{ij} \sigma_x^i \otimes \sigma_x^j + c_y^{ij} \sigma_y^i \otimes \sigma_y^j + c_z^{ij} \sigma_z^i \otimes \sigma_z^j \quad (6)$$

where (i, j) represents pairs of qubits with interaction.

Corollary The total fidelity $F(t)$ of a n -qubit system with pairwise interactions is invariable for any initial state and any prescribed standard quantum state if and only if the nonlocal system Hamiltonian H_I satisfied the condition that $c_x^{ij} = c_y^{ij} = c_z^{ij}$.

Proof This can be obtained by calculate $[H_I, \sum_{i=1}^n C_i]$ straightforwardly.

In the rest of this Letter, we apply *Information Conversation Law* in the quantum spin networks that have been used for short-distance quantum communication [9, 10, 19]. The unknown quantum state to be transmitted is $|\phi\rangle = \alpha|0\rangle + \beta|1\rangle$, located at the first qubit. Thus it is the standard quantum state. And the other qubits of this spin system are set to the ground state $|00\cdots 0\rangle$. Then the nearest-neighbor interactions are switched on. After time interval t , we can recover this standard quantum state from the i th qubit with a fidelity $F_i(t)$. From the view of quantum communications, the most important goal is to make the maximal fidelity of the n th qubit from which qubit we hope to recover the input state, i.e. $F_n(t)$, as high as possible, i.e. close to 1. Very recently, M. Christandl et al propose a protocol [10] that can achieve perfect transfer of any quantum state in a constant time. This scheme is theoretically rather beautiful, but from a practical point of view it is complex. It needs accurate control of coupling strengths and static magnetic fields. For example, over a distance n , it need to control $2n + 1$ parameters rigorously. Furthermore, the maximal coupling strength J_{\max} and the minimal coupling strength J_{\min} has the relation that $J_{\max}/J_{\min} \sim \sqrt{n}$. If n is large enough, then this condition may be not easy to be satisfied in practical experiments. Here we propose efficient and easy methods for achieving this goal based on *Information Conservation Law*. Without loss of generality, we average the pure standard quantum state over the whole Bloch-Sphere. We consider the Heisenberg coupling linear n -qubit chain model without using static magnetic fields. The Hamiltonian is given by

$$H_L = - \sum_{i=1}^{n-1} J_{i,i+1} \vec{\sigma}^i \cdot \vec{\sigma}^{i+1} \quad (7)$$

The fidelity of each qubit $F_i(t)$ surges with time approximately periodically. The period

is determined by the coupling strengths. According to *Corollary 2*, we know that the total fidelity $F(t)$ of all the qubits in this spin chain is conserved during the evolution. So there are obviously two type of methods for improving the maximal fidelity of the n th qubit based on *Information Conversation Law*. One is to increase $F(t)$ by choosing a more better initial state. The other is to reduce the total fidelity of all the qubits except the n th qubit. Two schemes of this type are proposed in the following.

I. We demonstrate the first method in the case of $n = 3$. The initial state is set as $|\psi\rangle = |\phi\rangle \otimes |0\rangle \otimes |0\rangle$ or $|\psi'\rangle = |\phi\rangle \otimes |\phi\rangle \otimes |0\rangle$. Then the fidelity of the destination qubit, that is the 3rd qubit, is $F_3(t)$ and $F'_3(t)$ correspondingly. It can be seen that $F'_3(t)$ is always larger than $F_3(t)$. Though this method is simple and may be not very practical, it demonstrates *Information Conversation Law* very well.

II. The starting point of the second type of method is to make the minimum points of the fidelities of the qubits except the n th qubit as close as possible.

(a). Consider the qubits except the n th qubit. Since $F_i(t)$ changes with time approximately periodically, we can adjust the coupling strengths of different pairs of qubits to make them not all the same. This adjustment is very simple and free without much restriction. The effect is to make the fidelities of these qubits come into being beats, so there will always exit some time t when the minimum points of the fidelities of these qubits are very close. Obviously at this time, the fidelity of the n th qubit is maximal and the maximum is improved. We show this result in the simple case of $n = 3$. See Fig. 1. We can see that the maximum fidelity of the 3rd qubit is much improved. In fact it can even be larger than 0.99.

This scheme is very practical in experiments. It is well known that it is difficult to construct a Heisenberg model system with all the same coupling strength between different pairs of qubits, which is employed in [9], due to experimental technology. Here we show that it is not a limitation in quantum communications through an open end spin chain based on *Information Conversation Law*. Actually, we can benefit from this fact. As long as the coupling strengths is fixed, we can determined the optimal time when the improved maximal fidelity is obtained through numerical methods. We need not to control the coupling strengths strictly.

(b). Another scheme is to use spin networks of special graph, for example using spin ring. This result has also been addressed in [9], e.g. quantum communications using a spin ring are more efficient than using a spin line over the same distance. Here we give an explicit




FIG. 1: $F_3(t) \sim t$ (blue line), $F'_3(t) \sim t$ (red line). We set the standard state $|\phi\rangle$, the initial state $|\psi\rangle = |\phi\rangle \otimes |0\rangle \otimes |0\rangle$. The parameters in H_I : $J_{1,2} = J_{2,3} = 0.6$ and $J_{1,2} = 0.6, J_{2,3} = 0.5$ respectively.

explanation based on *Information Conversation Law*.

For a ring with $2N$ qubits, the qubits except the source (black circle in Fig. 2) and destination qubit (blue circle) located at diametrical opposite sites can be separated into symmetric groups (red circles). The qubits of each symmetric group arrive the minimum points at the same time. So the decrease of the total fidelity qubits except the N th qubit is obviously larger than the corresponding one in the situation of using a spin line over the same distance. Of course the optimal fidelity of quantum communications is improved. For the extension of this result, we know that if using spin networks with more symmetric branches, as depicted in Fig. 2, higher efficiency can be obtained.

In conclusion, in this Letter we put forward a new concept: *Conservative Hamiltonians*. If a multi-qubit system's Hamiltonian is conservative then the total information about a prescribed standard quantum state is conserved during the evolution of the system. We also point out when the nonlocal Hamiltonians with pairwise interactions are *Conservative Hamiltonians*. This provides us a new perspective for the research of nonlocal Hamiltonians and nonlocal gates. It can also help us to obtain more in-depth comprehension about many problems involved fidelity in quantum computation and quantum information, e.g. no cloning theorem and capacities of quantum channels. We also apply the *Information*




FIG. 2: Spin graph with nearest-neighbor interactions (dash line) for quantum communications over distance $N = 2$. (a) Spin line; (b) Spin ring; (c) Spin graph with three symmetric branches.

Conversation Law in the situation of quantum communications through spin networks. Easy methods for improving the efficiency of this kind of quantum communications have been proposed. They does not involve too sophisticate and strict parameter control.

Acknowledgments

This work was funded by National Fundamental Research Program (2001CB309300), the Innovation funds from Chinese Academy of Sciences, Z.-W. Zhou acknowledges funds from National Natural Science Foundation of China (Grant No. 10204020).

-
- [1] C.H. Bennett and D.P. DiVincenzo, *Nature* 404, 247 (2000).
 - [2] L.K. Grover, *Proceedings, 28th, ACM Symposium on Theory of Computation*, 212, 1996.
 - [3] L.K. Grover, *Phys. Rev. Lett.* 79, 325 (1997).
 - [4] P.W. Shor, *quant-ph/9508027* (1995).
 - [5] S.C. Benjamin, *Phys. Rev. Lett.* 88, 017904 (2002). X.Zhou et al, *Phys. Rev. Lett.* 89, 197903 (2002). S.C. Benjamin, and S. Bose, *Phys. Rev. Lett.* 90, 247901 (2003).

- [6] A. Barenco et al., Phys. Rev. A 52, 3257 (1995).
- [7] G. Vidal, and C.M. Dawson, Phys. Rev. A 69, 010301 (2004).
- [8] Robert Raussendorf, and Hans J. Briegel, Phys. Rev. Lett. 86, 5188 (2001).
- [9] Sougato Bose, Phys. Rev. Lett. 91, 207901 (2003).
- [10] Matthias Christandl, Nilanjana Datta, Artur Ekerk and Andrew J. Landahl, quant-ph/0309131 (2003).
- [11] M. Horodecki and P. Horodecki, quant-ph/9708015.
- [12] M. Horodecki and R. Horodecki, Phys. Lett. A 244, 473 (1998).
- [13] R. Horodecki, M. Horodecki and P. Horodecki, quant-ph/0002021.
- [14] R. Horodecki, M. Horodecki and P. Horodecki, Phys. Rev. A. 63, 022310 (2001).
- [15] M. Horodecki, R. Horodecki, Aditi Sen De and Ujjwal Sen, quant-ph/0407038.
- [16] M. Horodecki, R. Horodecki, Aditi Sen De and Ujjwal Sen, quant-ph/0306044.
- [17] C.H. Bennett, J.I. Cirac et al., Phys. Rev. A 66, 012305 (2002). H.L. Haselgrove, M.A. Nielsen, and T.J. Osborne, Phys. Rev. A 68, 042303 (2003).
- [18] W.K. Wootters, W.H. Zurek, Nature, 299, 802 (1982). L.M. Duan, and G.C Guo, Phys. Lett. A 243, 261 (1998). L.M. Duan, and G.C Guo, Phys. Rev. Lett. 80, 4999 (1998).
- [19] V. Subrahmanyam, Phys. Rev. A 69, 034304 (2004).