

# A theory of electromagnetic fluctuations for metallic surfaces and van der Waals interactions between metallic bodies.

Giuseppe Bimonte\*

*Dipartimento di Scienze Fisiche,  
Università di Napoli Federico II,  
Complesso Universitario MSA,  
Via Cintia I-80126 Napoli, Italy;  
INFN, Sezione di Napoli, Napoli, ITALY*

(Dated: July 4, 2018)

A new general expression is derived for the fluctuating electromagnetic field outside a metal surface, in terms of its surface impedance. It provides a generalization to real metals of Lifshitz theory of molecular interactions between dielectric solids. The theory is used to compute the radiative heat transfer between two parallel metal surfaces at different temperatures. It is shown that a measurement of this quantity may provide an experimental resolution of a long-standing controversy about the effect of thermal corrections on the Casimir force between real metal plates.

PACS numbers: 12.20.Ds, 42.50.Lc

Keywords: Fluctuations, surface impedance, Casimir, superconductors.

One of the most intriguing predictions of Quantum Electrodynamics is the existence of irreducible fluctuations of the electromagnetic (e.m.) field in the vacuum. It was Casimir's fundamental discovery [1] to realize that the effects of this purely quantum phenomenon were not confined to the atomic scale, as in the Lamb shift, but would rather manifest themselves also at the macroscopic scale, in the form of a force of attraction between two parallel discharged metal plates at distance  $L$ . By modern experimental techniques the Casimir force has now been measured with an accuracy of a few percent (see Refs.[2] and Refs. therein). For a recent review on both theoretical and experimental aspects of the Casimir effect, see Ref.[3]. A few years after Casimir's seminal paper, E.M. Lifshitz developed the general theory of molecular attractive forces between dielectric bodies [4] based on Rytov's theory of e.m. fluctuations [5]. In this theory, the physical origin of the attraction resides in the fluctuating e.m. field which is always present in the interior of any absorbing medium, and extends beyond its boundaries partly in the form of propagating waves and partly as evanescent waves. The resulting expression of the attractive force automatically includes both non-retarded and retarded effects, as well as the effect of the bodies temperature. Moreover, for zero temperature and in the limit  $\epsilon(\omega) = \infty$ , it reproduces the force found by Casimir for two perfect mirrors.

The basic assumption in Lifshitz theory is that one can describe the propagation of e.m. fields inside the materials by means of a dielectric constant. As it is well known [6, 7], this is not always the case for metals, because when the penetration depth  $\delta$  of the e.m. field becomes comparable or larger than the free mean path of

the conduction electrons, there are non-local correlations in the material, and it is no longer possible to use the simple form of the wavevector derived from a dielectric response (anomalous skin-effect). It is well known, however that in a wide range of frequencies, including the anomalous region, the fields *outside* a metal can be accurately described by means of the *surface impedance*  $\zeta(\omega)$  introduced by M. A. Leontovich [7]. We recall that the surface impedance  $\zeta$  relates the tangential components  $E_t$  and  $B_t$  of the e.m. fields on the surface of the metal. For isotropic surfaces:

$$\mathbf{E}_t = \zeta(\omega) [\mathbf{B}_t \times \mathbf{n}], \quad (1)$$

where  $\mathbf{n}$  is the inward unit normal to the surface. For an ideal metal  $\zeta = 0$ , while for a good conductor  $|\zeta| \ll 1$ .

The purpose of this Letter is to develop the theory of e.m. fluctuations for metal surfaces. The main result is a new formula for the correlation functions of the random e.m. fields that are present *outside* a metal surface in thermal equilibrium, as a result of the fluctuating microscopic currents in the interior of the metal. A key feature is that the correlation functions are expressed in terms of the surface impedance, and therefore they apply to the anomalous region, as well as to superconductors (extreme anomalous effect).

Let the metal occupy the  $z < 0$  half-space. We consider the Fourier decomposition of the e.m. field outside the metal. For the electric field of TE modes, we write:

$$\vec{E}^{(TE)} = \frac{1}{(2\pi)^{3/2}} \int_0^\infty d\omega \int d^2k_\perp a(\omega, \vec{k}_\perp) \vec{e}_\perp e^{i(\vec{k}_\perp \cdot \vec{x} - \omega t)} + c.c., \quad (2)$$

where  $\vec{k}_\perp$  is the tangential component of the wave-vector  $\vec{k}$  and  $\vec{e}_\perp$  is a unit vector perpendicular to the plane formed by  $\vec{k}_\perp$  and the normal to the metal surface. The third component of the wave-vector  $k_z = \sqrt{\omega^2/c^2 - k_\perp^2}$  is defined such that  $Re(k_z) \geq 0$  and  $Im(k_z) \geq 0$ . In

---

\*Bimonte@na.infn.it%;

this way the external field appears as a superposition of *propagating waves* (p.w.) travelling away from the surface (for  $k_{\perp} < \omega/c$ ) and of *evanescent waves* (e.w.) (for  $k_{\perp} > \omega/c$ ) exponentially dying out away from the surface. Similarly, we write for the magnetic field of TM modes:

$$\vec{B}^{(TM)} = \frac{1}{(2\pi)^{3/2}} \int_0^{\infty} d\omega \int d^2k_{\perp} b(\omega, \vec{k}_{\perp}) \vec{e}_{\perp} e^{i(\vec{k} \cdot \vec{x} - \omega t)} + c.c. . \quad (3)$$

Our new formulae for the correlation functions of the fluctuating e.m. field are expressed in terms of the statistical averages of the products of the amplitudes  $a(\omega, \vec{k}_{\perp})$  and  $b(\omega, \vec{k}_{\perp})$ :

$$\langle a(\omega, \vec{k}_{\perp}) a^*(\omega', \vec{k}'_{\perp}) \rangle = \frac{4\pi \hbar \omega}{c} \coth\left(\frac{\hbar\omega}{2kT}\right) \frac{Re(\zeta)}{|1 + \zeta k_z|^2} \delta(\omega - \omega') \delta(\vec{k}_{\perp} - \vec{k}'_{\perp}), \quad (4)$$

$$\langle b(\omega, \vec{k}_{\perp}) b^*(\omega', \vec{k}'_{\perp}) \rangle = \frac{4\pi \hbar \omega}{c} \coth\left(\frac{\hbar\omega}{2kT}\right) \frac{Re(\zeta)}{|\zeta + k_z|^2} \delta(\omega - \omega') \delta(\vec{k}_{\perp} - \vec{k}'_{\perp}), \quad (5)$$

$$\langle a(\omega, \vec{k}_{\perp}) a(\omega', \vec{k}'_{\perp}) \rangle = \langle b(\omega, \vec{k}_{\perp}) b(\omega', \vec{k}'_{\perp}) \rangle = \langle a(\omega, \vec{k}_{\perp}) b^*(\omega', \vec{k}'_{\perp}) \rangle = \langle a(\omega, \vec{k}_{\perp}) b(\omega', \vec{k}'_{\perp}) \rangle = 0, \quad (6)$$

with  $k$  Boltzmann's constant. We obtained the above formulae by considering p.w. with frequencies  $\omega$  belonging to the domain of the normal skin effect, for which dielectric and surface impedance b.c. give coinciding results provided only that  $\epsilon'' \gg 1$ , with  $\epsilon''$  the imaginary part of  $\epsilon$ . In this region [7] one can take  $\zeta = \sqrt{\mu/\epsilon}$ , with  $\mu$  the magnetic permeability (we took  $\mu = 1$ ) and then, for p.w. in this domain, Eqs.(4-6) coincide to high accuracy with the analogous expressions derived from Lifshitz theory. We stress however that the validity of Eqs. (4-6) extends to the anomalous region, where Lifshitz theory is inapplicable, and therefore they constitute a new non-trivial generalization of that theory. Details will be given elsewhere.

As a first application of Eqs.(4-6), we have obtained a new derivation of the Casimir force between two metallic plates. Existing derivations based on surface impedance b.c., as found in the literature (see for example Ref.[8]), use a mode analysis to evaluate the Casimir free energy of the cavity, much in the spirit of the original Casimir's paper. The delicate thing in this approach is to justify the validity of the result in the case of dissipative mirrors, which requires consideration of an auxiliary electrodynamic problem (see the second of Ref. [8] for details). Using Eqs. (4-6), we can however obtain the Casimir force in much the same way used by Lifshitz, i.e. by computing the statistical average of the  $z$ -component of the Maxwell stress tensor inside the gap, resulting from fluctuating e.m. fields, at a point close to the surface of either plate. Dissipation poses no problem now, and is taken into account from the start. The computation is rather lengthy and gives the following result for the spectral density  $F_{\omega}$  of the attraction force  $F$  per unit

area ( $F = \int_0^{\infty} d\omega F_{\omega}$ ):

$$F_{\omega} = \coth\left(\frac{\hbar\omega}{2kT}\right) \frac{\hbar\omega^3}{2\pi^2 c^3} Re \int p^2 dp \sum_{\alpha} [(C_{\alpha}-1)^{-1} + 1/2], \quad (7)$$

where

$$C_{\alpha} = e^{-2ip\omega L/c} / (r_{\alpha}^{(1)} r_{\alpha}^{(2)}), \quad \alpha = TE, TM. \quad (8)$$

Here  $p = ck_z/\omega$  and the integration with respect to  $p$  is along the real axis from 1 to 0 (corresponding to p.w.) and thence along the imaginary axis to  $i\infty$  (corresponding to e.w.), while  $r_{\alpha}^{(1)}$  and  $r_{\alpha}^{(2)}$  are the impedance reflection coefficients for mirror 1 and 2 respectively:

$$r_{TM}^{(i)} = \frac{c k_z/\omega - \zeta_i}{c k_z/\omega + \zeta_i}, \quad r_{TE}^{(i)} = \frac{\zeta_i c k_z/\omega - 1}{\zeta_i c k_z/\omega + 1}. \quad (9)$$

Eqs. (7) and (8) are of the same form as those obtained by Lifshitz, apart from the expression of the reflection coefficients. After disregarding in Eq.(7) the terms with  $1/2$ , which give a divergent  $L$ -independent contribution, and after a rotation of the integration contours of  $\omega$  and  $p$ , our expression for  $F$  becomes identical to that in [8].

The second application of Eqs.(4-6) that we consider is new and much more interesting, as it relates to the presently controversial issue of the modification of the Casimir force arising from a non zero temperature of the mirrors, when the latter are treated as real metals. The debate was raised by the findings of Refs. [9], showing that the combined effect of temperature and finite conductivity leads to large deviations from the ideal metal case. This result was obtained within the framework of




FIG. 1: Plots of radiative heat transfer between two Al plates (in  $\text{erg}/(\text{s cm}^2)$ ) at temperatures  $T_1 = 323$  K and  $T_2 = 300$  K, as a function of separation (in microns), according to Lifshitz theory (dashed line) and surface impedance approach (continuous line).

Lifshitz theory, by using the Drude dielectric function  $\epsilon_D(\omega)$

$$\epsilon_D(\omega) = 1 - \Omega_p^2 / [\omega(\omega + i\gamma)] \quad (10)$$

to describe the metal. Finite conductivity is taken into account by allowing a non-vanishing value for the relaxation frequency  $\gamma$ . The results of Refs. [9] have been criticized by several authors, and supported by others. There is no space here to discuss the matter, and we address the reader to Refs. [8] and [10] for a discussion of the extended literature on this topic.

Of particular interest to us is the criticism of Ref.[9] in the second of Refs.[8], where it is argued that the dielectric model, *in combination with the Drude dielectric function*, fails to describe the thermal Casimir force *even in the domain of the normal skin effect*, and that also in this region the appropriate description is provided by impedance b.c.. Further arguments in favor of this point of view were found in [11], where the spectrum of finite temperature corrections to the Casimir force is studied, for metallic plates of finite conductivity. This study shows that the large deviations from the ideal metal case in [9] arise from *thermal TE e.w. of low frequencies* ( $\omega = 10^{10} - 10^{13}$  rad/s for  $L = 1\mu\text{m}$ ), for which the description of the plates as bulk dielectrics is

---


$$S = -\frac{4\hbar}{\pi^2 c^2} \int_0^\infty d\omega \omega^3 \left( \frac{1}{\exp(\hbar\omega/kT_1) - 1} - \frac{1}{\exp(\hbar\omega/kT_2) - 1} \right) \text{Re}(\zeta_1)\text{Re}(\zeta_2) \text{Re} \int dp p |p|^2 |e^{2ipL\omega/c}| \left( \frac{1}{B_{TE}} + \frac{1}{B_{TM}} \right) \quad (11)$$


---

where the contour of integration for  $p$  is the same as in

not valid. It is also shown there that if surface impedance b.c. are used, with  $\zeta = 1/\sqrt{\epsilon_D}$  (which is the expression valid in the domain of the normal skin effect), instead of the large *repulsive* thermal correction from TE e.w. found in [9], one obtains an *attractive* correction, of much smaller magnitude, while no appreciable differences are found both in the TM and in the TE p.w. sectors [16]. The conclusion of this study is that the conflicting statements on thermal corrections to the Casimir force, may be traced back to different magnitudes for the thermal TE e.w. correction, in the two approaches.

In view of these findings, it would be important to devise new experiments to probe the effectiveness of the two approaches to describe physical effects of thermal TE e.w., in the frequency range that is relevant for the Casimir effect. Optical data are of no help here, since they refer to p.w., and therefore give no information on e.w.. A key remark now is that the relevant e.w. are not vacuum fluctuations, as in the Casimir force at zero temperature, but rather real thermal excitations. Now, it is known that thermal e.w. play an important role in the heat transfer  $S$  between two closely spaced metal surfaces, at temperatures  $T_1$  and  $T_2$  [13]. It is therefore very interesting to see what is the prediction of impedance theory for  $S$ , and to compare it with the result from Lifshitz theory. The computation of  $S$  goes through the same steps as in Lifshitz' computation of the Casimir force [13]. One assumes that each plate separately is in local thermal equilibrium, and emits a fluctuating e.m. field according to our Eqs. (4-6), with  $T$  the temperature of the emitting plate. Since thermal e.m. fields from plates 1 and 2 are uncorrelated, the radiated power  $S$  can be obtained by subtracting incoherently the average Poynting vectors for the multiply reflected radiation fields originating from either plate. Apart from the fact that we obviously evaluate the average Poynting vector instead of the Maxwell stress-tensor, the only difference with respect to the computation of thermal corrections to the Casimir force is that the fluctuating fields from plate 1 and 2 are relative now to distinct temperatures  $T_1$  and  $T_2$ . Apart from this, these thermal fields have the same expression as in the Casimir force computation, at nonzero temperature. Our final formula for the power  $S$  (per unit area) has the form of a difference between two terms, one for each plate:

Eq.(7), and the quantities  $B_{TE/TM}$  are defined as:

$$B_{TE} = |(1+p\zeta_1)(1+p\zeta_2) - (1-p\zeta_1)(1-p\zeta_2) \exp(2ipL\omega/c)|^2, \quad (12)$$




FIG. 2: Plot of the difference  $s_{TE\ e.w.}^{(imp)}(\omega) - s_{TE\ e.w.}^{(Lif)}(\omega)$  between TE e.w. contributions to the spectra  $s^{(imp)}(\omega)$  (impedance theory) and  $s^{(Lif)}(\omega)$  (Lifshitz theory) of radiative heat transfer between two Al plates (in  $\text{erg}/(\text{rad cm}^2)$ ) at temperatures  $T_1 = 323$  K and  $T_2 = 300$  K, as a function of  $\log_{10}(\omega)$ .  $L = 0.3 \mu\text{m}$ .

$$B_{TM} = |(p + \zeta_1)(p + \zeta_2) - (p - \zeta_1)(p - \zeta_2) \exp(2ipL\omega/c)|^2. \quad (13)$$

In Fig. 1 we show plots of the radiated power (in  $\text{erg}/(\text{s cm}^2)$ ), for two plates of Al as a function of the separation  $L$  (in microns), for  $T_1 = 323$  K and  $T_2 = 300$  K. In parallel with the analysis of thermal corrections to the Casimir force described earlier, the figure compares the result from Lifshitz theory, Eq. (19) of Ref. [13], with the Drude dielectric function Eq. (10) (dashed line), with our Eq. (11), with  $\zeta(\omega) = 1/\sqrt{\epsilon_D(\omega)}$  (solid line). In both cases, we took  $\hbar\Omega_p = 11.5$  eV and  $\hbar\gamma = .05$  eV. We see that impedance theory and Lifshitz theory give rather different predictions for  $S$  (for  $L = 0.3 \mu\text{m}$ , impedance theory gives for  $S$  a value approximately 17 % larger than Lifshitz theory). We observe that in the considered submicron separation range, heat transfer occurs mainly via TE e.w.. The smaller contribution from p.w. and from TM e.w. is practically the same for the two theories, to an accuracy of a few parts in ten thousand, and therefore the difference between the two curves in Fig. 1 arises exclusively from TE e.w.. For larger separations the relative weight of e.w. with respect to p.w. decreases, and the two theories lead to very similar results. Already at one micron the results from Lifshitz theory and impedance theory differ only by four percent. We eval-

uated the contribution  $s_{TE\ e.w.}(\omega)$  from TE e.w. to the spectra  $s(\omega)$  of the radiated power ( $S = \int_0^\infty d\omega s(\omega)$ ) for the two theories and in Fig. 2 we plot the difference  $s_{TE\ e.w.}^{(imp)}(\omega) - s_{TE\ e.w.}^{(Lif)}(\omega)$ , as a function of  $\log_{10}(\omega)$ , for  $L = 0.3 \mu\text{m}$ . We see that two spectra differ significantly precisely in the low-frequency region which, according to [11] is at the origin of the controversial thermal corrections to the Casimir force. As pointed out in Ref.[11], in this frequency domain the penetration depth is of the same order as the mean free path of the conduction electrons. Therefore the dielectric model for the plates is not valid, and a more realistic description is provided by surface impedance b.c.. It is clear that an accurate measurement of  $S$  would provide strong indications of which approach is preferable, a knowledge which could then be used in the evaluation of the thermal Casimir effect. An experiment measuring the radiative heat transfer between two chromium plates is described in Ref. [14]. Even though the author of Ref. [14] claims to have obtained some evidence for proximity effects related to e.w., his measurements were performed mostly for plates separations larger than one micron, where we expect small differences between the two theories. Moreover, as the author does not quote the area of the plates, no comparison is possible with his data. It would therefore be desirable to repeat the measurements for submicron separations, and for metals that are used in Casimir experiments.

In conclusion, we have presented new formulae for the correlation functions of e.m. fluctuations present outside a metal. The formulae involve the surface impedance of the metal, and are therefore applicable in the anomalous region, as well as in the extreme anomalous case (superconductors), where Lifshitz theory is not valid. As an application, we have evaluated the radiative heat transfer between two metal plates at different temperatures and we have shown that a measurement of this quantity should provide enough information to settle experimentally recent controversies about the thermal Casimir effect. We plan to apply our results to superconducting cavities, that were proposed recently [15] as a tool to measure the variation of Casimir energy across the superconducting phase transition.

The author thanks G.L. Klimchitskaya and V.M. Mostepanenko for valuable suggestions, and acknowledges partial financial support by PRIN *SINTESI*.

[1] H.B.G. Casimir, Proc. K. Ned. Akad. Wet. Rev. **51**, 793 (1948).  
 [2] S.K. Lamoreaux, Phys. Rev. Lett. **78**, 5 (1997); U. Mohideen and A. Roy, ibid. **81**, 4549 (1998); G. Bressi, G. Carugno, R. Onofrio and G. Ruoso, ibid. **88** 041804 (2002); R.S. Decca, D. López, E. Fischbach, and D.E. Krause, ibid. **91**, 050402 (2003).  
 [3] M. Bordag, U. Mohideen and V.M. Mostepanenko, Phys. Rep. **353**, 1 (2001).

[4] E.M. Lifshitz, Sov. Phys. JETP **2**, 73 (1956); E.M. Lifshitz and L.P. Pitaevskii, *Landau and Lifshitz Course of Theoretical Physics: Statistical Physics Part II* (Butterworth-Heinemann, 1980.)  
 [5] S.M. Rytov, *Theory of Electrical Fluctuations and Thermal Radiation*, Publishing House, Academy of Sciences, USSR (1953).  
 [6] H. London, Proc. R. Soc. **A 176**, 522 (1940).  
 [7] L.D. Landau and E.M. Lifshitz, *Landau and Lifshitz*

*Course of Theoretical Physics: Electrodynamics of Continuous Media* (Oxford: Pergamon, 1960.)

- [8] V.B. Bezerra, G.L. Klimchitskaya and C. Romero, Phys. Rev. **A 65** 012111 (2001); B. Geyer, G.L. Klimchitskaya and V.M. Mostepanenko, *ibid.* **A 67** 062102 (2003); V.B. Svetovoy and M.V. Lokhanin, Phys. Rev. **A 67**, 022113 (2003).
- [9] M. Bostrom and B.E. Sernelius, Phys. Rev. Lett. **84**, 4757 (2000); B.E. Sernelius, *ibid.* **87**, 139102 (2001).
- [10] J.S. Høye, I. Brevik, J.B. Aarseth and K.A. Milton, Phys. Rev. **E 67**, 056116 (2003) and references therein.
- [11] J.R. Torgerson and S.K. Lamoreaux, Phys. Rev. **E 70**, 047102 (2004); S.K. Lamoreaux, Rep. Prog. Phys. **68**, 201 (2005).
- [12] G. Bimonte, *Comment on "Low-frequency character of the Casimir force between metallic films"*, in press on Phys. Rev. **E**, (2006).
- [13] D. Polder and M. Van Hove, Phys. Rev. **B 4**, 3303 (1971).
- [14] C.M. Hargreaves, Phys. Lett **A 30**, 491 (1969).
- [15] G.Bimonte, E. Calloni, G. Esposito, L. Milano and L. Rosa, Phys. Rev. Lett. **94**, 180402 (2005); G. Bimonte, E. Calloni, G. Esposito, and L. Rosa, Nucl. Phys. **B 726**, 441 (2005).
- [16] See Ref. [12], where a mistake in impedance b.c. for TE modes made in [11] is corrected.