

Still more on norms of completely positive maps

Stanislaw J. Szarek (Cleveland and Paris)

Let \mathcal{M}_n denote the space of $n \times n$ (real or complex) matrices and, for $A \in \mathcal{M}_n$ and $p \geq 1$, let $\|A\|_p := (\text{tr}(A^\dagger A)^{p/2})^{1/p}$ be the Schatten p -norm of A , with the limit case $p = \infty$ corresponding to the usual operator norm. Further, if $\Phi : \mathcal{M}_m \rightarrow \mathcal{M}_n$ is a linear map and $p, q \in [1, \infty]$, we consider

$$\|\Phi\|_{p \rightarrow q} := \max\{\|\Phi(\sigma)\|_q : \sigma \in \mathcal{M}_m, \|\sigma\|_p \leq 1\}, \quad (1)$$

i.e., the norm of Φ as an operator between the normed spaces $(\mathcal{M}_m, \|\cdot\|_p)$ and $(\mathcal{M}_n, \|\cdot\|_q)$. Such quantities were studied (in the context of quantum information theory) in [1], where the question was raised under what conditions (1) coincides with the *a priori* smaller norm

$$\|\Phi\|_{p \rightarrow q}^H := \max\{\|\Phi(\sigma)\|_q : \sigma \in \mathcal{M}_m, \sigma = \sigma^\dagger, \|\sigma\|_p \leq 1\} \quad (2)$$

of the restriction of Φ to the (real linear) subspace of *Hermitian* matrices and, in particular, whether this holds if Φ is *completely positive*. [Note that if Φ is just *positivity preserving*, it maps Hermitian matrices to Hermitian matrices.] The latter was subsequently confirmed in [2, 3], the first of which also contains an assortment of examples showing when such equalities may or may not hold (see also [4]). Here we provide one more proof. More precisely, we will show

Proposition *If Φ is 2-positive, then $\|\Phi\|_{p \rightarrow q} = \|\Phi\|_{p \rightarrow q}^H$. Moreover, similar equality holds if the domain and the range of Φ are endowed with any unitarily invariant norms.*

Recall that a norm $\|\cdot\|$ on \mathcal{M}_n is called *unitarily invariant* if $\|UAV\| = \|A\|$ for any $A \in \mathcal{M}_n$ and any $U, V \in U(n)$ (resp., $O(n)$ in the real case); see [5, 6]. This is equivalent to requiring that the norm of a matrix depends only on its *singular values* (called in some circles ‘‘Schmidt coefficients’’). Besides using a slightly weaker hypothesis and yielding a slightly more general assertion, the argument we present is self-contained and uses only definitions and elementary facts and concepts from linear algebra, of which the most sophisticated is the singular value decomposition. It may thus be argued that it is the ‘‘right’’ proof. [Note that an analysis of [3] and its references shows that in fact only 2-positivity is needed there, too.]

Proof For clarity, we will consider first the case when $p = 1$, i.e., when the domain of Φ is endowed with the trace class norm. In this case the extreme points of the respective unit balls (on which the maxima in (1) and (2) are necessarily achieved) are particularly simple: they are rank one operators. Accordingly, the question reduces to showing that

$$\max_{|u|=|v|=1} \|\Phi(|v\rangle\langle u|)\|_q \leq \max_{|u|=1} \|\Phi(|u\rangle\langle u|)\|_q, \quad (3)$$

where $u, v \in \mathbb{C}^m$ (or \mathbb{R}^m , depending on the context) and $|\cdot|$ is the Euclidean norm. Given such u, v , consider the block matrix $M_{u,v} = \begin{bmatrix} |u\rangle\langle u| & |u\rangle\langle v| \\ |v\rangle\langle u| & |v\rangle\langle v| \end{bmatrix} \in \mathcal{M}_{2m}$ and note that $M_{u,v} = |\xi\rangle\langle \xi|$ where $|\xi\rangle = (|u\rangle, |v\rangle) \in \mathbb{C}^r \oplus \mathbb{C}^r$ (in particular $M_{u,v} \geq 0$). Considering $M_{u,v}$ as an element of $\mathcal{M}_m \otimes \mathcal{M}_2$ and appealing to 2-positivity of Φ we deduce that $(\Phi \otimes Id_{\mathcal{M}_2})(M_{u,v}) = \begin{bmatrix} \Phi(|u\rangle\langle u|) & \Phi(|u\rangle\langle v|) \\ \Phi(|v\rangle\langle u|) & \Phi(|v\rangle\langle v|) \end{bmatrix} \geq 0$. The conclusion now follows from the following lemma (see, e.g., [6], Theorem 3.5.15; for completeness we include a proof at the end of this note).

Lemma *Let $A, B, C \in \mathcal{M}_r$ be such that the $2r \times 2r$ block matrix $M = \begin{bmatrix} A & B \\ B^\dagger & C \end{bmatrix}$ is positive semi-definite, and let $\|\cdot\|$ be a unitarily invariant norm on \mathcal{M}_r . Then $\|B\|^2 \leq \|A\| \|C\|$.*

The case of arbitrary $p \in [1, \infty]$ is almost as simple. First, for $\sigma \in \mathcal{M}_m$ with $\|\sigma\|_p \leq 1$ we consider the positive semi-definite matrix $M_\sigma = \begin{bmatrix} (\sigma\sigma^\dagger)^{1/2} & \sigma \\ \sigma^\dagger & (\sigma^\dagger\sigma)^{1/2} \end{bmatrix}$. [Positivity is seen, e.g., by writing down the singular value decompositions of the entries and expressing M_σ as a positive linear combination of matrices of the type $M_{u,v}$ considered above.] Since unitarily invariant norms depend only on singular values of a matrix, we have $\|(\sigma\sigma^\dagger)^{1/2}\|_p = \|(\sigma^\dagger\sigma)^{1/2}\|_p = \|\sigma\|_p \leq 1$. On the other hand, arguing as in the special case $p = 1$, we deduce from the Lemma that $\|\Phi(\sigma)\|_q^2 \leq \|\Phi((\sigma\sigma^\dagger)^{1/2})\|_q \|\Phi((\sigma^\dagger\sigma)^{1/2})\|_q \leq (\|\Phi\|_{p \rightarrow q}^H)^2$, and the conclusion follows by taking the maximum over σ . The proof for general unitarily invariant norms is the same. \square

Proof of the Lemma [Written for $\|\cdot\| = \|\cdot\|_q$, but the general case works in the same way.] Let $B = \sum_{j=1}^r \lambda_j |\varphi_j\rangle\langle\psi_j|$ be the Schmidt decomposition. Consider the orthonormal basis of \mathbb{C}^{2r} which is a concatenation of $(|\varphi_j\rangle)$ and $(|\psi_j\rangle)$. The representation of M in that basis is

$$M' := \begin{bmatrix} (\langle\varphi_j|A|\varphi_k\rangle)_{j,k=1}^r & \text{Diag}(\lambda) \\ \text{Diag}(\lambda) & (\langle\psi_j|C|\psi_k\rangle)_{j,k=1}^r \end{bmatrix},$$

where $\text{Diag}(\mu)$ is the diagonal matrix with the sequence $\mu = (\mu_j)$ on the diagonal. Given $j \in \{1, \dots, r\}$, the 2×2 matrix $\begin{bmatrix} \langle\varphi_j|A|\varphi_j\rangle & \lambda_j \\ \lambda_j & \langle\psi_j|C|\psi_j\rangle \end{bmatrix}$ is a minor of M' and hence positive semi-definite, and so $\lambda_j \leq \sqrt{\langle\varphi_j|A|\varphi_j\rangle\langle\psi_j|C|\psi_j\rangle} \leq (\langle\varphi_j|A|\varphi_j\rangle + \langle\psi_j|C|\psi_j\rangle)/2$. Consequently

$$\begin{aligned} \|B\|_q &= \left(\sum_j \lambda_j^q\right)^{1/q} \leq \left(\sum_j \langle\varphi_j|A|\varphi_j\rangle^q + \sum_j \langle\psi_j|C|\psi_j\rangle^q\right)^{1/q} / 2 \\ &\leq (\|A\|_q + \|C\|_q) / 2. \end{aligned} \quad (4)$$

The last inequality in (4) follows from the well-known fact that, for *any* square matrix $S = (S_{jk})$, $\|S\|_q \geq (\sum_j S_{jj}^q)^{1/q}$ (which in turn is a consequence of $(S_{jk}\delta_{jk})$, the diagonal part of S , being the average of $\text{Diag}(\varepsilon)S\text{Diag}(\varepsilon)$, where $\varepsilon = (\varepsilon_j)$ varies over all choices of $\varepsilon_j = \pm 1$). The bound from (4) is already sufficient to prove (3) (and the Proposition). To obtain the stronger statement from the Lemma we use the inequality $ab \leq \frac{1}{2}(ta + b/t)$ (for $t > 0$, instead of $ab \leq \frac{1}{2}(a + b)$) to obtain $\|B\|_q \leq \frac{1}{2}(t\|A\|_q + \|C\|_q/t)$, and then specify the optimal value $t = (\|C\|_q/\|A\|_q)^{1/2}$. Passing to a generic unitarily invariant norm requires just replacing everywhere $(\sum_j \mu_j^q)^{1/q}$ by $\|\text{Diag}(\mu)\|$; equalities such as $\|B\| = \|\text{Diag}(\lambda)\|$ or $\|A\| = \|(\langle\varphi_j|A|\varphi_k\rangle)_{j,k=1}^r\|$ just express the unitary invariance of the norm. \square

The author thanks K. Audenaert and M. B. Ruskai for comments on the first version of this note.

References

- [1] C. King and M. B. Ruskai, Comments on multiplicativity of maximal p -norms when $p = 2$, in *Quantum Information, Statistics and Probability*, ed. by O. Hirota, World Scientific 2004, see also arxiv.org e-print [quant-ph/0401026](#)
- [2] J. Watrous, Notes on super-operator norms induced by Schatten norms, *QIC* 5 (2005), 58–68.
- [3] K. Audenaert, A Note on the $p \rightarrow q$ norms of Completely Positive Maps, arxiv.org e-print [math-ph/0505085](#)
- [4] I. Devetak, M. Junge, C. King and M. B. Ruskai, Multiplicativity of completely bounded p -norms implies a new additivity result, arxiv.org e-print [quant-ph/0506196](#)
- [5] R. Schatten, *Norm Ideals of Completely Continuous Operators*. Springer-Verlag, Berlin, 1970.
- [6] R. A. Horn and C. R. Johnson, *Topics in matrix analysis*. Cambridge University Press, Cambridge, 1994.

Université Pierre et Marie Curie-Paris 6, UMR 7586-Institut de Mathématiques, Analyse Fonctionnelle, BC 186, 75252 Paris, France and
Case Western Reserve University, Department of Mathematics, Cleveland, Ohio 44106-7058, U.S.A.
Email: szarek@math.jussieu.fr