

Entanglement Entropy: Helicity versus Spin

Song He

Institute of Theoretical Physics, School of Physics, Peking University, Beijing, 100871, China

Shuxin Shao

Department of Physics, Beijing Normal University, Beijing, 100875, China

Hongbao Zhang*

Department of Astronomy, Beijing Normal University, Beijing, 100875, China

Department of Physics, Beijing Normal University, Beijing, 100875, China

CCAST (World Laboratory), P.O. Box 8730, Beijing, 100080, China

(Dated: October 8, 2018)

For a massive spin $\frac{1}{2}$ field, we present the reduced spin and helicity density matrix, respectively, for the same pure one particle state. Their relation has also been developed. Furthermore, we calculate and compare the corresponding entanglement entropy for spin and helicity within the same inertial reference frame. Due to the distinct dependence on momentum degree of freedom between spin and helicity states, the resultant helicity entropy is different from that of spin in general. In particular, we find that both helicity entanglement for a spin eigenstate and spin entanglement for a right handed or left handed helicity state do not vanish and their Von Neumann entropy has no dependence on the specific form of momentum distribution as long as it is isotropic.

PACS numbers: 03.67.Mn 03.65.Ge 03.65.Ud

Quantum information theory is usually formulated in the framework of non-relativistic quantum mechanics, since particles moving at relativistic speeds may not be needed to realize the promise of quantum information process such as quantum computation. However, relativity, especially special relativity plays a significant role in quantum entanglement and related quantum technology, such as teleportation. This point is obviously justified by quantum optics, which is well established on the basis of not only quantum theory but also special relativity in nature[1]. For example, most of EPR-type experiments have been performed by photon pairs[2, 3]. In addition, experiments of quantum teleportation have also been extensively carried out by photons[4, 5].

On the other hand, considerable efforts have also been expanded on the theoretical investigation of quantum information theory in relativistic framework, which has gone beyond from photons to electrons, and from explicit examples calculated in some specific cases to general framework formulated in relativistic quantum mechanics and even relativistic quantum field theory[6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36]. For review, please refer to [18]. However, for Dirac fields, previous discussions of relativistic quantum information theory are restricted primarily on quantum entanglement between spin and momentum degrees of freedom. Only recently has quantum entanglement between helicity and momentum been systematically formulated and numerically analyzed in relativistic framework for the first time

in [37]. This pioneering work not only sets a starting point for further worthwhile investigations along this line, but also acquires much importance from the perspective of high energy physics. It is the helicity rather than spin that is more often under both theoretical consideration and experimental detection in high energy physics, since the helicity has an advantage in providing a smooth transition to the massless case. Although both the helicity states and spin states can constitute the basis of Hilbert space of one particle, they differ in the way of unitary transformation under the action of Lorentz group. As a result, the entanglement properties for helicity differs remarkably from those for spin after we trace out the momentum degree of freedom. Especially, it is found that in the sharp momentum limit, unlike the vanishing spin entropy, at small velocities of the inertial observer the helicity entropy demonstrates a sudden jump onto a constant value, half of the entropy for the maximal entangled Bell states, which may be easily observed in high energy physics experiments.

As a further step along this direction, the purpose of this paper is to investigate the explicit difference between the reduced spin and helicity density matrix. Especially, for a pure one particle state, we find the resultant helicity entropy is different from that for spin with respect to the same inertial observer.

Start with a field with positive mass m and spin $\frac{1}{2}$, we can construct the spin states $|p, \sigma\rangle$ as a complete orthonormal basis for Hilbert space of one particle, where p is the four momentum of particle, and σ represents the spin along the z axis. Similarly, we can also choose the helicity states $|p; \lambda\rangle$ to form a complete orthonormal basis for the same Hilbert space with the helicity denoted by λ . The former is usually called spin representation,

*Current address is Perimeter Institute for Theoretical Physics, Waterloo, Ontario, N2L 2Y5, Canada.

and the latter is called helicity representation. Moreover, these spin states and helicity states are related by the representation transformation as[38]

$$|p; \lambda\rangle = D_{\sigma\lambda}[R(p)]|p, \sigma\rangle. \quad (1)$$

Here $R(p)$ is the rotation that carries the z axis into the direction \mathbf{p} , and D is the spin $\frac{1}{2}$ irreducible unitary representation of Lorentz group, given by

$$D[R(p)] = \begin{pmatrix} e^{-i\frac{\phi}{2}} & 0 \\ 0 & e^{i\frac{\phi}{2}} \end{pmatrix} \begin{pmatrix} \cos\frac{\theta}{2} & -\sin\frac{\theta}{2} \\ \sin\frac{\theta}{2} & \cos\frac{\theta}{2} \end{pmatrix} \quad (2)$$

with $\hat{\mathbf{p}} = (\sin\theta\cos\phi, \sin\theta\sin\phi, \cos\theta)$. Note that the spin states and helicity states consider here are observed within the same inertial reference frame.

For a pure one particle state, we can represent it as expansion of the spin states, i.e.,

$$|\psi\rangle = \sum_{\sigma=\pm\frac{1}{2}} \int d^3\mathbf{p} \psi(\sigma, \mathbf{p}) |p, \sigma\rangle \quad (3)$$

with the normalized condition

$$\sum_{\sigma=\pm\frac{1}{2}} \int d^3\mathbf{p} |\psi(\sigma, \mathbf{p})|^2 = 1. \quad (4)$$

It is noteworthy that this normalized state with a superposition of various momenta represents a more physical reality since a particle has no definite momentum in general, although for convenience the momentum eigenstates are extensively employed in textbooks on high energy physics and quantum field theory. Then the reduced spin density matrix associated with the above normalized state is obtained by tracing the momentum degree, i.e.,

$$\begin{aligned} \rho &= Tr_{\mathbf{p}}[|\psi\rangle\langle\psi|] = \int d^3\mathbf{p} \langle\psi|\mathbf{p}\rangle\langle\psi|\mathbf{p}\rangle \\ &= \sum_{\sigma, \tilde{\sigma}} \int d^3\mathbf{p} [\psi(\sigma, \mathbf{p})\psi^*(\tilde{\sigma}, \mathbf{p})|\sigma\rangle\langle\tilde{\sigma}|]. \end{aligned} \quad (5)$$

Here, we have used the orthonormal relation for the spin states.

Next, as mentioned above, $|\psi\rangle$ can also be expanded by the helicity states as

$$|\psi\rangle = \sum_{\lambda=\pm\frac{1}{2}} \int d^3\mathbf{p} \psi'(\lambda, \mathbf{p}) |p; \lambda\rangle. \quad (6)$$

According to the transformation relation between the spin states and helicity states Eq.(1), we have

$$\psi'(\lambda, \mathbf{p}) = D_{\lambda\sigma}^{-1}[R(p)]\psi(\sigma, \mathbf{p}), \quad (7)$$

which follows the reduced helicity density matrix as

$$\begin{aligned} \rho' &= \sum_{\lambda\tilde{\lambda}} \int d^3\mathbf{p} \\ &\{D_{\lambda\sigma}^{-1}[R(p)]\psi(\sigma, \mathbf{p})\psi^*(\tilde{\sigma}, \mathbf{p})D_{\tilde{\sigma}\tilde{\lambda}}[R(p)]|\lambda\rangle\langle\tilde{\lambda}|\}. \end{aligned} \quad (8)$$

As an example, consider a particle prepared with the spin in z direction, i.e., $\psi(-\frac{1}{2}, \mathbf{p}) = 0$, which implies that the corresponding spin entropy is zero. However, it does not mean that the helicity entropy also vanishes for this state. By Eq.(2) and Eq.(8), the reduced helicity density matrix can be explicitly written as

$$\rho' = \int d^3\mathbf{p} |\psi(\frac{1}{2}, \mathbf{p})|^2 \begin{pmatrix} \frac{1+\cos\theta}{2} & -\frac{\sin\theta}{2} \\ -\frac{\sin\theta}{2} & \frac{1-\cos\theta}{2} \end{pmatrix} \quad (9)$$

In particular, for simplicity but without loss of generality, let $\psi(\frac{1}{2}, \mathbf{p})$ be arbitrary except independent of the angle θ , then the reduced helicity density matrix can be calculated out as

$$\rho' = \begin{pmatrix} \frac{1}{2} & -\frac{\pi}{8} \\ -\frac{\pi}{8} & \frac{1}{2} \end{pmatrix}, \quad (10)$$

whose eigenvalues are easy to obtain as $\rho'_1 = \frac{1}{2} + \frac{\pi}{8}$ and $\rho'_2 = \frac{1}{2} - \frac{\pi}{8}$, respectively. Whence, the helicity entropy reads

$$S = -\left[\left(\frac{1}{2} + \frac{\pi}{8}\right)\log_2\left(\frac{1}{2} + \frac{\pi}{8}\right) + \left(\frac{1}{2} - \frac{\pi}{8}\right)\log_2\left(\frac{1}{2} - \frac{\pi}{8}\right)\right], \quad (11)$$

which is obviously positive.

Note that, in the case considered above, the helicity entropy does not depend on the specific form of wave function $\psi(\frac{1}{2}, \mathbf{p})$ in deed. Especially, if we choose $\psi(\frac{1}{2}, \mathbf{p})$ to be a Gaussian, i.e.,

$$\psi\left(\frac{1}{2}, \mathbf{p}\right) = \pi^{\frac{3}{4}} \tau^{-\frac{3}{2}} e^{-\frac{\mathbf{p}^2}{2\tau^2}}, \quad (12)$$

which is a minimum uncertainty state, the helicity entropy is independent of the distribution width parameter τ .

On the other hand, it is noteworthy that the above result does not show that the helicity entropy always exceeds the corresponding spin entropy for the same state. In order to confirm this assertion, we first combine Eq.(5) with Eq.(8) to obtain the reduced spin density matrix as

$$\begin{aligned} \rho &= \sum_{\sigma\tilde{\sigma}} \int d^3\mathbf{p} \\ &\{D_{\sigma\lambda}[R(p)]\psi'(\lambda, \mathbf{p})\psi'^*(\tilde{\lambda}, \mathbf{p})D_{\tilde{\lambda}\tilde{\sigma}}^{-1}[R(p)]|\sigma\rangle\langle\tilde{\sigma}|\}, \end{aligned} \quad (13)$$

if the corresponding reduced helicity density matrix reads

$$\rho' = \sum_{\lambda\tilde{\lambda}} \int d^3\mathbf{p} [\psi'(\lambda, \mathbf{p})\psi'^*(\tilde{\lambda}, \mathbf{p})|\lambda\rangle\langle\tilde{\lambda}|]. \quad (14)$$

Now consider instead a particle in the eigenstate of helicity with an eigenvalue $+\frac{1}{2}$ and isotropic momentum distributions such as Gaussian. Obviously, associated with this state, the helicity entropy vanishes. However, straightforward but lengthy calculations lead to the reduced spin density matrix as

$$\rho = \int d^3\mathbf{p} |\psi'(\frac{1}{2}, \mathbf{p})|^2 \begin{pmatrix} \frac{1+\cos\theta}{2} & \frac{e^{-i\phi}\sin\theta}{2} \\ \frac{e^{i\phi}\sin\theta}{2} & \frac{1-\cos\theta}{2} \end{pmatrix} = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{pmatrix}, \quad (15)$$

which follows that the spin entropy takes 1 in terms of binary logarithm. Similarly, this result does not rely on the specific momentum distribution of right handed helicity state except that isotropy is required here. In addition, this resultant spin entropy value implies that spin is maximally entangled with momentum for the prepared isotropic right handed helicity state.

In conclusion, although both the helicity states and spin states can constitute the basis of Hilbert space for one particle, they have significantly distinct dependence on momentum degree of freedom and thus demonstrate remarkably different properties of entanglement. Thus, even if in high energy physics experiments the prepared state is a direct product of a function of momentum and a function of spin, it is not a direct product of momentum and helicity, namely, the helicity and momentum appear to be entangled for this state, which means that the mea-

surement of momentum will cause the collapse of helicity degree of freedom in the inner space of the particle, and vice versa. This result may be of instructive significance to both proposal of experiment schemes and analysis of experiment data related to helicity and spin in high energy physics. In addition, it is found that both helicity entanglement for a spin eigenstate and spin entanglement for a right handed or left handed helicity states do not vanish and their Von Neumann entropy has no dependence on the specific form of momentum distribution as long as it is isotropic.

Stimulating communications on the Shining Star Science Forum in web(www.changhai.org), especially with Zhangqi Yin are much appreciated. Work by SH was supported by NSFC(Nos.10235040 and 10421003). SS was supported by LFS of BNU, and HZ was supported in part by NSFC(No.10533010).

-
- [1] D. F. Walls and G. J. Milburn, *Quantum Optics*(Springer-Verlag, Berlin, 1994).
- [2] A. Aspect, *Phys. Rev. Lett.* 49: 1804(1982).
- [3] G. Weihs *et al.*, *Phys. Rev. Lett.* 81: 5039(1998).
- [4] D. Bouwmeester *et al.*, *Nature* 390: 575(1997).
- [5] A. Furusawa *et al.*, *Science* 282: 706(1998).
- [6] M. Czachor, *Phys. Rev. A*55: 72(1997).
- [7] A. Peres and D. R. Terno, *J. Mod. Opt.* 49: 1255(2002).
- [8] A. Peres *et al.*, *Phys. Rev. Lett.* 88: 230402(2002).
- [9] P. M. Alsing and G. J. Milburn, quant-ph/0203051.
- [10] R. M. Gingrich and C. Adami, *Phys. Rev. Lett.* 89: 270402(2002).
- [11] A. Peres and D. R. Terno, *J. Mod. Opt.* 50: 1165(2003).
- [12] A. Peres and D. R. Terno, *Int. J. Quant. Inf.* 1: 225(2003).
- [13] J. Pachos and E. Solano, *Quant. Inf. Comput.* 3: 115(2003).
- [14] S. J. van Enk and T. Rudolph, *Quant. Inf. Comput.* 3: 423(2003).
- [15] M. Czachor and M. Wilczewski, *Phys. Rev. A*68: 010302(2003).
- [16] A. J. Bergou *et al.* *Phys. Rev. A*68: 042102(2003).
- [17] P. M. Alsing and G. J. Milburn, *Phys. Rev. Lett.* 91: 180404(2003).
- [18] A. Peres and D. R. Terno, *Rev. Mod. Phys.* 76: 93(2004).
- [19] C. Soo and C. C. Y. Lin, *Int. J. Quant. Inf.* 2: 183(2004).
- [20] P. M. Alsing *et al.*, *J. Optics.* B6: S834(2004).
- [21] Y. Shi, *Phys. Rev. D*70: 105001(2004).
- [22] W. T. Kim and E. J. Son, *Phys. Rev. A*71: 014102(2005).
- [23] M. Czachor, *Phys. Rev. Lett.* 94: 078901(2005).
- [24] I. Fuentes-Schuller and R. B. Mann, *Phys. Rev. Lett.* 95: 120404(2005).
- [25] P. Caban and J. Rembielinski, *Phys. Rev. A*72: 012103(2005).
- [26] P. Kok and S. L. Braunstein, *Int. J. Quant. Inf.* 4: 119(2006).
- [27] J. L. Ball *et al.*, *Phys. Lett.* A359: 550(2006).
- [28] L. Lamata *et al.*, *Phys. Rev. Lett.* 97: 250502(2006).
- [29] L. Lamata *et al.*, *Phys. Rev. A*73: 012335(2006).
- [30] T. F. Jordan *et al.*, *Phys. Rev. A*73: 032104(2006).
- [31] P. M. Alsing *et al.*, *Phys. Rev. A*74: 032326(2006).
- [32] Y. Shi, *Phys. Lett.* B641: 486(2006).
- [33] P. Caban and J. Rembielinski, *Phys. Rev. A*74: 042103(2006).
- [34] T. F. Jordan *et al.*, quant-ph/0608061.
- [35] Y. Ling *et al.*, *J. Phys.* A40: 9025(2007) .
- [36] G. Adesso *et al.*, quant-ph/0701074.
- [37] S. He *et al.*, *J. Phys.* A40: F857(2007).
- [38] S. Weinberg, in *Lectures on Particles and Field Theory*, edited by S. Deser and F. W. Ford((Prentice-Hall, Inc., Englewood Cliffs, N. J., 1964), vol. II of Lectures delivered at Brandeis Summer Institute in Theoretical Physics.