

Quantum Database Searching by a Single Query

DONG PYO CHI ^{*†} and JINSOO KIM ^{*‡}

15 July 1997

Abstract

In this paper we give a quantum mechanical algorithm that can search a database by a single query, when the number of solutions is more than a quarter. It utilizes modified Grover operator of arbitrary phase.

1 Introduction

For $N \in \mathbb{N}$, let $\mathbb{Z}_N = \{0, 1, \dots, N-1\}$ denote the additive cyclic group of order N and consider an arbitrary function $F : \mathbb{Z}_N \rightarrow \mathbb{Z}_2$. The database searching problem is to find some $i \in \mathbb{Z}_N$ such that $F(i) = 1$ under the assumption that such an i exists. We assume that the structure of F is unknown so that it is not possible to obtain a knowledge about F without evaluating it on \mathbb{Z}_N .

Let $t = |\{i \in \mathbb{Z}_N : F(i) = 1\}|$. There is a quantum mechanical algorithm to solve this problem in expected time of order $O(\sqrt{N/t})$, which is optimal up to a multiplicative constant [4, 2, 1]. Especially when $t = N/4$ is known, the original Grover algorithm in [4] can search a solution only by a single query [2]. It uses the π -phase, i.e., marking the states by multiplying $e^{\pi i} = -1$. When $t = N/2$, by changing this phase to $\pi/2$, that is, by marking the states by multiplying $e^{\pi i/2} = i$ and modifying the corresponding diffusion transform according to this phase, the solution can be found with certainty after a single iteration [3].

In this paper, we generalize Grover algorithm for arbitrary phase. When $t \geq N/4$, we give generalized conditional phase and diffusion transform

*Department of Mathematics, Seoul National University, Seoul 151-742, Korea.

†E-mail address: dpchi@math.snu.ac.kr

‡E-mail address: jskim@math.snu.ac.kr

depending on t , and then formulate a quantum mechanical algorithm that solves the database searching problem in a single query.

2 Grover Operator of Arbitrary Phase

Let $\mathcal{B}_N = \{|a\rangle\}_{a \in \mathbb{Z}_2^n} = \{|e_1\rangle, |e_2\rangle, \dots, |e_N\rangle\}$ be the standard basis of an n -qubit quantum register with $N = 2^n$ and \mathcal{H}_N be the corresponding Hilbert space, which represents the state vectors of a quantum system. Let \mathcal{H}_m be an m -dimensional subspace of \mathcal{H}_N spanned by a basis $\mathcal{B}_m = \{|e_{i_1}\rangle, |e_{i_2}\rangle, \dots, |e_{i_m}\rangle\}$. Let l be a positive integer such that $1 \leq l \leq m$.

For $\gamma \in \mathbb{R}$, the *conditional γ -phase transform* on a subspace \mathcal{H}_m , $\mathbf{S}_{F_l, \gamma}^{\mathcal{H}_m} : \mathcal{H}_m \rightarrow \mathcal{H}_m$ is defined by

$$\mathbf{S}_{F_l, \gamma}^{\mathcal{H}_m} |e_{i_k}\rangle = (e^{i\gamma})^{F_l(e_{i_k})} |e_{i_k}\rangle,$$

for $k = 1, 2, \dots, m$, where $i = \sqrt{-1}$. Let $\mathbf{S}_{l, \gamma}^{\mathcal{H}_m}$ denote $\mathbf{S}_{F_l, \gamma}^{\mathcal{H}_m}$, where $F_l(e_{i_k}) = \delta_{i_k i_l}$.

Let $\mathbf{W}_l^{\mathcal{H}_m}$ be any unitary transformation on \mathcal{H}_m satisfying

$$\mathbf{W}_l^{\mathcal{H}_m} |e_{i_l}\rangle = \frac{1}{\sqrt{m}} \sum_{k=1}^m |e_{i_k}\rangle.$$

For example, when m is a power of 2, in a suitably arranged basis we may set $\mathbf{W}_1^{\mathcal{H}_m}$ to be the Walsh-Hadamard transform. When m is not a power of 2, the approximate Fourier transform in [5] can be used.

For $\beta \in \mathbb{R}$, the *β -phase diffusion transform* on a subspace \mathcal{H}_m , $\mathbf{D}_\beta^{\mathcal{H}_m} : \mathcal{H}_m \rightarrow \mathcal{H}_m$ is defined by

$$\mathbf{D}_{\beta, ij}^{\mathcal{H}_m} = \langle e_i | \mathbf{D}_\beta^{\mathcal{H}_m} |e_j\rangle = \begin{cases} \frac{e^{i\beta} - 1}{m} & \text{when } |e_i\rangle, |e_j\rangle \in \mathcal{B}_m \text{ and } i \neq j, \\ 1 + \frac{e^{i\beta} - 1}{m} & \text{when } |e_i\rangle, |e_j\rangle \in \mathcal{B}_m \text{ and } i = j, \\ \delta_{ij} & \text{otherwise.} \end{cases}$$

If we rearrange the basis \mathcal{B}_N so that $\mathcal{B}_m = \{|e_1\rangle, |e_2\rangle, \dots, |e_m\rangle\}$ and represent $\mathbf{D}_\beta^{\mathcal{H}_m}$ by its matrix $\mathbf{D}_\beta^{\mathcal{H}_m} = (\mathbf{D}_{\beta, ij}^{\mathcal{H}_m})$, then we have

$$\begin{aligned} \mathbf{D}_\beta^{\mathcal{H}_m} &= \begin{bmatrix} \mathbf{W}_l^{\mathcal{H}_m} \mathbf{S}_{l, \beta}^{\mathcal{H}_m} \mathbf{W}_l^{\mathcal{H}_m \dagger} & 0 \\ 0 & I \end{bmatrix} \\ &= \begin{bmatrix} I + (e^{i\beta} - 1) P^{\mathcal{H}_m} & 0 \\ 0 & I \end{bmatrix}, \end{aligned}$$

where $P^{\mathcal{H}_m} = (P_{ij}^{\mathcal{H}_m})$ is a projection matrix in \mathcal{H}_m with $P_{ij}^{\mathcal{H}_m} = \frac{1}{m}$. Note that $\mathbf{S}_{F,\gamma}^{\mathcal{H}_m}$ and $\mathbf{D}_\beta^{\mathcal{H}_m}$ are unitary.

Let $\mathbf{G}_{F,\beta,\gamma}^{\mathcal{H}_m} : \mathcal{H}_N \rightarrow \mathcal{H}_N$ be the *Grover operator* of (β, γ) -phase in \mathcal{H}_m defined by

$$\mathbf{G}_{F,\beta,\gamma}^{\mathcal{H}_m} = \mathbf{D}_\beta^{\mathcal{H}_m} \mathbf{S}_{F,\gamma}^{\mathcal{H}_m}.$$

When $\beta = \gamma$ set $\mathbf{G}_{F,\gamma}^{\mathcal{H}_m} = \mathbf{G}_{F,\beta,\gamma}^{\mathcal{H}_m}$. For simplicity, we shall assume that $m = N$ and omit the superscript \mathcal{H}_m .

Let

$$\begin{aligned} A &= \{e_j \in \mathbb{Z}_2^n | F(e_j) = 1\}, \\ B &= \{e_j \in \mathbb{Z}_2^n | F(e_j) = 0\}, \end{aligned}$$

and let $t = |A|$. For $k, l \in \mathbb{C}$ such that $t|k|^2 + (N-t)|l|^2 = 1$, define

$$|\psi(k, l)\rangle = \sum_{e_j \in A} k|e_j\rangle + \sum_{e_j \in B} l|e_j\rangle.$$

Lemma 1 For $\beta, \gamma \in \mathbb{R}$, let

$$|\psi(k_j, l_j)\rangle = \mathbf{G}_{F,\beta,\gamma}^j |\psi(k_0, l_0)\rangle.$$

Then after applying $j+1$ Grover operator of (β, γ) -phase, we have

$$\begin{cases} k_{j+1} = \frac{(e^{i\beta} - 1)t + N}{N} e^{i\gamma} k_j + \frac{(e^{i\beta} - 1)(N-t)}{N} l_j, \\ l_{j+1} = \frac{(e^{i\beta} - 1)t}{N} e^{i\gamma} k_j + \frac{(e^{i\beta} - 1)(N-t) + N}{N} l_j. \end{cases} \quad (2.1)$$

Theorem 1 Assume that $k_0 = l_0$ and $\beta, \gamma \in [0, 2\pi]$. Then $l_1 = 0$ if and only if $\frac{N}{4} \leq t \leq N$ and $\beta = \gamma = \cos^{-1} \left(1 - \frac{N}{2t}\right)$. In this case, $k_1 = (e^{i\gamma} - 1)k_0$.

Proof. By (2.1), we get

$$\begin{cases} k_1 = \left[(e^{i\beta} - 1)(e^{i\gamma} - 1) \frac{t}{N} + e^{i\gamma} + e^{i\beta} - 1 \right] k_0, \\ l_1 = \left[(e^{i\beta} - 1)(e^{i\gamma} - 1) \frac{t}{N} + e^{i\beta} \right] l_0. \end{cases} \quad (2.2)$$

By considering the imaginary part of $e^{-i\beta\frac{l_1}{t_0}}$, the equation

$$\frac{t}{N} \{(1 - \cos \beta) \sin \gamma + (\cos \gamma - 1) \sin \beta\} = 0$$

is equivalent to

$$\frac{1 - \cos \beta}{\sin \beta} = \frac{1 - \cos \gamma}{\sin \gamma}. \quad (2.3)$$

Considering the real part of $e^{-i\beta\frac{l_1}{t_0}}$, from (2.3) it follows that the equation

$$\frac{t}{N} \{(1 - \cos \beta)(1 - \cos \gamma) + \sin \beta \sin \gamma\} - 1 = 0$$

is equivalent to

$$\cos \beta = \cos \gamma = 1 - \frac{N}{2t}.$$

Again by (2.3), we get $\beta = \gamma$ and $t \geq \frac{N}{4}$. Furthermore, by (2.2) we obtain $k_1 = (e^{\gamma} - 1)k_0$. This completes the proof. \square

For the case of π -phase in [4], $-\mathbf{D}_\pi = -I + 2P$ is an inversion about average operation and we have

$$-\mathbf{D}_\pi |\psi(k, l)\rangle = \left| \psi\left(-\frac{N-2t}{N}k + \frac{2(N-t)}{N}l, \frac{N-2t}{N}l + \frac{2t}{N}k\right) \right\rangle.$$

In this case, there is an explicit closed-form formula for k_j and l_j ;

$$\begin{cases} k_j = (-1)^j \frac{1}{\sqrt{t}} \sin((2j+1)\theta), \\ l_j = (-1)^j \frac{1}{\sqrt{N-t}} \cos((2j+1)\theta), \end{cases}$$

for $j = 0, 1, \dots$, where the angle θ is defined so that $\sin^2 \theta = \frac{t}{N}$ [2]. Especially when $t = N/4$, we have $l_1 = 0$.

Grover operator of $\frac{\pi}{2}$ -phase was used in [3]. Since

$$\mathbf{D}_{\frac{\pi}{2}} |\psi(k, l)\rangle = \left| \psi\left(\frac{(i-1)t+N}{N}k + \frac{(i-1)(N-t)}{N}l, \frac{(i-1)(N-t)+N}{N}l + \frac{(i-1)t}{N}k\right) \right\rangle,$$

when $t = \frac{N}{2}$, we have

$$\mathbf{G}_{F, \frac{\pi}{2}} |\psi(k, k)\rangle = |\psi((i-1)k, 0)\rangle.$$

By Theorem 1, When $t \in [N/4, N]$, we have

$$\mathbf{G}_{F, \gamma} |\psi(k_0, k_0)\rangle = |\psi((e^{\gamma} - 1)k_0, 0)\rangle,$$

where the phase γ is defined by $\gamma = \cos^{-1}\left(1 - \frac{N}{2t}\right)$.

References

- [1] C. H. Bennett, E. Bernstein, G. Brassard and U. Vazirani, *Strengths and weaknesses of quantum computing*, Los Alamos e-print quant-ph/9701001, 1997.
- [2] M. Boyer, G. Brassard, P. Høyer and A. Tapp, *Tight bounds on quantum searching*, Proceedings of the Fourth Workshop on Physics and Computation, New England Complex Systems Institute, 1996, pp. 36–43; Los Alamos e-print quant-ph/9605034, 1996.
- [3] G. Brassard and P. Høyer, *An exact quantum polynomial-time algorithm for Simon's problem*, Proceedings of the Fifth Israeli Symposium on Theory of Computing and Systems, 1997, (to appear); Los Alamos e-print quant-ph/9704027, 1997.
- [4] L. K. Grover, *Quantum mechanics helps in searching for a needle in a haystack*, Phys. Rev. Lett., (to appear); Los Alamos e-print quant-ph/9605043, 1996.
- [5] A. Yu. Kitaev, *Quantum measurements and the abelian stabilizer problem*, Los Alamos e-print quant-ph/9511026, 1995.