

Relativistically Extended Modification of the Schrödinger Equation

Waldemar Puszkarz*

*Department of Physics and Astronomy,
University of South Carolina,
Columbia, SC 29208*

(February 28, 1999)

Abstract

We propose a nonlinear modification of the Schrödinger equation that possesses the main properties of this equation such as the Galilean invariance, the weak separability of composite systems, and the homogeneity in the wave function. The modification is derived from the relativistic relation between the energy and momentum of free particle and as such it is the best relativistic extension of the Schrödinger equation that preserves the properties in question. The only change it effectively entails in the Schrödinger equation involves the conserved probability current. It is pointed out that it partially retains the linear superposition principle and can be used to model the process of decoherence.

*Electronic address: puszkarz@cosm.sc.edu

Since its discovery in 1926, the Schrödinger equation,

$$i\hbar\frac{\partial\Psi}{\partial t} = -\frac{\hbar^2}{2m}\Delta\Psi + V\Psi, \quad (1)$$

has been subjected to a very thorough examination, both experimental and theoretical. In the very same year, Madelung showed that one can represent this equation as a system of two nonlinear equations,

$$\hbar\frac{\partial R^2}{\partial t} + \frac{\hbar^2}{m}\vec{\nabla} \cdot (R^2\vec{\nabla} S) = 0, \quad (2)$$

$$\frac{\hbar^2}{m}\Delta R - \frac{\hbar^2}{m}R(\vec{\nabla} S)^2 - 2\hbar R\frac{\partial S}{\partial t} - 2RV = 0, \quad (3)$$

for the amplitude R and the phase S of the wave function $\Psi = Re^{iS}$. They form the so-called hydrodynamic representation of the Schrödinger equation and can be obtained as the imaginary and real part of this equation, correspondingly. The first of these equations is the continuity equation with the probability current

$$\vec{j}_{Sch} = \frac{\hbar^2}{m}R^2\vec{\nabla} S \quad (4)$$

and the probability density $\rho = R^2$. However, it is the other equation that is crucial for determining the stationary states of quantum-mechanical systems defined by the condition $\partial R^2/\partial t = 0$.

Let us recall that heuristically the free Schrödinger equation can be derived from the relativistic dispersion relation between the energy and momentum of a free particle

$$E = c\sqrt{p^2 + (mc)^2} = mc^2\sqrt{1 + \left(\frac{p}{mc}\right)^2}. \quad (5)$$

One obtains it by discarding the rest energy term from the leading approximation to (5),

$$E = mc^2 + \frac{p^2}{2m}, \quad (6)$$

and applying the first quantization procedure. To account for a wider range of physical situations one can subsequently add the potential term to the RHS of (6). This leads directly to (1).

It is the purpose of this paper to derive a relativistic extension of the Schrödinger equation which would stem from a closer approximation to (5) than the truncated first order approximation (6) and which in fact would be a nonlinear modification of the equation concerned. The interest in nonlinear modifications of the fundamental equation of quantum mechanics has recently been greatly stimulated by the work of Weinberg [1, 2] that proposed a relatively general framework for nonlinear modifications of quantum theory.¹

The Schrödinger equation is homogeneous of degree one in the wave function, however it is not entirely clear if this property is underlied by or entails any physically important principle. It is thus conceivable that this property is an accidental feature of the Schrödinger equation and therefore disposing of it in a nonlinear modification of this equation will most likely not cause any serious physical consequences. Nevertheless, as pointed out in [3], nonhomogeneous modifications of the

¹For a broader exposition of various aspects of nonlinear quantum mechanics see [3]. More elaborate, though still far from comprehensive a list of approaches and motivations to modify this equation can be found in [4].

Schrödinger equation suffer from the ambiguity in the definition of the energy functional. Although this occurs also in homogenous nonlinear variants of this equation, there exists a class of homogenous modifications for which the quantum-mechanical energy functional defined as the mean value of the corresponding Hamiltonian operator equals the energy defined within the Lagrangian framework as a certain conserved quantity [3]. This indicates that the homogeneity is necessary but not sufficient to guarantee the uniqueness of the energy functional. Weinberg has elevated the discussed property of the Schrödinger equation to one of the fundamental assumptions of his original generalization of the linear framework of quantum mechanics [1, 2]. It was his hope, particularly expressed in [2], that the homogeneity of nonlinear variants of this equation might entail their separability.² Even if this is true in his modification and was shown to be valid in some others [4, 5], one can find examples of nonlinear variants of the Schrödinger equation that contradict this thesis [6].

The separability of noninteracting subsystems [7, 4] has much more profound and better understood physical implications. Mathematically, it means that if $\Psi_1(\vec{x}_1, t)$ and $\Psi_2(\vec{x}_2, t)$ are wave functions for such subsystems-particles subjected to potentials $V_i(\vec{x}_i, t)$, ($i = 1, 2$) then it is their product $\Psi_1(\vec{x}_1, t)\Psi_2(\vec{x}_2, t)$ that represents the wave function for a system consisting of these particles in the presence of a potential

$$V(\vec{x}_1, \vec{x}_2, t) = V_1(\vec{x}_1, t) + V_2(\vec{x}_2, t). \quad (7)$$

The essential element here is that the subsystems are uncorrelated as expressed by the fact that the total wave function is a product of $\Psi_1(\vec{x}_1, t)$ and $\Psi_2(\vec{x}_2, t)$. The additive form of the total potential ensures that no interaction between the subsystems occurs, so they remain uncorrelated during the evolution.³ Such an interaction may however appear in nonlinear modifications of the Schrödinger equation even if the form of the potential itself does not imply that. This is due to a coupling that a nonlinear term usually causes between $\Psi_1(\vec{x}_1, t)$ and $\Psi_2(\vec{x}_2, t)$. As a result, even in the absence of forces the very existence of one of the particles affects the evolution of the other, clearly violating causality. Needless to say, this is not a physically sound situation. This particular feature of the Schrödinger equation distinguishes it from the Klein-Gordon equation. One arrives at the latter from (5) upon squaring it and employing the first quantization. The time derivative of this equation introduces a coupling between $\Psi_1(\vec{x}_1, t)$ and $\Psi_2(\vec{x}_2, t)$, rendering the equation inseparable in the sense discussed. Another physically significant difference between these equations is in the probability density which can be negative for the Klein-Gordon equation.

To shed more light on this issue let us demonstrate the separability of noninteracting systems for the Schrödinger equation in its hydrodynamic formulation. We are considering a physical system made up of two noninteracting subsystems in the sense of (7). We will show that one can write a solution of the Schrödinger equation for this system in the form of the product of wave functions for individual subsystems, that is $\Psi(x_1, x_2, t) = \Psi_1(x_1, t)\Psi_2(x_2, t) = R_1(x_1, t)R_2(x_2, t)\exp\{i(S_1(x_1, t) + S_2(x_2, t))\}$. The Schrödinger equation for the total system, assuming that the subsystems have the same mass m , reads now

$$\hbar \frac{\partial R_1^2 R_2^2}{\partial t} + \frac{\hbar^2}{m} \left\{ (\vec{\nabla}_1 + \vec{\nabla}_2) \cdot [R_1^2 R_2^2 (\vec{\nabla}_1 S_1 + \vec{\nabla}_2 S_2)] \right\} =$$

²As stated in [2]: “The problem of dealing with separated systems has led other authors to limit possible nonlinear terms in the Schrödinger equation to a logarithmic form;” and “The homogeneity assumption (2) makes this unnecessary.”

³The discussed separability is sometimes called the weak separability as opposed to the strong separability. The latter can be formulated also for entangled systems in some nonlinear generalizations of the Schrödinger equation [8].

$$\begin{aligned} & \hbar R_2^2 \frac{\partial R_1^2}{\partial t} + \hbar R_1^2 \frac{\partial R_2^2}{\partial t} + \frac{\hbar^2}{m} R_2^2 \vec{\nabla}_1 \cdot (R_1^2 \vec{\nabla}_1 S_1) + \frac{\hbar^2}{m} R_1^2 \vec{\nabla}_2 \cdot (R_2^2 \vec{\nabla}_2 S_2) = \\ & R_1^2 R_2^2 \hbar \left\{ \frac{1}{R_1^2} \frac{\partial R_1^2}{\partial t} + \frac{\hbar^2}{m} \frac{1}{R_1^2} \vec{\nabla}_1 \cdot (R_1^2 \vec{\nabla}_1 S_1) + \left[\hbar \frac{1}{R_2^2} \frac{\partial R_2^2}{\partial t} + \frac{\hbar^2}{m} \frac{1}{R_2^2} \vec{\nabla}_2 \cdot (R_2^2 \vec{\nabla}_2 S_2) \right] \right\} = 0 \end{aligned} \quad (8)$$

and

$$\begin{aligned} & \frac{\hbar^2}{m} (\Delta_1 + \Delta_2) R_1 R_2 - 2\hbar R_1 R_2 \frac{\partial (S_1 + S_2)}{\partial t} - \frac{\hbar^2}{m} R_1 R_2 (\vec{\nabla}_1 S_1 + \vec{\nabla}_2 S_2)^2 - \\ & (V_1 + V_2) R_1 R_2 = \frac{\hbar^2}{m} R_2 \Delta_1 R_1 + \frac{\hbar^2}{m} R_1 \Delta_2 R_2 - 2\hbar R_1 R_2 \frac{\partial S_1}{\partial t} - 2\hbar R_1 R_2 \frac{\partial S_2}{\partial t} \\ & + \frac{\hbar^2}{m} R_1 R_2 (\vec{\nabla}_1 S_1)^2 + \frac{\hbar^2}{m} R_1 R_2 (\vec{\nabla}_2 S_2)^2 - V_1 R_1 R_2 - V_2 R_1 R_2 = \\ & R_1 R_2 \left\{ \left[\frac{\hbar^2}{m} R \Delta_1 R_1 - 2\hbar \frac{\partial S_1}{\partial t} + \frac{\hbar^2}{m} (\vec{\nabla}_1 S_1)^2 - V_1 \right] + \left[\frac{\hbar^2}{m} \frac{\Delta_2 R_2}{R_2} - 2\hbar \frac{\partial S_2}{\partial t} + \right. \right. \\ & \left. \left. \frac{\hbar^2}{m} (\vec{\nabla}_2 S_2)^2 - V_2 \right] \right\} = 0. \end{aligned} \quad (9)$$

Implicit in the derivation of these equations is the fact that $\vec{\nabla}_1 f_1 \cdot \vec{\nabla}_2 g_2 = 0$, where f_1 and g_2 are certain scalar functions defined on the configuration space of particle 1 and 2, correspondingly. What we obtained is a system of two equations, each one consisting of terms (in square brackets) that pertain to only one of the subsystems. By dividing the first equation by $R_1^2 R_2^2$ and the second by $R_1 R_2$ one completes the separation of the Schrödinger equation and shows that indeed the product of wave functions of subsystems provides the wave function for the total system.

We would like the modified Schrödinger equation to possess the property of separability of composite noninteracting systems in the sense described and to be Galilean invariant. It should also ensure a non-negative probability density emerging in the continuity equation. Moreover, we require that the probability current of this equation be conserved. These properties are certainly of physical relevance. As we will see, our approach will result in a homogeneous equation by way of construction, exactly as the Schrödinger equation is rendered homogenous when derived in the manner outlined above.

To proceed, let us note that by expanding (5) in a series of p^2 and applying the first quantization we obtain terms $\lambda_c^{2n} \Delta^n \Psi$, where $\lambda_c = \frac{\hbar}{mc}$ is the Compton wavelength. Employing the Schrödinger-Madelung representation, we observe that the terms concerned contain growing with n a number of terms involving derivatives of R and S . The essence of our approach is to extract from them the terms that would satisfy our criteria. It is easy to check that the following recurrence relation emerges for $n \geq 1$

$$\begin{aligned} \Delta^n \Psi &= (A_n + iB_n) \exp(iS) = \left\{ [(\Delta - (\vec{\nabla} S)^2) A_{n-1} - (\Delta S + 2\vec{\nabla} S \cdot \vec{\nabla}) B_{n-1}] + \right. \\ & \left. i [(\Delta - (\vec{\nabla} S)^2) B_{n-1} + (\Delta S + 2\vec{\nabla} S \cdot \vec{\nabla}) A_{n-1}] \right\} \exp(iS), \end{aligned} \quad (10)$$

where $A_0 = R$ and $B_0 = 0$. The first part of this equality is trivially valid for $n = 0$ as well. It should be noted that by expanding (5) in a series of $\Delta^n \Psi$ one obtains nonseparable terms, the only exception being the leading $n = 1$ term that gives rise to the Schrödinger equation. Moreover, such an expansion would not be Galilean invariant. To ensure that both of these properties are

maintained in our modification, it is more convenient if not necessary to work in the Schrödinger-Madelung representation. To facilitate more general considerations, let us first examine in more detail the lowest order terms emerging from (10). These are

$$A_1 = \Delta R - (\vec{\nabla} S)^2 R = [\Delta - (\vec{\nabla} S)^2] R, \quad (11)$$

$$B_1 = (\Delta S + 2\vec{\nabla} S \cdot \vec{\nabla}) R = \frac{\vec{\nabla} \cdot (R^2 \vec{\nabla} S)}{R}, \quad (12)$$

and

$$A_2 = [\Delta - (\vec{\nabla} S)^2]^2 R - (\Delta S + 2\vec{\nabla} S \cdot \vec{\nabla}) \frac{\vec{\nabla} \cdot (R^2 \vec{\nabla} S)}{R}, \quad (13)$$

$$B_2 = [\Delta - (\vec{\nabla} S)^2] \frac{\vec{\nabla} \cdot (R^2 \vec{\nabla} S)}{R} + [\Delta S + 2\vec{\nabla} S \cdot \vec{\nabla}] [\Delta - (\vec{\nabla} S)^2] R. \quad (14)$$

Among them, only B_1 and ΔB_1 are separable, although not Galilean invariant. RB_1 is proportional to the divergence of the probability current of the Schrödinger equation, but the continuity equation (2) is Galilean invariant only as a result of cancellation of noninvariant contributions coming from its both terms. To see this, let us recall that under the Galilean transformation,

$$t = t', \quad \vec{x} = \vec{x}' + \vec{v}t, \quad (15)$$

the amplitude of wave function does not change, the phase changes according to

$$S(t, \vec{x}) = S'(t', \vec{x}') + m\vec{v} \cdot \vec{x}' + \frac{m}{2}\vec{v}^2 t', \quad (16)$$

whereas the operators that affect these quantities in (2-3) transform as

$$\vec{\nabla} = \vec{\nabla}', \quad \frac{\partial}{\partial t} = \frac{\partial}{\partial t'} - \vec{v} \cdot \vec{\nabla}'. \quad (17)$$

As one can deduce from the general recurrence rule (10) and the last two equations, only the terms ΔA_n and ΔB_n could produce Galilean invariant and separable terms from the terms that became separable in a preceding iteration, but were not necessarily Galilean invariant. The conditions of separability and Galilean invariance are stringent enough to select only one term that maintains separability in successive iterations of Δ and at the same time yields terms that are Galilean invariant. It is

$$\alpha = B_1 = R\Delta S + 2\vec{\nabla} S \cdot \vec{\nabla} R. \quad (18)$$

Let us note that even if $\vec{\nabla} S \cdot \vec{\nabla} R$ is separable, it is no longer so when acted upon by $R\Delta$ as it gives rise to nonseparable terms

$$R\Delta(\vec{\nabla} S \cdot \vec{\nabla} R) = \sum_i R\vec{\nabla} \cdot [\partial_i S \vec{\nabla} \cdot (\partial_i R) + \partial_i R \vec{\nabla} \cdot (\partial_i S)]. \quad (19)$$

The same applies to other terms of the form $R\Delta(\vec{\nabla} \Delta^{2n} S \cdot \vec{\nabla} R)$ ($n \geq 1$) that emerge in higher iterations. For this reason they will be discarded. Now, with the second term of (18) discarded, $R\Delta\alpha = \vec{\nabla} \cdot (R^2 \vec{\nabla}^3 S) + R\Delta R\Delta S$, but since $R\Delta R\Delta S$ is not separable, it should be excluded from our

construction as should other terms of the form $R\Delta R\Delta^n S$ ($n \geq 1$) appearing in subsequent iterations. Repeating this procedure for $R\Delta^n \alpha$ ($n \geq 1$) one observes that it effectively produces $\vec{\nabla} \cdot (R^2 \vec{\nabla} \Delta^n S)$.

The multiplication of all terms of the expansion in question by R is required in order to arrive at the correct form of the continuity equation. For the term B_1 this has already been pointed out. Namely, it is $RB_1 = R\alpha$ and not B_1 itself that gives rise to the divergence of the probability current. In the same manner, the terms $R\Delta^n \alpha$ ($n \geq 1$) contribute to the divergence of a new probability current in the modified continuity equation. We see that the only change our construction of the relativistically extended Schrödinger equation brings about is in this current, or, what amounts to the same, in the continuity equation of the Schrödinger-Madelung system (2-3). One can write this equation as

$$\hbar \frac{\partial R}{\partial t} = mc^2 \lambda_c^2 \sum_{n=0}^{\infty} c_n (\lambda_c^2 \Delta)^n \alpha, \quad (20)$$

where c_n are the coefficients of the expansion. Multiplying both sides of it by R one arrives at

$$\frac{\partial R^2}{\partial t} = \frac{2\hbar}{m} \sum_n \lambda_c^{2n} c_n R \Delta^n \alpha = \frac{2\hbar}{m} \sum_n \lambda_c^{2n} c_n \vec{\nabla} \cdot (R^2 \vec{\nabla} \Delta^n S) = \frac{2\hbar}{m} \vec{\nabla} \cdot \left[R^2 \vec{\nabla} \sum_{n=0}^{\infty} \lambda_c^{2n} c_n \Delta^n S \right]. \quad (21)$$

The last sum simplifies to $G(\lambda_c^2 \Delta)S$, where

$$G(x) = \frac{\sqrt{1-x^2} - 1}{x^2} = -2 \frac{\sin^2(\frac{\sin^{-1} x}{2})}{x^2}, \quad (22)$$

the second equality being a straightforward consequence of relations $\sqrt{1-x^2} = \cos(\sin^{-1} x)$ and $\cos y - 1 = -2 \sin^2 \frac{y}{2}$. Eventually, the continuity equation reads

$$\frac{\partial R^2}{\partial t} + \vec{\nabla} \cdot \vec{j}_{RM} = 0, \quad (23)$$

where

$$\vec{j}_{RM} = -\frac{2\hbar}{m} R^2 \vec{\nabla} (G(\lambda_c^2 \Delta)S). \quad (24)$$

Alternatively, one can write the modified Schrödinger equation as

$$i\hbar \frac{\partial \Psi}{\partial t} = \frac{\hbar^2}{2m} \left[-\Delta - \frac{i}{\rho} (\vec{\nabla} \cdot \vec{j}_{RM}) + \frac{i}{\rho} (\vec{\nabla} \cdot \vec{j}_{Sch}) \right] \Psi + V\Psi, \quad (25)$$

which is equivalent to

$$i\hbar \frac{\partial \Psi}{\partial t} = -\frac{\hbar^2}{2m} \left[\Delta + \frac{i}{\rho} \vec{\nabla} \cdot [\rho \vec{\nabla} (2G(\lambda_c^2 \Delta) - 1)S] \right] \Psi + V\Psi. \quad (26)$$

In general, the only notable exception to this rule being the modification of Białynicki-Birula and Mycielski, the nonlinear modifications of the Schrödinger equation do not have the classical limit in the sense of the Ehrenfest theorem. The discussed variant of the modification presented is not exempted from this either. The nonlinear terms it introduces lead to some corrections to the Ehrenfest relations. We will now work out these corrections. For a general observable A one finds that

$$\frac{d}{dt} \langle A \rangle = \frac{d}{dt} \langle A \rangle_L + \frac{d}{dt} \langle A \rangle_{NL}, \quad (27)$$

where the nonlinear contribution is due to $H_{NL} = H_R + iH_I$, H_R and H_I representing the real and imaginary part of H_{NL} , respectively. The brackets $\langle \rangle$ denote the mean value of the quantity embraced. Specifying A for the position and momentum operators one obtains the general form of the modified Ehrenfest relations [6]

$$m \frac{d}{dt} \langle \vec{r} \rangle = \langle \vec{p} \rangle + I_1, \quad (28)$$

$$\frac{d}{dt} \langle \vec{p} \rangle = - \langle \vec{\nabla} V \rangle + I_2, \quad (29)$$

where

$$I_1 = \frac{2m}{\hbar} \int dV \vec{r} \rho H_I, \quad (30)$$

$$I_2 = \int dV \rho \left(2H_I \vec{\nabla} S - \vec{\nabla} H_R \right). \quad (31)$$

In the derivation of the last formula it was assumed that $\int dV \vec{\nabla} (\rho H_I) = 0$ which for homogeneous modifications as the one in question seems to be well justified [6]. For the extension concerned these integrals are found to be

$$I_1 = -\hbar \int dV \vec{r} \vec{\nabla} \cdot \left[\rho \vec{\nabla} \left(2G(\lambda_c^2 \Delta) - 1 \right) S \right], \quad (32)$$

$$I_2 = -\frac{\hbar^2}{m} \int dV \vec{\nabla} \cdot \left[\rho \vec{\nabla} \left(2G(\lambda_c^2 \Delta) - 1 \right) S \right] \vec{\nabla} S. \quad (33)$$

The presented modification is the best relativistic approximation to the Schrödinger equation that shares with it the properties of separability for composed noninteracting systems, the Galilean invariance, and the homogeneity in the wave function. Moreover, the probability density of equation (25) (or (26)) is a non-negative function unlike that of the Klein-Gordon equation that originates within the same framework. This is what distinguishes the modification discussed from other nonlinear variants of the Schrödinger equation [7, 9, 5] that encompass the same properties, but make additional assumptions or employ principles beyond those specified here. These essentially are necessary to keep the number of new terms at a reasonably manageable level. Such a problem does not arise here as, in principle, there is only one new term, which certainly adds to the attractiveness of this proposal. What is even more, our modification does not introduce any new constants. This makes it truly unique in this respect: all other nonlinear modifications of the Schrödinger equation do involve some new parameters.

The modification in question supports stationary states of the linear Schrödinger equation for which $S = -Et/\hbar + const$, without affecting their energy E . Some other characteristic solutions to it include the plane wave, but interestingly enough also ordinary Gaussian wave packets and coherent states. Since both of these are the result of superposing more elementary wave functions, each of which is a solution to the nonlinear equation, this means that similarly as the Schrödinger equation our modification allows, although only partially, for the linear superposition principle. It is easy to understand how this comes about here. The third spatial derivative of the phase of a wave-packet for simplicity in one dimension ($\hbar = 1$),

$$S = \frac{m t x^2}{2(t^2 + t_0^2)} - \frac{1}{2} \arctan \frac{t}{t_0}, \quad (34)$$

vanishes as do its higher order derivatives. The current correction of (26) starts with $\vec{\nabla}^3 S$ in the expansion, which means that in the case of wave packets the new term does not affect the Schrödinger equation at all. The same holds true for coherent states for which $\Delta S = 0$. The nonlinear modifications endowed with the discussed property that we chose to call the weak nonlinearity are rare (see [5, 10] for the only other known examples of this kind).

As noted, the modification proposed employs only the most general and universal assumptions. Because of that, it seems to be perfectly suited to describe phenomena of some universal nature. As pointed out in [11], one of the most important phenomena in the quantum realm, the process of decoherence responsible for rendering classical features of the ultimately quantum world, can be modeled by Hamiltonians involving imaginary terms. This is exactly the case of our modification. It introduces only such terms and it does so in a minimal model-independent manner: the imaginary terms that play the role of “optical” potentials emerge naturally, being an integral part of the modification and do not have to be put by hand. An attractive consequence of this is that the process in question can be thought of as caused by relativistic nonlinear corrections. One can hope that the modification proposed will provide a good model to investigate this process in greater detail.

Acknowledgments

I would like to thank Professor Paweł O. Mazur for bringing to my attention the work of Professor A. Staruszkiewicz that started my interest in nonlinear modifications of the Schrödinger equation and Dr. G. A. Goldin for his interest in this paper. This work was partially supported by the NSF grant No. 13020 F167 and the ONR grant R&T No. 3124141.

References

- [1] S. Weinberg, *Ann. Phys. (N. Y.)* **194**, 336 (1989).
- [2] S. Weinberg, *Phys. Rev. Lett.* **62**, 485 (1989).
- [3] W. Puszkarz, *Energy Ambiguity in Nonlinear Quantum Mechanics*, preprint quant-ph/9802001.
- [4] G. A. Goldin and G. Svetlichny, *J. Math. Phys.* **35**, 3322, (1995).
- [5] W. Puszkarz, *Higher Order Modification of the Schrödinger Equation*, preprint quant-ph/9710007.
- [6] W. Puszkarz, *Nonlinear Phase Modification of the Schrödinger Equation*, preprint quant-ph/9710010.
- [7] I. Białynicki-Birula and J. Mycielski, *Ann. Phys. (N. Y.)* **100**, 62 (1976).
- [8] M. Czachor, *Phys. Rev. A* **57**, 4122 (1998).
- [9] H.-D. Doebner and G. A. Goldin, *Phys. Lett.* **162A**, 397 (1992); *J. Phys. A* **27**, 1771 (1994).
- [10] A. Staruszkiewicz, *Acta Phys. Pol.* **B14**, 907 (1983).
- [11] M. B. Mensky, *Phys. Lett. A* **196**, 159 (1995).