## Fabrication of Magnetic Charge from Excited States of H-atom

## T. PRADHAN

Institute of Physics, Bhubaneswar-751005, INDIA

PACS - 14.80 Hv

## Abstract

It is shown that the excited states of hydrogen atom in a uniform electric field (Stark States) posess magnetic charge whose magnitude is given by a Dirac-Saha type relation:

$$\frac{eg}{\hbar c} = \sqrt{3}n$$

An experiment is proposed to fabricate such states and to detect their magnetic charge.

Dirac [1] incorporated magnetic charge (monopole) in electrodynamics by introducing a string singularity in the vector potential and showed that its strength is given by the relation

$$\frac{eg}{\hbar c} = \frac{n}{2}, \ n = 1, 2, 3...$$
(1)

This relation was independently obtained by Saha [2] and Wilson [3] by quantizing the angular momentum of a two-body system consisting of a magnetic point charge and an electric point charge. Existence of magnetic charge is also a feature of certain non-abelian gauge theories [4]. Although numerous experimental searches [5] have been undertaken to detect them, none have been found so far. In this note we show that it is possible to fabricate magnetically charged states from the excited states of the hydrogen atom by putting it in a uiform external electrostatic field and detect their magnetic charge by a simple experiment.

If magnetic charges exist, Maxwell equations take the form:

$$\vec{\nabla}.\vec{E} = \rho^{(e)}, \quad \vec{\nabla} \times \vec{B} - \frac{\partial \vec{E}}{\partial t} = \vec{J}^{(e)}$$

$$\vec{\nabla}.\vec{B} = \rho^{(m)}, \quad \vec{\nabla} \times \vec{E} + \frac{\partial \vec{B}}{\partial t} = -\vec{J}^{(m)}$$
(2)

where the superscripts (e) and (m) on  $\vec{J}$  and  $\rho$  stand for electric and magnetic densities. Our contention is that analogous to the relation

$$\vec{\mu} = \frac{1}{2} \int d^3 r (\vec{r} \times \vec{J}^{(e)}(\vec{r}) \tag{3}$$

between magnetic dipole moment  $\vec{\mu}$  and electric current  $\vec{J}^{(e)}$ , there a the relation

$$\vec{d} = \frac{1}{2} \int d^3 r [\vec{r} \times \vec{J}^{(m)}(\vec{r})] \tag{4}$$

between the permanent electric dipole moment  $\vec{d}$  and the magnetic current  $\vec{J}^{(m)}$ . In quantum mechanics this relation takes the form

$$<\vec{d}>=\frac{1}{2}\int d^{3}r \;\psi^{*}(\vec{r})[\vec{r}\times\vec{J}^{(m)}(\vec{r})]\psi(\vec{r})$$
 (5)

Since  $[\vec{r} \times \vec{J}^{(m)}(\vec{r})]$  has odd parity,  $\langle \vec{d} \rangle$  will vanish since in general  $\psi(\vec{r})$  has definite parity. However, excited states of hydrogen atom do not have definite parity on account of l-degeneracy. These states have permanent electric dipole moment.

It may be noted here that the electric dipole moment is usually defined as

$$\vec{d} = \int d^3 r \ \vec{r} \rho^{e)}(\vec{r}) \tag{6}$$

From Maxwell equations (2), this can be written as

$$d_i = \int d^3r \ r_i \partial_j E_j = -\int d^3r \ E_i + \int d^3r \ \partial_j (r_i E_j) \tag{7}$$

The second term on the right can be converted into a surface integral which for the surface at infinity would vanish on account of the fact that the expectation value of electric field  $\vec{E}$  produced by the charge distribution of the hydrogen atom vanishes exponentially at infinity. Similarly the definition (4) for the electric dipole moment can be written as

$$d_{i} = \frac{1}{2} \int d^{3}r(\vec{r} \times \vec{J}^{(m)}) = \frac{1}{2} \int d^{3}r \ r_{j}(-\partial_{i}E_{j} + \partial_{j}E_{i}) = -\int d^{3}rE_{i} + \frac{1}{2} \int d^{3}r[\partial_{j}(r_{j}E_{i}) - \partial_{i}(r_{j}E_{j})]$$

$$(8)$$

in the static limit i.e.  $\frac{\partial B}{\partial t} = 0$ . Since the surface integrals vanish for hydrogen atom, we have the identity

$$\int d^3r \ \vec{r}\rho^{(e)}(\vec{r}) = \frac{1}{2} \int d^3r [\vec{r} \times \vec{J}^{(m)}(\vec{r})]$$
(9)

Physically what this means is that the motion of the electron in the hydrogen atom which gives rise to a permanent electric dipole moment can be considered to have generated a magnetic current in terms of which this dipole moment can be expressed.

; From parity and dimensional considerations the magnetic current  $\vec{J}^{(m)}(\vec{r})$  can be written as

$$\vec{J}^{(m)}(\vec{r}) = g \frac{\vec{L}}{mcr} \tag{10}$$

where  $\vec{L}$  is the orbital angular momentum and g is the magnetic charge. The quantum machenical electric dipole moment operator in that case can be written as

$$\vec{d} = \frac{1}{2} \frac{\vec{r} \times \vec{L}}{mcr} = -\frac{i\hbar g}{2 \times 2mcr} (\vec{r} \times \overline{\nabla})$$

$$where \ a\overline{\nabla}b = a(\vec{\nabla}b) - (\vec{\nabla}a)b$$
(11)

When the hydrogen atom is placed in a static uniform electric field  $(0, 0, \mathcal{E})$  the interaction energy operator  $H_z$  has the form

$$H_I = \vec{d}.\vec{\mathcal{E}} = \frac{-ig\lambda\mathcal{E}}{2\times2} [x^2 + y^2)\frac{\overline{\partial}}{\partial z} - z(x\frac{\overline{\partial}}{\partial x} - y\frac{\overline{\partial}}{\partial y})]$$
(12)

where  $\lambda = \frac{\hbar}{mc}$  is the compton wavelength of the electron. In parabolic co-ordinates

$$H_I = \frac{-ig\lambda\mathcal{E}}{2} \frac{\xi\eta}{\xi+\eta} \left(\frac{\overline{\partial}}{\partial\xi} - \frac{\overline{\partial}}{\partial\eta}\right)$$
(13)

The Schroedinger equation takes the form

$$(H_0 + H_I)\Phi = E\Phi = (E_0 + \Delta E)\Phi \tag{14}$$

with

$$H_0 = \frac{-\hbar^2}{2m} \left\{ \frac{4}{\xi + \eta} \left[ \frac{\partial}{\partial \xi} \left( \xi \frac{\partial}{\xi} \right) + \frac{\partial}{\partial \eta} \left( \eta \frac{\partial}{\partial \eta} \right) \right] + \frac{1}{\xi \eta} \frac{\partial^2}{\partial \varphi^2} \right\} - \frac{2e^2}{\xi + \eta}$$
(15)

In order to solve eqn(14) we take

$$E = E_0 + \Delta E, \ \Phi = \psi exp(\frac{ieg}{\hbar c}F - iM\varphi) \quad F = \log\frac{\xi\eta}{a_0^2}$$
(16)

where  $a_0$  ground state Bohr radius, M is magnetic quantum number and  $\psi$  is real. To order g

$$H_0 \Phi = E_0 \Phi \tag{17}$$

becomes

$$-\frac{\hbar^{2}}{2m}\left(\frac{4}{\xi+\eta}\right)\left[\frac{\partial}{\partial\xi}\left(\xi\frac{\partial\psi}{\partial\xi}+\frac{\partial}{\partial\eta}\left(\eta\frac{\partial\psi}{\partial\eta}\right)\right]+\frac{ieg}{\hbar c}\left[\frac{\partial}{\partial\xi}\left(\xi\frac{\partial F}{\partial\xi}\right)+\frac{\partial}{\partial\eta}\left(\eta\frac{\partial F}{\partial\eta}\right)\right]$$
$$+\frac{2ieg}{\hbar c}\left[\frac{\partial\psi}{\partial\xi}\left(\xi\frac{\partial F}{\partial\xi}\right)+\frac{\partial\psi}{\partial\eta}\left(\eta\frac{\partial F}{\partial\eta}\right)\right]+\frac{\hbar^{2}M^{2}}{2m}\frac{\psi}{\xi\eta}-\frac{-2e^{2}\psi}{\xi+\eta}\psi=E_{0}\psi$$
(18)

With

$$F = \log \frac{\xi \eta}{a_0^2} , \quad \frac{\partial F}{\partial \xi} = \frac{1}{\xi}, \quad \frac{\partial}{\partial \xi} (\xi \frac{\partial F}{\partial \xi}) = 0$$
(19)

equation (18) becomes

$$-\frac{\hbar^2}{2m}(\frac{4}{\xi+\eta})\left[\frac{\partial}{\partial\xi}(\xi\frac{\partial\psi}{\partial\xi}+\frac{\partial}{\partial\eta}(\eta\frac{\partial\psi}{\partial\eta})\right]+\frac{2ieg}{\hbar c}(\frac{\partial\psi}{\partial\xi}+\frac{\partial\psi}{\partial\eta})\right]+\frac{\hbar^2M^2\psi}{2m\xi\eta}-\frac{2c^2\psi}{\xi+\eta}=E_0\psi$$
 (20)

Since  $E_0$  is obtained by multiplying both sides of this equation on the left by  $\psi^*$  and taking volume integral and since  $\psi^* = \psi$ 

$$\int_0^\infty d\xi \psi^* \frac{\partial \psi}{\partial \xi} = \frac{1}{2} \int_0^\infty d\xi \frac{\partial(\psi\psi)}{\partial \xi} = 0$$

equation (20) is equivalent to

$$-\frac{\hbar^2}{2m}(\frac{4}{\xi+\eta})\left[\frac{\partial}{\partial\xi}\left(\xi\frac{\partial\psi}{\partial\xi}+\frac{\partial}{\partial\eta}(\eta\frac{\partial\psi}{\partial\eta})\right]+\frac{\hbar^2M^2}{2m}\frac{\psi}{\xi\eta}-\frac{2e^2\psi}{\xi+\eta}=E_0\psi\tag{21}$$

which is the standard H-atom Schroedinger equation in the absence of external field  $\mathcal{E}$ .

¿From eqn(13) (14) and (16) we have, to orger  $g^2$ ,

$$\Delta E = \langle \Phi \mid H_I \mid \Phi \rangle = \frac{ig\lambda\mathcal{E}}{8} \int d\xi d\eta \xi \eta \psi (\frac{\overline{\partial}}{\partial\xi} - \frac{\overline{\partial}}{\partial\eta}) \psi - \frac{eg^2\lambda\mathcal{E}}{4\hbar c} \int d\xi d\eta \psi (\frac{\partial F}{\partial\xi} - \frac{\partial F}{\partial\eta}) \psi$$
(22)

which on use of (19) gives

$$\Delta E = \frac{eg^2\lambda\mathcal{E}}{4\hbar c} \int d\xi d\eta \psi(\xi - \eta)\psi = \frac{eg^2\lambda}{2\hbar c}\epsilon(\frac{n_1 - n_2}{n})$$
(23)

on account of vanishing of the first term on r.h.s. of eqn(22). The conventional approach with the definition (6) for electric dipole moment, gives

$$\Delta E = \frac{3\eta \mathcal{E}}{2} (n_1 - n_2) \frac{\hbar^2}{me} \tag{24}$$

In view of the identify (9) we equate (23) and (24) and obtain

$$\frac{eg}{\hbar c} = \sqrt{3}n\tag{25}$$

which differs from Dirac's result by a numerical factor.

It is worth noting that the wave function  $\Phi$  in eqn(16) has a logarithmic singularity either for  $\xi = r + z = 0$  or for  $\eta = r - z = 0$ . The former corresponds to singularity extending from 0 to  $-\infty$  along the negative z-axis and the latter extending from 0 to  $+\infty$  along the positive z-axis which is reminiscent of the Dirac string.

The linear combination of degenerate excited states of hydrogen atom that posses permanent electric dipole moment and have magnetic charge are just those discussed in this paper i.e. states with the parabolic co-ordinate. These states form when the hydrogen atom is placed in an uniform static electric field. Magnetic charged states can therefore be experimentally fabricated by passing hydrogen atoms selectively excited to a specified n quantum number through a region of uniform electrostatic field. For convenience one can excite the atoms to the n = 2 state by absorption of a laser beam of  $\lambda = 2430A$  which can be generated by frequency doubling of an intense dye laser beam at  $\lambda = 4860 A$  [6]. The two states of the beam with quantumnumbers ( $n_1 = 0$ ,  $n_2 = 1$ , m =0) and ( $n_1 = 1, n_2 = 0, m = 0$ ) possesing equal and opposite magnetic charge get bent in opposite directions due to the action of the electrostatic field. The two separated magnetic charged particles in uniform magnetostatic field. The two separated magnetic charged particle beams can be passed through superconducting rings [4] causing flow of electric current therein which can be detected by SQUID (Superconducting quantum interpreference device).

In view of the fact that the non-degenerate ground state  $\psi_o$  of the hydrogen atom can have no magnetic charge, it would appear that conservation of magnetic charge will be violated when the atom makes a radiative transition to (form) the ground state from (to) an excited state. However, a closer analysis shows that this is not so. Due to interaction of the atom with the radiation field, the states acquire time dependence as per below.

$$\psi_n(t) = [Cos \ \omega_n t] \ \psi_n(0) + \ [Sin \ \omega_n t] \psi_0(0)$$
  
$$\psi_0(t) = [Cos \ \omega_n t] \ \psi_0(0) - \ [Sin \ \omega_n t] \psi_n(0)$$
  
(26)

where  $\omega_n$  is the frequency of radiation resulting from the transition between states  $|n\rangle$ and  $|0\rangle$ . It therefore follows that

$$g_n(t) = g_n \ Cos^2 \omega_n \ t$$

$$g_0(t) = g_n \ Sin^2 \omega_n \ t$$
(27)

so that

$$g_n(t) + g_0(t) = g_n \tag{28}$$

remains constant.

## References

- 1. P.A. M. Dirac, Proc. Roy. Soc. A133 60 (1931), Phys. Rev. 74 817 (1948).
- 2. M.N. Saha, Ind. J. Phys. 10 141 (1936), Phys. Rev. 75 1968 (1949).
- G't Hooft, Nucl. Phys. **B79** 276 (1974).
   A.M. Polyakov, JETP Lett. **20** 194 (1974).
- E. Amaldi in "Old and New Problems in Elementary Particles" ed. G. Puppi (Acad. Press, New York 1968).
  - A.S. Goldhabor and J. Smith, Rep. Prog. Phys. **38** 731 (1975).
  - B. Cabrera, Phys. Rev. Lett. 48 1378 (1982)
  - B. Cabrera etal. Phys. Rev. Lett. **51** 1993 (1983).
  - T. Hara et.al. Phys. Rev. Lett. 56 553 (1986).
  - T. Gentile et.al. Phys. Rev. **D35** 1081 (1987).
  - The MACRO collaboration, Nucl. Instrum. Methods. Phys. Res. Sec. A **264** 18 (1988).
- 5. N. Cabibbo and E. Ferrari, Nuovo Cimento 23 1147 (1962).
  C.R. Hagen, Phys. Rev. 140B 804 (1965).
  D. Zwanziger, Phys. Rev. D3 880 (1971).
  - W. Baker and F. Graziani, Phys. Rev. **D18** 3849 (1978) and Am. J. Phys. **46** 1111 (1978).
  - D. Singleton, Am. J. Phys. **64** 452 (1996).
- 6. T.W. Hansch, Physics Today **30** 34 (1977).