A priori Probabilities of Separable Quantum States

Paul B. Slater

ISBER, University of California, Santa Barbara, CA 93106-2150 e-mail: slater@itp.ucsb.edu, FAX: (805) 893-7995

(November 26, 2024)

Życzkowski, Horodecki, Sanpera, and Lewenstein (ZHSL) recently proposed a "natural measure" on the N-dimensional quantum systems, but expressed surprise when it led them to conclude that for $N = 2 \times 2$, disentangled (separable) systems are more probable (0.632 ± 0.002) in nature than entangled ones. We contend, however, that ZHSL's (rejected) intuition has, in fact, a sound theoretical basis, and that the *a priori* probability of disentangled 2×2 systems should more properly be viewed as (considerably) less than 0.5. We arrive at this conclusion in two quite distinct ways, the first based on classical and the second, quantum considerations. Both approaches, however, replace (in whole or part) the ZHSL (product) measure by ones based on the volume elements of *monotone* metrics, which in the classical case amounts to adopting the Jeffreys' prior of Bayesian theory. Only the quantum-theoretic analysis — which yields the smallest probabilities of disentanglement — uses the *minimum* number of parameters possible, that is $N^2 - 1$, as opposed to $N^2 + N - 1$ (although this "over-parameterization", as recently indicated by Byrd, should be avoidable). However, despite substantial computation, we are not able to obtain precise estimates of these probabilities and the need for additional (possibly supercomputer) analyses is indicated — particularly so for higher-dimensional quantum systems (such as the 2×3 ones, we also study here).

PACS Numbers 03.67.-a, 02.50.-r, 89.70.+c, 02.40.Ky

Keywords: Quantum entanglement, separable states, monotone metrics, Jeffreys' prior, Haar measure, partial transposition

I. INTRODUCTION

In a recent paper [1], Życzkowski, Horodecki, Sanpera and Lewenstein (ZHSL) [1] sought to estimate "how many entangled (disentangled) states exist among all quantum states". They gave three principal reasons for their study: (1) to answer the question "is the world *more classical* or *more quantum*?"; (2) to know, for the purposes of numerical simulation, "to what extent entangled quantum systems may be considered as typical"; and (3) "to investigate how frequently certain nonseparable states, 'peculiarly' admitting time reversal in one subsystem, arise". In response to the first query, ZHSL concluded — to their "surprise" [1, p. 889] — that although the (higher-dimensional) "world" is, in general, more quantum than classical, this is not so for the 2×2 quantum systems. We contend here, however, that alternative analyses based on the concept of *monotone* metrics on classical and quantum systems [2], lead to the elimination of this exception to their general rule.

In their investigation, ZHSL obtained a variety of both analytical and numerical bounds on the volumes of the sets of separable states for various dimensions, using what they asserted was a "natural measure" on the space of density matrices (cf. [3]). In the first analytical part of this communication (sec. II), we indicate an essential degree of arbitrariness in the choice of measure by ZHSL, and its consequences for the results they have reported (cf. [4]). We then argue in favor of a specific alternative — well-founded on statistical principles — which leads to a markedly smaller probability of encountering a disentangled (separable) state. (The numerical results we report are for the 2×2 , 2×3 and 3×3 quantum systems.) Then, in sec. III, we study the use of methods more fundamentally quantum-theoretic in nature — requiring us to develop a quite distinct set of procedures than those used by ZHSL and followed in sec. II. (Due to the associated large computational demands, we have primarily limited our analyses to the 2×2 systems, but in sec IIIG we do, in fact, initiate a parallel investigation of the 2×3 systems.) We obtain for each of more than thirty pairs of parameters, determining the fineness of approximating square grids and three-dimensional simplicial decompositions, a set of three probabilities of disentanglement (Table I), each probability being based on a distinct form of monotone metric [6,7]. The sets are intended to determine a range of values within which any suitable candidate (meeting underlying natural criteria of *monotonicity*) for the "true" a priori probability of disentanglement must lie. Essentially all such probabilities we obtain turn out to be considerably smaller than both the result of ZHSL $(0.632 \pm .002)$ and the alternative to it (≈ 0.35) we promote in sec. II, based on the Jeffreys prior of Bayesian theory. However, the need for additional computational work is indicated in order to sharpen the estimates reported in secs. IIIA IIIB and IIIC, as well as to extend our general approach to higher-dimensional

quantum systems. (We are reminded, to some degree, of the computational/combinatorial challenges of lattice gauge theory [5].) In sec. IIIE, we switch from the explicit enumeration approach (based on regular grids and simplicial decompositions) to a randomization methodology (such as ZHSL employed in their studies) — but also find this to be highly computationally demanding, since we must search in a high-dimensional parameter space for those particular points corresponding to density matrices.

A conservative evaluation of the accumulated evidence of the multiple quantum-theoretic analyses we report (Tables I II III) indicates that the *a priori* probability of disentanglement for the 2×2 systems should be regarded — using any of a continuum of possible acceptable standards, in particular that provided by the *minimal* monotone (Bures) metric — as *no more* than eleven percent. As to a *lower* bound, on the other hand, on the probability of separability, it remains an unsettled issue as to whether or not the *maximal* monotone metric should be viewed as furnishing a bound strictly greater than zero.

In our concluding remarks (sec. IV), we draw attention to an interesting recent analysis of M. Byrd (personal communication), bearing upon the issue of whether or not the use of "over-parameterizations" by ZHSL and (following them) by us in sec. II, can be averted.

II. SEMICLASSICAL STATISTICAL ANALYSES OF 3 × 3, 2 × 3 AND 2 × 2 QUANTUM SYSTEMS

ZHSL [1] used as a measure on the space of $N \times N$ density matrices the product of the Haar measure for the unitary group U(N) and the uniform distribution on the (N-1)-dimensional simplex spanned by the N eigenvalues of the density matrix. Now, we see no basis (within the semiclassical framework adopted by ZHSL) for questioning the use of the Haar measure. However, the selection of the uniform distribution on the simplex appears not to be so compelling, as it lacks as convincing a rationale as the group-theoretic argument for the Haar measure (cf. [8]). Also, we must point out that the analyses of ZHSL are based on "over-parametrizations", since $N^2 + N - 1$ parameters are used, while the convex set of $N \times N$ density matrices is only $(N^2 - 1)$ -dimensional in nature. Though we adhere to this over-parameterization in the analyses of this section, in sec. III we revert to the more natural and conventional form.

The uniform distribution on the (N-1)-dimensional simplex $(p_1 + \ldots + p_N = 1; p_i \ge 0)$ can be considered to be that specific member of the (continuous) family of Dirichlet probability distributions [9,10, sec. 7.7],

$$\frac{\Gamma(\nu_1 + \dots + \nu_N)}{\Gamma(\nu_1) \dots \Gamma(\nu_N)} p_1^{\nu_1 - 1} \dots p_{N-1}^{\nu_{N-1} - 1} (1 - p_1 - \dots - p_{N-1})^{\nu_N - 1}; \quad \nu_1 > 0, \dots, \nu_N > 0$$
(1)

which has all its N parameters (ν 's) set equal to unity. The family of Dirchlet distributions is *conjugate*, in that if one selects a *prior* distribution belonging to it, then, through the application of Bayes' rule to observations drawn from a multinomial distribution, one arrives at a *posterior* distribution which is also within the family.

A. Jeffreys' prior

Of strongest interest, however, for our purposes here, is that the principle of reparameterization invariance (based on the Fisher information [11,12]) leads to the special case (the Jeffreys' prior) in which (1) has all its N parameters set equal to one-half [13, eq. (3.7)] [14] and not unity, as for the uniform distribution. "The main intuitive motivation for Jeffreys' priors is not their invariance, which is certainly a necessary, but in general far from sufficient condition to determine a sensible reference prior; what makes Jeffreys' priors unique is that they are uniform measures in a particular metric which may be defended as the 'natural' choice for statistical inference" [15]. By way of illustration, Kass [14, sec. 2] (cf. [16]), using the transformations $p_i = 2z_i^2$, demonstrates how the Jeffreys' prior for the trinomial model (N = 3) on the two-dimensional simplex, can be (making use of spherical polar coordinates) transformed to the uniform distribution on the positive-octant portion of the two-dimensional sphere, $z_1^2 + z_2^2 + z_3^2 = 4$ of radius 2. (Braunstein and Milburn [17] show that for two-level quantum systems, statistical distinguishability is just the [Bures/minimal monotone] metric on the surface of the unit sphere in four dimensions. In contrast, the space of *n*-level quantum systems is "not a space of constant curvature for n > 2 and not even a locally symmetric space. The physical meaning of this fact seems to be an interesting open question" [18] (cf. [19,20]).)

Clarke and Barron [21,22] (cf. [23]) have established that Jeffreys' priors (the normalized volume elements of Fisher information metrics) asymptotically maximize Shannon's mutual information between a sample of size n and the parameter, and that Jeffreys' prior is the unique continuous prior that achieves the asymptotic minimax

risk when the loss function is the Kullback-Leibler distance between the true density and the predictive density. (The possibility of extending the "universal coding" results of Clarke and Barron to the *quantum* domain, has been investigated in [24] (cf. [25]).) Clarke [26] asserts that "Jeffreys' prior can be justified by four distinct arguments". In addition, Balasubramanian [27] "cast parametric model selection as a disordered statistical mechanics on the space of probability distributions" and "derived and discussed a novel interpretation of Jeffreys' prior as the uniform prior on the probability distributions indexed by a parametric family" (cf. [28]).

Now, it is of interest to note that in the limit in which the N parameters (ν 's) of the Dirichlet distribution (1) all go to zero, the distribution becomes totally concentrated on the pure states of the N-dimensional quantum system. Since ZHSL showed that in "the subspace of all pure states, the measure of separable states is equal to zero" [1, p. 886], we would anticipate, making use of a continuity argument, that the measure or volume of the set of separable states would increase if all N parameters of (1) were fixed at one-half (Jeffreys' prior), but still be less than if they were all taken to be equal to unity, *etc.* ("The purer a quantum state is, the smaller its probability of being separable" [1, p. 891].)

B. The case of 3×3 quantum systems

We have, in fact, tested these last contentions regarding competing measures of separability, through numerical means, first, generating a set of three thousand random 9×9 unitary matrices (N = 9), following the (Hurwitz/Euler angle) prescription given in [29, eqs. (3.1)-(3.5)]. From it, we produced (in the manner of ZHSL [1, eq. (34)]) three sets of three thousand 9×9 density matrices: one set based on the selection $\nu_1 = \ldots = \nu_9 = \frac{1}{2}$; another for $\nu_1 = \ldots = \nu_9 = 1$ (as, in effect, done in [1]); and a third for $\nu_1 = \ldots = \nu_9 = \frac{3}{2}$. (Random realizations of the Dirichlet distributions were generated based on the fact that they can be considered to be joint distributions of [univariate] gamma distributions [9,10]. The three thousand instances we obtain are obviously far fewer in number than the "several millions" ZHSL [1] apparently employed as a general rule in their series of analyses of quantum systems of various dimensions. This is primarily due to our full reliance on MATHEMATICA, while ZHSL — as K. Życzkowski wrote in a personal communication — employed FORTRAN routines for random number generation. Nevertheless, as noted immediately below, Życzkowski, using his speedier routines, has confirmed the main aspects of our analysis. Additionally, the quantum-theoretic analyses of sec. III require a quite different set of algorithms, and it is far from clear whether our use there of MATHEMATICA is in any way relatively inefficient.)

Then, we determined whether all (nine of) the eigenvalues of the *partial transpositions* of the random density matrices (viewing them as $(3 \times 3) \times (3 \times 3)$ density matrices, in the manner of (21) in [30]) were positive (as they must be in the separable case) or not. For the $\nu = \frac{1}{2}$ (Jeffreys' prior) scenario, eighty-three of the three thousand density matrices had this positivity property, while considerably more (six hundred and two) possessed it for $\nu = 1$ and still more (one thousand two hundred and ninety-six) for $\nu = \frac{3}{2}$. Thus, we note an approximate decrease by a factor of $\frac{83}{602} \approx .138$ in the upper bound on our suggested probability of encountering a separable state *vis-á-vis* the analysis of ZHSL.

C. The cases of 2×3 and 2×2 quantum sytems

We, then, conducted parallel analyses to those in sec. II B for the 2×3 and 2×2 systems. (For both such systems, but not higher-dimensional ones, such as the 3×3 , the positivity of the partial transposition is a sufficient, as well as necessary condition for separability [30]. So, we will be estimating probabilities themselves, rather than upper bounds on them.) In both cases, we now employed ten thousand realizations. In the 2×3 case, we found 1,309 separable states, using $\nu = \frac{1}{2}$, and 4,135 for $\nu = 1$, as well as 6,357 for $\nu = \frac{3}{2}$. (Our statistic of .4135 needs to be compared with that of .384 ± .002 of ZHSL — which, as noted, was based on a much larger sample.) For the 2×2 systems, the analogous results were: 3,633; 6,564; and 7,946. So, we would conclude, in this analytical framework, that the proportion of separable states among the 2×2 quantum systems should be taken to be approximately .36 — which is well *below* the demarcation point of .5, *above* which ZHSL found their result of .632 ± .002 (roughly comparable to ours of .6564) to (counterintuitively) lie.

K. Życzkowski has kindly repeated the analyses reported above for the case $\nu = \frac{1}{2}$ (that is, Jeffreys' prior), using 200,000 random realizations for each of the three scenarios. The probabilities of separability he obtained were (all digits being significant he states): .022 (for the 3 × 3 systems); .122 (2 × 3 systems); and .352 (2 × 2 systems). These should be compared with our results (based on considerably smaller samples) of .0277, .1309 and .3633, respectively.

These various numerical results are, thus, quite supportive of our arguments and help to fulfill the first objective of this letter of showing the dependence of estimates of the volume of the set of separable states on the particular choice of (symmetric) Dirichlet distribution on the (N-1)-dimensional simplex spanned by the N eigenvalues of the $N \times N$ density matrix (ρ). We note again that ZHSL [1, p. 889] expressed "surprise that the probability that a mixed state $\rho \in H_2 \times H_2$ is separable exceeds fifty percent". Thus, they would have apparently been not so confounded if the uniform distribution on the three-simplex of eigenvalues had been replaced by the Jeffreys' prior, since its use yields a more modest percentage of approximately thirty-five.

Let us also note that our suggested modification $(\nu = \frac{1}{2})$ of the ZHSL measure $(\nu = 1)$ would appear to find some support in a recent paper concerned with a parameterization (sharing certain features with that of ZHSL) of the $N \times N$ density matrices [8]. Its authors consider the N eigenvalues to be parameterized by the "squared components" of the (N - 1)-sphere (rather than coordinates in the (N - 1)-dimensional simplex, as in ZHSL). As noted above, in relation to [14], the Jeffreys' prior is simply the uniform distribution on such a *sphere* — while ZHSL used instead the uniform distribution on the *simplex*.

III. QUANTUM-THEORETIC STATISTICAL ANALYSES OF 2×2 AND 2×3 SYSTEMS

In [4, sec. II.C], we presented evidence that certain statistical features of the product measure employed by ZHSL [1] were not reproducible through the use of any of the possible (continuum of) *monotone* metrics. A similar conclusion appears to hold if one replaces the uniform distribution in the ZHSL product measure (as we have done above in sec. II) by any other member of the family of Dirichlet distributions (1). Since Petz and Sudár [2] have argued that monotone metrics are the quantum analogues of the (classically unique) Fisher information metric, it would seem highly desirable to replace the product measures so far employed by ones based directly on the volume elements of such metrics. (In [31], efforts were reported to integrate the volume elements of the *minimal* and *maximal* monotone metrics over the convex sets of 3×3 and 4×4 density matrices.) In so doing, we would avoid the nonparsimonious "over-parameterization" mentioned at the outset of sec. II. (However, it will be incumbent upon us to develop a quite distinct set of computational methods than those used by ZHSL and applied in sec. II.)

We have, in fact, conducted such a series of analyses for the 2×2 quantum systems, based on a MATHEMATICA program containing *two* parameters of choice, n_1 and n_2 . The parameter n_1 determines the fineness of a regular decomposition of the three-dimensional simplex — the points of which correspond now to the *diagonal* entries of ρ , and not the *eigenvalues*, as in sec. II and the work of ZHSL [1] and Boya *et al* [8]. (Of course, both the eigenvalues and diagonal entries of a density matrix are nonnegative and sum to unity. To compute the coordinates of the simplicial coordinates, we followed an algorithm for the next *composition* of an integer N into K parts, given in [32, chap. 5], taking K = 4 for our purposes, and then dividing each of the $\binom{N+3}{N}$ compositions generated by N.) The reciprocal of the parameter n_2 is the distance between adjacent points of a regular square grid — having its extreme points/corners at $(\frac{1}{2}, \frac{1}{2}), (\frac{1}{2}, -\frac{1}{2}), (-\frac{1}{2}, -\frac{1}{2})$ and $(-\frac{1}{2}, \frac{1}{2})$ — imposed on a circle of radius one-half centered at the origin of the complex plane. (The off-diagonal entries of a density matrix can not exceed one-half in absolute value, so they must lie within this circle.)

For specific values of n_1 and n_2 , within but challenging our computational capabilities, we generated the associated three-dimensional simplicial decompositions and twelve-dimensional uniform lattices (the six-fold Cartesian product of the imposed two-dimensional square grid — six, of course, corresponding to the number of pairs of off-diagonal entries). Then, we explicitly enumerated all those points in the fifteen-dimensional product space parameterizing the 4×4 density matrices of mixed states (that is, yielding matrices having all strictly positive eigenvalues, noting that the additional Hermiticity and trace requirements are automatically satisfied by construction). We would reject any density matrices of pure states (the totality of which form a six-dimensional subspace [33]) that happened to be generated, since our measures (see immediately below) are singular on them, as well as more generally, degenerate density matrices, those density matrices not being of full rank (and hence having zero determinant). However, the possibility remains — in particular, since we will be computing (nonrobust) averages — that the behavior of the measures for a relatively few nearly degenerate states, can strongly influence the results (cf. Tables I and II).

By our purposeful design, the explicitly enumerated points are uniformly distributed (using the conventional parameterization) in the fifteen-dimensional convex set of 4×4 density matrices. We took several significant steps in our MATHEMATICA ("backtrack" [32, chap. 27]) program to cut down on the (potentially huge) search spaces, by utilizing the requirement that all the principal minors of a density matrix must be nonnegative [34, Thm. 7.2.5] [37]. (A sufficient condition only, of possible interest, would be that the matrix is diagonally-dominant [34].) Also, in our later, larger analyses, we exploited certain permutational symmetries.

We, then, employed an *ansatz* of ours [31], building upon a result of Dittmann [18, p. 76] (pertaining to the Bures or minimal monotone metric) regarding the spectrum of the sum of the operators of left and right multiplication of matrices cf. [38, p. 112]. Utilizing it, we assigned as a weight to each density matrix (ρ) generated (the eigenvalues of which are denoted by λ_i), the volume elements of certain monotone metrics of particular interest. These elements we took to be of the form $[\Pi_{i,j=1}^4 f(i,j)]^{\frac{1}{2}}$, where (the "Morozova-Chentsov" function [2] [39, eq. (3)]) f(i,j) is equal to $\frac{2}{(\lambda_i+\lambda_j)}$ in the minimal monotone case, $\frac{(\lambda_i+\lambda_j)}{2\lambda_i\lambda_j}$ in the maximal monotone case, and for the Kubo–Mori/Bogoliubov (KMB) metric (associated with the relative entropy) [7,40,41], $\frac{\log \lambda_i - \log \lambda_j}{\lambda_i - \lambda_j}$. (In this last case, if $\lambda_i = \lambda_j$, we take $f(i, j) = \lambda_i^{-1}$. It is interesting to note that the inverses of these Morozova-Chentsov functions are simply wellknown indicators of *central tendency*, such as the arithmetic mean, the logarithmic mean, and the harmonic mean [42]. Choosing a particular monotone metric is, therefore, akin to selecting such an indicator.) We also checked if ρ satisfied the partial transposition condition, necessary for separability.

We now report several sets of results in this regard — but let us first make some important observations (taking into account that the determinant of a matrix is equal to the product of its eigenvalues). Using the formulas just given, one can show [43] that for an $N \times N$ density matrix (ρ) , the volume element of the minimal monotone metric (and also the KMB-metric) is directly proportional to $(\det \rho)^{-\frac{1}{2}}$, while for the maximal monotone metric, the volume element is directly proportional to $(\det \rho)^{\frac{1-2N}{2}}$. So, the divergence near the boundary of degenerate states $(\det \rho = 0)$ of the volume element of the maximal monotone metric is much more severe than for the other two metrics under investigation. In fact, in our previous studies [6,31], we have concluded that the integral of the volume element of the maximal monotone metric over the convex set of 2×2 density matrices does *not* converge (in contrast to those for the minimal monotone and KMB-metrics). (For N > 2, however, the issue of convergence appears to be unsettled.) So, it would seem that — unless one chooses to remove from consideration (as was done in [6], for inferential purposes) those states the degeneracy of which exceeds some prescribed level [6] (cf. sec. III D) — one can not, in fact, define a probability distribution based on the maximal monotone metric. We have been able, however, for N = 3, by taking the limit of a certain ratio, to obtain associated *marginal* probability distributions using the maximal monotone metric [31].) Also, the maximal monotone metric is of substantial interest, in that it has been characterized as the most *noninformative* of the monotone metrics [6,7].

The motivating hypothesis for pursuing the analyses immediately below is that one should be able to find values of the parameters n_1 and n_2 large enough (say, m_1 and m_2), so that the associated probabilities of disentanglement are within some ϵ of each other for any choices of n_1 and n_2 which dominate both m_1 and m_2 . (It is useful to bear in mind, however, that there is a qualitative difference between analyses based on even or odd values of n_2 , as will be indicated.) This would indicate a convergence of these probabilities in the continuum limit as n_1 and n_2 each become indefinitely large.

A. The case $n_1 = 23$, $n_2 = 7$

The choice of $n_2 = 7$ leads to a square grid, having thirty-two points — serving as trial off-diagonal entries — lying within the circle of radius one-half. (The particular arrangement of the lattice points, then, mandates that those density matrices we will be able to construct will have off-diagonal entries of modulus no less than $\frac{1}{7\sqrt{2}} \approx .101015$. This, in turn, implies that the product of any pair of diagonal entries of the density matrices will not be less than this value.) The numbers of density matrices we were, then, able to construct were 1,340,928. Of these, 356,096 passed the transposition test for separability. Applying the weights based on the three monotone metrics considered, we obtained prior probabilities of encountering separable states of

$$p_{min} = .111102, \quad p_{KMB} = .0873186, \quad p_{max} = .0846153.$$
 (2)

Of course, these three values are all considerably less than both the ZHSL statistic of $.632 \pm .002$ and the preferred one (of the three given) of sec. II based on the Jeffreys' prior, that is, $\approx .35$.

We have also computed the "degree of entanglement", $\sum_{i=1}^{4} \lambda'_i - 1$ (which must lie between zero and unity) for all the (1,340,928) density matrices and averaged the results with the same set of three weights as used to obtain (2). The outcomes were

$$d_{min} = .18206, \quad d_{KMB} = .208022, \quad d_{max} = .248457.$$
 (3)

The corresponding value obtained by ZHSL for the 2×2 systems was considerably smaller, that is .057 [1, App. B]. (ZHSL remarked that this quantity seemed to saturate at approximately 0.10 for large systems.)

B. The case $n_1 = 19$, $n_2 = 8$

Since the parameter n_2 is now an even integer, the origin (0,0) of the complex plane becomes, by our particular mode of construction, one of the forty-nine intersection points of the square grid lying within the circle of trial values for the off-diagonal entries. There are, then, no nontrivial lower bounds imposed on the moduli of these entries, as there are for odd values of n_2 — such as *seven* in the immediately preceding analysis of sec. III A. We obtained 4,443,408 density matrices, of which 1,284,816 satisfied the separability criterion. Use of the volume elements of the three selected monotone metrics as weights resulted in

$$p_{min} = .147968, \quad p_{KMB} = .123283, \quad p_{max} = .0554999, \tag{4}$$

and

$$d_{min} = .18686, \quad d_{KMB} = .184149, \quad d_{max} = .17713.$$
 (5)

C. Additional (nontruncated) analyses

Continuing along the same lines as secs. III A and III B, we have conducted analyses for additional choices of n_1 and n_2 . (It is interesting to note that unit increases in n_2 are relatively more costly computationally than in n_1 .) We report our accumulated set of results in Table I. The analyses are listed in increasing order of the total number of density matrices generated. (During the course of conducting these analyses, we were able to undertake larger-sized studies, corresponding to those listed at the bottom of the table, by taking advantage of certain inherent permutational symmetries. An analogous assertion can be made in regard to Table IV.)

As a general rule, the probability of disentanglement is greatest for the minimal monotone metric, although still markedly less than the ZHSL result (0.632 ± 0.002) or that of sec. II (≈ 0.35) based on the Jeffreys' prior. The stability of the results, on the other hand, is least for the maximal monotone metric (in particular, notoriously so, for the case, $n_1 = 30$, $n_2 = 7$ — but see sec. III E). This instability may be explainable by the fact that the volume element of the maximal monotone metric, as previously noted, is not normalizable (to form a probability distribution) over the convex sets of $N \times N$ density matrices (in particular, for N = 4), being highly singular near the degenerate states (while the volume elements of the minimal and KMB metrics, though, still singular, are markedly less so, and are apparently normalizable, extrapolating from the 2×2 case). So, one might rely upon either the minimal monotone metric should, at least for fine enough grids and simplicial decompositions, dominate estimates based on any other member of the *continuum* of monotone metrics. Following the arguments of Petz and Sudar [2], we contend that any estimates not based on such metrics (such as the "over-parameterized" results of ZHSL [1] — cf. [8] — and those of sec. II here) fail to meet certain natural requirements and should, thus, be taken *cum grano salis*.

Let us also note a specific relation between the minimal monotone (Bures) metric and the results of ZHSL. The scalar curvature of this metric has recently been shown to attain its minimum $-\frac{(5N^2-4)(N^2-1)}{2}$ — for the totally mixed N-dimensional (tracial) state (corresponding to the $N \times N$ diagonal density matrix having all its nonzero entries equal to $\frac{1}{N}$), and to diverge on the degenerate states, those not of full rank [19]. Now, in their analysis, ZHSL concluded both that all states in a small enough neighborhood of the totally mixed state are separable, and that the "purer a quantum state is, the smaller its probability of being separable" [1]. Braunstein *et al* have given "a constructive proof that all mixed states of N qubits in a sufficiently small neighborhood of the maximally mixed state are separable" [35], while Vidal and Tarrach have also reached the same conclusion [36].

D. Reanalysis of the anomalous $n_1 = 30$, $n_2 = 7$ case based on truncation of states near to degeneracy

We have also considered the possibility of introducing a third parameter of choice, that is det ρ — in addition to n_1 and n_2 — into our computations. It would control the level of degeneracy below which we reject for further consideration (due to the singular behavior of the volume elements of the monotone metrics), the (nearly degenerate) density matrices our explicit enumeration method of sec. III generates. (This third parameter has been implicitly zero in sec. III.) We have, in fact, conducted three additional analyses for the case $n_1 = 30$, $n_2 = 7$, for which we previously obtained results of a peculiar nature (Table I). The largest possible value the determinant of a 4×4 density matrix can possess is $(\frac{1}{4})^4 = \frac{1}{256} = .00390625$. In the first analysis, we rejected all those density matrices with determinants less than $\frac{1}{256} \cdot 10^{-4}$, in the second, $\frac{1}{256} \cdot 10^{-3}$ and in the third, $\frac{1}{256} \cdot 10^{-2}$. We report these results in Table II. It appears then (based on the smallest nonzero threshold, that is $\frac{1}{256} \cdot 10^{-4}$) that the previously reported (zero-

It appears then (based on the smallest nonzero threshold, that is $\frac{1}{256} \cdot 10^{-4}$) that the previously reported (zero-threshold) anomalous behavior for the maximal and KMB-monotone metrics was attributable to some set (the number of which we are not certain) of near-degenerate states which, in fact, passed the partial transposition test for separability.

E. Analyses based on randomized searches

In the previous quantum-theoretic statistical analyses of this section, we employed systematic explicit enumeration methods to generate 4×4 density matrices, which we then tested for separability. We embarked on such a course after initial computations indicated that it was extremely difficult to locate the four-by-four density matrices (in the ambient fifteen-dimensional parameter space) using *random* search methods, in the fashion of ZHSL [1] and sec. II of this paper. Nevertheless, at a later point, we chose to intensively pursue such a strategy.

In almost eight hundred and fifty million *ab initio* searches, we succeeded in obtaining (only) sixty-one density matrices — of which, twelve turned out to be separable. Realizing that the "hit-rate" would be enhanced if instead of searching for possible off-diagonal entries in the circle of radius one-half in the complex plane, we also conducted analyses (though at the risk of introducing possible biases) based on radii of one-third and one-fourth, as well (and also, in a supplementary analysis, five-twelfth). The results are reported in Table III. (As in sec. II, standard deviations were *not* determined, so no specific assessment of the number of significant digits in the probabilistic results is immediately available.) They are, then, arguably, generally consistent with the sets of smaller probabilities reported in Table I, in particular, for those based on the largest number of generated density matrices (corresponding to the bottom rows of the table), in which we naturally repose the greatest confidence.

F. Probabilities as a function of the participation ratio

In Fig. 1, we show (using bins of width .05), relying upon the analysis for the 2×2 case $n_1 = 22$, $n_2 = 10$, the conditional probability (P_{sep}) of separability based on the minimal monotone metric, for a given participation ratio R (defined as the reciprocal of the trace of the square of the density matrix [1, eq. (17)]). In Fig. 2, we show its counterpart based on the KMB-metric. (These two figures — both having an unexplained "anomalous blip" in the interval [1.65, 1.7] — are the monotone metric analogues of Fig. 2(b) of [1]. It is encouraging, however, that the "blip" does not seem to appear in analogous plots for other values of n_1 and n_2 . The value of R, in the $N = 2 \times 2$ case, must lie between 1 and 4. If $R \geq 3$, the density matrix must be separable [1, eq. (18)].)



FIG. 1. Conditional probability — based on the minimal monotone (Bures) metric — for $N = 2 \times 2$ of finding a separable state, given a certain range (of width .05) of the participation ratio R, for the scenario $n_1 = 22, n_2 = 10$



FIG. 2. Conditional probability — based on the KMB-metric for $N = 2 \times 2$ — of finding a separable state, given a certain range (of width .05) of the participation ratio R, for the scenario $n_1 = 22$, $n_2 = 10$

Życzkowski [44], drawing upon a long list of open problems he presents, considers one of the "most relevant" to be the question of "whether the dependence of the conditional probability on the participation ratio, obtained for product measures, holds also for the measures based on the monotone metrics". He has hypothesized the existence of certain universal/metric-independent features in this regard, that is, he proposes that all "reasonable" metrics should yield similar such plots. (He has, in fact, superimposed the figures here upon those previously generated by him, and found a strong degree of resemblance between them.)

In Figs. 3 and 4, we show (again for the case $n_1 = 22$, $n_2 = 10$ of Table I) the probabilities of our generating a density matrix (either separable or inseparable) based on the minimal monotone and KMB-metrics, respectively. (These are the monotone metric counterparts of Fig. 2(a) of [1]. Since we find more probability concentrated at smaller values of R than did ZHSL, these two figures help us to understand why we obtain smaller *overall* probabilities of separability — in particular, less than .5 in the $N = 2 \times 2$ case — than their "surprising" result of 0.632 ± 0.002 .)



FIG. 3. Probability — based on the minimal monotone (Bures) metric — for $N = 2 \times 2$ of finding a quantum state (whether separable or not), given a certain range (of width .05) of the participation ratio R, for the scenario $n_1 = 22$, $n_2 = 10$



FIG. 4. Probability — based on the KMB-metric – for $N = 2 \times 2$ of finding a quantum state (whether separable or not), given a certain range (of width .05) of the participation ratio R, for the scenario $n_1 = 22, n_2 = 10$

G. The case of 2×3 systems

In Table IV, we report initial findings (of an, unfortunately, rather unstable nature) for the 2×3 systems — parallel to those given in Table I for the 2×2 systems. (We note that the probability of separability obtained by ZHSL [1] for the 2×3 systems was $.384 \pm .002$ and, with the alternative use of the Jeffreys' prior in sec. II C, .122.) There are now six diagonal entries (associated with n_1) and fifteen pairs of off-diagonal entries (associated with n_2) to consider, so computational demands are substantially increased. We, of course, expect the probabilities of separability to be *less* than the corresponding ones in the 2×2 case reported in Table I, and this is certainly the case for the most extensive analysis ($n_1 = 20, n_2 = 8$).

The somewhat counterintuitive observation that for $n_1 = 14$, the choice of $n_2 = 8$ leads to many more generated density matrices than for $n_2 = 9$, is comprehensible in that only for even values of n_2 are no nonzero lower bounds placed on the possible absolute values of off-diagonal entries. We were not able for any $n_2 = 7$ scenario — due to memory limitations — to find a large enough n_1 , for which any density matrices at all were generated. We also possess no immediate explanation for the equality of the three p's and d's for the three cases involving $n_2 = 9$. (The relatively large probabilities for the scenario, $n_1 = 24$, $n_2 = 6$ may be attributable to a "number-theoretic" effect, given that 24 is exactly divisible by 6.)

A randomization approach, such as we pursued in sec. III E for the 2×2 systems, would clearly yield even fewer density matrices for a given number of independent searches than there (Table III).

IV. CONCLUDING REMARKS

We have presented in this study, two forms of evidence (one essentially classical and the other, quantum-theoretic in nature) that the specific choice of (product) measure of ZHSL [1] led them to substantially overestimate the extent to which quantum systems — in particular, for $N = 2 \times 2$ and 2×3 — should be considered to possess, in some natural *a priori* sense, the property of separability or disentanglement. The preponderance of evidence adduced indicates that the probability of separability for the $N = 2 \times 2$ systems, based on the minimal monotone (Bures) metric is no greater than eleven percent — and if one views the evidence somewhat less conservatively, perhaps less than ten percent. In turn, estimates founded on any other member of the continuum of monotone metrics should be lower still. For instance, for the Kubo-Mori/Bogoliubov metric [7,40,41], an estimate of nine percent would seem conservatively high. Apparently, the *maximal* monotone metric — the volume element of which possesses a high degree of singularity on the degenerate states (det $\rho = 0$), associated with its conjectured nonnormalizability over the fifteen-dimensional convex set of states — must, in some (perhaps limiting) sense, furnish a *lower* bound on the probability of separability. This bound would have to be strictly greater than zero, if the related arguments made in [1,35,36,44], in fact, apply.

Let us also note that in continuing work, pertaining to [24], we have found a quite interesting distinguished role (that of yielding both the *minimax* and *maximin* in universal quantum coding) for a monotone metric that has not apparently previously been noted. Its associated Morozova-Chentsov function,

$$e(\lambda_i^{\lambda_i}/\lambda_j^{\lambda_j})^{\frac{1}{\lambda_j-\lambda_i}},\tag{6}$$

is simply the reciprocal of the *exponential* or *identric* mean [42] of λ_i and λ_j . The behavior of the related monotone metric appears to be quite close to that of the minimal (Bures) monotone metric.

The need for additional computational work (possibly utilizing supercomputers) is indicated, in regard to what we contend are the theoretically superior (properly parameterized) quantum-theoretic analyses of sec. III, in order to more closely pinpoint estimates. Such analyses could be based on finer simplicial decompositions for the trial diagonal entries (that is, higher values of n_1) and finer square grids for the trial off-diagonal entries (that is, higher values of n_2), than those reported in sec. III, and/or possibly randomization procedures, as in sec. III E.

At several points in this paper, we have indicated that the analyses of ZHSL [1] were "over-parameterized", in that $N^2 + N - 1$ parameters were employed, rather than $N^2 - 1$, as is clearly most natural for the $N \times N$ density matrices (which form an $N^2 - 1$ -dimensional convex set). However, in a personal communication, M. Byrd has asserted that this bothersome feature could be avoided, since N - 1 Euler angles (in addition to the "phase", as is well known) can be seen to, in fact, vanish in the ZHSL-type representation [1, eq. (25)] of a density matrix in the product form U'DU. Byrd has been able to explicitly show this in the case N = 3, based on the Euler angle parameterization given in [45] (the angles c and ϕ vanishing) and contends that analogous phenomena must hold for N > 3, as well (cf. [8]). (This vanishing does not appear to occur with the particular Euler angle parameterizations [associated with Hurwitz] used by ZHSL, given in [29,46]. Byrd suggests that this is because the "diagonal matrices that make up the maximal torus" do not appear on the end, while if they did, they would commute with the diagonalized density matrix.) This highly interesting line of thought would suggest that the analyses of ZHSL, Życzkowski [44] and those of sec. II here (but, of course, not those "properly parameterized" ones of sec. III) should be repeated in such a more parameter-wise economical framework, and the new results compared with those previously obtained, to see whether any differences are found (cf. [47]).

ACKNOWLEDGMENTS

I would like to express appreciation to the Institute for Theoretical Physics for computational support in this research, and to K. Życzkowski for several comments, as well as providing the simulations reported at the end of sec. II, pointing out to me [16], and suggesting that it would be of considerable interest to "bin" the results here, thus, leading to the four included figures. I am also grateful to M. Byrd for responding to my queries regarding the issue of "over-parameterization" in the work of ZHSL.

- [1] K. Życzkowski, P. Horodecki, A. Sanpera, and M. Lewenstein, Phys. Rev. A 58, 883 (1998).
- [2] D. Petz and C. Sudár, J. Math. Phys. 37, 2662 (1996).
- [3] W. K. Wootters, Found. Phys. 20, 1365 (1990).
- [4] P. B. Slater, $SU(N) \times S_m$ -Invariant Eigenspaces of $N^m \times N^m$ Mean Density Matrices, quant-ph/9802019.
- [5] C. Davies, New Scient. 158, 32 (1998).
- [6] P. B. Slater, Phys. Lett. A 247, 1 (1998).
- [7] D. Petz and G. Toth, Lett. Math. Phys. 27, 205 (1993).
- [8] L. J. Boya, M. Byrd, M. Mims, and E. C. G. Sudarshan, Density Matrices and Geometric Phases for n-state systems, quant-ph/9810084.
- [9] S. S. Wilks, *Mathematical Statistics*, (Wiley, New York, 1962).
- [10] T. S. Ferguson, Ann. Statist. 1, 209 (1973).
- [11] B. R. Frieden and B. H. Soffer, Phys. Rev. E 52, 2274 (1995).
- [12] B. R. Frieden, Probability, Statistical Optics, and Data Testing, (Berlin, Springer-Verlag, 1991).
- [13] J. O. Berger and J. M. Bernardo, Biometrika 79, 25 (1992).
- [14] R. E. Kass, Statist. Sci. 4, 188 (1989).
- [15] J. M. Bernardo, Statist. Sci. 4, 227 (1989).
- [16] P. L. Antonelli and C. Strobeck, Adv. Appl. Prob. 9 (1977), 238.
- [17] S. L. Braunstein and G. J. Milburn, Phys. Rev. A 51, 1820 (1995).
- [18] J. Dittmann, Sem. Sophus Lie 3, 73 (1993).
- [19] J. Dittmann, The Scalar Curvature of the Bures Metric on the Space of Density Matrices, quant-ph/9810012.

- [20] F. Hiai, D. Petz, and G. Toth, Statist. Scient. Math. Hung. 32, 235 (1996).
- [21] B. S. Clarke and A. R. Barron, IEEE Trans. Inform. Th. 36, 453 (1990).
- [22] B. S. Clarke and A. R. Barron, J. Statist. Plann. Infer. 41, 37 (1994).
- [23] J. Rissanen, IEEE Trans. Inform. Th. 42, 40 (1996).
- [24] C. Krattenthaler and P. B. Slater, Asymptotic Redundancies for Universal Quantum Coding, quant-ph/9612043.
- [25] R. Jozsa, M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. 81, 1714 (1998).
- [26] B. S. Clarke, J. Amer. Statist. Assoc. 91, 173 (1996).
- [27] V. Balasubramanian, Neural Comput. 9, 349 (1997).
- [28] V. Balasubramanian, in Maximum Entropy and Bayesian Methods, edited by K. M. Hanson and R. N. Silver (Kluwer, Dordrecht, 1996), p. 277
- [29] K. Życzkowski and M. Kuś, J. Phys. A 27, 4235 (1994).
- [30] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Lett. A 223, 1 (1996).
- [31] P. B. Slater, Statistical Thermodynamics Based on the Volume Elements of the Noninformative Maximal Monotone Metrics on Quantum States: Investigations of the Spin-1 and Spin-³/₂ Cases, quant-ph/9802019.
- [32] H. Wilf and A. Nijenhuis, Combinatorial Algorithms for Computers and Calculators, (Academic, New York, 1978).
- [33] V. E. Mkrtchian and V. O. Chaltykian, Opt. Commun. 3, 239 (1987).
- [34] R. A. Horn and C. A. Johnson, *Matrix Analysis*, (Cambridge Univ. Press, Cambridge, 1985).
- [35] S. L. Braunstein, C. M. Caves, R. Jozsa, N. Linden, S. Popescu, and R. Schack, Separability of Very Noisy Mixed States and Implications for NMR Quantum Computing, quant-ph/9811018.
- [36] G. Vidal and R. Tarrach, Robustness of Entanglement, quant-ph/9806094.
- [37] F. J. Bloore, J. Phys. A 9, 2059 (1976).
- [38] R. Bhatia, *Matrix Analysis*, (Springer-Verlag, New York, 1997).
- [39] D. Petz, Lin. Alg. Applics. 244, 81 (1996).
- [40] D. Petz, J. Math. Phys. 35, 780 (1994).
- [41] R. Balian, Y. Alhassid, and H. Reinhardt, Phys. Rep. 131, 1 (1986).
- [42] F. Qi, Proc. R. Soc. Lond. A 454, 2723 (1998).
- [43] P. B. Slater, J. Phys. A 29, L601 (1996).
- [44] K. Życzkowski, On the Volume of the Set of Mixed Entangled States. II, quant-ph/9902050.
- [45] M. Byrd, J. Math. Phys. 39, 6125 (1998).
- [46] M. Poźniak, K. Zyczkowski and M. Kuś, J. Phys. A 31, 1059 (1998).
- [47] P. B. Slater, Comment on "Volume of the Set of Separable States", quant-ph/9904014.

TABLE I. Quantum-theoretic statistical analyses based on the minimal, KMB and maximal monotone metrics. The parameters n_1 and n_2 determine the resolution of simplicial decompositions and square grids for trial diagonal and off-diagonal entries, respectively. The variable p denotes the associated probability of disentanglement and d, the averaged degree of entanglement. The only generated density matrices which have been omitted from consideration are those which correspond precisely to degenerate states, that is, det $\rho = 0$. The results are tabulated in increasing order of the total number of density matrices generated — given in the third column.

n_1	n_2	ρ	$\rho_{separable}$	p_{min}	p_{KMB}	p_{max}	d_{min}	d_{KMB}	d_{max}
23	7	1,340,928	356,096	.111102	.0873186	.0846153	.18206	.208022	.248457
35	6	$1,\!425,\!216$	$467,\!424$.193939	.178191	.0980763	.250552	.229161	.154696
30	7	2,919,680	806,400	.119669	.17601	.749588	.184696	.170555	.00940062
45	6	3,033,084	$987,\!484$.220936	.224241	.293623	.248278	.243704	.144297
4	12	4,228,817	$1,\!634,\!577$.249824	.232509	.309404	.117852	.118964	.0886535
19	8	4,443,408	$1,\!284,\!816$.147968	.123283	.0554999	.18686	.184149	.17713
8	10	$4,\!645,\!163$	$1,\!230,\!411$.11469	.0982187	.179838	.18989	.192975	.143422
35	7	4,673,024	$1,\!286,\!656$.0927196	.0594478	.152368	.220529	.26694	.156327
13	9	$5,\!540,\!864$	$1,\!341,\!440$.0756821	.0361165	.00173733	.208862	.252169	.349782
6	11	6,161,152	1,703,808	.114669	.0859676	.155728	.187623	.205257	.119427
10	10	7,103,372	$2,\!232,\!836$.195802	.188155	.243187	.131533	.130076	.103806
23	8	8,026,372	$2,\!286,\!148$.109203	.0747959	.000112443	.24179	.260264	.296032
50	7	$13,\!522,\!176$	3,705,472	.10125	.072653	.687396	.206883	.230803	.0422876
19	9	$16,\!603,\!136$	4,096,000	.0920722	.083066	.0820477	.186361	.188775	.190464
$\overline{7}$	12	$27,\!658,\!276$	$6,\!897,\!940$.0779479	.0553514	.31028	.221055	.256207	.0959608
35	8	28,582,224	$8,\!104,\!555$.140105	.122355	.199223	.22233	.228248	.118103
15	10	29,328,236	8,006,017	.120671	.0963975	.198144	.194026	.210798	.134718
11	11	36,913,664	9,049,600	.104142	.0891733	.164511	.177627	.179959	.122263
25	9	37,671,424	9,440,000	.0837076	.0343513	.0027555	.202037	.229187	.198646
8	12	40,487,643	$10,\!493,\!443$.109191	.0873665	.16817	.185359	.201147	.134193
27	9	47,381,504	$11,\!798,\!016$.0900729	.0619535	.169748	.197141	.219559	.124459
6	13	47,815,680	$12,\!558,\!464$.100372	.0710627	.112995	.19102	.205754	.149789
18	10	52,099,496	13,733,736	.104203	.0821789	.0834828	.205028	.215253	.198921
29	9	58,888,704	$14,\!645,\!792$.0886606	.054769	.00175691	.190217	.201052	.150746
9	12	$58,\!900,\!696$	$14,\!677,\!208$.0933132	.0726818	.176224	.20039	.218327	.131068
13	11	$60,\!453,\!376$	$14,\!847,\!232$.0848635	.0569472	.062366	.193203	.213676	.208112
19	10	$61,\!584,\!896$	$16,\!090,\!832$.0865202	.052442	.155109	.226729	.245587	.127568
30	9	$65,\!276,\!416$	$16,\!252,\!736$.0892746	.0685755	.0839835	.20221	.239333	.0959485
32	9	79,412,992	19,729,792	.0856388	.0562649	.123084	.200964	.232005	.141376
21	10	$83,\!685,\!188$	$21,\!982,\!132$.0993573	.062451	.138054	.20348	.228876	.114666
15	11	92,920,832	$22,\!811,\!392$.0848161	.0550803	.147779	.193381	.211392	.152707
34	9	94,713,344	$23,\!606,\!336$.0913722	.0638694	.131152	.197612	.230137	.2171
22	10	96,084,402	$25,\!272,\!244$.106789	.0808784	.0769463	.205559	.23114	.187346

TABLE II. Reanalyses of anomalous results (Table I) for $n_1 = 30$, $n_2 = 7$, using varying thresholds of degeneracy, as indexed by det ρ , below which the generated density matrices are rejected from further consideration.

threshold on $\det\rho$	ρ	$\rho_{separable}$	p_{min}	p_{KMB}	p_{max}	d_{min}	d_{KMB}	d_{max}
0	2,919,680	806,400	.119669	.17601	.749588	.184696	.170555	.00940062
$\frac{1}{256} \cdot 10^{-4}$	2,913,536	801,792	.102275	.0664226	.00698098	.189535	.201332	.0983051
$\frac{-10}{256} \cdot 10^{-3}$	2,856,704	$796,\!672$.123362	.108123	.386157	.180807	.191661	.101308
$\frac{\frac{230}{1}}{256} \cdot 10^{-2}$	$2,\!381,\!312$	$724,\!864$.180938	.171253	.260637	.14053	.140938	.101575

TABLE III. Results of *random* searches of the fifteen-dimensional parameter space, using differing values of the radius (r) of the circle in the complex plane centered at (0,0), from which the possible off-diagonal entries of the 4×4 density matrices are chosen.

r	searches	ρ	$\rho_{separable}$	p_{min}	p_{KMB}	p_{max}	d_{min}	d_{KMB}	d_{max}
1/2	847,500,000	49	12	.0711773	.0548709	.0269786	.197979	.2155	.252218
5/12	387,900,000	175	47	.098809	.0408168	.00218239	.1736	.18024	.194678
1/3	351,700,000	2,438	619	.0853483	.0654313	.180815	.201774	.22259	.147024
1/4	$74,\!300,\!000$	15,701	3,912	.0846071	.0364071	.00148944	.17873	.20229	.290831

TABLE IV. Analogues for the 2×3 systems of the results of Table I

n_1	n_2	ρ	$ ho_{separable}$	p_{min}	p_{KMB}	p_{max}	d_{min}	d_{KMB}	d_{max}
8	8	7,581	5,205	.643091	.654085	.678443	.0254769	.024673	.0230053
11	6	35,268	9,828	.000171812	.000192414	.000212583	.592094	.592009	.591929
14	6	$149,\!607$	59,727	.368812	.425491	.46159	.147718	.118212	.10094
15	6	158,522	59,185	.147933	.152787	.128344	.299006	.281777	.235139
13	9	245,760	8,448	.034375	.034375	.034375	.185994	.185994	.185994
9	8	$235,\!616$	$65,\!840$.234146	.262858	.266667	.0803952	.0654659	.0637942
15	9	245,760	768	.003125	.003125	.003125	.182635	.182635	.182635
14	9	$368,\!640$	9,600	.0260417	.0260417	.0260417	.188924	.188924	.188924
16	6	$370,\!479$	$131,\!055$.00584908	.0172682	.359634	.581471	.558051	.119815
17	6	$557,\!304$	164,264	.000248636	.000243731	.000240557	.59097	.591328	.591453
10	8	$579,\!186$	150,090	.173755	.174011	.174992	.184237	.194057	.195405
12	8	$1,\!352,\!182$	303,022	.0463408	.0872936	.19522	.383656	.307713	.117593
11	8	$1,\!593,\!588$	520,068	.299328	.309032	.326253	.0995446	.0968566	.0940759
22	6	$2,\!875,\!965$	$974,\!818$.0160034	.125932	.0721558	.562473	.374128	.259626
24	6	$3,\!870,\!989$	1,360,213	.192395	.19933	.314609	.292782	.268782	.140023
13	8	4,408,872	881,397	.0551892	.0659333	.256791	.294217	.259817	.0915806
14	8	6,073,071	882,781	.000948857	.000644427	.00952389	.61686	.476343	.359429
16	8	$7,\!373,\!379$	$1,\!609,\!392$.103614	.121313	.203796	.204274	.183523	.119163
26	6	$7,\!696,\!926$	$2,\!514,\!130$.1132	.111881	.0999924	.368272	.289389	.229179
15	8	$15,\!603,\!746$	$2,\!109,\!650$.000899908	.00162305	.00114651	.63942	.599576	.461553
17	8	$19,\!413,\!528$	$3,\!311,\!159$.054504	.0562656	.0398168	.323892	.298912	.259056
18	8	29,075,408	$5,\!061,\!131$.081246	.0737912	.0958513	.2304	.215317	.181838
20	8	45,002,652	7,755,473	.00401427	.0131804	.0525791	.606695	.487055	.206391

List of Tables

Quantum-theoretic statistical analyses based on the minimal, KMB and maximal monotone metrics.	
The parameters n_1 and n_2 determine the resolution of simplicial decompositions and square grids for	
trial diagonal and off-diagonal entries, respectively. The variable p denotes the associated probability	
of disentanglement and d , the averaged degree of entanglement. The only generated density matrices	
which have been omitted from consideration are those which correspond precisely to degenerate states,	
that is, det $\rho = 0$. The results are tabulated in increasing order of the total number of density matrices	
generated — given in the third column.	12
Reanalyses of anomalous results (Table I) for $n_1 = 30$, $n_2 = 7$, using varying thresholds of degeneracy,	
as indexed by det ρ , below which the generated density matrices are rejected from further consideration.	12
Results of random searches of the fifteen-dimensional parameter space, using differing values of the	
radius (r) of the circle in the complex plane centered at $(0,0)$, from which the possible off-diagonal	
entries of the 4×4 density matrices are chosen	13
Analogues for the 2×3 systems of the results of Table I $\ldots \ldots $	13
	Quantum-theoretic statistical analyses based on the minimal, KMB and maximal monotone metrics. The parameters n_1 and n_2 determine the resolution of simplicial decompositions and square grids for trial diagonal and off-diagonal entries, respectively. The variable p denotes the associated probability of disentanglement and d , the averaged degree of entanglement. The only generated density matrices which have been omitted from consideration are those which correspond precisely to degenerate states, that is, det $\rho = 0$. The results are tabulated in increasing order of the total number of density matrices generated — given in the third column

List of Figures

1	Conditional probability — based on the minimal monotone (Bures) metric — for $N = 2 \times 2$ of finding	
	a separable state, given a certain range (of width $.05$) of the participation ratio R , for the scenario	
	$n_1 = 22, n_2 = 10$	7
2	Conditional probability — based on the KMB-metric for $N = 2 \times 2$ — of finding a separable state,	
	given a certain range (of width .05) of the participation ratio R, for the scenario $n_1 = 22, n_2 = 10$	8
3	Probability — based on the minimal monotone (Bures) metric — for $N = 2 \times 2$ of finding a quantum	
	state (whether separable or not), given a certain range (of width $.05$) of the participation ratio R , for	
	the scenario $n_1 = 22, n_2 = 10$	8
4	Probability — based on the KMB-metric – for $N = 2 \times 2$ of finding a quantum state (whether separable	
	or not), given a certain range (of width .05) of the participation ratio R, for the scenario $n_1 = 22, n_2 = 10$	9