## Information, Relative Entropy of Entanglement and Irreversibility

L. Henderson and V. Vedral

Centre for Quantum Computation, Clarendon Laboratory, University of Oxford, Parks Road OX1 3PU

## Abstract

Previously proposed measures of entanglement, such as entanglement of formation and assistance, are shown to be special cases of the relative entropy of entanglement. The difference between these measures for an ensemble of mixed states is shown to depend on the availability of classical information about particular members of the ensemble. Based on this, relations between relative entropy of entanglement and mutual information are derived.

In quantifying entanglement, a number of measures have been proposed. For bipartite pure states,  $\rho_{AB}$ , the Von Neumann entropy of the reduced density matrix of either subsystem,  $\rho_A$  or  $\rho_B$ , has been found to be a good and unique measure, [1],[2]. Relative entropy of entanglement has been proposed as a measure which extends to mixed states, [3], [4]. Loosely speaking, it quantifies how 'far' an entangled state is from the set of disentangled states. Entanglement of mixed states has also been characterised by the 'entanglement of formation', [5], [6], and by the 'entanglement of distillation', [5]. Rather surprisingly, use of entanglement in mixed states is not reversible in the sense that all the entanglement required to construct a particular mixed state cannot be distilled out again, so the entanglement of formation is greater than the entanglement of distillation, [4]. In this paper, we clarify the role of classical information about the identity of particular members of an ensemble of mixed states, and show that the loss of such information is responsible for the difference between the entanglement of formation and the entanglement of distillation. We provide a unifying frame-work for entanglement measures by showing how previously proposed measures are special cases of the relative entropy of entanglement. This gives a strong physical argument for using quantum relative entropy as a unique way to understand entanglement in general.

Suppose that Alice and Bob share a state described by the density matrix  $\rho_{AB}$ . The state  $\rho_{AB}$  has an infinite number of different decompositions  $\varepsilon = \{ |\psi_{AB}^i\rangle \langle \psi_{AB}^i|, p_i \}$ , into pure states  $|\psi_{AB}^i\rangle$ , with probabilities  $p_i$ , [7]. We denote the mixed state  $\rho_{AB}$  written in decomposition  $\varepsilon$  by

$$\rho_{AB}^{\varepsilon} = \sum_{i} p_{i} \left| \psi_{AB}^{i} \right\rangle \left\langle \psi_{AB}^{i} \right| \tag{1}$$

The entanglement of formation is the average entanglement of the pure states, minimised over all decompositions, [6]:

$$E_F(\rho_{AB}) = \min_{\varepsilon} \sum_i p_i S(\rho_B^i)$$
(2)

Here  $\rho_B^i$  is the reduced density matrix for subsystem A of the pure state  $|\psi_{AB}^i\rangle\langle\psi_{AB}^i|$ . The physical importance of entanglement of formation lies in the fact that it is possible to convert an ensemble of m maximally entangled singlets into a smaller number, n, of non-maximally entangled states,  $\rho_{AB}^{\otimes n}$ , using only local operations and classical communication, [5], and entanglement of formation is the asymptotic conversion ratio,  $\frac{m}{n}$ , in the limit of infinitely many copies.

The 'entanglement of distillation',  $E_D(\rho_{AB})$ , is the number of maximally entangled singlets per copy of  $\rho_{AB}$  which can be distilled from an asymptotically large ensemble of copies of  $\rho_{AB}$  by a purification procedure involving only local operations and classical communication, [5]. For a mixed state, it is lower than or equal to the entanglement of formation, [4].

Relative entropy of entanglement of the mixed state is defined as

$$E_{RE}(\rho_{AB}) = \min_{\sigma_{AB} \in D} S(\rho_{AB} || \sigma_{AB})$$

where  $S(\rho||\sigma) = Tr(\rho \log \rho - \rho \log \sigma)$  is the quantum relative entropy, [4]. The minimum is taken over D, the set of completely disentangled or 'separable' states. A state is separable if it can be written as a convex combination of product states  $\sigma = \sum_i p_i \sigma_A^i \otimes \sigma_B^i$ , with  $\sum_i p_i = 1$ . The relative entropy of entanglement provides an upper bound for the distillable entanglement, [4]. The known relationships between the different measures of entanglement for mixed states are therefore  $E_D(\rho_{AB}) \leq E_{RE}(\rho_{AB}) \leq E_F(\rho_{AB})$ . Equality holds for pure states, where all the measures reduce to the Von Neumann entropy,  $S(\rho_A) = S(\rho_B)$ . We will give a straight-forward argument for the second inequality later in the paper. Protocols for formation and distillation of pure and mixed entangled states have been introduced [1], [5]. We first briefly review the pure state procedures, and then go on to discuss the role of classical information and relative entropy of entanglement in the mixed state case.

The basis of formation is that Alice and Bob would like to create an ensemble of n copies of the non-maximally entangled state,  $\rho_{AB}$ , using only local operations, classical communication, and a number of maximally entangled pairs. It is customary to consider the process of formation which consumes the least entanglement, since the only 'cost' in communication is due to the use of entanglement resources, or sending information down a quantum channel, and classical communication costs nothing. Distillation is the reverse process, where Alice and Bob share an ensemble of n copies of the non-maximally entangled state,  $\rho_{AB}$ , and would like to extract the largest number of maximally entangled pairs using only local operations and classical communications.

Formation of an ensemble of n non-maximally entangled *pure* states,  $\rho_{AB} = |\psi_{AB}\rangle \langle \psi_{AB}|$  is achieved by the following protocol. Alice first prepares the states she would like to share with Bob locally. She then uses Schumacher compression, [8], to compress these states into  $nS(\rho_B)$ states. One particle of each pair is then teleported to Bob using  $nS(\rho_B)$ maximally entangled pairs. Bob decompresses the states he receives and so ends up sharing n copies of  $\rho_{AB}$  with Alice. The entanglement of formation is therefore  $E_F(\rho_{AB}) = S(\rho_B)$ . For pure states, this process requires no classical communication in the asymptotic limit, [9]. The reverse process of distillation is accomplished using the Schmidt projection method, [1], which allows  $nS(\rho_B)$  maximally entangled pairs to be distilled in the limit as n becomes very large. Again, this process involves no classical communication between the separated parties. Therefore pure states are fully inter-convertible in the asymptotic limit.

The situation for mixed states is more complex. When any mixed state, denoted by Eq.(1), is created, it is necessarily part of an extended system whose state is pure. We will consider the pure states  $|\psi_{AB}^i\rangle$  in the mixture to be correlated to orthogonal states  $|m_i\rangle$  of a memory M. The extended system is in the pure state

$$|\psi_{MAB}\rangle = \sum_{i} \sqrt{p_i} |m_i\rangle |\psi^i_{AB}\rangle$$

If we have no access to the memory system, we trace over it to obtain the mixed state in Eq.(1). We will see that the amount of entanglement involved in the different entanglement manipulations of mixed states depends on the accessibility of the information in the memory at different stages. Note that a unitary operation on  $|\psi_{MAB}\rangle$  will convert it into another pure state  $|\phi_{MAB}\rangle$  with the same entanglement, [10],

$$|\phi_{MAB}\rangle = \sum_{j} \sqrt{q_j} |n_j\rangle |\phi_{AB}^j\rangle \tag{3}$$

Tracing over the memory in this case gives another decomposition,  $\zeta = \{ |\phi_{AB}^{j}\rangle \langle \phi_{AB}^{j} |, q_{j} \}$ , of  $\rho_{AB}$  into pure states

$$\rho_{AB}^{\zeta} = \sum_{j} q_{j} \left| \phi_{AB}^{j} \right\rangle \left\langle \phi_{AB}^{j} \right| \tag{4}$$

The reduction of the pure state, (3), to the mixed state, (4), may be regarded as due to a projection-valued measurement on the memory with operators  $\{E_j = |n_j\rangle \langle n_j|\}$ .

Consider first the protocol of formation by means of which Alice and Bob come to share an ensemble of n mixed state  $\rho_{AB}$ . Alice first creates the mixed states locally by preparing a collection of n states in a particular decomposition,  $\varepsilon = \{ |\psi_{AB}^i\rangle \langle \psi_{AB}^i|, p_i \}$  by making  $np_i$  copies of each pure state  $|\psi_{AB}^i\rangle$ . At the same time a memory system entangled to the pure states is generated, which keeps track of the identity of each member of the ensemble. Note that as long as we consider Alice's entire environment, the state of subsystems A and B together with the memory may always be taken to be pure. Later, we will consider the situation in which Alice's memory is decohered. There are then three ways for her to share these states with Bob. First of all, she may simply compress subsystem B to  $nS(\rho_B)$  states, and teleport these to Bob using  $nS(\rho_B)$  maximally entangled pairs. The choice of which subsystem to teleport is made so as to minimise the amount of entanglement required, so that  $S(\rho_B) \leq S(\rho_A)$ . The teleportation in this case would require no classical communication in the asymptotic limit, just as for pure states, [9]. The state of the whole system which is created by this process is an ensemble of pure states  $|\psi_{MAB}\rangle$ , where subsystems M and A are on Alice's side and subsystem B is on Bob's side. In terms of entanglement resources, however, this process is not the most efficient way for Alice to send the states to Bob. She may do it more efficiently by using the memory system of  $|\psi_{MAB}\rangle$  to identify blocks of  $np_i$  members in each pure state  $|\psi_{AB}^i\rangle$ , and applying compression to each block to give  $np_i S(\rho_B^i)$ states. Then the total number of maximally entangled pairs required to teleport these states to Bob is  $n \sum_{i} p_i S(\rho_B^i)$ , which is clearly less than  $nS(\rho_B)$ , by concavity of the entropy. The amount of entanglement required clearly depends on the decomposition of the mixed state  $\rho_{AB}$ . However, in order to decompress these states, Bob must also be able to identify which members of the ensemble are in which state. Therefore Alice must also send him the memory system. She now has two options. She may either teleport the memory to Bob, which would use more entanglement resources. Or she may communicate the information in the memory classically, with no further use of entanglement. When Alice uses the minimum entanglement decomposition,  $\varepsilon = \{|\psi_{AB}^i\rangle \langle \psi_{AB}^i|, p_i\}$ , this process, originally introduced by Bennett *et al.*, [5], makes the most efficient use of entanglement, consuming only the entanglement of formation of the mixed state,  $E_F(\rho_{AB}) = \sum_i p_i S(\rho_B^i)$ . We may think of the classical communication between Alice and Bob in one of two equivalent ways. Alice may either measure the memory locally to decohere it, and then send the result to Bob classically, or she may send the memory through a completely decohering quantum channel. In this case, the interaction with the channel is given by

$$\begin{aligned} |\psi_{MAB}\rangle \left|c\right\rangle &= \sum_{i} \sqrt{p_{i}} |m_{i}\rangle |\psi_{AB}^{i}\rangle \left|c\right\rangle \\ &\longrightarrow \sum_{i} \sqrt{p_{i}} |\psi_{AB}^{i}\rangle \left|m_{i}\rangle \left|c_{i}\right\rangle \end{aligned}$$

where  $|c\rangle$  is the initial state of the channel and  $\{|c_i\rangle\}$  are orthogonal channel states. Since Alice and Bob have no access to the channel, the state of the whole system which is created by this process is the mixed state

$$\rho_{ABM}^{\varepsilon} = \sum_{i} p_{i} |\psi_{AB}^{i}\rangle \langle \psi_{AB}^{i}| \otimes |m_{i}\rangle \langle m_{i}|$$
(5)

where Bob is classically correlated to the AB subsystem. Bob is then able to decompress his states using the memory to identify members of the ensemble.

Once the collection of n pairs is shared between Alice and Bob, it is converted into an ensemble of n mixed states  $\rho_{AB}$  by destroying access to the memory which contains the information about the state of any particular member of the ensemble. It is the loss of this information which is responsible for the fact that entanglement of distillation is lower than than entanglement of formation<sup>1</sup>. Distillation is not carried out by people like Alice and Bob who have access to the memory, but by people who have just received the ensemble of n mixed states  $\rho_{AB}$ with no further information. If Alice and Bob were to carry out the distillation, they could obtain as much entanglement from the ensemble

<sup>&</sup>lt;sup>1</sup>The relation between classical information and distillable entanglement was previously discussed by Eisert *et al.*, [11], in a different context.

as was required to form it. In the case where Alice and Bob share an ensemble of the pure state  $|\psi_{MAB}\rangle$ , they would simply apply the Schmidt projection method, [1]. The relative entropy of entanglement gives the upper bound to distillable entanglement,

$$E_{RE}(|\psi_{(MA):B}\rangle\langle\psi_{(MA):B}|) = S(\rho_B) \tag{6}$$

which is the same as the amount of entanglement required to create the ensemble of pure states, as described above. Here MA and B are spatially separated subsystems on which joint operations may not be performed. In our notation, we use a colon to separate the local subsystems.

On the other hand, if Alice used the least entanglement for producing an ensemble of the mixed state  $\rho_{AB}$ , together with classical communication, the state of the whole system is an ensemble of the mixed state  $\rho_{ABM}^{\epsilon}$ , and the process is still reversible. Because of the classical correlation to the states  $|\psi_{AB}^i\rangle$ , Alice and Bob may identify blocks of members in each pure state  $|\psi_{AB}^i\rangle$ , and apply the Schmidt projection method to them, giving  $np_i S(\rho_B^i)$  maximally entangled pairs, and hence a total entanglement of distillation of  $\sum_i p_i S(\rho_B^i)$ . The relative entropy of entanglement again quantifies the amount of distillable entanglement from the state  $\rho_{ABM}^{\epsilon}$  and is given by

$$E_{RE}(\rho_{A:(BM)}^{\varepsilon}) = \min_{\sigma_{ABM} \in D} S(\rho_{ABM}^{\varepsilon} || \sigma_{ABM})$$

The disentangled state which minimises the relative entropy is  $\sigma_{ABM} = \sum_i p_i \sigma_{AB}^i \otimes |m_i\rangle \langle m_i|$ , where  $\sigma_{AB}^i$  is obtained from  $|\psi_{AB}^i\rangle \langle \psi_{AB}^i|$  by deleting the off-diagonal elements in the Schmidt basis. This is the minimum because the state  $\rho_{MAB}$  is a mixture of the orthogonal states  $|m_i\rangle |\psi_{AB}^i\rangle$ , and for a pure state  $|\psi_{AB}^i\rangle$ , the disentangled state which minimises the relative entropy is  $\sigma_{AB}^i$ . The minimum relative entropy of the extended system is then

$$S(\rho_{ABM}^{\varepsilon}||\sigma_{ABM}) = \sum_{i} p_{i}S(\rho_{B}^{i})$$

This relative entropy,  $E_{RE}(\rho_{A:(BM)}^{\varepsilon})$ , has previously been called 'entanglement of projection', [12], because the measurement on the memory projects the pure state of the full system into a particular decomposition. The minimum of  $E_{RE}(\rho_{A:(BM)}^{\varepsilon})$  over all decompositions is equal to the entanglement of formation of  $\rho_{AB}$ . However, Alice and Bob may choose to create the state  $\rho_{AB}$  by using a decomposition with higher entanglement than the entanglement of formation. The maximum of  $E_{RE}(\rho_{A:(BM)}^{\varepsilon})$  over all possible decompositions is called the 'entanglement of assistance' of  $\rho_{AB}$ , [13]. Because  $E_{RE}(\rho_{A:(BM)}^{\varepsilon})$  is a relative entropy, it is invariant under local operations and non-increasing under general operations, properties which are conditions for a good measure of entanglement, [4]. However, unlike  $E_{RE}(\rho_{AB})$  and  $E_F(\rho_{AB})$ , it is not zero for completely disentangled states. In this sense, the relative entropy of entanglement,  $E_{RE}(\rho_{A:(BM)}^{\varepsilon})$ , defines a class of entanglement measures interpolating between the entanglement of formation and entanglement of assistance. Note that an upper bound for the entanglement of assistance,  $E_A$ , can be shown using concavity, [13], to be  $E_A(\rho_{AB}) \leq \min[S(\rho_A), S(\rho_B)]$ . This bound can also be shown from the fact that the distillable entanglement from any decomposition,  $E_{RE}(\rho_{A:(BM)}^{\varepsilon}) \leq E_A(\rho_{AB})$  cannot be greater than the entanglement of the original pure state.

We may also derive relative entropy measures that interpolate between the relative entropy of entanglement and the entanglement of formation by considering non-orthogonal measurements on the memory. First of all, the fact that the entanglement of formation is in general greater than the upper bound for entanglement of distillation, emerges as a property of the relative entropy, namely that it cannot increase under the local operation of tracing one subsystem, [10],

$$E_F(\rho_{AB}) = \min_{\sigma_{ABM} \in D} S(\rho_{ABM} || \sigma_{ABM}) \ge \min_{\sigma_{AB} \in D} S(\rho_{AB} || \sigma_{AB})$$

In general, the loss of the information in the memory may be regarded as a result of an imperfect classical channel. This is equivalent to Alice making a non-orthogonal measurement on the memory, and sending the result to Bob. In the most general case,  $\{E_i = A_i A_i^+\}$  is a POVM performed on the memory. The decomposition corresponding to this measurement is composed of mixed states,  $\xi = \{q_i, Tr_M(A_i\rho_{MAB}A_i^+)\},\$ where  $q_i = Tr(A_i \rho_{MAB} A_i^+)$ . The relative entropy of entanglement of the state  $\rho_{MAB}^{\xi}$ , when  $\xi$  is a decomposition of  $\rho_{AB}$  resulting from a nonorthogonal measurement on M, defines a class of entanglement measures interpolating between the relative entropy of entanglement and the entanglement of formation of the state  $\rho_{AB}$ . In the extreme case where the measurement gives no information about the state  $\rho_{AB}$ ,  $E_{RE}(\rho_{A;(BM)}^{\varepsilon})$ becomes the relative entropy of entanglement of the state  $\rho_{AB}$  itself. In between, the measurement gives partial information. We note that instead of an imperfect measurement, we may regard the memory itself as imperfect, in the sense that the memory states are non-orthogonal,  $\langle m_i | m_i \rangle \neq 0$  for  $i \neq j$ .

Now we will relate the loss of entanglement to the loss of information in the memory. As we have argued so far, there are two stages at which distillable entanglement is lost. The first is in the conversion of the pure state  $|\psi_{MAB}\rangle$  into a mixed state  $\rho_{ABM}$ . This happens because Alice uses a *classical* channel to communicate the memory to Bob. The second is due to the loss of the memory, M, taking the state  $\rho_{ABM}$  to  $\rho_{AB}$ . The amount of information lost may be quantified by the difference in mutual information between the respective states. Mutual information is a measure of correlations between the memory M and the system AB, giving the amount of information about AB which may be obtained from a measurement on M. The quantum mutual information between M and AB is defined as

$$I_Q(\rho_{M:(AB)}) = S(\rho_M) + S(\rho_{AB}) - S(\rho_{MAB})$$

The quantum mutual information of the pure state  $|\psi_{MAB}\rangle$  is  $I_Q(|\psi_{M:(AB)}\rangle \langle \psi_{M:(AB)}|) = 2S(\rho_{AB})$ , and of the mixed state in Eq.(5) is  $I_Q(\rho_{M:(AB)}) = S(\rho_{AB})$ . Therefore the mutual information loss in the first stage is  $\Delta I_Q = S(\rho_{AB})$ . There is a corresponding reduction in the relative entropy of entanglement, from the entanglement of the original pure state,  $E_{RE}(|\psi_{(MA):B}\rangle \langle \psi_{(MA):B}|)$ , to the entanglement of the mixed state  $E_{RE}(\rho_{A:(BM)}^{\varepsilon})$  for all decompositions  $\varepsilon$  arising as the result of an orthogonal measurement on the memory. We now show that when the mutual information loss is added to the relative entropy of entanglement of the original pure state,  $E_{RE}(|\psi_{(MA):B}|)$ , the result is greater than the relative entropy of entanglement of the original pure state, which occurs when  $E_{RE}(\rho_{A:(BM)}^{\varepsilon}) = E_F(\rho_{AB})$ :

$$E_{RE}(\left|\psi_{(MA):B}\right\rangle\left\langle\psi_{(MA):B}\right|) \le E_F(\rho_{AB}) + S(\rho_{AB}) \tag{7}$$

The proof goes as follows. Let  $\varepsilon = \{ |\psi_{AB}^i\rangle \langle \psi_{AB}^i|, p_i \}$  be the minimal entanglement decomposition giving rise to the entanglement of formation, see Eq.(2). Then,

$$S(\rho_{AB}) = \sum_{i} p_{i} S(|\psi_{AB}^{i}\rangle \langle \psi_{AB}^{i}|||\rho_{AB})$$
$$\geq \sum_{i} p_{i} S(\rho_{B}^{i}||\rho_{B})$$
$$= S(\rho_{B}) - \sum_{i} p_{i} S(\rho_{B}^{i})$$

The inequality results from the fact that the relative entropy does not increase under the local operation of tracing subsystem A, [10]. Using Eq.(6), and the fact that for this decomposition,  $E_F(\rho_{AB}) = \sum_i p_i S(\rho_B^i)$ , gives inequality (7).

A similar result may be proved for the second loss, due to loss of the memory. After this, the mutual information between the memory and AB of the state  $\rho_{AB}$  is zero. Therefore the mutual information lost in losing the memory is again  $\Delta I_Q = S(\rho_{AB})$ . The relative entropy of entanglement is reduced from  $E_{RE}(\rho_{A:(BM)}^{\varepsilon})$ , for any decomposition  $\varepsilon$  resulting from an orthogonal measurement on the memory, to  $E_{RE}(\rho_{AB})$ , the relative entropy of entanglement of the state  $\rho_{AB}$  with no memory. We show that when the mutual information loss is added to  $E_{RE}(\rho_{AB})$ , the result is greater than  $E_{RE}(\rho_{A:(BM)}^{\varepsilon})$ . In this case, the result is strongest for  $E_{RE}(\rho_{A:(BM)}^{\varepsilon}) = E_A(\rho_{AB})$ :

$$E_A(\rho_{AB}) \le E_{RE}(\rho_{AB}) + S(\rho_{AB}) \tag{8}$$

Let  $\zeta = \{ |\phi_{AB}^i\rangle \langle \phi_{AB}^i|, q_i \}$  be the maximal entanglement decomposition giving rise to the entanglement of assistance. Then

$$E_{RE}(\rho_{AB}) + S(\rho_{AB}) - E_{RE}(\rho_{A:(BM)}^{\zeta}) = -\min_{\sigma_{AB}\in D} Tr(\rho_{AB}\ln\sigma_{AB}) - \sum_{i} q_{i}S(\rho_{B}^{i}) = \sum_{i} q_{i}(-\min_{\sigma_{AB}\in D} \langle \phi_{AB}^{i}|\ln\sigma_{AB} | \phi_{AB}^{i} \rangle - S(\rho_{B}^{i})) \geq \sum_{i} q_{i}(\min_{\sigma_{AB}^{i}\in D} S(|\phi_{AB}^{i}\rangle \langle \phi_{AB}^{i}|||\sigma_{AB}^{i}) - S(\rho_{B}^{i})) = 0$$

The inequality holds because  $\sigma_{AB}$  is the disentangled state which minimises the relative entropy of the state  $\rho_{AB}$ , but may not minimise the relative entropy for each of the component pure states,  $|\phi_{AB}^i\rangle$ . Notice that if  $\rho_{AB}$  is a pure state, then  $S(\rho_{AB}) = 0$ , and equality holds.

Inequalities (7) and (8) provide lower bounds for  $E_F(\rho_{AB})$  and  $E_{RE}(\rho_{AB})$  respectively. They are of a form typical of irreversible processes in that restoring the information in M is not sufficient to restore the original correlations between M and AB. In particular, they express that the loss of entanglement between Alice and Bob at each stage must be accompanied by an even greater reduction in mutual information between the memory and subsystems AB. This raises the interesting open question of whether the inequalities (7) and (8) may be generalised to a relation of the kind

$$E_{RE}(\rho_{A:(BM)}) \le E_{RE}(\chi_{A:(BM)}) + I_Q(\rho_{M:(AB)}) - I_Q(\chi_{M:(AB)})$$
(9)

for any two entangled states,  $\rho$  and  $\chi$ , where  $\chi$  is obtained from  $\rho$  by any operation on the memory. This would give the physically reasonable property that loss of the information in M about AB is always greater than the loss of entanglement between the separated subsystems.

In summary, there are numerous decompositions of any bipartite mixed state into a set of states  $\rho_i$  with probability  $p_i$ . The average entanglement of states in each decomposition is given by the relative entropy of entanglement of the system extended by a memory whose orthogonal states are classically correlated to the states of the decomposition. This correlation records which state  $\rho_i$  any member of an ensemble of mixed states  $\rho_{AB}^{\otimes n}$  is in. It is available to parties involved in formation of the mixed state, but is not accessible to parties carrying out distillation. When the classical information is fully available, different decompositions give rise to different amounts of distillable entanglement, the highest being entanglement of assistance and the lowest, entanglement of formation. When access to the classical record is reduced, the amount of distillable entanglement is reduced. In the limit where no information is available, the distillable entanglement is given by the relative entropy of entanglement of the state  $\rho_{AB}$  itself, without the extension of the classical memory. Our work shows that relative entropy of entanglement provides a unifying measure for all cases, elucidating the role of classical information and the appearance of irreversibility in manipulations of mixed state entanglement.

## References

- C. H. Bennett, H. J. Bernstein, S. Popescu, and B. Schumacher, Phys. Rev. A 53, 2046 (1996).
- [2] S. Popescu and D. Rohrlich, Phys. Rev. A 56, R3319 (1997).
- [3] V. Vedral, M. B. Plenio, M. A. Rippin, and P. L. Knight, Phys. Rev. Lett. 78, 2275 (1997).
- [4] V. Vedral and M. B. Plenio, Phys. Rev. A 57, 1619 (1998).
- [5] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Phys. Rev. A 54, 3824 (1996).
- [6] W. K. Wootters, Phys. Rev. Lett. 80, 2245 (1998).
- [7] L. P. Hughston, R. Jozsa, and W. K. Wootters, Phys. Lett. A 183, 14 (1993).
- [8] B. Schumacher, Phys. Rev. A **51**, 2738 (1995).
- [9] H. Lo and S. Popescu, quant-ph/9902045 (1999).
- [10] G. Lindblad, Comm. Math. Phys. 40, 147 (1975).
- [11] J. Eisert *et al.*, quant-ph/ 9907021 (1999).
- [12] R. Garisto and L. Hardy, quant-ph/9808007 (1998).
- [13] D. P. DiVincenzo *et al.*, quant-ph/9803033 (1998).