

# Explanation of Quantum Mechanics

Gang Zhao

Department of Physics, Nankai University, Tianjing 300071, P.R. China

E-mail: zhaog@nankai.edu.cn

September 24, 2018

## Abstract

By assuming that the kinetic energy, potential energy, momentum, and some other physical quantities of a particle exist in the field out of the particle, the Schrödinger equation is an equation describing field of a particle, but not the particle itself.

That there are energy and momentum in electromagnetic field is an indisputable fact. According to Maxwell equation, density of electromagnetic energy and momentum in vacuum respectively is

$$W = \frac{1}{2}(\epsilon_0 E^2 + \frac{1}{\mu_0} B^2) \quad (1)$$

$$\mathbf{p} = \epsilon_0 \mathbf{E} \times \mathbf{B} \quad (2)$$

This means the energy and momentum of electromagnetic field exist in whole space. The Coulomb interaction of charges can be described in terms of field, too. Assume the distance of two charges  $Q_1, Q_2$  is  $\vec{r}$ , hence

$$\mathbf{F} = -\nabla U = -\nabla \int \frac{1}{2} \epsilon_0 \mathbf{E}^2 d\tau = -\nabla \int \frac{1}{2} \epsilon_0 \mathbf{E}_1 \cdot \mathbf{E}_2 d\tau = \frac{1}{4\pi\epsilon_0} Q_1 Q_2 \frac{\vec{r}}{r^3} \quad (3)$$

Note that the potential energy  $U$  is a spatial integration, the energy distributes in whole space. The energy in any volume element  $d\tau$  that contributes to potential energy is  $\frac{1}{2} \epsilon_0 E_1 \cdot E_2$ , where  $E_1, E_2$  are electric amplitudes of  $Q_1, Q_2$ , respectively.

Since the electromagnetic energy, momentum, potential energy of a particle exist in field, we make a crazy, but reasonable assumption: the kinetic energy, momentum, potential energy, and some other physical quantities of a particle are taken by the field (real field) out of the particle (energy distributes in space), but not be taken by the particle itself. Hence, if we know the movement equation of field (real field) out of a particle, we can

calculate the kinetic energy, momentum, potential energy, and some other physical quantities of the particle. So we should find the equation of field. The easiest and most natural way is to assume the Schrödinger equation  $i\hbar \frac{\partial}{\partial t} \psi = H\psi$  is the equation describing the movement of field.

In classical physics, the equation  $\frac{d}{dt} \psi = \alpha \nabla^2 \psi$  can be used as heat field equation [1], that we assume Schrödinger equation is the equation of field is reasonable.

As to the movement of a particle, we can accept L. De Broglie's idea [2] that particle moves led by wave, since we admit that particle and field are parts of a body.

#### References

1. Zhen-Qiu Lu, *Equations of Classical and Modern Mathematical Physics*, Shenhai, 1991.
2. L. De Broglie, *La physique quantique restera-telle indéterministe?* Gauthier-Villars, Paris, 1953.