Casorati Determinant Solutions for the Discrete Painlevé III Equation

Kenji Kajiwara

Department of Electrical Engineering, Doshisha University, Tanabe, Kyoto 610-03, Japan Yasuhiro Ohta Department of Applied Mathematics, Faculty of Engineering, Hiroshima University, 1-4-1 Kagamiyama, Higashi-Hiroshima, Hiroshima 724, Japan

and

Junkichi Satsuma

Department of Mathematical Sciences, University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo 153, Japan

Abstract

The discrete Painlevé III equation is investigated based on the bilinear formalism. It is shown that it admits the solutions expressed by the Casorati determinant whose entries are given by the discrete Bessel function. Moreover, based on the observation that these discrete Bessel functions are transformed to the q-Bessel functions by a simple variable transformation, we present a q-difference analogue of the Painlevé III equation.

1 Introduction

Recently, several interesting ordinary difference equations have been proposed as discrete analogues of the Painlevé equations. Among them, the discrete Painlevé $I(dP_I)$ and $II(dP_{II})$ are naturally derived in calculating the partition functions of some models in the theory of two-dimensional quantum gravity[1,2,3]. In other words, they have close relationship with the theory of orthogonal polynomials. This example implies the significance of discrete Painlevé equations. Moreover, they are expected to possess rich mathematical structures. Besides this derivation, the discrete Painlevé equations can be obtained in several context[4,5,6,7,8].

It is known that the discrete Painlevé equations admit several properties analogous to continuous ones, such as the existence of Lax pair, Bäcklund transformation, coalescence cascade and singularity confinement(SC) property[5,6,7,9,10,11,12]. However, are these properties sufficient to regard the equations really as the discrete Painlevé equations?

In the continuous case, the Painlevé equations are "integrated" in the sense that they are reduced to linear integral equations by means of inverse scattering transform from their Lax pairs[13]. In the discrete case, however, the dP_I equation is the only known example which is "integrated" based on its Lax pair.

In the continuous case, series of exact solutions which are expressed by determinants are obtained by using the Bäcklund transformation[14]. However, no attempt has been done to construct such solutions for the discrete case.

The coalescence cascade is a property of the Painlevé equations which is related to the confluence of singularities of linear ordinary differential equations in the theory of monodromy preserving deformation. Hence, this property is considered to have no direct relationship to the structure of their solution spaces.

The SC is a concept which was extracted as a common property from the class of integrable ordinary difference equations proposed by Quispel et al.[15] In this context, "integrable" means the possession of conserved quantities. Grammaticos et al. have employed this property as an integrability criterion of difference equations[8]. By applying this test to non-autonomous equations, they have proposed several discrete Painlevé equations[5]. Since the SC might be regarded as the discrete analogue of the Painlevé property, it provides us with a useful and powerful tool to detect "good" equations. However, we do not know whether the equations which pass the SC are really "integrable" in the above sense.

The Painlevé equations are characterized by the property that they do not possess movable singularities except for poles. It is also a significant property that they admit series of exact solutions expressed by determinants for special values of parameters, although their solutions are transcendental in general. We believe that the existence of such solutions is a direct reflection of the "integrability" of equations. From this viewpoint, we consider that the preservation of the structure of solution space is the most important for the discretization of Painlevé equations.

In the previous paper, we have shown that the dP_{II} admits a class of solutions expressed by the Casorati determinants of discrete Airy function[16]. They reduce to the Wronskian solutions of P_{II} in the continuous limit. In this paper, we investigate the solutions of special function type for the discrete Painlevé III (dP_{III}) equation. We will show that they are expressed in terms of the Casorati determinants of the discrete Bessel functions. The dP_{III} occupies a particular place among the discrete Painlevé equations, since this is the simplest equation which is derived only through the SC. Hence, the existence of such solutions may be a good check of the validity of SC.

Moreover, we will show that the discrete Bessel functions in our result are transformed to the q-Bessel functions by a simple independent variable transformation. Based on this observation, we present a q-difference analogue of the P_{III} .

In section 2, we present the explicit form of exact solutions for the dP_{III} . The derivation is discussed in detail by means of the bilinear formalism in section 3. In section 4, we present a *q*-difference analogue of the P_{III} . Finally in section 5, we give concluding remarks.

2 Solutions of dP_{III}

In this section, we investigate the solutions of $dP_{III}[5]$,

$$w_{n+1}w_{n-1} = \frac{\alpha w_n^2 + \beta \lambda^n w_n + \gamma \lambda^{2n}}{w_n^2 + \delta w_n + \alpha} , \qquad (1)$$

where α , β , γ , δ , and λ are parameters. Before going to the discrete case, let us remind of the special function type solutions for the P_{III} equation,

$$\frac{d^2u}{dx^2} = \frac{1}{u} \left(\frac{du}{dx}\right)^2 - \frac{1}{x}\frac{du}{dx} + \frac{1}{x}(\alpha u^2 + \beta) + \gamma u^3 + \frac{\delta}{u} , \qquad (2)$$

where α , β , γ and δ are parameters. It is known that eq.(2) admits the solutions expressed by the τ function[14],

$$\tau_{N}^{\nu} = \begin{vmatrix} J_{\nu} & D & J_{\nu} & \cdots & D^{N-1} J_{\nu} \\ D & J_{\nu} & D^{2} & J_{\nu} & \cdots & D^{N} & J_{\nu} \\ \vdots & \vdots & \ddots & \vdots \\ D^{N-1} J_{\nu} & D^{N} & J_{\nu} & \cdots & D^{2N-2} & J_{\nu} \end{vmatrix},$$
(3)

where $D = x \frac{d}{dx}$ and J_{ν} is the Bessel function of degree ν . The dependent variable u in eq.(2) is expressed as

$$u = \left(\log\frac{\tau_N^{\nu+1}}{\tau_{N+1}^{\nu}}\right)_x + \frac{\nu+N}{x} .$$
 (4)

Equation (4) gives the particular solutions of eq.(2) with the parameters,

$$\alpha = 2(N - \nu)$$
, $\beta = 2(\nu + N + 1)$, $\gamma = 1$, $\delta = -1$. (5)

To obtain the particular solutions of dP_{III} eq.(1), we start from the discrete Riccati equation[17]. In this case, it is easily shown that if w_n satisfies

$$w_{n+1} = -\frac{aw_n + \lambda^n}{w_n + d} , \qquad (6)$$

where a and d are constants, then w_n gives a solution of eq.(1). Assuming the fractional form,

$$w_n = \frac{F_n}{G_n} , \qquad (7)$$

and substituting this into eq.(6), we get

$$w_n = \frac{G_{n+1}}{G_n} - d , \qquad (8)$$

$$G_{n+2} + (a-d)G_{n+1} + (\lambda^n - ad)G_n = 0.$$
(9)

On the analogy of the continuous case, it is expected that eq.(9) is a discrete version of the Bessel equation. However, it is not trivial to identify eq.(9) with the Bessel equation. Instead of eqs.(8) and (9), we consider

$$w_n = \frac{J_\nu(n+1)}{J_\nu(n)} - q^\nu,$$
(10)

$$J_{\nu}(n+2) - (q^{\nu} + q^{-\nu})J_{\nu}(n+1) + \{1 + (1-q)^2 q^{2n}\}J_{\nu}(n) = 0.$$
(11)

We can show that if $J_{\nu}(n)$ satisfies eq.(11), then w_n in eq.(10) gives the solution of dP_{III} eq.(1) with the parameters,

$$\alpha = -1, \quad \beta = (q^{\nu} - q^{-\nu-2})(1-q)^2, \quad \gamma = \frac{(1-q)^4}{q^2}, \quad \delta = q^{\nu} - q^{-\nu}, \quad \lambda = q^2 .$$
(12)

Continuous limit of eq.(11) is given by putting $q = 1 + \epsilon$, $n = \frac{z}{\epsilon}$ and taking the limit of $\epsilon \to 0$, which yields

$$\frac{d^2 J_{\nu}}{dz^2} + (e^{2z} - \nu^2) J_{\nu} = 0 .$$
(13)

Equation (13) reduces to the Bessel equation in ordinary form if we put $x = e^{z}$. In this sense, eq.(11) is regarded as the discrete Bessel equation. We note that the same results for this simplest solution is also derived in ref.[12].

The critical point is that the above result can be extended to the $N \times N$ determinant solution. We here give only the results, leaving the derivation in the next section. We consider the τ function,

$$\tau_N^{\nu}(n) = \begin{vmatrix} J_{\nu}(n) & J_{\nu}(n+1) & \cdots & J_{\nu}(n+N-1) \\ J_{\nu}(n+2) & J_{\nu}(n+3) & \cdots & J_{\nu}(n+N+1) \\ \vdots & \vdots & \ddots & \vdots \\ J_{\nu}(n+2N-2) & J_{\nu}(n+2N-1) & \cdots & J_{\nu}(n+3N-3) \end{vmatrix},$$
(14)

where $J_{\nu}(n)$ satisfies the following contiguity relations,

$$J_{\nu}(n) - q^{-\nu} J_{\nu}(n+1) = -(1-q)q^n J_{\nu+1}(n) , \qquad (15)$$

$$J_{\nu}(n) - q^{\nu} J_{\nu}(n+1) = (1-q)q^n J_{\nu-1}(n) .$$
(16)

We notice that the discrete Bessel equation (11) is derived ¿from above contiguity relations. Then we can show that

$$w_n = \frac{\tau_{N+1}^{\nu}(n+1)\tau_N^{\nu+1}(n)}{\tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)} - q^{\nu+N} , \qquad (17)$$

gives the solutions of the dP_{III} ,

$$w_{n+1}w_{n-1} = \frac{\alpha w_n^2 + \beta q^{2n} w_n + \gamma q^{4n}}{w_n^2 + \delta w_n + \alpha},$$
(18)

with the parameters,

$$\begin{aligned} \alpha &= -q^{4N}, & \beta &= (q^{\nu+N} - q^{-\nu-N-2}) q^{8N} (1-q)^2 , \\ \gamma &= q^{2(6N-1)} (1-q)^4 , & \delta &= (q^{\nu-N} - q^{-\nu+N}) q^{2N} . \end{aligned}$$
 (19)

It is remarkable that the τ function has a similar asymmetric structure to that of dP_{II}[16]. Namely, the index of elements of the determinant, n, increases by two in one direction while by one in another direction.

3 Derivation of Results

In this section, we show that the τ function (14) actually gives the solution of eq.(18) with the parameters (19), through the dependent variable transformation (17). First, we give the bilinear difference equations satisfied by the τ function (14). Secondly, we show that eq.(18) follows from the bilinear difference equations.

3.1 Bilinear Difference Equations

The bilinear difference equations for the τ function (14) are given as

$$\tau_{N+1}^{\nu}(n) \ \tau_N^{\nu+1}(n+1) - q^{-\nu-N} \tau_{N+1}^{\nu}(n+1) \ \tau_N^{\nu+1}(n) = -(1-q)q^{n+2N} \tau_{N+1}^{\nu+1}(n) \ \tau_N^{\nu}(n+1) \ ,$$
(20)

$$\tau_{N+1}^{\nu+1}(n) \ \tau_N^{\nu}(n+1) - q^{\nu-N+1} \tau_{N+1}^{\nu+1}(n+1) \ \tau_N^{\nu}(n) = (1-q)q^{n+2N} \tau_{N+1}^{\nu}(n) \ \tau_N^{\nu+1}(n+1) \ ,$$
(21)

$$\tau_{N+1}^{\nu}(n) \ \tau_N^{\nu+1}(n+3) - q^{-\nu-N} \tau_{N+1}^{\nu}(n+1) \ \tau_N^{\nu+1}(n+2) = -(1-q)q^n \tau_{N+1}^{\nu+1}(n) \ \tau_N^{\nu}(n+3) \ ,$$
(22)

$$\tau_{N+1}^{\nu+1}(n) \ \tau_N^{\nu}(n+3) - q^{\nu-N+1} \tau_{N+1}^{\nu+1}(n+1) \ \tau_N^{\nu}(n+2) = (1-q)q^n \tau_{N+1}^{\nu}(n) \ \tau_N^{\nu+1}(n+3) \ ,$$
(23)

$$\tau_{N+1}^{\nu}(n+2) \ \tau_{N}^{\nu+1}(n+1) - q^{2N}(q^{\nu-N} + q^{-\nu+N})\tau_{N+1}^{\nu}(n+1) \ \tau_{N}^{\nu+1}(n+2) + q^{4N}\{1 + (1-q)^2 q^{2n}\}\tau_{N+1}^{\nu}(n) \ \tau_{N}^{\nu+1}(n+3) = 0 \ .$$
(24)

In the following, we give a detail of the proof that eqs.(20)-(24) hold for the τ function (14).

(1) Difference Formulas

The Plücker relations are the quadratic identities for determinants some of whose columns are shifted. Bilinear equations are obtained from the relations. Therefore the first step is to construct several formulas which relate the "shifted determinants". For usual integrable systems, such as the KP, Toda lattices and discrete KP, the entries of their determinant representation of τ functions satisfy quite simple linear differential or difference equations, and hence we can easily construct the differential or difference formulas for the τ functions (see, for example, [18,19,20,21]). On the other hand, in this case, the entries of the τ functions satisfy linear, but variable-coefficient difference equations as shown below, which makes the calculations somewhat technical.

We now consider the τ function (14) provided that the element $J_{\nu}(n)$ satisfies the contiguity relations (15) and (16). Let us introduce a notation for simplicity,

$$\tau_N^{\nu}(n) = |0_{\nu}, 1_{\nu}, \cdots, N - 2_{\nu}, N - 1_{\nu}|, \qquad (25)$$

where k_{ν} stands for the column which begins with $J_{\nu}(n+k)$,

$$k_{\nu} = \begin{pmatrix} J_{\nu}(n+k) \\ J_{\nu}(n+k+2) \\ J_{\nu}(n+k+4) \\ \vdots \end{pmatrix} .$$
(26)

Here the height of the column k_{ν} is N. However in the following, we use the same symbol k_{ν} for the bigger determinant $\tau_{N+1}^{\nu}(n)$ because there is no possibility of confusion. So the size of k_{ν} should be read appropriately case by case. By adding (i+1)-th column multiplied by $-q^{-\nu}$ to *i*-th column and using eq.(15) for $i = 1 \sim (N-1)$, we have

$$\begin{aligned} \tau_N^{\nu}(n) &= \{-(1-q)\}^{N-1} q^{\frac{(2n+N-2)(N-1)}{2}} \\ &\times \begin{vmatrix} J_{\nu+1}(n) & \cdots & J_{\nu+1}(n+N-2) & J_{\nu}(n+N-1) \\ q^2 J_{\nu+1}(n+2) & \cdots & q^2 J_{\nu+1}(n+N) & J_{\nu}(n+N+1) \\ \vdots & \cdots & \vdots & \vdots \\ q^{2N-2} J_{\nu+1}(n+2N-2) & \cdots & q^{2N-2} J_{\nu+1}(n+3N-4) & J_{\nu}(n+3N-3) \end{vmatrix} \end{aligned} (27) \\ &= \{-(1-q)\}^{N-1} q^{\frac{(2n+3N-2)(N-1)}{2}} \\ &\times \begin{vmatrix} J_{\nu+1}(n) & \cdots & J_{\nu+1}(n+N-2) & J_{\nu}(n+N-1) \\ J_{\nu+1}(n+2) & \cdots & J_{\nu+1}(n+N) & q^{-2} J_{\nu}(n+N+1) \\ \vdots & \cdots & \vdots & \vdots \\ J_{\nu+1}(n+2N-2) & \cdots & J_{\nu+1}(n+3N-4) & q^{-2N+2} J_{\nu}(n+3N-3) \end{vmatrix} \\ &= \{-(1-q)\}^{N-1} q^{\frac{(2n+3N-2)(N-1)}{2}} & |0_{\nu+1}, \cdots, N-2_{\nu+1}, N-1'_{\nu}| , \end{aligned}$$

where k'_{ν} stands for the column,

$$k'_{\nu} = \begin{pmatrix} J_{\nu}(n+k) \\ q^{-2}J_{\nu}(n+k+2) \\ q^{-4}J_{\nu}(n+k+4) \\ \vdots \end{pmatrix} .$$
(28)

Hence we obtain a difference formula,

$$|0_{\nu}, \cdots, N - 2_{\nu}, N - 1'_{\nu-1}| = \{-(1-q)\}^{-N+1}q^{-\frac{(2n+3N-2)(N-1)}{2}} \quad \tau_N^{\nu-1}(n).$$
(29)

Let us construct another difference formula. In eq.(27), multiplying N-th column by $q^{-\nu}$, adding (N-1)-th column multiplied by $-(1-q)q^{n+N-2}$ to N-th column and using eq.(15), we have

$$q^{-\nu}\tau_{N}^{\nu}(n) = \{-(1-q)\}^{N-1}q^{\frac{(2n+N-2)(N-1)}{2}} \\ \times \begin{vmatrix} J_{\nu+1}(n) & \cdots & J_{\nu+1}(n+N-2) & J_{\nu}(n+N-2) \\ q^{2}J_{\nu+1}(n+2) & \cdots & q^{2}J_{\nu+1}(n+N) & J_{\nu}(n+N) \\ \vdots & \cdots & \vdots & \vdots \\ q^{2N-2}J_{\nu+1}(n+2N-2) & \cdots & q^{2N-2}J_{\nu+1}(n+3N-4) & J_{\nu}(n+3N-4) \end{vmatrix}$$

$$= \{-(1-q)\}^{N-1} q^{\frac{(2n+3N-2)(N-1)}{2}} \times \begin{vmatrix} J_{\nu+1}(n) & \cdots & J_{\nu+1}(n+N-2) & J_{\nu}(n+N-2) \\ J_{\nu+1}(n+2) & \cdots & J_{\nu+1}(n+N) & q^{-2}J_{\nu}(n+N) \\ \vdots & \cdots & \vdots & \vdots \\ J_{\nu+1}(n+2N-2) & \cdots & J_{\nu+1}(n+3N-4) & q^{-2N+2}J_{\nu}(n+3N-4) \end{vmatrix} \\= \{-(1-q)\}^{N-1} q^{\frac{(2n+3N-2)(N-1)}{2}} |0_{\nu+1}, \cdots, N-2_{\nu+1}, N-2'_{\nu}|.$$

Hence we get a difference formula,

$$|0_{\nu}, \cdots, N - 2_{\nu}, N - 2'_{\nu-1}| = \{-(1-q)\}^{-N+1} q^{-\frac{(2n+3N-2)(N-1)}{2}-\nu+1} \quad \tau_N^{\nu-1}(n) \;. \tag{30}$$

Similar calculations give the following formulas by using eq.(16) instead of (15),

$$|0_{\nu}, \cdots, N - 2_{\nu}, N - 1_{\nu+1}'| = (1 - q)^{-N+1} q^{-\frac{(2n+3N-2)(N-1)}{2}} \tau_N^{\nu+1}(n) , \qquad (31)$$

$$|0_{\nu}, \cdots, N - 2_{\nu}, N - 2'_{\nu+1}| = (1 - q)^{-N+1} q^{-\frac{(2n+3N-2)(N-1)}{2} + \nu + 1} \quad \tau_N^{\nu+1}(n) \;. \tag{32}$$

So far we have constructed the difference formulas of τ for "horizontal shifts". Next we consider formulas for "vertical shifts" (shifts of rows in eq.(14)). For notational convenience, we take the transpose of τ ,

$$\tau_{N}^{\nu}(n) = \begin{vmatrix} J_{\nu}(n) & J_{\nu}(n+2) & \cdots & J_{\nu}(n+2N-2) \\ J_{\nu}(n+1) & J_{\nu}(n+3) & \cdots & J_{\nu}(n+2N-1) \\ \vdots & \vdots & \cdots & \vdots \\ J_{\nu}(n+N-1) & J_{\nu}(n+N+1) & \cdots & J_{\nu}(n+3N-3) \end{vmatrix}$$
$$= |\hat{0}_{\nu}, \hat{2}_{\nu}, \cdots, 2\widehat{N-4_{\nu}}, 2\widehat{N-2_{\nu}}| .$$
(33)

Here \hat{k}_{ν} stands for the column,

$$\hat{k}_{\nu} = \begin{pmatrix} J_{\nu}(n+k) \\ J_{\nu}(n+k+1) \\ J_{\nu}(n+k+2) \\ \vdots \end{pmatrix} .$$
(34)

Using the contiguity relations (15) and (16), we can verify

$$\hat{2}_{\nu} - q^{2\nu} (1 + (1 - q)^2 q^{2n}) \, \hat{0}_{\nu}
= (1 - q) q^{n+3\nu+1} \begin{pmatrix} (1 + q^{-2\nu}) J_{\nu+1}(n+1) \\ q^{-1} \{ (1 + q^{-2\nu+2}) J_{\nu+1}(n+2) - (1 - q^2) q^{-\nu-1} J_{\nu+1}(n+1) \} \\ q^{-2} \{ (1 + q^{-2\nu+4}) J_{\nu+1}(n+3) - (1 - q^4) q^{-\nu-1} J_{\nu+1}(n+2) \} \\ \vdots \end{pmatrix}. (35)$$

In eq.(33), we first add *j*-th column multiplied by $-q^{2\nu}(1+(1-q)^2q^{2n+4j-4})$ to (j+1)-th column for $j = (N-1) \sim 1$ and use eq.(35). Secondly, we multiply *i*-th row by $q^{i-1}/(1+q)^2q^{2n+4j-4}$

 $q^{-2\nu+2(i-1)}$) for $i = 1 \sim N$. Finally adding *i*-th row multiplied by $(1-q^{2i})q^{-\nu-1}/(1+q^{-2\nu+2i})$ to (i+1)-th row for $i = 1 \sim (N-1)$, we obtain

$$\begin{aligned} &|\hat{0}_{\nu}', \hat{1}_{\nu+1}, \hat{3}_{\nu+1}, \cdots, 2\widehat{N-3}_{\nu+1}| \\ &= \{(1-q)q^{n+3\nu-1}\}^{-N+1}q^{-\frac{N(N-1)}{2}} \prod_{k=0}^{N-1} \frac{1}{1+q^{-2\nu+2k}} \quad \tau_N^{\nu}(n) , \end{aligned}$$
(36)

where

=

$$\hat{k}_{\nu}' = \begin{pmatrix} K_{\nu}(n+k)_{0} \\ K_{\nu}(n+k)_{1} \\ K_{\nu}(n+k)_{2} \\ \vdots \end{pmatrix},$$
(37)

and $K_{\nu}(n)_i$ is defined by the following recursion relation,

$$\begin{cases} K_{\nu}(n)_{i} = \frac{1}{1+q^{-2\nu+2i}}(q^{i}J_{\nu}(n+i)+(1-q^{2i})q^{-\nu-1}K_{\nu}(n)_{i-1}), & \text{for } i \geq 1, \\ K_{\nu}(n)_{0} = \frac{1}{1+q^{-2\nu}}J_{\nu}(n). \end{cases}$$
(38)

Let us derive another difference formula. We multiply 1st column of $\tau_N^{\nu}(n)$ in eq.(33) by $-q^{2\nu}(1+(1-q)^2q^{2n})$ and add 2nd column to 1st column. Then we follow the same procedure as above calculation, that is, add *j*-th column multiplied by $-q^{2\nu}(1+(1-q)^2q^{2n+4j-4})$ to (j+1)-th column for $j = (N-1) \sim 2$, multiply *i*-th row by $q^{i-1}/(1+q^{-2\nu+2(i-1)})$ for $i = 1 \sim N$, and add *i*-th row multiplied by $(1-q^{2i})q^{-\nu-1}/(1+q^{-2\nu+2i})$ to (i+1)-th row for $i = 1 \sim (N-1)$. Then we obtain

$$\begin{aligned} |\hat{1}_{\nu+1}, \hat{2}'_{\nu}, \hat{3}_{\nu+1}, \cdots, 2\widehat{N-3}_{\nu+1}| \\ &= \{(1-q)q^{n+3\nu-1}\}^{-N+1}q^{-\frac{N(N-1)}{2}}\prod_{k=0}^{N-1}\frac{1}{1+q^{-2\nu+2k}} \\ &\times (-q^{2\nu})\{1+(1-q)^2q^{2n}\} \quad \tau_N^{\nu}(n) . \end{aligned}$$
(39)

We have constructed six difference formulas (29), (30), (31), (32), (36) and (39). In the following, using these difference formulas for τ function, we prove that the Plücker relations reduce to the bilinear difference equations (20)-(24).

(2) Bilinear Identities

A. Equations (20) and (21)

First we derive eq.(20). Consider the identity of determinant,

$$0 = \begin{vmatrix} 0_{\nu} & 1_{\nu} & \cdots & N - 1_{\nu} & \emptyset & N_{\nu} & N'_{\nu-1} & \phi_1 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 1_{\nu} & \cdots & N - 1_{\nu} & N_{\nu} & N'_{\nu-1} & \phi_1 \end{vmatrix}, \quad (40)$$

where k_{ν} and k'_{ν} denote columns defined in eqs.(26) and (28), \emptyset means an empty matrix of relevant size and

$$\phi_1 = \begin{pmatrix} 0\\ \vdots\\ 0\\ 1 \end{pmatrix} \} N + 1 .$$
(41)

Applying the Laplace expansion on the right-hand side, we have

$$0 = |0_{\nu}, \dots, N - 1_{\nu}, N'_{\nu-1}| \times |1_{\nu}, \dots, N - 1_{\nu}, N_{\nu}, \phi_{1}| - |1_{\nu}, \dots, N - 1_{\nu}, N_{\nu}, N'_{\nu-1}| \times |0_{\nu}, \dots, N - 1_{\nu}, \phi_{1}| - |0_{\nu}, \dots, N - 1_{\nu}, N_{\nu}| \times |1_{\nu}, \dots, N - 1_{\nu}, N'_{\nu-1}, \phi_{1}| .$$

$$(42)$$

We note that eq.(42) is nothing but one of the Plücker relations. This identity is rewritten in terms of τ by using eqs. (25), (29) and (30) as

$$\tau_{N+1}^{\nu-1}(n) \ \tau_N^{\nu}(n+1) - q^{-\nu-N+1} \tau_{N+1}^{\nu-1}(n+1) \ \tau_N^{\nu}(n) = -(1-q)q^{n+2N} \tau_{N+1}^{\nu}(n) \ \tau_N^{\nu-1}(n+1) \ ,$$
(43)

which is essentially the same as eq.(20). Equation (21) is derived from the same identity as eq.(42) except that $N'_{\nu-1}$ is replaced by $N'_{\nu+1}$ and that eqs.(31) and (32) are used instead of eqs.(29) and (30).

B. Equations (22) and (23)

We next derive eq.(22). Replacing ϕ_1 in eq.(42) by

$$\phi_2 = \begin{pmatrix} 1\\0\\\vdots\\0 \end{pmatrix} , \qquad (44)$$

we have

$$0 = |0_{\nu}, \cdots, N - 1_{\nu}, N'_{\nu-1}| \times |1_{\nu}, \cdots, N - 1_{\nu}, N_{\nu}, \phi_{2}| - |1_{\nu}, \cdots, N - 1_{\nu}, N_{\nu}, N'_{\nu-1}| \times |0_{\nu}, \cdots, N - 1_{\nu}, \phi_{2}| - |0_{\nu}, \cdots, N - 1_{\nu}, N_{\nu}| \times |1_{\nu}, \cdots, N - 1_{\nu}, N'_{\nu-1}, \phi_{2}| ,$$
(45)

which can be rewritten in terms of τ again. Here we give notes on the treatise of the determinant including the column ϕ_2 . As an example, let us consider $|1_{\nu}, \dots, N-1_{\nu}, N'_{\nu-1}, \phi_2|$. Expanding by (N+1)-th column, we have

$$|1_{\nu}, \cdots, N - 1_{\nu}, N'_{\nu-1}, \phi_2|$$

$$= \begin{vmatrix} J_{\nu}(n+1) & \cdots & J_{\nu}(n+N-1) & J_{\nu-1}(n+N) & 1 \\ J_{\nu}(n+3) & \cdots & J_{\nu}(n+N+1) & q^{-2}J_{\nu-1}(n+N+2) & 0 \\ \vdots & \cdots & \vdots & \vdots & \vdots \\ J_{\nu}(n+2N+1) & \cdots & J_{\nu}(n+3N-1) & q^{-2N}J_{\nu-1}(n+3N) & 0 \end{vmatrix}$$
$$= (-1)^{N} \begin{vmatrix} J_{\nu}(n+3) & \cdots & J_{\nu}(n+N+1) & q^{-2}J_{\nu-1}(n+N+2) \\ \vdots & \cdots & \vdots & \vdots \\ J_{\nu}(n+2N+1) & \cdots & J_{\nu}(n+3N-1) & q^{-2N}J_{\nu-1}(n+3N) \end{vmatrix}$$
$$= (-1)^{N}q^{-2} \quad |3_{\nu}, 4_{\nu}, \cdots, N+1_{\nu}, N+2'_{\nu-1}| , \qquad (46)$$

which indicates that an additional factor $(-1)^N q^{-2}$ is needed. Taking this into account, we obtain

$$\tau_{N+1}^{\nu-1}(n) \ \tau_N^{\nu}(n+3) - q^{-\nu-N+1} \tau_{N+1}^{\nu-1}(n+1) \ \tau_N^{\nu}(n+2) = -(1-q)q^n \tau_{N+1}^{\nu}(n) \ \tau_N^{\nu-1}(n+3) \ , \tag{47}$$

which is essentially the same as eq.(22). Similarly, eq.(23) is derived from the same identity as eq.(45) except that the column $N'_{\nu-1}$ is replaced by $N'_{\nu+1}$.

C. Equation (24)

Finally we derive eq.(24). We consider the identity of determinant,

where \hat{k}_{ν} , \hat{k}'_{ν} and ϕ_1 are defined in eqs.(34), (37) and (41), respectively. Applying the Laplace expansion to the right-hand side of eq.(48), we have

$$0 = |\hat{2}'_{\nu}, \hat{3}_{\nu+1}, \cdots, 2\widehat{N-1}_{\nu+1}, 2\widehat{N+1}_{\nu+1}| \times |\hat{1}_{\nu+1}, \hat{3}_{\nu+1}, \cdots, 2\widehat{N-1}_{\nu+1}, \phi_1| + |\hat{1}_{\nu+1}, \hat{2}'_{\nu}, \hat{3}_{\nu+1}, \cdots, 2\widehat{N-1}_{\nu+1}| \times |\hat{3}_{\nu+1}, \cdots, 2\widehat{N-1}_{\nu+1}, 2\widehat{N+1}_{\nu+1}, \phi_1| - |\hat{1}_{\nu+1}, \hat{3}_{\nu+1}, \cdots, 2\widehat{N-1}_{\nu+1}, 2\widehat{N+1}_{\nu+1}| \times |\hat{2}'_{\nu}, \hat{3}_{\nu+1}, \cdots, 2\widehat{N-1}_{\nu+1}, \phi_1| .$$
(49)

Using eqs.(33), (36) and (39), we get

$$\tau_{N+1}^{\nu}(n+2) \ \tau_N^{\nu+1}(n+1) - q^{2\nu+2N} \{1 + (1-q)^2 q^{2n}\} \tau_{N+1}^{\nu}(n) \ \tau_N^{\nu+1}(n+3)$$

= $(1-q)q^{n+\nu+N+1}(q^{2N}+q^{2\nu}) \tau_{N+1}^{\nu+1}(n+1) \ \tau_N^{\nu}(n+2) \ .$ (50)

Combining eqs.(50) and (20), we obtain eq.(24). Thus we have proved that the τ function (14) really satisfies the bilinear difference equations (20)-(24).

3.2 Derivation of dP_{III} from Bilinear Difference Equations

We now show that eq.(18) actually follows from the bilinear difference equations (20)-(24). We first note that eq.(18) is factorized as

$$w_{n+1}w_{n-1} = -\frac{\left(q^{2N}w_n - (1-q)^2 q^{2n+\nu+7N}\right) \left(q^{2N}w_n + (1-q)^2 q^{2n-\nu+5N-2}\right)}{\left(w_n + q^{\nu+N}\right) \left(w_n - q^{-\nu+3N}\right)} .$$
(51)

Let us rewrite the factors in eq.(51). Equation (17) gives

$$w_n = \frac{\tau_{N+1}^{\nu}(n+1)\tau_N^{\nu+1}(n) - q^{\nu+N}\tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)}{\tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)} , \qquad (52)$$

which is rewritten by using eq.(20) as

$$w_n = (1-q)q^{n+\nu+3N} \frac{\tau_{N+1}^{\nu+1}(n)\tau_N^{\nu}(n+1)}{\tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)} .$$
(53)

By the definition, it is obvious that

$$w_n + q^{\nu+N} = \frac{\tau_{N+1}^{\nu}(n+1)\tau_N^{\nu+1}(n)}{\tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)} .$$
(54)

Using eq.(24), we get

$$w_n - q^{-\nu+3N} = \frac{\tau_{N+1}^{\nu}(n+1)\tau_N^{\nu+1}(n) - (q^{\nu+N} + q^{-\nu+3N})\tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)}{\tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)} = -q^{4N} \{1 + (1-q)^2 q^{2(n-1)}\} \frac{\tau_{N+1}^{\nu}(n-1)\tau_N^{\nu+1}(n+2)}{\tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)} .$$
 (55)

From eqs.(53)-(55), we obtain

$$w_{n+1}w_{n-1}(w_n + q^{\nu+N})(w_n - q^{-\nu+3N})$$

$$= -(1-q)^2 q^{2n+2\nu+10N} \{1 + (1-q)^2 q^{2(n-1)}\}$$

$$\times \frac{\tau_{N+1}^{\nu+1}(n+1)\tau_N^{\nu}(n+2)\tau_{N+1}^{\nu+1}(n-1)\tau_N^{\nu}(n)}{\tau_{N+1}^{\nu}(n)^2 \tau_N^{\nu+1}(n+1)^2}.$$
(56)

The factor $q^{2N}w_n - (1-q)^2 q^{2n+\nu+7N}$ is rewritten as follows,

$$q^{2N}w_n - (1-q)^2 q^{2n+\nu+7N} = (1-q)q^{n+\nu+5N} \frac{\tau_{N+1}^{\nu+1}(n)\tau_N^{\nu}(n+1) - (1-q)q^{n+2N}\tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)}{\tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)} = (1-q)q^{n+2\nu+4N+1} \frac{\tau_{N+1}^{\nu+1}(n+1)\tau_N^{\nu}(n)}{\tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)},$$
(57)

where we have used eq.(53) in the first line and eq.(21) in the second line, respectively. Finally the factor $q^{2N}w_n + (1-q)^2 q^{2n-\nu+5N-2}$ is rewritten as

$$q^{2N}w_n + (1-q)^2 q^{2n-\nu+5N-2}$$

$$= (1-q)q^{n+\nu+5N} \frac{\tau_{N+1}^{\nu+1}(n)\tau_N^{\nu}(n+1) + (1-q)q^{n-2\nu-2}\tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)}{\tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)}$$

$$= (1-q)q^{n+\nu+5N} \times \left\{ \left(q^{-\nu+N-1}\tau_{N+1}^{\nu+1}(n-1)\tau_N^{\nu}(n+2) - (1-q)q^{n-\nu+N-2}\tau_{N+1}^{\nu}(n-1)\tau_N^{\nu+1}(n+2) \right) + (1-q)q^{n-\nu+N-2} \left(\tau_{N+1}^{\nu}(n-1)\tau_N^{\nu+1}(n+2) + (1-q)q^{n-1}\tau_{N+1}^{\nu+1}(n-1)\tau_N^{\nu}(n+2) \right) \right\} / \tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)$$

$$= (1-q)q^{n+6N-1} \left\{ 1 + (1-q)^2 q^{2(n-1)} \right\} \frac{\tau_{N+1}^{\nu+1}(n-1)\tau_N^{\nu}(n+2)}{\tau_{N+1}^{\nu}(n)\tau_N^{\nu+1}(n+1)}, \qquad (58)$$

where we have used eq.(53) in the first line and eqs.(22) and (23) in the second line, respectively. From eqs.(57) and (58), we get

$$-\left(q^{2N}w_n - (1-q)^2 q^{2n+\nu+7N}\right) \left(q^{2N}w_n + (1-q)^2 q^{2n-\nu+5N-2}\right)$$

= $-(1-q)^2 q^{2n+2\nu+10N} \{1 + (1-q)^2 q^{2(n-1)}\}$
 $\times \frac{\tau_{N+1}^{\nu+1}(n+1)\tau_N^{\nu}(n)\tau_{N+1}^{\nu+1}(n-1)\tau_N^{\nu}(n+2)}{\tau_{N+1}^{\nu}(n)^2 \tau_N^{\nu+1}(n+1)^2},$ (59)

which is the same as eq.(56). Thus we have proved that the dP_{III} eq.(51) follows from the bilinear difference equations (20)-(24).

4 q-Difference Analogue of P_{III}

We have discussed the solution of dP_{III} , and shown that it is expressed by the discrete Bessel function. It should be noted, however, that eq.(11) is essentially the same as the *q*-Bessel equation,

$$J_{\nu}(q^2x) - (q^{\nu} + q^{-\nu})J_{\nu}(qx) + \{1 + (1-q)^2x^2\}J_{\nu}(x) = 0.$$
(60)

Actually, eq.(11) yields eq.(60) by the replacement $q^n \to x$. This fact implies that the dP_{III} is a q-discrete system rather than a discrete system in essence. More precisely, let us consider the following equation instead of the dP_{III} (18),

$$w(qx)w(q^{-1}x) = \frac{\alpha w(x)^2 + \beta x^2 w(x) + \gamma x^4}{w(x)^2 + \delta w(x) + \alpha} .$$
(61)

The solutions of eq.(61), with the parameters being the same as eq.(19), are given by

$$w(x) = \frac{\tau_{N+1}^{\nu}(qx)\tau_N^{\nu+1}(x)}{\tau_{N+1}^{\nu}(x)\tau_N^{\nu+1}(qx)} - q^{\nu+N} , \qquad (62)$$

and the τ function $\tau_N^{\nu}(x)$ is expressed as

$$\tau_N^{\nu}(x) = \begin{vmatrix} J_{\nu}(x) & J_{\nu}(qx) & \cdots & J_{\nu}(q^{N-1}x) \\ J_{\nu}(q^2x) & J_{\nu}(q^3x) & \cdots & J_{\nu}(q^{N+1}x) \\ \vdots & \vdots & \cdots & \vdots \\ J_{\nu}(q^{2N-2}x) & J_{\nu}(q^{2N-1}x) & \cdots & J_{\nu}(q^{3N-3}x) \end{vmatrix} .$$
(63)

Here $J_{\nu}(x)$ is the q-Bessel function of degree ν .

We next mention briefly the Lax pair for eq.(61). The auxiliary linear system for dP_{III} is given by [9,12]

$$\Phi_n(qh) = L_n(h)\Phi_n(h) , \qquad (64)$$

$$\Phi_{n+1}(h) = M_n(h)\Phi_n(h) , \qquad (65)$$

whose compatibility condition is written as

$$M_n(qh)L_n(h) = L_{n+1}(h)M_n(h) . (66)$$

If we choose suitable 4×4 matrices as L and M[12], eq.(66) yields dP_{III} (18). The linear system for eq.(61) is written as q-difference system both in x and h,

$$\Phi(x;qh) = L(x;h)\Phi(x;h) , \qquad (67)$$

$$\Phi(qx;h) = M(x;h)\Phi(x;h), \tag{68}$$

whose compatibility condition yields

$$M(x;qh)L(x;h) = L(qx;h)M(x;h)$$
 . (69)

Let the matrices L and M be

$$L(x;h) = \begin{pmatrix} \kappa & \frac{c}{w(x)} + \kappa & \frac{c}{w(x)} & 0\\ 0 & \frac{q^2\lambda}{x^2} & q^2\frac{w(q^{-1}x) + \lambda}{x^2} & q^2\frac{w(q^{-1}x)}{x^2}\\ \frac{w(x)}{h^4x^2} & 0 & \mu & \frac{w(x)}{x^2} + \mu\\ \frac{1}{h^4}\left(\frac{c}{q^4w(q^{-1}x)} + \frac{\lambda}{x^2}\right) & \frac{c}{h^4q^4w(q^{-1}x)} & 0 & \frac{\lambda}{x^2} \end{pmatrix} , (70)$$

$$M(x;h) = \begin{pmatrix} \frac{(\kappa x^2 - q^2 \lambda)w(x)}{q^2 \lambda w(x) + cx^2} & x^2 \frac{\kappa w(x) + c}{q^2 \lambda w(x) + cx^2} & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & \frac{\mu x^2 - \lambda}{w(x) + \lambda} & \frac{w(x) + \mu x^2}{w(x) + \lambda}\\ \frac{1}{h^4} & 0 & 0 & 0 \end{pmatrix}, \quad (71)$$

respectively, where c, κ, λ and μ are arbitrary constants. Then we have from eq.(69),

$$w(qx)w(q^{-1}x) = \frac{c \ (q^2\lambda w(x) + cx^2)(w(x) + \mu x^2)}{q^2(\kappa w(x) + c)(w(x) + \lambda)} ,$$
(72)

which is essentially the same as eq.(61).

Finally, we comment on the limit to P_{III} . We put w = (1-q)xu and take the limit $q \to 1$. Then eq.(61) with the parameters (19) reduces to the P_{III} eq.(2) with the parameters (5). Moreover, the τ function (63) reduces to that of P_{III} (3) up to a trivial factor and the dependent variable transformation (62) yields eq.(4).

In summary,

- 1. Equation (61) reduces to the P_{III} equation in the limit $q \rightarrow 1$.
- 2. Equation (61) has particular solutions expressed by the determinant whose entries are given by the q-Bessel functions.
- 3. The auxiliary linear system and the Lax equation for eq.(61) are written as q-difference equations.

These facts imply that eq.(61) is one candidate for the q-difference analogue of P_{III} .

5 Concluding Remarks

In this paper, we have investigated the solutions of special function type for the dP_{III} equation, and shown that they are expressed in terms of the Casorati determinants of the discrete analogue of the Bessel functions. The solutions are reduced to the Wronskian solutions of the P_{III} in the continuous limit. Hence we conclude that the dP_{III} is a good discretization of the P_{III} .

The discrete Bessel functions are transformed to the q-Bessel functions by a simple independent variable transformation. Based on this observation, we have presented a q-difference analogue of the P_{III}, whose particular solutions are expressed by the Wronskitype determinants of the q-Bessel functions. Moreover, we have mentioned that the Lax equation and its auxiliary linear problem are written as q-difference equations. We expected that the τ function of the dP_{III} satisfies the bilinear form of the *q*-cylindrical Toda molecule equation which we proposed in ref.[22] since that of the P_{III} satisfies the cylindrical Toda molecule equation[14]. However, we found it is not the case because of the asymmetry of structure of the τ function. Actually the τ function of the dP_{III} satisfies a slightly modified version of the *q*-cylindrical Toda molecule equation which also reduces to the cylindrical Toda molecule equation in the limit $q \rightarrow 1$. Similar asymmetric structure of the τ function has also been obtained for the solutions of the dP_{II}[16]. We do not know yet whether such asymmetry is essential or not for the discrete Painlevé equations.

It is an interesting problem to study whether the other discrete Painlevé equations proposed in refs.[6,7,9,10,11,12] admit similar determinant solutions or not. The existence of rational solutions together with their determinant structures is also an interesting subject to be investigated.

Acknowledgment: We are grateful to Prof. Grammaticos and Prof. Ramani for fruitful discussions and useful comments.

References

- [1] E. Brézin and V. A. Kazakov, Phys. Lett. B **236** (1990) 144.
- [2] V. Periwal and D. Shevitz, Phys. Rev. Lett. **64** (1990) 1326.
- [3] A. R. Its, A. V. Kitaev and A. S. Fokas, Russ. Math. Surveys 45, No. 6 (1990) 155.
- [4] F. W. Nijhoff and V. G. Papageorgiou, Phys. Lett. A 153 (1991) 337.
- [5] A. Ramani, B. Grammaticos and J. Hietarinta, Phys. Rev. Lett. 67 (1991) 1829.
- [6] B. Grammaticos and A. Ramani, in Applications of Analytic and Geometric Methods to Nonlinear Differential Equations, ed. P. A. Clarkson (NATO ASI Ser. C413, Kluwer, 1993) 299.
- [7] A. S. Fokas, B. Grammaticos and A. Ramani, J. Math. Anal. Appl. 180 (1993) 342.
- [8] B. Grammaticos, A. Ramani and V. Papageorgiou, Phys. Rev. Lett. 67 (1991) 1825.
- [9] V. G. Papageorgiou, F. W. Nijhoff, B. Grammaticos and A. Ramani, Phys. Lett. A 164 (1992) 57.

- [10] A. Ramani and B. Grammaticos, J. Phys. A **25** (1992) L633.
- [11] K. M. Tamizhmani, B. Grammaticos and A. Ramani, Lett. Math. Phys. **29** (1993) 49.
- [12] B. Grammaticos, F. W. Nijhoff, V. Papageorgiou, A. Ramani and J. Satsuma, Phys. Lett. A 185 (1994) 446.
- [13] M. J. Ablowitz, A. Ramani and H. Segur, J. Math. Phys. **21** (1980) 715.
- [14] See for example, K. Okamoto, Ann. Mat. Pura Appl. 146 (1987) 337; Japan. J. Math.
 13 (1987) 47; Math. Ann. 275 (1986) 221; Funkcial. Ekvac. 30 (1987) 305.
- [15] G. R. W. Quispel, J. A. G. Roberts and C. J. Thompson, Physica D 34 (1989) 183.
- [16] K. Kajiwara, Y. Ohta, J. Satsuma, B. Grammaticos and A. Ramani, J. Phys. A 27 (1994) 915.
- [17] R. Hirota, J. Phys. Soc. Jpn. 46 (1979) 312.
- [18] Y. Ohta, J. Satsuma, D. Takahashi and T. Tokihiro, Prog. Theor. Phys. Suppl. No. 94 (1988) 210.
- [19] R. Hirota, Y. Ohta and J. Satsuma, Prog. Theor. Phys. Suppl. No. 94 (1988) 59.
- [20] Y. Ohta, R. Hirota, S. Tsujimoto and T. Imai, J. Phys. Soc. Jpn. **62** (1993) 1872.
- [21] Y. Ohta, K. Kajiwara, J. Matsukidaira and J. Satsuma, J. Math. Phys. 34 (1993) 5190.
- [22] K. Kajiwara, Y. Ohta and J. Satsuma, Phys. Lett. A **180** (1993) 249.