Integrable systems and Riemann surfaces of infinite genus

Martin U. Schmidt¹

Institut für Theoretische Physik Freie Universität Berlin Arnimallee 14 D-14195 Berlin

Contents

Supported by Forschungsinstitut für Mathematik (ETH Zürich) and by DFG, SFB 288 "Differentialgeometrie und Quantenphysik"

[A Borel summability](#page-81-0) 81

[B Another reality condition](#page-85-0) 85

Abstract.

To the spectral curves of smooth periodic solutions of the n-wave equation the points with infinite energy are added. The resulting spaces are considered as generalized Riemann surfcae. In general the genus is equal to infinity, nethertheless these Riemann surfaces are similar to compact Riemann surfaces. After proving a Riemann Roch Theorem we can carry over most of the constructions of the finite gap potentials to all smooth periodic potentials. The symplectic form turns out to be closely related to Serre duality. Finally we prove that all non-linear PDE's, which belong to the focussing case of the non-linear Schrödinger equation, have global solutions for arbitrary smooth periodic inital potantials.

1 Introduction

The investigation of the Korteweg-de Vries equation initiated the development of many new ideas on integrable systems². A large class of integrable systems was discovered, which turned out to be closely related to the theory of Riemann surfaces. In 1976 McKean and Trubowitz [[MK-T-1\]](#page-91-0) succeeded to establish a one to one correspondence between periodic smooth solutions of the Korteweg-de Vries equation and divisors of Riemann surfaces of infinite genus. In 1980 Adler and van Moerbeke[[A-vM](#page-90-0)] and afterwards Reyman and Semenov-Tian-Shansky[[R-S-T](#page-91-0)] generalized ideas of Krichever [\[Kr-1\]](#page-90-0). In the finite dimensional case they showed that the Lagrangian submanifolds are connected components of the Picard group³ of compact Riemann surfaces. Moreover, the hamiltonian flows turned out to be given by the action of one dimensional subgroups of the Picard group on these Lagrangian submanifolds (see e.g. [\[R-S-T\]](#page-91-0)). It was clear that the same is true at least for the so called algebraic geometric solutions of the corresponding infinite dimensional systems. In consideration of the results of[[MK-T-1\]](#page-91-0) and [[MK-T-2\]](#page-91-0) it is natural to expect that this correspondence could be generalized to non-algebraic geometric solutions. Unfortunately there does not exist a Riemann surface with an infinite dimensional Picard group. Either the Riemann surface is compact and the corresponding Picard group is finite dimensional, or the Riemann surface is not compact and the corresponding Picard group is trivial.

McKean and Trubowitz overcame this problem by using another space associated to a Riemann surface instead of a connected component of the Picard group. The first homology group has a natural embedding into the dual space of the holomorphic forms. This embedding may be described by the integration of the holomorphic forms over representative 1-cycles. For compact Riemann surfaces the Jacobian variety, the connected component of the identity of the Picard group, is isomorphic to the quotient of the dual space of the holomorphic forms divided by the first homology group. This quotient turns out to be a compact abelian group. For Riemann surfaces of infinite genus the natural generalization of this quotient is the dual of the Hilbert space of holomorphic forms divided by the closure of the first homology group. In fact, in[[MK-T-1\]](#page-91-0) it is proven, that the real part of this quotient is a compact abelian group. Moreover, this group can be identified with all divisors corresponding to the potentials out of the isospectral sets. For compact Riemann surfaces it is possible to define the solutions directly as functions on these compact abelian groups: Multivalued holomorphic functions on these groups are called theta functions. The Its Matveev formula [\[I-M](#page-90-0)] gives solutions of the Korteweg-de Vries equation in terms of these theta functions. McKean and Trubowitz were able to construct such theta functions on the infinite dimensional quotient space of square integrable holomorphic forms divided by the first homology group of some Riemann surface of infinite genus. In [\[MK-T-2](#page-91-0)] the Its Matveev formula is proven even in this situation. Now in general for Riemann surfaces of infinite genus there are two possibilities: Either we use these

²We commend the review[[D-K-N](#page-90-0)].

³The Picard group of a Riemann surface is defined to be the set of all equivalence classes of holomorphic line bundles together with the multiplication induced by the tensor product of line bundles. In this article we will often use divisors to describe line bundles. This is possible if the line bundle admits a meromorphic cross section. For Riemann surfaces in the correct sense this is always true (see e.g. [\[Fo,](#page-90-0) 29.17]). We do not know whether this is true for the generalized Riemann surfaces considered in this article. But the line bundles we are interested in admit always meromorphic cross sections.

theta functions⁴, or we characterize those divisors, which correspond to the potentials. In the first approach the correspondence between line bundles and potentials may be omitted. We choose the second approach, which refers back to the Picard group: Due to Abel's Theorem for compact Riemann surfaces the elements of the quotient space of holomorphic forms divided by the first homology group are in one to one correspondence with the equivalence classes of divisors of degree zero. The methods of [\[MK-T-1](#page-91-0)] and[[MK-T-2\]](#page-91-0) suggest a modification of the equivalence relation in the case of non-compact Riemann surfaces of infinite genus. Only those meromorphic functions have a divisor equivalent to the zero divisor in this modified sense, which are bounded in some sense near infinity.

This paper gives a slightly different and more geometric approach to the Picard group of Riemann surfaces of infinite genus. Similar to the algebraic geometric case we add to the spectral curve some points corresponding to the value $\lambda = \infty$ of the spectral parameter. In every neighbourhood of the form $|\lambda| > 1/\epsilon$ of these points there are in general infinitely many branchpoints. Hence the resulting space is not a Riemann surface in the correct sense. Now we define a base of neighbourhoods of such points, such that in some neighbourhood of these points there are no branchpoints. Together with the usual topology of the spectral curve this gives a topology on the resulting space. Typically these neighbourhoods are of the following form:

$$
\left\{\lambda^{-1}\in\mathbb{C}\,\bigg|\,\,|\lambda^{-1}|<\epsilon,\,\,|\lambda^{-1}-a_n|>\frac{c_n}{\epsilon}\text{ for all }n\in\mathbb{N}\right\},\
$$

with some $\epsilon > 0$, some sequence $(a_n)_{n \in \mathbb{N}}$, which converges to zero, and some positive sequence $(c_n)_{n\in\mathbb{N}}$, which converges much faster to zero, than the sequence $(|a_n|)_{n\in\mathbb{N}}$.

In figure 1 we draw a schematic picture of such a neighbourhood in the $1/\lambda$ -plane. Now we call a function f on such a set holomorphic, if

- (i) f is holomorphic on the interior of this set with respect to the usual topology of \mathbb{C} ,
- (ii) f and all derivatives of f extend to continuous functions on the whole set with respect to the topology defined by these neighbourhoods.

⁴To the authors knowledge this is done in [\[F-K-T](#page-90-0)].

This definition makes it possible to carry over almost all concepts of the theory of Riemann surfaces to these generalized Riemann surfaces. For example the sheaves of holomorphic and meromorphic functions, and the sheaves of holomorphic and meromorphic forms are defined in a natural way, respectively. Furthermore, the concept of holomorphic line bundles and of divisors may be generalized to this situation.

These generalized Riemann surfaces turn out to be similar to compact Riemann surfaces, although they are not compact. For example all global holomorphic functions are constant, and their Picard groups are very large. A large part of the article will be devoted to an analysis of these Picard groups. It turns out that the smooth periodic solutions of the generalized nonlinear Schrödinger equation are in one to one correspondence with some part of the Picard groups.

In some sense the nonlinear Schrödinger equation is the simplest case of those integrable systems, which may be described by some Lax equation or some zero curvature equation⁵. In this article we restrict attention to the generalization of the nonlinear Schrödinger equation to semisimple Lie groups introduced in $[F-K]$. It will not suffice to look at the nonlinear Schrödinger equation only, because a simple count of dimensions in the infinite dimensional case suggests that the systems corresponding to higher groups than $SL(2,\mathbb{C})$ are not integrable⁶. Indeed, the number of series of integrals are equal to the rank of the group and the number of fields is equal to the dimension of the group minus the rank of the group. This article shows that the generalized nonlinear Schrödinger equation is quite similar to the nonlinear Schrödinger equation in the case of periodic boundary conditions⁷. We believe that almost all methods and results of this paper may be generalized to other integrable systems, as for example the generalized sine Gordon and the sinh Gordon equation. The case of the two dimensional Kadomtsev-Petviashvili equation seems to be more complicated. Some of the ideas of Appendix [B](#page-85-0) may be helpful.

It is possible to modify these methods in order to cover all potentials, which are elements of some Sobolev spaces, instead of the Fréchet space of smooth periodic potentials. For this purpose the topology of the Riemann surface will have to be changed. Furthermore, condition (ii) of the definition of holomorphic functions on these neighbourhoods described above has to be replaced by

(ii)' f and all derivatives up to some fixed order extend to continuous functions on the whole set.

In this article only the smooth case is covered. Then the resulting Riemann surfaces seem to be more similar to usual Riemann surfaces.

Let us now give a short summary of the article. In [\[H-S-S](#page-90-0)] it is shown that the Floquet matrix⁸ may be diagonalized formally by a matrix valued formal power series in λ^{-1} . In the second section we prove that this power series is an asymptotic expansion of a holomorphic diagonalization. The domain of this asymptotic expansion is chosen to be as large as possible.

 5 Thenonlinear Schrödinger equation is the standard example of [[F-T](#page-90-0)].

 6 Comparewith [[B-S\]](#page-90-0).

⁷On the line this might be different. Then all Riemann surfaces are singular. For higher groups all isospectral sets may decompose into uncountable many components with respect to the action of the Picard group. Hence there may be additional integrals of motion (see Example [10.7](#page-80-0)). Due to Proposition [10.1](#page-74-0) this is impossible for $GL(2,\mathbb{C})$ and $SL(2,\mathbb{C})$.

⁸This matrix is the same as the monodromy matrix.

This domain is used in the third section to define the topology of the completed spectral curve. The resulting space is a Riemann surface in the sense described above. Moreover, we prove that all global holomorphic functions are constant (Lemma [3.3](#page-15-0)) and that the total residue of meromorphic differential forms is zero (Theorem [3.5](#page-17-0)). Finally the divisor of a meromorphic function is shown to be an infinite sum of finite divisors of degree zero (Corollary [3.6](#page-17-0)). With the exception of a finite part of the divisor each of these finite divisors are located in one of the domains, which are excluded from the neighbourhoods of points corresponding to $\lambda = \infty$ as described above. Hence the degree of a divisor turns out to be a sequence of degrees of finite divisors indexed by the excluded domains.

In the fourth section the eigen vectors of the monodromy are shown to define holomorphic line bundles over this Riemann surface. It turns out that the Riemann surfaces corresponding to some potentials are singular. This case is treated separately in the tenth section. The degree of the eigen bundle is determined in Theorem [4.6.](#page-23-0) Finally it is shown that the Riemann surface together with the eigen bundle completely determines the potential.

In order to classify all line bundles, which are equivalent to the eigen bundle of some potential, a Riemann-Roch Theorem for the generalized Riemann surfaces is proven in the fifth section. A condition on divisors is given, such that the space of global sections is finite dimensional and the space of holomorphic forms with values in the dual line bundle is finite dimensional, too (Theorem [5.5](#page-29-0)). In a general situation all integral divisors of a specified degree fulfil this condition (Corollary [5.11\)](#page-34-0). Hence the topology defined above essentially determines the location properties of these divisors.

Now the classification of all eigen bundles of the potentials is given in Theorem [6.6.](#page-37-0) Furthermore the one to one correspondence between potentials and divisors defines a homeomorphism with respect to a suitable topology on the set of these divisors.

The seventh section is a short excursion on Darboux coordinates⁹. These Darboux coordinates are given by the values of the spectral parameter λ and the logarithm of the values of the Floquet multiplier μ (eigenvalues of the monodromy) at all points of the divisor, which describes the eigen bundles corresponding to the potentials. In case of the Korteweg-de Vries equation this was proven in[[P-T](#page-91-0), Theorem 2.8 and Theorem 3.5]. This book gives a comprehensive picture of these coordinates. In the finite dimensional case the analogous result was recently proven in[[A-H-H\]](#page-89-0).

To the authors knowledge the main result of the eighth section is new even in the finite dimensional case. Theorem [8.5](#page-55-0) shows that the symplectic form is given by the Serre duality. Loosely speaking the tangent space may be identified with the direct sum of the first cohomology group of the sheaf of holomorphic functions on the spectral curve and the space of holomorphic forms on the spectral curve. The first summand is naturally isomorphic to the Lie algebra of the Picard group and therefore also isomorphic to the tangent space along the corresponding Lagrangian submanifold. The second summand is isomorphic to any maximal isotropic subspace of the tangent space transversal to the tangent space along the Lagrangian submanifold. Then the symplectic form defines a non-degenerate pairing between these two spaces. Theorem [8.5](#page-55-0) shows that this pairing is the same as the pairing given by Serre duality¹⁰. This relation

⁹In Example [10.3](#page-77-0) it is shown that in general they are not global coordinates.

 10 For compact Riemann surfaces Serre duality is isomorphic to a natural symplectic form of the Riemann surface. In fact, the wedge product and integration over the Riemann surface together defines a natural symplectic form on the first cohomology group of the deRham complex and the decomposition into the direct sum of the

might be the central point of the connection between Riemann surfaces and integrable systems. Theorem [8.5](#page-55-0) proves in addition that these systems are completely integrable in a weak sense: the tangent space along the Lagrangian submanifolds is shown to be maximal isotropic with respect to the symplectic form.

In the ninth section we formulate a reality condition. In case of the nonlinear Schrödinger equation this yields the self-focussing nonlinear Schrödinger equation¹¹. Summing up Corollary [9.11](#page-69-0) and Corollary [10.6](#page-79-0) leads to the following picture: All isospectral sets are homoemorphic to finite unions of groups of the form

 $(\mathbb{R}/\mathbb{Z})^{\text{defective}}$ times a finite dimensional abelian Lie group.

Here $\mathcal{I}_{\text{effective}}$ is some countable set. Moreover, all tangent vectors along the isospectral set correspond to a hamiltonian flow on this isospectral set. In particular all partial differential equations, which describe such hamiltonian flows corresponding to local integrals of motion (see [\[H-S-S\]](#page-90-0) and [\[Sch\]](#page-91-0)), are shown to have global solutions¹².

In Appendix A we show that the formal power series, which diagonalizes the monodromy, is Borel summable if and only if the potential is analytic. Hence the asymptotic expansion of the eigen vectors completely determines the potential (and therefore also the meromorphic eigenvector function) if and only if the potential is analytic.

In Appendix B we include another reality condition in our approach. In case of the nonlinear Schrödinger equation this gives the non-focussing nonlinear Schrödinger equation¹³. It turns out that the two reality conditions are related to the two covering maps of the Riemann surface induced by the spectral parameter and the Floquet multiplier, respectively.

In some footnotes we mark those parts of the article, which may be passed over, if only a rough comprehension is intended.

2 An asymptotic expansion

Let p be a diagonal matrix $p = diagonal(p_1, \ldots, p_n)$, such that all p_i are distinct and let $q(x)$ be a smooth function into the $n \times n$ -matrices, which is periodic with period 1. Let us first consider the fundamental solution of the auxiliary problem

$$
L \cdot g(x, \lambda, q) = 0, \quad g(0, \lambda, q) = 1,
$$

with the Lax operator $L = \frac{d}{dx} + q(x) + \lambda p$. The following well known lemma will give us a power series expansion of g.

Lemma 2.1 Let $g(x)$ be the unique fundamental solution of the homogenous differential equation

$$
\left(\frac{d}{dx} + a(x)\right)g(x) = 0, \ \ g(0) = 1.
$$

space of hoplomorphic and antiholomorphic forms is a decomposition into Lagrangian subspaces with respect to this symplectic form. Due to Dolbeault's Theorem this decomposition of the first cohomology group of the deRham complex is isomorphic the direct sum of the space of holomorphic forms and the first cohomology group of the sheaf of holomorphic functions and this isomorphism transforms the symplectic form into Serre duality.

¹¹The methods of $[MK-T-1]$ do not cover this case.

 12 By pure analytic methods this is proven in [\[Bo](#page-90-0)] for some of these partial differential equations.

¹³The methods of [\[MK-T-1](#page-91-0)] can be carried over to this case.

Then $f(x) = g(x) \left(\int_0^x g^{-1}(t) b(t) dt + f_0 \right)$ is the unique solution of the inhomogeneous differential equation

$$
\left(\frac{d}{dx} + a(x)\right) f(x) = b(x), \quad f(0) = f_0.
$$

We omit the easy proof. \Box

Now we make an ansatz for the fundamental solution $g(x)$ with $a(x) = g(x) + p\lambda$.

$$
g(x, \lambda, q) = \sum_{0}^{\infty} \gamma_n(x, \lambda, q)
$$
 (1)

with
$$
\gamma_0(x,\lambda) = \exp(-x\lambda p)
$$
 and $\left(\frac{d}{dx} + \lambda p\right) \gamma_{n+1}(x,\lambda,q) = -q(x)\gamma_n(x,\lambda,q).$

Due to Lemma [2.1](#page-6-0) we obtain the recursion relation

$$
\gamma_{n+1}(x,\lambda,q) = -\int_0^x \gamma_0(x-t,\lambda)q(t)\gamma_n(t,\lambda,q)dt,
$$

and finally the explicit solution $\gamma_n(x, \lambda, q) =$

$$
= (-1)^n \int_{0 \le t_1 \le \dots \le t_n \le x} \gamma_0(x - t_n, \lambda) q(t_n) \gamma_0(t_n - t_{n-1}, \lambda) \dots q(t_1) \gamma_0(t_1, \lambda) dt_1 \dots dt_n.
$$
 (2)

On the space of potentials q we make use of the natural scalar product

 $\langle \tilde{q}, q \rangle = \int_0^1 \text{tr}(\tilde{q}(t)q^*(t)) dt$ and the corresponding norm $||q||$. Although, in the end we are only interested in smooth periodic potentials, we consider the fundamental solution g for arbitrary q in the Hilbert space H corresponding to the above scalar product. The space of $n \times n$ -matrices is endowed with the Banach norm of operators of the Hilbert space \mathbb{C}^n .

Theorem 2.2 The formal power series (1) for $g(x, \lambda, q)$ converge uniformly on bounded subsets of $[0,1] \times \mathbb{C} \times \mathcal{H}$ to the unique solution of the auxiliary problem. For each fixed x and q, $g(x, \cdot, q)$ is an entire function.

This result compares with Theorem 1 of[[P-T](#page-91-0)] in the case of Hill's equation. Proof: For $\gamma_0(x,\lambda)$ we start with the obvious bound

 $\|\gamma_0(x, \lambda)\| = \exp(x \cdot \sup\{\Re(\lambda p_i)|i = 1, \ldots, n\}).$

Inserting this estimate into (2) gives

$$
\|\gamma_n(x,\lambda,q)\| \le \|\gamma_0(x,\lambda)\| \frac{1}{n!} \left(\int_0^x \|q(t)\| dt\right)^n
$$

For the last factor we give a bound in terms of $||q||$:

$$
\int_0^x \|q(t)\| dt \le \sup \left\{ \left| \langle q, \tilde{q} \rangle \right| \, \left| \|\tilde{q}\|^2 = x \right. \right\} = \|q\| \sqrt{x}.
$$

In fact this is true because on the space of $n \times n$ -matrices we have

$$
||A|| \le \sup \{|tr(AB)| | tr(BB^*) = 1\}.
$$

Inserting this bound into([1\)](#page-7-0) finally gives

$$
g(x, \lambda, q) \le \exp\left(x \cdot \sup\left\{\Re(\lambda p_i) \mid i = 1, ..., n\right\} + ||q||\sqrt{x}\right).
$$

This shows the convergence of $g(x, \lambda, q)$. The uniqueness is well known. All summands of the expansion [\(1](#page-7-0)) are entire functions with respect to λ for fixed x and q. Hence $g(x, \cdot, q)$ is entire, too. too. \Box

Now let \tilde{q} be a small perturbation of $q : ||q - \tilde{q}|| < \epsilon$. Then formula [\(2\)](#page-7-0) implies

$$
\|\gamma_n(x,\lambda,q) - \gamma_n(x,\lambda,\tilde{q})\| \leq n\epsilon \sqrt{x} \|\gamma_0(x,\lambda)\| \frac{1}{n!} \left((\|q\| + \epsilon)\sqrt{x} \right)^{n-1} \leq
$$

$$
\leq \epsilon \sqrt{x} \|\gamma_0(x,\lambda)\| \frac{1}{(n-1)!} \left((\|q\| + \epsilon)\sqrt{x} \right)^{n-1}.
$$

This proves the

Corollary 2.3 If q and \tilde{q} satisfy $||q - \tilde{q}|| < \epsilon$, the following estimate holds:

$$
||g(x, \lambda, q) - g(x, \lambda, \tilde{q})|| \le \epsilon \sqrt{x} ||\gamma_0(x, \lambda)|| \exp \left((||q|| + \epsilon)\sqrt{x} \right).
$$

Let us now recall, how the fundamental solution transforms under gauge transformations. If $q(x)$ is the fundamental solution of

$$
\left(\frac{d}{dx} + a(x)\right)g(x) = 0, \ g(0) = 1
$$

and if $h(x)$ is a differentiable function into $GL(n, \mathbb{C})$, then $\tilde{g}(x) = h^{-1}(x)g(x)h(0)$ is the fundamental solution of

$$
\left(\frac{d}{dx} + \tilde{a}(x)\right)\tilde{g}(x) = 0, \quad \tilde{g}(0) = \mathbf{1} \quad \text{with} \quad \tilde{a}(x) = Ad(h^{-1}(x))a(x) + h^{-1}(x)\frac{dh(x)}{dx}.
$$

In the case, in which both $a(x)$ and $h(x)$ are periodic with period 1, the Floquet matrix $g(1)$ transforms to $\tilde{g}(1) = Ad(h^{-1}(0))g(1)$. This shows that whenever a periodic $h(x)$ 'diagonalizes' the operator $\frac{d}{dx} + a(x)$, which means that $\tilde{a}(x)$ is a diagonal matrix for all x, then $h(0)$ diagonalizesthe Floquet matrix $g(1)$. The next theorem from [[H-S-S\]](#page-90-0) presents a formal diagonalization of the foregoing Lax operator. The rest of this sections concerns the analytic content of this formal diagonalization.

Theorem 2.4 Let q be a smooth periodic potential. Then there exist two series

$$
a_1(x), a_2(x), \ldots
$$
 of offdiagonal matrices and
 $b_0(x), b_1(x), \ldots$ of diagonal matrices, respectively,

such that $a_{m+1}(x)$ and $b_m(x)$ are differential polynomials in $q(x)$ with derivatives of order m at most and the following equality for formal power series of λ^{-1} holds:

$$
L\left(1+\sum_{m=1}^{\infty}a_m(x)\lambda^{-m}\right) = \left(1+\sum_{m=1}^{\infty}a_m(x)\lambda^{-m}\right)\left(\frac{d}{dx}+p\lambda+\sum_{m=0}^{\infty}b_m(x)\lambda^{-m}\right). \tag{3}
$$

 \Box

In the particular case where q is only M times differentiable, there exists an $\epsilon > 0$ and $c > 0$, such that for all $|\lambda|^{-1} < \epsilon$, $x \in [0, 1]$

$$
\left\| \left(1 + \sum_{m=1}^{M+1} a_m(x) \lambda^{-m} \right)^{-1} L \left(1 + \sum_{m=1}^{M+1} a_m(x) \lambda^{-m} \right) - \frac{d}{dx} + p\lambda + \sum_{m=0}^{M} b_m(x) \lambda^{-m} \right\| < \frac{c}{|\lambda|^{M+1}}.
$$
\n(4)

Proof:We solve inductively the ansatz ([3](#page-8-0)) in all powers of λ^{-1} . For the power λ^{0} we obtain the equation

$$
q(x) = [a_1(x), p] - b_0(x). \tag{5}
$$

Since all diagonal entries of p are distinct, $ad(p)$ is invertible on the space of all offdiagonal matrices and the equation has a unique solution. For the power λ^{-M} we obtain the equation

$$
\frac{da_M(x)}{dx} + q(x)a_M(x) - \sum_{m=1}^M a_m(x)b_{M-m}(x) = [a_{M+1}(x), p] + b_M(x). \tag{6}
$$

These equations give inductively a unique solution of([3\)](#page-8-0) with the desired properties. For fixed M it is obvious that for all λ out of some neighbourhood of infinity and for all $x \in$ [0, 1] $(1 + \sum_{m=1}^{M} a_m(x)\lambda^{-m})$ has a uniformly bounded inverse. On the other hand the solution of (3) (3) (3) implies that there exists a \tilde{c} , such that for the same domain

$$
\left\|\left(1+\sum_{m=1}^{M+1}a_m(x)\lambda^{-m}\right)^{-1}L\left(1+\sum_{m=1}^{M+1}a_m(x)\lambda^{-m}\right)-\frac{d}{dx}+p\lambda+\sum_{m=0}^{M}b_m(x)\lambda^{-m}\right\|<\frac{\tilde{c}}{|\lambda|^{M+1}}.
$$

This completes the proof of the theorem. \Box

Itwas shown in [[H-S-S\]](#page-90-0) that these formal power series are convergent, if and only if $q(x)$ is an algebraic geometric potential. In Appendix [A](#page-81-0) we present a proof that these power series are Borel summable, if and only if $q(x)$ is an analytic potential. But for general smooth potentials these power series seem not to be convergent in any sense. Formally these power series diagonalize the Floquet matrix at $\lambda = \infty$. We are now going to prove the main statement of this section that indeed these power series are an asymptotic expansion of the diagonalization of the Floquet matrix near $\lambda = \infty$. Later on we will see that in some sense it is even a Taylor expansion of this diagonalization. Let us first define the domain of the asymptotic expansion.

Definition 2.5 For any $l \in \mathbb{N}_0$, $\epsilon \in \mathbb{R}^+$ let $O_{l,\epsilon}$ be the set

$$
O_{l,\epsilon} = \left\{\lambda \in \mathbb{C} \,\,\bigg|\, |\lambda|^{-1} < \epsilon, \,\, |\lambda - k\pi\sqrt{-1}| > \frac{1}{\epsilon(|k|\pi)^l} \,\, \text{for all } k \in \mathbb{Z} \setminus \{0\} \right\}^{14}.
$$

For a fixed M times continuously differentiable potential q let $exp(p_i(\lambda))$ be the *i*-th eigenvalue of the Floquet matrix of the Lax operator on the right hand side of (3) :

$$
p_{i,M}(\lambda) = -p_i \lambda - \sum_{m=0}^{M} \lambda^{-m} \int_0^1 \left(b_m(x) \right)_{ii} dx. \tag{7}
$$

¹⁴For $l = 0$ we will always assume $\epsilon > 2/\pi$. This ensures that all these sets contain circles around infinity.

Finally let $U_{M,\epsilon}$ be the set given by

$$
U_{M,\epsilon} = \left\{ \lambda \in \mathbb{C} \; \left| \; \frac{p_{i,M}(\lambda) - p_{j,M}(\lambda)}{2} \in O_{M,\epsilon} \text{ for all } i \neq j \in \{1,\ldots,n\} \right. \right\}.
$$

The polynomials $p_i(\lambda)$ and the sets $U_{M,\epsilon}$ depend on the potential q. But for large λ , which corresponds to small ϵ , $p_i(\lambda)$ is nearly equal to $p_i\lambda$. Hence $U_{M,\epsilon}$ consists of all sufficient large λ with the exception of small domains near $\lambda = \frac{2n\pi\sqrt{-1}}{n-m}$ $\frac{n\pi\sqrt{-1}}{p_i-p_j}, n \in \mathbb{Z}$ $\frac{n\pi\sqrt{-1}}{p_i-p_j}, n \in \mathbb{Z}$ $\frac{n\pi\sqrt{-1}}{p_i-p_j}, n \in \mathbb{Z}$. With the help of ([7\)](#page-9-0) it would be easy to give an asymptotic expansion of the exact localization of these excluded domains. Later on we will see that in the case of Hill's equation, this reproduces the well known asymptotic expansion of the periodic and antiperiodic eigenvalues (compare e.g. [[M-W](#page-91-0)], [\[Ho\]](#page-90-0) .)

Theorem 2.6 Let q be an M times continuously differentiable periodic potential. Then three is a holomorphic matrix valued function h on some $U_{l,\epsilon}$, which diagonalizes the Floquet matrix $g(1, \lambda, q)$ for all $l \leq M$. Moreover this matrix valued function can be chosen to be of the form $h = 1 + a$ holomorphic offdiagonal matrix valued function. Then this function h and the eigenvalues μ_1, \ldots, μ_n of the Floquet matrix may be expanded asymptotically and uniformly on some $U_{l,\epsilon}$:

$$
\|h(\lambda) - \left(1 + \sum_{m=1}^{M-l} a_m(0)\lambda^{-m}\right)\| < \frac{\delta}{|\lambda|^{M-l+1}} 15,
$$

$$
|\mu_i - \exp(p_{i,M}(\lambda))| < \frac{c}{|\lambda|^{M+1}} |\exp(p_{i,M}(\lambda))| \quad i = 1, ..., n,
$$

with some $\delta, c > 0^{16}$.

In order to prove this theorem we need some lemmata.

Lemma 2.7 Let B be a diagonal matrix $B = diagonal(\beta_1, \ldots, \beta_n)$ such that β_1 is the eigenvalue of maximal length, which implies $|\beta_1| = ||B||$. In addition consider a matrix A of the form $A =$ $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with respect to the decomposition $\mathbb{C}^n = \mathbb{C} \oplus \mathbb{C}^{n-1}$. Moreover we assume the $\emph{following estimates to be valid for some small $\epsilon>0$, $0<\delta\leq\frac{1}{4}$}$ $\frac{1}{4}$:

$$
||A - B|| \le \epsilon ||B|| \text{ and } |\beta_1 - \beta_i| \ge \frac{\epsilon}{\delta} |\beta_1| \text{ for all } i \ne 1.
$$

Then there exists exactly one eigenvalue α of A obeying the estimate $|\alpha - \beta_1| \leq \epsilon |\beta_1|$. Furthermore there exists exactly one matrix of the form $\begin{pmatrix} 1 & w \\ & 1 \end{pmatrix}$ \boldsymbol{v} 1 \setminus , which satisfies

$$
\left(\begin{array}{cc} a & b \\ c & d \end{array}\right) \left(\begin{array}{cc} 1 & w \\ v & \mathbf{1} \end{array}\right) = \left(\begin{array}{cc} 1 & w \\ v & \mathbf{1} \end{array}\right) \left(\begin{array}{cc} \alpha & 0 \\ 0 & cw + d \end{array}\right).
$$

Finally the assumptions imply the following inequalities:

$$
||v|| \le \frac{\delta}{1-2\delta}, \quad ||w|| \le \frac{\delta}{1-2\delta} \quad and \quad ||\begin{pmatrix} \alpha & 0\\ 0 & cw+d \end{pmatrix} - B|| \le \epsilon \frac{1-\delta}{1-2\delta} ||B||.
$$

¹⁵Actually we will prove that for all $1 \leq l \leq M$ and all $\delta > 0$ there exists an $\epsilon > 0$, such that this estimate holds uniformly on $U_{l,\epsilon}$. Furthermore, for $l = 0$ this estimate holds with some $\delta > 0$ uniformly on $U_{0,4/\pi}$ (compare with footnote [17](#page-13-0)).

¹⁶The rest of this section may be passed over. It contains the proof of this theorem.

Proof: Set $A(z) = B + z(A - B)$. Now we make the ansatz $A(z)v(z) = \alpha(z)v(z)$, with

$$
\alpha(z) = \beta_1 + \sum_{k=1}^{\infty} \alpha_k z^k, \ v(z) = \sum_{k=0}^{\infty} v_k z^k, \ v_0 = \begin{pmatrix} 1 \\ 0 \\ \vdots \end{pmatrix}.
$$

For the K-th power of z we obtain the equation: $Bv_K + (A - B)v_{K-1} = \sum_{k=0}^{K} \alpha_k v_{K-k}$. All these equations have a unique solution with $v_0^t v_K = 0$ for all $K > 0$, $\alpha_K = v_0^t (B - A) v_{K-1}$ and $(B - \beta_1 \mathbf{1})v_K = (B - A)v_{K-1} + \sum_{k=1}^K \alpha_k v_{K-k}$. Using the assumptions we can give bounds for $|\alpha_K|$ and $||v_K||$: $|a_k| < \epsilon |B|$ $\|a_k\|$ $\|a_0\|$

$$
|v_K| \le \frac{\delta}{\epsilon |\beta_1|} \left(\epsilon |\beta_1| \cdot \|v_{K-1}\| + \sum_{k=1}^{K-1} \epsilon |\beta_1| \cdot \|v_{K-1}\| \cdot \|v_{K-k}\| \right) \le \delta \sum_{k=0}^{K-1} \|v_k\| \cdot \|v_{K-k-1}\|.
$$

The equation $\gamma(z) = \delta z \gamma^2(z) + 1$ has one solution $\gamma(z) = 1/(2\delta z) \left(1 - \sqrt{1 - 4\delta z}\right)$ with $\gamma(0) = 1$. The Taylor expansion of this function converges on the domain $|z| \leq 1/(4\delta)$ and is bounded on this domain by $|\gamma(z)| \leq 2$. This proves that $\alpha = \alpha(1)$ and $v(1)$ exist and are bounded by

$$
|\alpha - \beta_1| \le 2\epsilon |\beta_1|, \quad ||v(1) - v_0|| \le \frac{1}{2\delta} \left(1 - 2\delta - \sqrt{1 - 4\delta} \right).
$$

Let us now improve these estimates. Since α is an eigenvalue of A we have

$$
\inf\{|\alpha-\beta_i| \mid i=1,\ldots,n\} \leq \epsilon |\beta_1|
$$

Together with $|\alpha - \beta_1| \le 2\epsilon |\beta_1|$ and $|\beta_1 - \beta_i| \ge 4\epsilon |\beta_1|$ for all $i = 2, \ldots, n$ this implies $|\alpha - \beta_1| \le$ $\epsilon|\beta_1|$. The following ansatz

$$
\left(\begin{array}{cc} a & b \\ c & d \end{array}\right) \left(\begin{array}{cc} 1 & w \\ v & 1 \end{array}\right) = \left(\begin{array}{cc} 1 & w \\ v & 1 \end{array}\right) \left(\begin{array}{cc} \alpha & 0 \\ 0 & cw + d \end{array}\right)
$$

is equivalent to the three equations

$$
a + bv = \alpha, \quad aw + b = wcw + wd,
$$

$$
c + dv = v\alpha
$$

Since α is an eigenvalue of A , $v = (\alpha \mathbf{1} - d)^{-1}c$ is a solution of the two equations involving w. The third equation is equivalent to

$$
\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & w \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & w \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \alpha & 0 \\ c & cw + d \end{pmatrix} \text{ and to}
$$

$$
\begin{pmatrix} 1 & -w \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} \alpha & 0 \\ c & cw + d \end{pmatrix} \begin{pmatrix} 1 & -w \\ 0 & 1 \end{pmatrix}.
$$

Again since α is an eigenvalue, $w = b(d - \alpha \mathbf{1})^{-1}$ is a solution. The assumptions also guarantee the following three estimates:

$$
||d-diagonal(\beta_2,\ldots,\beta_n)|| \leq \epsilon |\beta_1|,
$$

$$
\|\operatorname{diagonal}(\beta_2, \dots, \beta_n)v - \beta_1 v\| \ge \frac{\epsilon}{\delta} \|\beta_1 v\| \text{ for all } v \in \mathbb{C}^{n-1},
$$

$$
\|\beta_1 \mathbf{1} - \alpha \mathbf{1}\| \le \epsilon |\beta_1|.
$$
This implies
$$
\|dv - \alpha v\| \ge \frac{\epsilon |\beta_1|(1 - 2\delta)}{\delta} \|v\| \text{ and furthermore } \|(d - \alpha \mathbf{1})^{-1}\| \le \frac{\delta}{\epsilon (1 - 2\delta) |\beta_1|}
$$

Therefore
$$
\|v\| \le \frac{\delta}{1 - 2\delta} \text{ and } \|w\| \le \frac{\delta}{1 - 2\delta} \text{ and finally}
$$

$$
\left\| \begin{pmatrix} \alpha & 0 \\ 0 & cw + d \end{pmatrix} - B \right\| \le \epsilon |\beta_1| + \left\| \begin{pmatrix} 0 & 0 \\ 0 & cw \end{pmatrix} \right\| \le \epsilon |\beta_1| \left(1 + \frac{\delta}{1 - 2\delta}\right).
$$

All other eigenvalues α' of A are also eigenvalues of $cw + d$. Hence they obey the estimate $\inf \{ |\alpha' - \beta_i| \mid i = 2, \ldots, n \} \leq \frac{3}{2} \epsilon |\beta_1|$, which is a contradiction to $|\alpha' - \beta_1| \leq \epsilon |\beta_1|$. This proves that α is unique. \Box

Lemma 2.8 Let B be the diagonal matrix $B = diagonal(\beta_1, \ldots, \beta_n)$ and A another $n \times n$ -matrix such that the following estimates hold for small $\epsilon, \delta \in \mathbb{R}^+$ and all $i = 1, \ldots, n$:

$$
\left\|\bigwedge^i(A)-\bigwedge^i(B)\right\| \leq \epsilon \left\|\bigwedge^i(B)\right\| \text{ and } |\beta_i-\beta_j| \geq \frac{\epsilon}{\delta} \sup\{|\beta_i|, |\beta_j|\}.
$$

Then there exists an invertible matrix h of the form $1 +$ offdiagonal satisfying $Ah = h \cdot$ $diagonal(\alpha_1, \ldots, \alpha_n)$ such that the following inequalities are valid:

$$
||h-1|| = o(\delta) \text{ and } |\alpha_i - \beta_i| = o(\epsilon)|\beta_i|.
$$

Proof: First we rearrange lines and columns in a way, such that $\prod_{j=1}^{i} \beta_j$ is an eigenvalue of maximal length of $\wedge^i(B)$. Then we make use of the last lemma and, furthermore claim that

$$
\left\|\bigwedge^i (cw+d) - \bigwedge^i (diagonal(\beta_2,\ldots,\beta_n))\right\| = o(\epsilon) \left\|\bigwedge^i (diagonal(\beta_2,\ldots,\beta_n))\right\|.
$$

In order to prove this claim we note that the matrix

$$
\bigwedge^{i+1} \left((1 - wv) \begin{pmatrix} 1 & w \\ v & 1 \end{pmatrix}^{-1} \right) = \bigwedge^{i+1} \begin{pmatrix} 1 & -w \\ -v & (1 - wv)1 + vw \end{pmatrix}
$$
 has the form $\begin{pmatrix} 1 & \cdot \\ \cdot & \cdot \end{pmatrix}$

with respect to the natural decomposition \wedge^{i+1} ($\mathbb{C} \oplus \mathbb{C}^{n-1}$) $\cong \wedge^{i}$ (\mathbb{C}^{n-1}) $\oplus \wedge^{i+1}$ (\mathbb{C}^{n-1}). Hence the matrix

$$
\bigwedge^{i+1} \left(\begin{array}{cc} 1 & -w \\ -v & (1 - wv)\mathbf{1} + vw \end{array} \right) \left(\bigwedge^{i+1} (A) - \bigwedge^{i+1} (B) \right)
$$

has the form

$$
\left(\begin{array}{cc}\alpha\,\Lambda^i(cw+d)-\beta_1\,\Lambda^i\,(diagonal(\beta_2,\ldots,\beta_n))&\cdot\\ \cdot&\cdot\end{array}\right)
$$

.

Finally the bound

$$
\left\|\bigwedge^i(cw+d)-\bigwedge^i(diagonal(\beta_2,\ldots,\beta_n))\right\|\leq
$$

.

$$
\leq \frac{1}{|\beta_1|} \left\| \bigwedge^{i+1} \left(\begin{array}{cc} 1 & -w \\ -v & (1-wv)\mathbb{1} + vw \end{array} \right) \right\| \cdot \left\| \bigwedge^{i+1} (A) - \bigwedge^{i+1} (B) \right\| + \frac{|\alpha - \beta_1|}{|\beta_1|} \left\| \bigwedge^i (cw + d) \right\|
$$

and the last lemma proves the claim. The inductive use of this claim and the last lemma now proves the lemma. \Box

The assumptions of Lemma [2.8](#page-12-0) contain two estimates. In the application we have in mind the first is just the content of Corollary [2.3](#page-8-0), but to ensure the second estimate we need two further lemmata.

Lemma 2.9 For all $\delta > 0$ and $l \in \mathbb{N}$ there exists an $\epsilon > 0$, such that for all $\lambda \in O_{l,\epsilon}$

$$
|\sinh(\lambda)| \ge \frac{1}{\delta |\lambda|^l} \exp\left(|\Re(\lambda)|\right)^{17}.
$$

Proof: First we use the following bound from below: $|\sinh(\lambda)| \geq \exp(|\Re(\lambda)|) |\sin(\Im(\lambda))|/2$. To each $\delta > 0$ there exists an $\epsilon > 0$, such that for all $\sqrt{-1} \Im(\lambda) \in O_{l,\epsilon}$ $|\sin(\Im(\lambda))| \geq 2\delta^{-1} |\lambda|^{-l}$, and therefore the above estimate also holds. On the other hand $|\sinh(\lambda)|$ is again bounded from below by

$$
|\sinh(\lambda)| \ge \frac{1}{2} (1 - \exp(-2|\Re(\lambda)|)) \exp(|\Re(\lambda)|).
$$

Hence the proposed estimate holds also in the domain $(1 - \exp(-2|\Re(\lambda)|)) \geq 2\delta^{-1}|\lambda|^{-l}$. The union of the domains of both kinds clearly contains some $O_{l,\epsilon}$.

Lemma 2.10 Let $\beta_1(\lambda)$ and $\beta_2(\lambda)$ be two functions of the form

$$
\beta_1(\lambda) = \exp(p_1\lambda + polynomial(\lambda^{-1})) = \exp(p_1(\lambda)),
$$

$$
\beta_2(\lambda) = \exp(p_2\lambda + polynomial(\lambda^{-1})) = \exp(p_2(\lambda))
$$

Then for all $\delta > 0$ and $l \in \mathbb{N}$ there exists an $\epsilon > 0$, such that for all $\frac{p_1(\lambda)-p_2(\lambda)}{2} \in O_{l,\epsilon}$

$$
|\beta_1(\lambda) - \beta_2(\lambda)| \ge \frac{1}{\delta |\lambda|^l} \sup \{ |\beta_1(\lambda)|, |\beta_2(\lambda)| \}.
$$

Proof: We have

$$
\exp\left(p_1(\lambda)\right) - \exp\left(p_2(\lambda)\right) = 2 \exp\left(\frac{p_1(\lambda) + p_2(\lambda)}{2}\right) \sinh\left(\frac{p_1(\lambda) - p_2(\lambda)}{2}\right),
$$

$$
\sup\left\{|\exp\left(p_1(\lambda)\right)|, |\exp\left(p_2(\lambda)\right)|\right\} \left|\exp\left(-\frac{p_1(\lambda) + p_2(\lambda)}{2}\right)\right| = \exp\left(\left|\Re\left(\frac{p_1(\lambda) - p_2(\lambda)}{2}\right)\right|\right)
$$

Hence the last lemma gives for each $\delta > 0$ an $\epsilon > 0$, such that for all $\frac{p_1(\lambda)-p_2(\lambda)}{2} \in O_{l,\epsilon}$

$$
|\exp(p_1(\lambda)) - \exp(p_2(\lambda))| \ge \frac{\exp(\sup \{ \Re(p_1(\lambda)), \Re(p_2(\lambda)) \})}{\delta} \left| \frac{p_1(\lambda) - p_2(\lambda)}{2} \right|^{-l}.
$$

¹⁷This estimate is proven in [\[P-T](#page-91-0), Lemma 2.1] for $l = 0, \delta = 4$ and $\epsilon = 4/\pi$.

.

For $p_1 - p_2$ not being zero, there surely exists some neighbourhood of $\lambda = \infty$ and some $c > 0$ such that

$$
\left|\frac{p_1(\lambda)-p_2(\lambda)}{2}\right|^{-l} \ge c|\lambda|^{-l}.
$$

This proves the lemma. \Box

Now we are ready to give the Proof of Theorem [2.6:](#page-10-0) We want to apply Lemma [2.8.](#page-12-0) Two estimates are assumed in this lemma. These estimates are shown to be fulfilled one after another. First we make an observation. If g is the fundamental solution of the differential equation

$$
\left(\frac{d}{dx} + a(x)\right)g(x) = 0, \ \ g(0) = \mathbf{1},
$$

then $\Lambda^{i}(g)$ is the fundamental solution of the differential equation

$$
\left(\frac{d}{dx} + d\bigwedge^i (a(x))\right) \bigwedge^i (g(x)) = 0, \ \bigwedge^i (g(0)) = 1.
$$

Here $d \bigwedge^i (A)$ is defined to be $(A \wedge \mathbf{1} \wedge ... \wedge \mathbf{1}) + ... + (\mathbf{1} \wedge \mathbf{1} \wedge ... \wedge A)$. We also remark that if $||A - B|| \leq \epsilon$, then $||d \Lambda^{i}(A) - d \Lambda^{i}(B)|| \leq i\epsilon$. If we take the two Lax operators of inequality [\(4](#page-9-0)) in Theorem [2.4](#page-8-0) and the i -th exterior powers, respectively, these inequalities ensure the validity of the assumption of Corollary [2.3](#page-8-0) with $\epsilon = c|\lambda|^{-M-1}$. Corollary 2.3 for its part ensures the first estimate of the assumption in Lemma [2.8](#page-12-0) with $\epsilon = c|\lambda|^{-M-1}$ for some $c > 0$ (not necessarily the same as in Theorem [2.4\)](#page-8-0). Now the second estimate of the assumption of Lemma [2.8](#page-12-0) is guaranteed by Lemma [2.10,](#page-13-0) if δ in Lemma [2.8](#page-12-0) is changed to $\delta c |\lambda|^{l-M-1}$ and the same ϵ as in Theorem [2.6.](#page-10-0) Now the first estimate of Lemma [2.8](#page-12-0) ensures that the matrix, which diagonalizes thefirst Lax operator in the inequality ([4\)](#page-9-0) exists and is uniformly bounded by $\delta c |\lambda|^{l-M-1}$ with some constant c (not necessary the same as above) on the domain of Lemma [2.10](#page-13-0). The second estimate of Lemma [2.8](#page-12-0) proves the second estimate of Theorem [2.6](#page-10-0). \Box

3 The Riemann surface

In this section we will introduce the Riemann surface corresponding to the Floquet matrix. For the moment it is defined as the curve given in terms of the eigenvalue equation of the Floquet matrix

$$
R(\lambda, \mu) = \det \left(\mu \mathbf{1} - g(1, \lambda, q) \right) = 0.
$$

After normalization this is an open n-fold covering of the complex plane $\lambda \in \mathbb{C}$ and therefore an open smooth Riemann surface (see for example Theorem 8.9[[Fo](#page-90-0)]). We intend to establish a one to one correspondence between some line bundles and potentials. This construction has already been carried out in the case of algebraic geometric potentials[[Kr-1\]](#page-90-0) , [\[R-S-T\]](#page-91-0) . On open Riemann surfaces all line bundles are trivial (Theorem 30.3[[Fo](#page-90-0)]), so we have to impose some decay condition near $\lambda = \infty$. We want to do this with the help of the asymptotic expansion of the last section. Indeed, similar to the algebraic geometric case, we may add covering points of $\lambda = \infty$ to the Riemann surface, if we concede them an exceptional position. In this section we want to provide a first impression of this strange behaviour of the Riemann surface near

 $\lambda = \infty$. It can be summarized in the statement that almost everything holds as in the context of compact Riemann surfaces.

Definition 3.1 From now on we fix a smooth periodic potential q. Theorem [2.6](#page-10-0) shows that for an arbitrary $l \in \mathbb{N}_0$ over some $U_{l,\epsilon}$ the Riemann surface is an unbranched n-fold covering, and that there is a natural way to index the sheets with the numbers $i = 1, \ldots, n$ corresponding to the eigenvalue μ_i of the Floquet matrix. Each of these sheets can of course uniquely be extended to those small excluded domains of $U_{l,\epsilon}$, where only other sheets have branchpoints. To each extended sheet over a domain near $\lambda = \infty$ we add the point $\lambda = \infty$ and take the extended sets of $U_{l,\delta}$, $\delta \leq \epsilon$, as a base of neighbourhoods of this new point: For each $i = 1, \ldots, n$ these neighbourhoods are the liftings into the i-th sheet of the sets

$$
\left\{\lambda \in \mathbb{C} \left| \frac{p_{i,l}(\lambda) - p_{j,l}(\lambda)}{2} \in O_{l,\delta} \text{ for all } j \neq i \right\} \bigcup \left\{\infty\right\}.
$$

(Compare with Definition [2.5\)](#page-9-0). This base of neighbourhoods of the n covering points over $\lambda = \infty$ together with the usual topology over $\mathbb C$ defines the topology τ_l^{18} . Because $O_{l,\epsilon}$ is contained in $O_{m,\epsilon}$ for $\epsilon < 1, l \leq m$, these topologies are ordered: $\tau_0 \supset \tau_1 \supset \ldots$. Finally we define τ_{∞} as the finest topology, which is coarser than all these topologies. This n-fold covering over $\mathbb{C} \cup \{\infty\}$ together with the topology τ_{∞} will be called the Riemann surface Y. Taking the sets $U_{l,\epsilon} \cup \{\infty\}$ as a base of neighbourhoods of $\lambda = \infty$ there is also a sequence of topologies $\tau_0 \supset \tau_1 \supset ... \supset \tau_\infty$ on $\mathbb{C} \cup \{\infty\}$. The space $\mathbb{C} \cup \{\infty\}$ together with the topology τ_{∞} will be called X.

Remark 3.2 We will see in the next section that the Riemann surface Y may be an unbranched covering over some of these excluded domains. For the analytic investigation of the Riemann surface Y it would be appropriate not to exclude such domains from the neighbourhoods of the covering points of infinity. But it could happen that a singular Riemann surface, namely with singularities in such unbranched coverings over excluded domains, corresponds to the potential q in a way, which is established in the next section. The topology defined above fits in the most singular case, namely the algebraic curve defined by the equation $R(\lambda, \mu) = 0$.

Obviously X and Y are Hausdorff spaces, but no topological manifolds. The covering map $\pi : Y \mapsto \mathbb{P}_1$ is continuous, but in general the map $\pi : Y \mapsto X$ is not continuous. A function on X and Y, respectively is called holomorphic, if it is holomorphic on $\mathbb C$ and $\pi^{-1}(\mathbb C)$, respectively and if all derivatives of f can be extended continuously to the whole of X and Y , respectively. Analogously meromorphic functions are meromorphic functions on $\mathbb C$ and $\pi^{-1}(\mathbb C)$, respectively, such that on some neighbourhood of infinity and $\pi^{-1}(\infty)$ respectively $\lambda^{-1}f$ is holomorphic for some $l \in \mathbb{N}$. Now it should be evident how to define the sheaves \mathcal{O}_X and \mathcal{O}_Y of holomorphic functions and \mathcal{M}_X and \mathcal{M}_Y of meromorphic functions.

Lemma 3.3 For each open set U of \mathbb{P}_1 , the sections of the sheaf \mathcal{O}_X restricted to U coincide with the holomorphic functions on U in the usual sense. In particular $H^0(X, \mathcal{O}_X) = \mathbb{C}$ and $H^0(Y, \mathcal{O}_Y) = \mathbb{C}.$

¹⁸For $l = 0$ we assume that $\epsilon > 2/\pi$ (compare with footnote [14](#page-9-0)).

Proof: By the maximum modulus Theorem [\[Co](#page-90-0)] the sections of \mathcal{O}_X restricted to some U and all their derivatives are continuous with respect to the topology of \mathbb{P}_1 . Hence they are holomorphic in the usual sense. All holomorphic functions on \mathbb{P}_1 are constant, so that $H^0(X, \mathcal{O}_X) = \mathbb{C}$ follows. Due to the same argument all the elementary symmetric functions of a holomorphic function on Y with respect to the covering $\pi : Y \mapsto \mathbb{P}_1$ [\[Fo\]](#page-90-0) have to be constant and therefore the holomorphic functions on Y also have to be constant. the holomorphic functions on Y also have to be constant.

This is the first analogy of our Riemann surface with compact Riemann surfaces. Before we proceed to further analogies, let us join on to the last section.

Lemma 3.4 A function holomorphic on some $U_{l,\epsilon}$ can be extended to a function holomorphic on $U_{l,\epsilon} \cup \{\infty\}$, iff it can be expanded asymptotically uniformly on some $U_{L,\delta}$ for each $L > l$ in the following manner

$$
\left| f(\lambda) - \sum_{m=0}^{M} a_m \lambda^{-m} \right| < \frac{c_M}{|\lambda|^{M+1}} \text{ for all } M \in \mathbb{N}.
$$

Proof: If f is holomorphic on some $U_{l,\epsilon}$, it has by definition a Taylor expansion at $\lambda = \infty$. This Taylor expansion obviously gives an asymptotic expansion of the desired form. Conversely let f have such an asymptotic expansion. For each $\lambda_0 \in U_{L,\delta'}$ with $\delta' < \delta$ there is a ball $B(\lambda_0^{-1}, c|\lambda_0|^{-L-2}) = \left\{ \lambda \mid |\lambda^{-1} - \lambda_0^{-1}| < c|\lambda_0|^{-L-2} \right\}$ inside $U_{L,\delta}$ with some $c > 0$ depending on L and δ but not on λ_0 . Now with the help of Cauchy's estimate [\[Co](#page-90-0)] the asymptotic expansion leads to an asymptotic expansion of $f' = \frac{df}{d\lambda^{-1}}$ on $U_{L,\delta'}$ of the form

$$
\left| f'(\lambda) - \sum_{m=0}^{M} m a_m \lambda^{-m+1} \right| < \frac{c_M}{c |\lambda|^{M-L+3}}
$$

and therefore

$$
\left| f'(\lambda) - \sum_{m=0}^{M} m a_m \lambda^{-m+1} \right| < \frac{\tilde{c}_M}{|\lambda|^{M-L+3}}
$$

with some new constants \tilde{c}_M . This proves that $f'(\lambda)$ is continuous with respect to the topology τ_L and has an asymptotic expansion similar to f. The repeated application of this claim to higher and higher derivatives shows that all derivatives can continuously be extended with respect to all topologies τ_L . This proves the claim. \Box There are similar statements about the holomorphic functions on Y with completely analogous

proofs.

We emphasize that the inverse of a meromorphic function is not necessarily also a meromorphic function. In fact, the inverse of a meromorphic function would be meromorphic, if we could exclude that this meromorphic function has a zero of infinite order at infinity or at some of the covering points of infinity, respectively. However, this is not possible and we will provide an example of a function with these properties at the end of this section. Now it is quite obvious that the divisor of an invertible meromorphic function consists of a finite part far away from infinity, of sequences of finite divisors inside the small excluded domains of the neighbourhood of infinity and of a finite contribution at infinity. More precisely, to each meromorphic function there exists a neighbourhood of infinity and $\pi^{-1}(\infty)$ respectively, such that the restriction of this function to this neighbourhood has poles or zeros at most at infinity. Let us choose a

subset of this neighbourhood out of the base given above in Definition [3.1](#page-15-0), defining a sequence of excluded domains. In the sequel we will often make some statements about these excluded domains of this base of neighbourhoods. For this purpose we abbreviate the multiple index (i, j, k) used in Definition [3.1](#page-15-0) by ι :

$$
\iota \in \mathcal{I}, \text{ with } \mathcal{I} = \{ (i, j, k) \mid k \in \mathbb{Z}, \ i < j \in \{1, \dots, n\} \}.
$$

Moreover, the absolute value of ι is defined as $|\iota| = |(i, j, k)| = |k|$. This index will be used as follows. Given any neighbourhood $U_{l,\epsilon}$ as in Definition [2.5,](#page-9-0) the index $\iota = (i, j, k) \in \mathcal{I}$ specifies the excluded domain of $U_{l,\epsilon}$ defined as the union of the *i*-th and *j*-th sheet in Y over the set of λ 's satisfying

$$
\left|\frac{p_{i,l}(\lambda)-p_{j,l}(\lambda)}{2}-k\pi\sqrt{-1}\right|\leq \frac{1}{\epsilon(|k|\pi)^l}.
$$

By definition, if $k = 0$, the excluded domains form the whole of Y. Excluded domains having non empty intersection will be identified. Thus the set $\mathcal I$ does not really label the excluded domains. For the sake of simplicity we do not use a more precise notation. The actual precise meaning will be clear from the context.

Holomorphic and meromorphic differential forms on X and Y , respectively, are defined near infinity as holomorphic and meromorphic functions, respectively, times $d\lambda^{-1}$ and otherwise as usual.

Theorem 3.5 For each meromorphic differential form, the sequence $(r_{\iota})_{\iota \in \mathcal{I}}$ of the corresponding total residues of the excluded domains defines an element of the Fréchet space

$$
l_{\mathcal{I}}^{\infty} = \left\{ (\alpha_{\iota})_{\iota \in \mathcal{I}} \; \left| \; \sum_{\iota \in \mathcal{I}} |\alpha_{\iota}| \cdot |\iota|^{l} < \infty \; \text{ for all } l \in \mathbb{N}_{0} \right. \right\}
$$

Hence the total residue is well defined and is equal to zero.

Proof: The sequence of residues of every form $\omega = \left(\sum_{m=M_1}^{M_2} a_m \lambda^{-m}\right) d\lambda$ is asymptotically equal to zero for arbitrary integers $M_1 \leq M_2$. If we use the Cauchy integral representation of the total residue, the asymptotic expansion of the meromorphic differential form analogous to the asymptotic expansion of Lemma [3.4](#page-16-0) shows the convergence of the sequence of total residues to be faster than any inverse power of $|\lambda|$. Since the absolute value of the index ι is bounded by some constant times the absolute value of λ inside the excluded domain with index $\iota : |\iota| < c|\lambda|$, the claim about the asymptotic behaviour follows. The last statement is now obvious. the claim about the asymptotic behaviour follows. The last statement is now obvious.

Corollary 3.6 Let f be a meromorphic function, whose inverse is also a meromorphic function. Then the sequence of total degrees $(d_i)_{i\in\mathcal{I}}$ of the excluded domains is asymptotically equal to zero: $d_i = 0$ for all $|i| \geq N$, with some positive integer N. Therefore the total degree of f is well defined and is equal to zero.

Proof: Due to the assumption on f, $f^{-1}df$ is a meromorphic differential form. Therefore the residue of this form is a integer multiple of $2\pi\sqrt{-1}$ and the claim follows from the preceeding theorem. \Box

Remark 3.7 As a consequence of this corollary the equivalence classes of divisors decompose into components in analogy to the connected components of the Picard group of compact Riemann surfaces. But they are not only labeled by the total degree as in the case of compact Riemann surfaces: For every divisor, which in general is defined as a cross section of the multiplicative sheaf $\mathcal{M}^*_X/\mathcal{O}^*_X$ and $\mathcal{M}^*_Y/\mathcal{O}^*_Y$ respectively, there exists a neighbourhood of infinity and all covering points of infinity, respectively, such that the restriction to this neighbourhood may have only contributions at infinity and all covering points of infinity respectively. Hence we can associate to each divisor a sequence of total degrees in the excluded domains indexed by $\iota \in \mathcal{I}$:

 $deg: \text{ divisors } \to \text{ integer valued sequences indexed by } \iota \in \mathcal{I}, \ D \mapsto (\deg_{\iota}(D))_{\iota \in \mathcal{I}}.$

If the absolute value of ι is small, deg_{ι} depends on the choice of $U_{l,\epsilon}$ defining the excluded domain with index ι. Moreover, as mentioned before, there corresponds only one degree to excluded domains having a non empty intersection, namely the total degree of the union of these excluded domains. If $\deg_{\iota}(D) = \deg_{\iota}(D')$ for all $|\iota| \geq N$, with some positive integer N, the degrees of D and D' are called asymptotically equal. If in addition $\sum_{|\iota| \leq K} \deg_{\iota}(D) - \deg_{\iota}(D') = 0$ for all $K \geq N$ we call the degrees of two such divisors asymptotically and totally equal. Hence the degrees of two linear equivalent divisors D and D' are asymptotically and totally equal. Consequently the components of the equivalence classes of divisors are labeled by equivalence classes of integer valued sequences indexed by $\iota \in \mathcal{I}$, which are asymptotically and totally equal.

Having exhibited some analogies between our Riemann surfaces X and Y and ordinary compact Riemann surfaces, we will turn to the question whether there actually exist meromorphic functions with infinitely many poles. Before we answer this question in the next section for the surface Y, we will now give the answer for the surface X^{19} .

Example 3.8 Let $(\alpha_i)_{i\in\mathbb{N}}$ and $(\beta_i)_{i\in\mathbb{N}}$ be two sequences of $\mathbb{C} \subset X = \mathbb{P}_1$ satisfying the following conditions:

- (i) $|\alpha_{i+1}| \geq |\alpha_i|, |\beta_{i+1}| \geq |\beta_i|.$
- (ii) Both sequences have no accumulation points in X.
- (iii) For all large i, α_i and β_i are contained in a common excluded domain.
- (iv) $|\alpha_i|^l > i$, $|\beta_i|^l > i$ for $i \geq I$, with some positive integers I and l.

Due to (ii) each excluded domain contains only a finite number of α 's and β 's. Condition (iv) ensures that this number is bounded by some power of $|\lambda|$ with λ being a point in the excluded domain. In consideration of the topology of X these conditions ensure that the sequence $(\alpha_i - \beta_i)_{i \in \mathbb{N}}$ defines an element of the Fréchet space l_{∞} . Here l_{∞} is

$$
l_{\infty} = \left\{ (\alpha_{\iota})_{\iota \in \mathbb{N}} \; \left| \; \sum_{\iota \in \mathbb{N}} |\alpha_{\iota}| \cdot |\iota|^{l} < \infty \; \text{ for all } \, l \in \mathbb{N}_{0} \right. \right\}
$$

Now with help of the well known test of convergence for infinite products[[Co](#page-90-0)] it is straightforward to prove that $f(\lambda) = \prod_{i \in \mathbb{N}} (\lambda - \alpha_i) / (\lambda - \beta_i)$ is a meromorphic function on X.

¹⁹These two examples may be passed over.

4 THE DUAL EIGEN BUNDLE 19

Example 3.9 For all $\iota \in \mathcal{I}$ let $\alpha_{\iota}(\lambda)/\beta_{\iota}(\lambda)$ be a rational function on \mathbb{P}_1 , which has a zero at infinity and poles only inside the image under π of the excluded domain corresponding to the index ι of some $U_{l,\epsilon}$ for all $l \in \mathbb{N}$. Let $\|\cdot\|_{l,\epsilon,\iota}$ be the supremum norm on the boundary of the domain excluded from $U_{l,\epsilon}$ and corresponding to the index ι . Then it is quite easy to prove that

$$
f(\lambda) = \sum_{\iota \in \mathcal{I}} \frac{\alpha_{\iota}(\lambda)}{\beta_{\iota}(\lambda)}
$$

is a meromorphic function on X, whenever

$$
\sum_{\iota\in\mathcal{I}}\left\|\frac{\alpha_\iota(\lambda)}{\beta_\iota(\lambda)}\right\|_{l,\epsilon,\iota}<\infty\;\;\text{for all}\;\;l\in\mathbb{N}\;\;\text{with some}\;\;\epsilon>0\;\;\text{depending on}\;\;l.
$$

This even implies that for all $l, k \in \mathbb{N}$ and the same ϵ as before \sum ι∈I $\alpha_{\iota}(\lambda)$ $\beta_{\iota}(\lambda)$ $\bigg\|_{l,\epsilon,\iota} |\iota|^k < \infty.$

Finally we want to present the promised counterexample of a meromorphic function, whose inverse is not a meromorphic function. Let $g(x)$ be a smooth periodic function, with a Taylor expansion at $x = 0$ identically equal to zero. Then the coefficients of the Fourier expansion $g(x) = \sum_{j\in\mathbb{Z}} \gamma_j \exp\left(2\pi j\sqrt{-1}x\right)$ satisfy the relations $\sum_{j\in\mathbb{Z}} j^l \gamma_j = 0$ for all $l \in \mathbb{N}$. Now we define

$$
f(\lambda) = \sum_{j \in \mathbb{Z}} \frac{\gamma_j \lambda}{\lambda - j} = \sum_{j \in \mathbb{Z}} \frac{\gamma_j}{1 - j/\lambda},
$$

which according to Example 3.9 is a meromorphic function on some appropriately chosen X. But all derivatives of f at infinity vanish:

$$
f^{(l)}(\lambda) = \sum_{j \in \mathbb{Z}} \frac{(j)^l \gamma_j l!}{(1 - j/\lambda)^{l+1}}.
$$

Hence the inverse of f is not meromorphic. On the other hand f is not identically zero, if g is not identically zero. With the choice

$$
g(x) = \exp\left(\frac{1}{(z-1)^2}\right), \ \ z = \exp\left(2\pi\sqrt{-1}x\right)
$$

we obtain a meromorphic function $f \neq 0$, whose inverse is not meromorphic.

4 The dual eigen bundle

In this section we will associate a line bundle $E(q)$ over the Riemann surface Y to the potential q. This correspondence $q \mapsto E(q)$ will be injective, which means that although there are many potentials, which correspond to the same Riemann surface Y , there is only one, which in addition corresponds to a given line bundle. Moreover, we will be able even to reconstruct the potential from the line bundle. This inversion is called the inverse scattering map or the inverse spectral transform. Indeed, in the case of Hill's equation for example, the line bundle can be described in terms of the corresponding Dirichlet or Neumann eigenvalues; both of them essentially define equivalent divisors corresponding to the line bundle (for the precise statements seee.g. $[MK-T-1]$ $[MK-T-1]$.

The matrix $h(\lambda)$ in Theorem [2.6](#page-10-0) combines all eigenvectors of the Floquet matrix. Therefore it can also be used to describe the eigen bundle. But in case the potential has a trivial Taylor expansion at $x = 0$, the offdiagonal part of h has a trivial asymptotic expansion at infinity. This would face us with the problem of poles of infinite order. To avoid this we conjugate the Lax operator and therefore also the Floquet matrix with the matrix

 $h_0 =$ 1 \sqrt{n} $\begin{pmatrix} 1 & 1 & \dots & 1 \end{pmatrix}$ $\overline{}$ 1 s . . s^{n-1} 1 s^{n-1} . $s^{(n-1)(n-1)}$ \setminus $\overline{}$ $s \text{ a primitive } n\text{-th root of unity},$

i.e. $L \mapsto Ad(h_0)L$, $g(1, \lambda, q) \mapsto F(\lambda, q) = Ad(h_0)g(1, \lambda, q)$ and $h(\lambda) \mapsto h_0h(\lambda)$.

Lemma 4.1 Both equations

$$
\pi^* \left((F(\lambda, q)) v = v \mu \quad and \tag{8}
$$

$$
w\pi^*(F(\lambda, q)) = \mu w \tag{9}
$$

have solutions being vector valued meromorphic functions $v =$ $\sqrt{ }$ $\overline{}$ v_1 . . . v_n \setminus $\Big\}, w = (w_1, \ldots, w_n)$ on

Y. These solutions are unique under the additional normalization condition $v_i = 1$ and $w_i = 1$ for any fixed $i = 1, \ldots, n$.

Proof: From the definition of the Riemann surface we know that $\det (\mu \mathbb{1} - F(\lambda, q)) = 0$. Therefore for any $\lambda \in \mathbb{C}$ there exist solutions v and w as above with

$$
(\mu \mathbf{1} - F(\lambda, q)) v = 0 \text{ and } w(\mu \mathbf{1} - F(\lambda, q)) = 0.
$$

Explicitly such solutions v and w can be given in terms of polynomials of the entries of $F(\lambda, q)$ and μ . Therefore at least formally we can explicitly write down solutions of the desired form in terms of rational functions in the entries of $F(\lambda, q)$ and μ . If the denominator does not vanish identically, this gives a meromorphic solution on $\pi^{-1}(\mathbb{C})$. In the proof of Theorem [2.6](#page-10-0) it was shown that there exist values of λ , for which all eigenvalues are distinct. This implies the uniqueness of the solutions of the desired form. Furthermore Theorem [2.6](#page-10-0) gives a quotient of two asymptotic expansions for the solutions. The leading term of the denominator is given by some entry of h_0 and is therefore not zero. Hence we obtain an asymptotic expansion near infinity of the solutions, guaranteeing the formal solutions on $\pi^{-1}(\mathbb{C})$ to be meromorphic. Now Lemma [3.4](#page-16-0) proves the claim. \Box

This lemma shows that equations (8) and (9) define holomorphic maps from Y to \mathbb{P}_{n-1} and therefore holomorphic line bundles on Y. The line bundles described by v and w are called the eigen bundle and the transposed eigen bundle of q and denoted by $E(q)$ and $E^t(q)$ respectively.

4 THE DUAL EIGEN BUNDLE 21

It is more convenient to work with the corresponding dual bundles $E^*(q)$ and $E^{t*}(q)$. We want to use divisors to describe these line bundles: Set $D(q)$ to be the negative divisor of v with $v_1 = 1$ and $D^t(q)$ to be the negative divisor of w with $w_1 = 1$. They are both integral divisors. In general a divisor of a matrix valued meromorphic function is defined by the highest order of the poles and by the lowest order of the zeros of all its entries respectively. Finally let us introduce the sheaf \mathcal{O}_q of holomorphic functions on \mathbb{P}_1 with values in the space of the $n \times n$ matrices, which commute with $F(\cdot, q)$. Outside of the branchpoints, each diagonalization of F also diagonalizes every section of \mathcal{O}_q . Hence the eigenvalues of sections of \mathcal{O}_q can be considered asholomorphic functions on Y , at least outside the branchpoints, and with the help of [[Fo](#page-90-0), Theorem 8.2] everywhere on Y. This establishes a sheaf homomorphism $\varepsilon_q : \mathcal{O}_q \to \pi_*(\mathcal{O}_Y)$ of sheaves of rings on \mathbb{P}_1 . The map ε_q is injective, because the diagonalization of F can be used to invert ε_q outside of the branchpoints.

Definition 4.2 The potential q is called non-singular, if ε_q is an isomorphism of sheaves and singular, if ε_q is not surjective. If q is singular the image of \mathcal{O}_q defines a singular Riemann surface Y' with normalization $p: Y \to Y'$, such that the following diagram commutes

and the induced map $\varepsilon'_q : \mathcal{O}_q \to \pi'_*(\mathcal{O}_{Y'})$ is an isomorphism of sheaves. Here $\mathcal{O}_{Y'}$ is the structure sheaf of Y' .

The commutative diagram ensures that Y' can only have covering points over the same base point as multiple points. This is in fact the only possibility for singular points: a branchpoint of the eigenvalues of F , which is removed on Y . But we want to emphasize that there may exist branchpoints of the eigenvalues of F , which also are removed on Y' . Hence we have in general three branching divisors $b_{\text{analytic}} \leq b_{\text{effective}} \leq b_{\text{algebraic}}$ on Y. Here the analytic branching divisor is defined by the branching order of the map π . The effective branching divisor is in some sense the branching divisor of the singular Riemann surface Y'. It will be defined in Section [10.](#page-73-0) Finally the algebraic branching divisor is defined by the zeros of $\frac{\partial R(\lambda,\mu)}{\partial \mu}$. The algebraic branching divisor is equal to the effective branching divisor corresponding to the most singular curve, which is defined by $R(\lambda, \mu) = 0$. From now, unless stated explicitly, we restrict ourselves to the simplest case $b_{\text{analytic}} = b_{\text{effective}} = b_{\text{algebraic}}$. The general case will be treated in Section [10](#page-73-0).

The global sections $v_1 = 1, v_2, \ldots, v_n$ and $w_1 = 1, w_2, \ldots, w_n$ define sheaf homomorphisms $\phi_v: \mathcal{O}_{\mathbb{P}_1}^n \to \pi_*(\mathcal{O}_{D(q)})^{20}, (f_1, \ldots, f_n) \mapsto \sum_{i=1}^n f_i v_i$ $\phi_w: \mathcal{O}_{\mathbb{P}_1}^{\tilde{n}} \to \pi_*(\mathcal{O}_{D^t(q)}), (f_1, \ldots, f_n) \mapsto \sum_{i=1}^n f_i w_i.$

Theorem 4.3 ϕ_v and ϕ_w are isomorphisms. In particular v_1, \ldots, v_n and w_1, \ldots, w_n span the space of global sections of $\mathcal{O}_{D(q)}$ and $\mathcal{O}_{D^t(q)}$, respectively. Moreover, v_1, \ldots, v_n and w_1, \ldots, w_n are uniquely defined by $D(q)$ and $D^t(q)$, respectively and by their values at all covering points of infinity.

²⁰For all divisors D the sheaf \mathcal{O}_D is defined to be the sheaf of meromorphic functions f satisfying locally $(f) \geq -D$.

4 THE DUAL EIGEN BUNDLE 22

Proof: The proof is given for v. It is the same for w. Clearly the image of ϕ_v is an \mathcal{O}_q module. Hence it is also an $\pi_*(\mathcal{O}_Y)$ module. But $\mathcal{O}_{D(q)}$ is generated by v_1, \ldots, v_n as an \mathcal{O}_Y module. Therefore ϕ_v is surjective. At each point λ_0 with distinct eigenvalues μ_1, \ldots, μ_n of $F(\lambda_0, q)$, the values of v constitute a basis of \mathbb{C}^n . Hence ϕ_v is injective in a neighbourhood of such a point. Then ϕ_v is an isomorphism. The values of v at all covering points of infinity are the columns of h_0 . There they also form a basis. This verifies the last claim. \Box

The dependence on the values at infinity suggests a modification of the notion of linear equivalence: Modified principal divisors are defined as divisors of meromorphic functions, which are equal to 1 at all covering points of infinity. This also excludes meromorphic functions with zeros of infinite order. Geometrically this corresponds to the identification of all the covering points of infinity and therefore makes Y singular. But for reasons of simplicity we use the notion of modified linear equivalence. A line bundle in this modified sense is a line bundle on Y together with an identification of all its fibers over the covering points of infinity. Theorem [4.3](#page-21-0) shows that $D(q)$ is the unique integral divisor, which is modified equivalent to $D(q)$, or equivalently, that $E^*(q)$ has a unique cross section, which respects the identification over all covering points of infinity. The v_i 's are uniquely determined as the quotients of cross sections with specified values at the covering points divided by this unique cross section.

Counting Lemma 4.4 In the sense of Remark [3.7](#page-17-0) the branching divisor b has asymptotically and totally the degree $(2, 2, \ldots)$.

We want to remind the reader that although the number of branchpoints is of course infinite, this has a precise meaning: All but finitely many branchpoints are pairwise located in excluded domains indexed by $\iota \in \mathcal{I}$ with the exception of finitely many indices. Furthermore the number of excluded branchpoints is twice the number of excluded indices.

Proof: The branchpoints are the zeroes of the discriminant of $R(\lambda, \mu)$, considered as a polynomialin μ . In [[P-T](#page-91-0)] it is shown that Lemma [2.9](#page-13-0) is true even for $\delta = 4, l = 0$ and $\epsilon = 4/\pi$. Hence Lemma [2.10](#page-13-0) is also true for $\delta = 2, l = 0$ and $\epsilon = 4/\pi$: For $\lambda \in O_{0,\epsilon}$

$$
|\exp(p_1\lambda) - \exp(p_2\lambda)| \ge \frac{1}{2} \sup \{ |\exp(p_1\lambda)|, |\exp(p_2\lambda)| \}.
$$

On the other hand for $|\lambda|$ large enough Theorem [2.6](#page-10-0) gives:

$$
|\mu_1 - \exp(p_1\lambda)| < \frac{c}{|\lambda|} |\exp(p_1\lambda)|
$$
 and $|\mu_2 - \exp(p_2\lambda)| < \frac{c}{|\lambda|} |\exp(p_2\lambda)|$.

These two estimates give for $|\lambda|$ large enough and $\lambda \in O_{0,\epsilon}$:

$$
\left|(\mu_1-\mu_2)^2-(\exp(p_1\lambda)-\exp(p_2\lambda))^2\right|<|\exp(p_1\lambda)-\exp(p_2\lambda)|^2.
$$

Thesame is of course true for $\mu_i, \mu_j, \quad i \neq j \in \{1, \ldots, n\}$. Now Rouché's Theorem [[Co](#page-90-0)] proves the claim. \Box

Any two solutions of [\(8](#page-20-0)) and [\(9](#page-20-0)) can be used to define a matrix valued meromorphic function on the Riemann surface $Y: P = v(w \cdot v)^{-1}w$. If f and g are meromorphic functions on Y, and $\tilde{v} = vf$ and $\tilde{w} = gw$, then $\tilde{v}(\tilde{w} \cdot \tilde{v})^{-1}\tilde{w} = v(w \cdot v)^{-1}w$. Therefore P does not depend on the special choice of v and w . In the definition of P we may in particular assume that locally v and w have no poles and zeros. This function P has some nice properties:

 ${\bf Lemma \ 4.5 \ \ (i)} \ \ P^2 = P$

(ii) $P\pi^*(F) = \pi^*(F)P = \mu F$

(iii) The sum over the sheets of P is equal to 1 .

(iv) The divisor of P is $-b$.

Proof: (i) and (ii) directly follow from the definition. Now let v^1, \ldots, v^n and w^1, \ldots, w^n be the values of v and w, respectively at all the covering points of some λ_0 not being a branchpoint: set μ_1, \ldots, μ_n for the distinct eigenvalues of $F(\lambda_0, q)$. The following relations ensure the $v's$ and the w's to form two bases of \mathbb{C}^n : $Fv^i = v^i\mu_i$, $w^iF = \mu_iw^i$. This implies $w^iv^j = 0$ if $i \neq j$, and the v's and the w's are therefore up to a factor dual bases of \mathbb{C}^n . Hence we have $\sum_{i=1}^n v^i (w^i v^i)^{-1} w^i = 1$, which proves (iii). The negative divisor of P must be integral, because P only can have poles. In fact we may assume that v and w have neither poles nor zeroes, and only the denominator wv can vanish. The proof of (iii) shows that the poles of P are exactly the branchpoints and even more precisely, that the divisor of P is $-b$. Indeed considering the proof of (iii) we know that for y, y' in a small neighbourhood of some branchpoint y_0 , the number of zeros of the function $w(y)v(y')$, if we fix either y or y', is equal to the branching order. This proves (iv). \Box

Theorem 4.6 The divisors $D(q)$ and $D^t(q)$ have asymptotically and totally degree $(1, 1, \ldots)$ in the sense of Remark [3.7](#page-17-0). Moreover, if v and w are solutions of [\(8](#page-20-0))and ([9\)](#page-20-0) with $v_1 = 1 = w_1$, the following equation for divisors holds:

$$
D(q) + D^{t}(q) + (w \cdot v) = b.
$$
 (10)

Finally, the function wv induces an equivalence relation in the modified sense.

Proof: The divisor of $v \cdot w$ is equal to $-D(q)-D^{t}(q)$. Hence equation (10) is a direct consequence of (iv) from Lemma [4.5.](#page-22-0) At each covering point of infinity, $w \cdot v$ is equal to n. This proves the last statement. Now we claim that $D(q)$ and $D^t(q)$ have asymptotically and totally the same degree. With this claim the first statement of the theorem follows from Corollary [3.6](#page-17-0) and the counting Lemma²¹. In order to prove the claim we need another lemma. By π we also denote the induced homomorphism from the group of divisors of Y into the group of divisors of X.

Lemma 4.7 Let $g \in GL(n, \mathcal{M}_X)$ be the unique solution of $gv = w^t$. Then one has the following relation between divisors:

$$
(\det(g)) = \pi (D(q)) - \pi (D^{t}(q)).
$$

Proof: Due to Theorem [4.3](#page-21-0) the support of the divisor of $\det(g)$ is contained in the image of the union of the support of $D(q)$ and $D^t(q)$ under π . For every meromorphic function f on Y, there exists a unique $n \times n$ -matrix valued function g_f on X, such that $g_f v = v f$ and $g_f w^t = w^t f$, and the determinant of g_f is equal to the *n*-th elementary symmetric function of f with respect to the covering map π . Theorem [4.3](#page-21-0) even shows that the supports of the divisors of $\det(g) \det(g_f)$ and $\det^{-1}(g_f) \det(g)$ are contained in the image of the union of the support of $(f) + D(g)$ and

²¹Implicit in Section [5](#page-26-0) we will give another proof. In fact, the proof of the implication (i)⇒ (ii) of Theorem [5.5](#page-29-0) implies the first statement.

4 THE DUAL EIGEN BUNDLE 24

 $D^{t}(q)$ and the union of the support of $D(q)$ and $(f) + D^{t}(q)$ respectively. This proves the claim. \Box

Completion of the proof of Theorem [4.6:](#page-23-0) The function g of the last lemma is a meromorphic function on X, which is equal to $h_0^{-1}h_0^{-1}$ at infinity. Hence there exists a neighbourhood ofinfinity, such that $|\det(g) - 1| < 1$ in this neighbourhood. Now Rouché's Theorem [[Co](#page-90-0)] implies that $D(q)$ and $D^t(q)$ have asymptotically and totally the same degree. Indeed, if there exist disjoint excluded domains, whose images under π have non empty intersection, the *n*-fold covering π decomposes near these excluded domains into coverings containing actual only one excluded domain. The application of the last lemma to each of these coverings proves the claim even in this case. \Box

Due to equation [\(10\)](#page-23-0) the equivalence class of one of the divisors $D(q)$ and $D^{t}(q)$ determines the other equivalence class. In the remainder of this section we present a way to reconstruct the potential q from the divisor $D(q)$. This will be done in three steps:

- step 1: Theorem [4.3](#page-21-0) shows that the sections $v_1 = 1, v_2, \ldots, v_n$ of the sheaf $\mathcal{O}_{D(q)}$ are uniquely determined by $D(q)$.
- step 2: Theorem [2.4](#page-8-0) and Theorem [2.6](#page-10-0) enable us to obtain the complete Taylor expansion of $q(x)$ at the point $x = 0$ from the asymptotic expansion of v.
- step 3: For each $x \in \mathbb{R}/\mathbb{Z}$, let T_x be the shift by x on the space of potentials: $(T_xq)(y) = q(x+y)$. The Riemann surface Y will not change under the shift and the line bundle $E(T_xq)$ will be described in terms of the line bundle $E(q)$.

The first step has already been carried out.

In the second step we have to restrict the space of potentials:

Assumption 4.8 The diagonal part of the potential is equal to a fixed constant.

This fits with the group theoretical treatment. In fact, with this assumption the space of potentials forms a coadjoint orbit and therefore possesses a natural symplectic structure. This observation and a corresponding hamiltonian formulation is well known and can be found in e.g. $[F-T]$ $[F-T]$, $[Sch]$.

Equation([5](#page-9-0)) gives a formula for $q(0)$ in terms of the asymptotic expansion of v. One may obtain similar formulas for the higher order coefficients of the Taylor expansion of q with the help of([6\)](#page-9-0). Some details may be found in Appendix [A](#page-81-0).

Finally we turn to the third step. First we have to ensure that the shift T_x does not change the Riemann surface. For this purpose let $g(x)$ be the fundamental solution of the differential equation $\left(\frac{d}{dx} + a(x)\right)g(x) = 0$, $g(0) = 1$ with some periodic $a(x)$. Then the Floquet matrix of $T_x a$ is given by $g(x+1)g^{-1}(x) = Ad(g(x))g(1)$. Hence the eigenvalues of the Floquet matrices of all the T_xa coincide. In particular to all the potentials T_xq there corresponds the same Riemann surface.The main tool of this step is given by Floquet theory $[Fr]$ $[Fr]$:

If the Floquet matrix $g(1)$ of the above differential equation has a logarithm: $g(1) = \exp(\alpha)$, then the gauge transformation with the periodic differentiable matrix

 $h(x) = q(x) \exp(-x\alpha)$ transforms $a(x)$ into the constant α . For the application of this observation in the present context we need a lemma.

4 THE DUAL EIGEN BUNDLE 25

Lemma 4.9 On some neighbourhood of all covering points of infinity, $\ln(\mu)$ is a meromorphic function.

Proof: For ϵ small enough, the second estimate of Theorem [2.6](#page-10-0) implies an asymptotic expansion uniformly on $U_{M,\epsilon}$: $|\ln(\mu_i) - p_{i,M}(\lambda)| < c|\lambda|^{-M-1}$ with some constant depending on M. Of course only one branch of $\ln(\mu)$ satisfies this estimate. Together with Lemma [3.4](#page-16-0) this guarantees the existence of the meromorphic function $\ln(\mu)$.

Now an easy application of Floquet theory provides solutions of the equations([8\)](#page-20-0) and [\(9\)](#page-20-0) for the potential T_xq of the form:

$$
v(x,q) = \pi^* \left(h_0 g(x, \cdot, q) h_0^{-1} \right) v(q) \mu^{-x}
$$

$$
w(x,q) = \mu^x w(q) \pi^* \left(h_0 g^{-1}(x, \cdot, q) h_0^{-1} \right).
$$

Here μ^x is an abbreviation of $\exp(x \cdot \ln(\mu))$. Now we claim that for all $x \in \mathbb{R}$ these solutions are holomorphic vector valued functions on the neighbourhood described in the foregoing lemma, which take the same values as $v(q)$ and $w(q)$, respectively, at all covering points of infinity. Let $v(T_xq)$ and $w(T_xq)$ be the corresponding normalized solutions:

$$
v(T_xq) = \frac{v(x,q)}{v_1(x,q)}
$$

$$
w(T_x,q) = \frac{w(x,q)}{w_1(x,q)}.
$$

Then the functions $v_1(x, q)$ and $w_1(x, q)$ are the unique solutions of the differential equations

$$
\frac{\partial v_1(x,q)}{\partial x} = -\left(\left(\pi^*(h_0(p\lambda + q(x))h_0^{-1})v(T_xq) \right)_1 + \ln(\mu) \right) v_1(x,q), \ \ v_1(0,q) = 1 \text{ and}
$$

$$
\frac{\partial w_1(x,q)}{\partial x} = \left(\left(w(T_x q) \pi^*(h_0 (p \lambda + q(x)) h_0^{-1}) \right)_1 + \ln(\mu) \right) w_1(x,q), \quad w_1(0,q) = 1, \text{ respectively.}
$$

These solutions are given by

$$
v_1(x,q) = \exp\left(\int_0^x \left(\left(\pi^*(h_0(p\lambda + q(t))h_0^{-1})v(T_tq)\right)_1 + \ln(\mu)\right)dt\right) \text{ and}
$$

$$
w_1(x,q) = \exp\left(\int_0^x \left(\left(w(T_tq)\pi^*(h_0(p\lambda + q(t))h_0^{-1})\right)_1 + \ln(\mu)\right)dt\right), \text{respectively.}
$$

Now the asymptotic expansion of Theorem [2.6](#page-10-0), whose coefficients are calculated in Theorem [2.4](#page-8-0), shows that both functions under the integrals of are holomorphic functions on the neighbourhood described in the previous lemma. Furthermore, due to Assumption [4.8](#page-24-0) they vanish at all covering points of infinity. This proves the claim.

Thedefinition of these solutions of the equations ([8](#page-20-0)) and ([9](#page-20-0)) for the potential T_xq may be transformed to

$$
\pi^*(h_0g(x,\cdot,q)h_0^{-1})v(q) = v(x,q)\mu^x
$$
and

$$
w(x,q)\pi^*(h_0g(x,\cdot,q)h_0^{-1})\mu^xw(q),
$$
 respectively.

It is well known that $\det(g(x, \lambda, q)) = \exp(\int_0^x \text{tr}(p\lambda + q(t)) dt)$. Hence the map $\lambda \to g(x, \cdot, q)$ defines an entire function from $\mathbb C$ to $GL(n,\mathbb C)$. Thus the left hand sides define line bundles on $\pi^{-1}(\mathbb{C})$ isomorphic to the restriction of $E(q)$ and $E^t(T_xq)$, respectively. Due to Lemma [4.9](#page-24-0), the function μ^x is the cocycle of a holomorphic line bundle over Y with respect to the covering $\pi^{-1}(\mathbb{C})$ and the neighbourhood given by Lemma [4.9](#page-24-0). Let us denote this line bundle by $L(x)$. Then the foregoing claim implies that

$$
E(q) \simeq E(T_x q) \otimes L(x),
$$

$$
E^t(T_x q) \simeq E^t(q) \otimes L(x).
$$

After Theorem [4.3](#page-21-0) we introduced a modified notion for line bundles. In order to finish the reconstruction of the potential out of the line bundle in the modified sense, we remark that the claim proves these isomorphisms even to be isomorphisms in the modified sense. Since obviously $L^*(x) \simeq L(-x)$, this proves the

Theorem 4.10 There are two isomorphisms of line bundles in the modified sense for all $x \in \mathbb{R}$:

$$
E^*(T_xq) \simeq E^*(q) \otimes L(x),
$$

$$
E^{t^*}(T_xq) \simeq E^{t^*}(q) \otimes L(-x).
$$

For all $x \in \mathbb{Z}$, $L(x)$ is trivial, because μ and $1/\mu$ are holomorphic functions on $\pi^{-1}(\mathbb{C})$. The line bundles on the right hand sides are well defined for all $x \in \mathbb{C}$, but the left hand sides in general make no sense for $x \notin \mathbb{R}$. Only for analytic potentials the line bundles on the left hand side are defined for some $x \notin \mathbb{R}$. This observation provides an idea of the largeness even of the components of the Picard group $H^1(Y, \mathcal{O}_Y^*)$. From Remark [3.7](#page-17-0) we recall that the Picard group has as many components, as there are classes of asymptotically and totally equivalent integer valued sequences indexed by $\iota \in \mathcal{I}$. Now we are only interested in a small part of one component, namely the one corresponding to the sequence $(1, 1, \ldots)$. The next section will provide more details.

5 The Riemann-Roch Theorem

Our next goal is to classify the divisors, which correspond to some potential q . For this purpose we need as the main tool an appropriate version of the Riemann-Roch Theorem adapted to the present situation. This is of interest in its own right and the only subject of this section.

We start with a discussion, which will serve as a motivation. On a compact Riemann surface Y the Riemann-Roch Theorem (see e.g.[[Fo](#page-90-0)]) is given by the relation

$$
\dim H^0(Y, \mathcal{O}_D) - \dim H^1(Y, \mathcal{O}_D) = 1 - g + \deg(D),
$$

where g is the genus of Y, and D is any divisor on Y. If Y is an n-fold covering of the Riemann sphere, with branching divisor b, the genus can be calculated with the help of the Riemann-Hurwitz formula (see e.g.[[Fo](#page-90-0)]):

$$
g = \frac{\deg(b)}{2} - n + 1.
$$

 \Box

In our case $deg(b)$ is infinite, and the meaning of the Riemann-Roch Theorem is not clear. But if there is a finite interpretation of the expression $deg(D) - deg(b)/2$, then the Riemann-Roch Theorem still makes sense. Now Theorem [4.6](#page-23-0) shows that the divisors, we are interested in, are exactly of this kind. Also in Theorem [4.3](#page-21-0) the dimension of $H^0(Y, \mathcal{O}_{D(q)})$ is calculated to be equal to n. If we insert this into the Riemann-Roch Theorem, we would obtain $\dim H^1(Y, \mathcal{O}_{D(q)}) = 0$ $\dim H^1(Y, \mathcal{O}_{D(q)}) = 0$ $\dim H^1(Y, \mathcal{O}_{D(q)}) = 0$. At this point the first difficulty arises: Equation ([10](#page-23-0)) shows that there exists an exact sequence of homomorphisms of sheaves

$$
0 \to \mathcal{O}_{D(q)} \to \mathcal{O}_b \to \mathcal{O}_{D^t(q)}/\mathcal{O}_Y \to 0.
$$

If $\dim H^1(Y, \mathcal{O}_{D(q)})$ were to be equal to zero, the exact cohomology sequence of the short foregoing exact sequence would imply the short exact sequence:

$$
0 \to H^0(Y, \mathcal{O}_{D(q)}) \to H^0(Y, \mathcal{O}_b) \to H^0(Y, \mathcal{O}_{D^t(q)}/\mathcal{O}_Y) \to 0.
$$

This means that to any sequence of complex numbers indexed by the points of the divisor $D^{t}(q)$, there exists a form which is almost holomorphic and takes these values at the index points. (i.e. only with poles of order at most two at all covering points of some element of \mathbb{P}_1) This can't be true (see Theorem [9.5](#page-65-0)). The contradiction is related to the conclusion of the last section. Indeed, as we explained, we are only interested in an 'admissible' part of one component of the Picard group. Therefore we have to restrict the first cohomology groups to some 'admissible' parts. One way to avoid this difficulty is to rephrase the classical formulation of the Riemann-Roch Theorem:

$$
\dim H^0(Y, \mathcal{O}_D) - \dim H^0(Y, \Omega_{-D}) = 1 - g + \deg(D),
$$

whichdue to Serre duality (see e.g. [[Fo\]](#page-90-0)) is equivalent to the up-to-date version. There is an alternative approach, which in some sense compactifies the Riemann surface Y : We use the inverse image topology of the covering map to calculate the sheaf cohomology. For any sheaf $\mathcal F$ on Y define $\tilde H^i(Y,\mathcal F) = H^i(\mathbb{P}_1,\pi_*(\mathcal F))$. Clearly $\tilde H^0(Y,\mathcal F) = H^0(Y,\mathcal F)$. The behaviour of Čech cohomology under refinement (see e.g. [\[Fo](#page-90-0)]) shows that $\tilde{H}^1(Y,\mathcal{F})$ is in fact a subgroup of $H^1(Y,\mathcal{F}).$

Definition 5.1 A divisor D on Y is called of Riemann-Roch type, if

- (i) the support of D is contained in $\pi^{-1}(\mathbb{C})$.
- (ii) for some neighbourhood U of infinity in \mathbb{P}_1 , there exist cross sections u_1, \ldots, u_n of \mathcal{O}_D on $\pi^{-1}(U)$, such that the map $\phi_u : \mathcal{O}_{\mathbb{P}_1}^n \to \pi_*(\mathcal{O}_D)$, $(f_1, \ldots, f_n) \mapsto \sum_{i=1}^n f_i u_i$ is an isomorphism of sheaves on U.

The restriction of (i) may be dropped, but then we have to pay attention to the possibility of zeros of infinite order. On the other hand the condition (i) fits with the modification of linear equivalence in the last section. From this definition and Lemma [3.3](#page-15-0) it is quite obvious that for each divisor of Riemann-Roch type, $\pi_*(\mathcal{O}_D)$ is isomorphic to the sheaf of sections of a holomorphic vector bundle on \mathbb{P}_1 . In view of the Riemann-Roch Theorem for holomorphic vector bundles this suggests the following:

Riemann-Roch Theorem 5.2 Each divisor of Riemann-Roch type D has asymptotic degree $(1, 1, \ldots)$ in the sense of Remark [3.7.](#page-17-0) Hence $\deg(D) - \deg(b)/2$ is a well defined integer. Moreover both modified cohomology groups are finite dimensional and the following formula is valid:

$$
\dim \tilde{H}^0(Y, \mathcal{O}_D) - \dim \tilde{H}^1(Y, \mathcal{O}_D) = n + \deg(D) - \frac{\deg(b)}{2}.
$$

Proof: In condition (ii) we can impose that u_i has the same value as v_i of Lemma [4.1,](#page-20-0) with $v_1 = 1$, at all covering points of infinity. The same argument as used in the proof of Theorem [4.6](#page-23-0) shows that the divisors D and $D(q)$ have asymptotically the same degree. If in addition the u_i are even meromorphic on the whole of Y , so that the total degrees of the divisors of the vector valued function (u_1, \ldots, u_n) and of $D(q)$ coincide. But $\pi_*(\mathcal{O}_D)$ defines a holomorphic vector bundle on \mathbb{P}_1 , which always has n independent meromorphic sections [\[Gu\]](#page-90-0). Hence we can even assume that the u_i are meromorphic on \mathbb{P}_1 . Now the claim follows from the Riemann-Roch Theorem for holomorphic vector bundles $[G\mathrm{u}]$.

From the proof of Lemma [4.5](#page-22-0) we conclude that $\pi_*(\mathcal{O}_{D^t(q)})$ is natural isomorphic to the sheaf of sections of the vector bundle dual to the vector bundle associated to $\pi_*(\mathcal{O}_{D(q)})$. More generally, for each divisor D of Riemann-Roch type $b-D$ is also of Riemann-Roch type and the associated vector bundles are dual to each other. On the other hand the sheaf of holomorphic forms on Y is isomorphic to the tensor product of \mathcal{O}_b with the pullback of the sheaf of holomorphic forms on \mathbb{P}_1 . Now the Serre duality for holomorphic vector bundles on \mathbb{P}_1 [\[Gu](#page-90-0)] implies Serre duality for divisors of Riemann-Roch type on Y :

Serre duality Theorem 5.3 For each divisor of Riemann-Roch type $D \Omega_{-D}$ corresponds also to a divisor of Riemann-Roch type $D' \sim b - D - 2\pi^{-1}(\lambda)$ with some $\lambda \in \mathbb{P}_1$. Moreover there is a natural non-degenerate pairing between $\tilde{H}^0(Y, \mathcal{O}_D)$ and $\tilde{H}^1(Y, \Omega_{-D})$.

The definition of divisors of Riemann-Roch type essentially postpones the main problem, namely to find sufficiently many divisors of Riemann-Roch type. In the rest of this section we try to characterize some divisors of Riemann-Roch type through their location. In particular we are looking for sufficient geometric conditions which in addition are easy to handle.

Let us choose an arbitrary $U_{l,\epsilon}$. For each $\iota \in \mathcal{I}$ we define $\pi : Y_{l,\epsilon,\iota} \mapsto \mathbb{P}_1$ to be the *n*-fold covering, which is obtained from Y after removing all branching points outside of the domain excluded from $U_{l,\epsilon}$ with index ι . Let $D_{l,\epsilon,\iota}$ be the divisor of $Y_{l,\epsilon,\iota}$, whose support is contained in the excluded domain with index ι , and whose restriction to this excluded domain coincides with the restriction of D to this excluded domain. For large $|\iota|$ this definition does not depend on the choice of $U_{l,\epsilon}$. If we are only interested in such ι , we will sometimes omit the indices l and ϵ .

Definition 5.4 A divisor D of Y is called admissible, if

- (i) the support of D is contained in $\pi^{-1}(\mathbb{C})$.
- (ii) the direct image sheaf $\pi_*(\mathcal{O}_{D_i})$ is isomorphic to $\mathcal{O}_{\mathbb{P}_1}^n$ for all $\iota \in \mathcal{I}$, which correspond to excluded domains in the inverse image of some neighbourhood of infinity under the covering map π .

 \Box

The Riemann-Roch Theorem and the Riemann-Hurwitz formula together ensure $\deg(D_{\iota})$ to be equal to half the branching order of Y_t . Hence all admissible divisors have asymptotic degree $(1, 1, \ldots)$. Let D be any admissible divisor. For all sufficiently large ι there exist n unique cross sections $v_{1,\iota}, \ldots, v_{n,\iota}$ of $\mathcal{O}_{D_{\iota,\iota,\iota}}$, which take the same values as v_1, \ldots, v_n , with the normalization $v_1 = 1$, at the covering points of infinity. For all λ outside the domain excluded from $U_{l,\epsilon}$ with index ι we define the $n \times n$ -matrix $h_{l,\epsilon,\iota}(D,\lambda)$ as $(h_{l,\epsilon,\iota}(D,\lambda))_{i,j}$ = the value of $v_{i,\iota}$ at the covering point in the j-th sheet over λ . Therefore at infinity $h_{l,\epsilon,l}(D,\lambda)$ is equal to h_0 . Now we can state the main theorem of this section.

Theorem 5.5 The following statements are equivalent:

- (i) The divisor D is of Riemann-Roch type.
- (ii) The divisor D is admissible and $\sum_{\iota} ||h_{l,\epsilon,\iota}(D,\cdot) h_0||_{l,\epsilon,\iota} < \infty$ for all $l \in \mathbb{N}$ with some ϵ depending on l.

We defined the norm $\|\cdot\|_{l,\epsilon,\iota}$ in Example [3.9](#page-19-0) as the supremum norm on the boundary of the domain excluded from $U_{l,\epsilon}$ with index ι . The proof²² of this theorem is divided into two steps concerning the two implications. In each step we will first state some lemmata, which are needed in the following main part of the actual step. Let us begin with the implication (i) \Rightarrow (ii).

Lemma 5.6 Let $D(\lambda_0, R)$ be the circle $\{\lambda \in \mathbb{C} \mid |\lambda - \lambda_0| = R\}$ and let

 $g : D(\lambda_0, R) \to GL(n, \mathbb{C}), \ \lambda \mapsto g(\lambda)$ be an analytic map such that $||g(\lambda) - 1|| < 1$ for all $\lambda \in D(\lambda_0, R)$. This map g defines a cocycle of the covering $\{\lambda \in \mathbb{P}_1 \mid |\lambda - \lambda_0| \leq R\}$ $\{\lambda \in \mathbb{P}_1 \mid |\lambda - \lambda_0| \ge R\}$. Then the corresponding holomorphic vector bundle on \mathbb{P}_1 is trivial. Moreover, the trivialization can be chosen to be in accordance with a given identification of the fiber over infinity with \mathbb{C}^n .

Proof: We remind the reader of the biholomorphic map of \mathbb{P}_1 , which transforms $D(\lambda_0, R)$ into $D(0, 1)$ and fixes infinity. Let H be the Hilbert space $L^2(S^1, \mathbb{C}^n)$. We have the decomposition $H = H_+ \oplus H_-$ into boundary values of holomorphic maps from the interior of $D(0, 1)$ to \mathbb{C}^n and holomorphic maps defined outside of $D(0, 1)$, which are equal to zero at infinity, respectively. In[[P-S](#page-91-0), Proposition (6.3.1)] it is shown that $LGL(n,\mathbb{C})$ is a subgroup of $GL_{res}(H_+ \oplus H_-)$. Moreoverit is proven in [[P-S](#page-91-0), Theorem $(8.1.2)$ and Proposition $(8.4.1)$] that g admits a Birkhoff factorization $g = g_+g_-,$ if $g =$ $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ can be written as $\begin{pmatrix} a - bd^{-1}c & b \\ 0 & d \end{pmatrix}$ 0 d \bigwedge 1 0 $d^{-1}c$ 1 \setminus with respect to the decomposition $H = H_+ \oplus H_-$. Hence the vector bundle is trivial, if d is invertible. The assumption implies sup $\big\{\|g(\lambda)-1\| \mid \lambda \in D(0,1)\big\} < 1$ and therefore $\left(\begin{array}{cc} a & b \\ c & d \end{array}\right) - 1$ < 1 . This proves that d is invertible. In the Birkhoff factorization above $g_-\$ is equal to 1 at infinity, and the induced trivialization of the vector bundle is in accordance with a given identification of the fiber over infinity with \mathbb{C}^n . ✷

Proof of the implication (i) \Rightarrow (ii) of Theorem 5.5: Let D be a divisor of Riemann-Roch type. Then there exist n cross sections $v_1(D), \ldots, v_n(D)$ of \mathcal{O}_D over the inverse image of a

 22 The proof of this theorem may be passed over to the first paragraph in front of equation [\(13](#page-34-0)).

neighbourhood of infinity under π , which induce an isomorphism between the direct image sheaf $\pi_*(\mathcal{O}_D)$ and $\mathcal{O}_{\mathbb{P}_1}^n$. They can be chosen to take the same values as v_1, \ldots, v_n at the covering points of infinity. On some neighbourhood of infinity in X these cross sections define a holomorphic $n \times n$ -matrix valued function: $h(D, \lambda)_{i,j} =$ the value of $v_i(D)$ at the covering point in the j-th sheet over λ . Let $D(\lambda_t, R_t)$ be any circle around a domain excluded from $U_{l,\epsilon}$ with index ι , such that all other excluded domains sit outside of this circle. The holomorphic vector bundle of Lemma [5.6](#page-29-0) defined by the cocycle $g(\lambda) = h(D, \lambda)$ corresponds to the direct image sheaf $\pi_*(\mathcal{O}_{D_{l,\epsilon,\iota}})$. More precisely, the Birkhoff factorization of $h(D, \cdot)$ with respect to the circle $D(\lambda_i, R_i)$ is given by $h(D, \cdot) = (h(D, \cdot)h_{l, \epsilon, \iota}^{-1}(D, \cdot)h_0) \cdot (h_0^{-1}h_{l, \epsilon, \iota}(D, \cdot))$. Then Lemma [5.6](#page-29-0) proves that the divisor D is admissible, since $h_0^{-1}h(D, \lambda)$ is continuous and equal to the identity matrix at infinity. Moreover, $h(D, \lambda)$ has an asymptotic expansion:

$$
\left\| h(D, \lambda) - \left(h_0 + \sum_{m=1}^M a_m \lambda^{-m} \right) \right\| < \frac{c_M}{|\lambda|^{M+1}} \text{ for all } M \in \mathbb{N} \text{ uniformly on } U_{l, \epsilon}
$$

If ϵ is small enough, then $h_0 + \sum_{m=1}^M a_m \lambda^{-m}$ is a holomorphic map from $U_{l,\epsilon}$ into $GL(n,\mathbb{C})$ and furthermore \parallel $(h_0 + \sum_{m=1}^{M} a_m \lambda^{-m})^{-1}$ \leq 2 on this set. Thus we have

$$
\left\| \left(h_0 + \sum_{m=1}^M a_m \lambda^{-m} \right)^{-1} h(D, \lambda) - 1 \right\| < \frac{2c_M}{|\lambda|^{M+1}} \text{ for all } M \in \mathbb{N} \text{ uniformly on } U_{l, \epsilon}.
$$

The proof of Lemma [5.6](#page-29-0) even shows that

$$
\|g_{-}(\lambda)-1\| \le \frac{\|g(\lambda)-1\|}{1-\|g(\lambda)-1\|} \text{ for all } \lambda \in D(\lambda_0, R).
$$

Hence we obtain the estimate

$$
\left\|h_0^{-1}h_{l,\epsilon,\iota}(D,\lambda)-1\right\|_{l,\epsilon,\iota}\leq \frac{\tilde{c}_M}{|\lambda|^{M+1}}
$$

for all $M \in \mathbb{N}$ and for all indices ι , which correspond to domains excluded from $U_{l,\epsilon}$. This is true for all $l \in \mathbb{N}$ with some ϵ depending on l. This implies (ii).

Now we want to prove the implication (ii) \Rightarrow (i). Let us first explain the strategy of the proof. Let D be a divisor with the assumed properties. We will see that there exist unique cross sections $v_1(D), \ldots, v_n(D)$ of \mathcal{O}_D over $\pi^{-1}(U)$, with some neighbourhood U of infinity on \mathbb{P}_1 , which takes the same values as v_1, \ldots, v_n at infinity. Their elementary symmetric functions with respect to the covering map π [[Fo\]](#page-90-0) are of the form $c + \sum \alpha_i(\lambda)/\beta_i(\lambda)$ as in Example [3.9](#page-19-0) at the end of Section 2. This implies that D is of Riemann-Roch type. To establish this claim we will use perturbative methods. If we remove all the branchpoints and simultaneously all the points of the divisor D the resulting surface is isomorphic to $(\mathbb{P}_1)^n$ and \mathcal{O}_D becomes isomorphic to $(\mathcal{O}_{\mathbb{P}_1})^n$. Thus the claim is trivial in this case. In a first step we deform the Riemann surface Y to the compact Riemann surface Y_t over some neighbourhood of the branchpoints out of one excluded domain. By Definition [5.4](#page-28-0) the space of cross sections of $\mathcal{O}_{D_{\iota}}$ is isomorphic to the space of cross sections of $(\mathcal{O}_{\mathbb{P}_1})^n$. These deformations of the Riemann surface Y perturb each other.

In a second step we present a way to fit together all these deformations. The main tool will be provided by the elementary symmetric functions, because they determine functions on Y in terms of functions on \mathbb{P}_1 . It will turn out that these perturbations deform the isomorphism $\pi_*(\mathcal{O}_{D_i}) \simeq \mathcal{O}_{\mathbb{P}_1}$ into another isomorphism $\pi_*(\mathcal{O}_D) \simeq \mathcal{O}_{\mathbb{P}_1}$.

Let us omit the index ι in the first step. In the next lemma $\pi : Y \to \mathbb{P}_1$ is assumed to be a compact n-fold covering of \mathbb{P}_1 and D is assumed to be a divisor of Y, such that $\pi_* (\mathcal{O}_D)$ is isomorphic to $\mathcal{O}_{\mathbb{P}_1}^n$. If we impose that a cross section of \mathcal{O}_D takes the values x_1, \ldots, x_n at the covering points of infinity, this cross section is uniquely defined. Then the elementary symmetric functions of this cross section are rational functions, with poles only at the base points of the integral part of the divisor. The values at infinity are of course the usual elementary symmetric functions of x_1, \ldots, x_n . Hence for every choice of x_1, \ldots, x_n we have rational functions $c_1(\lambda, x_1, \ldots, x_n), \ldots, c_n(\lambda, x_1, \ldots, x_n)$ on \mathbb{P}_1 , such that

$$
f^{n} + f^{n-1}c_{1}(\lambda, x_{1}, \dots, x_{n}) + \dots + c_{n}(\lambda, x_{1}, \dots, x_{n}) = 0
$$
\n(11)

defines the unique cross section f of \mathcal{O}_D , which takes the values x_1, \ldots, x_n at the covering points of infinity. Each of these elementary symmetric functions has the form

$$
c_i(\lambda, x_1, \dots, x_n) = e_i(x_1, \dots, x_n) + \frac{\alpha_i(\lambda, x_1, \dots, x_n)}{\beta(\lambda)},
$$
\n(12)

where $e_i(x_1, \ldots, x_n)$ is the usual *i*-th elementary symmetric function of x_1, \ldots, x_n , $\beta(\lambda)$ is the polynomial whose zeroes are given by the base points of the integral part of the divisor and $\alpha_i(\lambda, x_1, \ldots, x_n)$ are polynomials in λ of degree less than the degree of $\beta(\lambda)$, whose coefficients are homogenous polynomials in x_1, \ldots, x_n .

Deformation Lemma 5.7 Let Y and D fulfil the conditions given above. Let U be an open subset of \mathbb{P}_1 , which contains a ball of radius $R > 0$ around all the base points of the integral part of the divisor. Moreover, let B be a ball contained in the intersection of all balls with radius R/2 around these base points, and let $\|\cdot\|_{\partial B}$ be the supremum norm on the boundary of B. Moreover, choose some values x_1, \ldots, x_n , such that $|x_i - x_j| > c > 0$ for all $i \neq j$. Assume $\|\alpha_i(\cdot, x_1, \ldots, x_n)/\beta\|_{\partial B}$ to be small enough for all $i = 1, \ldots, n$, and let f_1, \ldots, f_n be functions on U, such that $|f_i - x_i| < \epsilon$, with some $\epsilon < c/4$. Then there exists a unique cross section \tilde{f} of \mathcal{O}_D over $\pi^{-1}(U)$, such that the regular parts of the elementary symmetric functions \tilde{c}_i of \tilde{f} with respect to the covering π are equal to the usual elementary symmetric functions of f_1, \ldots, f_n :

$$
\tilde{c}_i = e_i(f_1, \ldots, f_n) + \frac{\tilde{\alpha}_i}{\beta}
$$
, with some polynomial $\tilde{\alpha}_i$ of degree smaller than the degree of β .

On the *i*-th sheet the section \tilde{f} is moreover assumed to be nearly equal to f_i away from the ball B. In this sense \tilde{f} is a deformation of f_1, \ldots, f_n on the trivial covering $\mathbb{P}_1^n \to \mathbb{P}_1$.

In order to prove this lemma we need two more lemmata.

Lemma 5.8 Let U be an open subset of \mathbb{P}_1 , which contains a ball of radius $R > 0$ around some points $\lambda_1, \ldots, \lambda_n \in \mathbb{C}$. Let $Hol(A)$ be the Banach space of holomorphic functions on U, which extend continuously to the closure A of U. Then the linear map

$$
Hol(A) \to Hol(A), \ \ g \mapsto reg\left(\frac{g}{\prod_{l=1}^d (\lambda - \lambda_l)}\right) \ \ is \ bounded \ in \ norm \ by \ (2/R)^d.
$$

Proof: We use induction in d. It is quite obvious that the regular part of $g(\lambda)(\lambda-\lambda_0)^{-1}\prod_{l=1}^d(\lambda-\lambda_l)$ $(\lambda_l)^{-1}$ is equal to the regular part of $\tilde{g}(\lambda) \prod_{l=1}^d (\lambda - \lambda_l)^{-1}$, where $\tilde{g}(\lambda) = (g(\lambda) - g(\lambda_0)) / (\lambda - \lambda_0)$. This function \tilde{g} extends to a continuous function on A. With the help of the maximum modulus Theorem [\[Co\]](#page-90-0) we obtain the bound $\|\tilde{g}\| \le \|g\|/R$. This proves the start of the induction and also each induction step also each induction step.

Lemma 5.9 In the situation of the preceeding lemma let B be a ball containing the points $\lambda_1, \ldots, \lambda_d$, which for his part is contained in the intersection of all balls with radius R/3 around the points $\lambda_1, \ldots, \lambda_d$. Again let $\|\cdot\|_{\partial B}$ be the supremum norm on the boundary of B. Moreover, let α be a polynomial of degree less than d, and $\beta(\lambda) = \prod_{l=1}^{d} (\lambda - \lambda_l)$. Then the linear map

$$
Hol(A) \to Hol(A), \ \ g \mapsto reg\left(\frac{g(\lambda)\alpha(\lambda)}{\beta(\lambda)}\right) \ \ is \ bounded \ in \ norm \ by \ 2\|\alpha/\beta\|_{\partial B}.
$$

Proof: The rational function $\alpha(\lambda)/\beta(\lambda)$ can be written as

$$
\frac{\alpha(\lambda)}{\beta(\lambda)} = \sum_{j=1}^d \frac{\gamma_j}{\prod_{l=j}^d (\lambda - \lambda_l)} \text{ with } \gamma_j = \frac{1}{2\pi\sqrt{-1}} \int_{\partial B} \frac{\alpha(\lambda)}{\beta(\lambda)} \prod_{l=j+1}^d (\lambda - \lambda_l) d\lambda.
$$

Here $|\gamma_j|$ is bounded by $\|\alpha/\beta\|_{\partial B} (R/3)^{d-j+1}$. The preceeding lemma now gives the estimate $||g\alpha/\beta|| \le ||g|| \, ||\alpha/\beta||_{\partial B} \sum_{j=1}^d (2/3)^{d-j+1} \le 2||g|| \, ||\alpha/\beta||_{\partial B}.$

Proof of the Deformation Lemma [5.7](#page-31-0): Let u_1, \ldots, u_n be the unique cross sections of \mathcal{O}_D , whose matrix of values at all covering points of infinity is equal to the identity matrix. For all $\tilde{f}_1,\ldots,\tilde{f}_n\in Hol(A)^n$ let $\tilde{c}_1,\ldots,\tilde{c}_n$ be the elementary symmetric functions with respect to the covering π of $\tilde{f} = \sum_{i=1}^n \tilde{f}_i u_i$. Due to the last lemma the map

$$
Hol(A)^n \to Hol(A)^n, \quad (\tilde{f}_1, \ldots, \tilde{f}_n) \mapsto (reg(\tilde{c}_1), \ldots, reg(\tilde{c}_n))
$$

is holomorphic. If $\tilde{f}_1, \ldots, \tilde{f}_n$ are all constant, they are mapped onto the ordinary elementary symmetric functions of these constants. But on the open subset of $Hol(A)^n$ with $|\tilde{f}_i - x_i| < 2\epsilon$, the map

 $(\tilde{f}_1, \ldots, \tilde{f}_n) \mapsto$ the usual elementary symmetric functions of $(\tilde{f}_1, \ldots, \tilde{f}_n)$

is invertible, if c is larger than 4ϵ . According to the last lemma the difference of these two maps is small. Hence we can apply the inverse function Theorem[[P-T\]](#page-91-0), and the above map is invertible for $|\tilde{f}_i - x_i| < 2\epsilon$. This proves the claim.

Now let D be any admissible divisor of Y. As we mentioned before, the surfaces $Y_{l,\epsilon,\iota}$ and the divisors $D_{l,\epsilon,\iota}$ fit into the situation described in the deformation Lemma whenever $|\iota|$ is large enough. In Example [3.9](#page-19-0) we introduced the supremum norm on the boundary of these excluded domains $\|\cdot\|_{l,\epsilon,\iota}$. Let $C_{\iota}(D, l, \epsilon) > 0$ be the smallest constant, such that

$$
\left\|\frac{\alpha_{i,\iota}(\cdot,x_1,\ldots,x_n)}{\beta_{\iota}(\cdot)}\right\|_{l,\epsilon,\iota} \leq C_{\iota}(D,l,\epsilon) \text{ for all } |x_1| \leq 1,\ldots, |x_n| \leq 1 \text{ and } i=1,\ldots,n.
$$

Proposition 5.10 Let D be an admissible divisor of Y. Then D is of Riemann-Roch type, if $\sum_{\iota} C_{\iota}(D, l, \epsilon) < \infty$ for all $l \in \mathbb{N}$ with some $\epsilon > 0$ depending on l.

Proof: Let x_1, \ldots, x_n be complex numbers, such that $|x_i - x_j| > c$. First we show that there exists a unique cross section of \mathcal{O}_D over the inverse image under π of some neighbourhood of infinity on \mathbb{P}_1 , which takes the values x_1, \ldots, x_n at all covering points of infinity. For the elementary symmetric functions of this cross section we make the ansatz:

$$
\tilde{c}_i(\lambda, x_1, \dots, x_n) = e_i(x_1, \dots, x_n) + \sum_{\iota \in \mathcal{I}'} \frac{\alpha_{i,\iota}(\lambda)}{\beta_{\iota}(\lambda)}
$$
 (compare with (12) and Example 3.9).

The sum contains only those indices, which correspond to excluded domains over the neighbourhood of infinity on \mathbb{P}_1 . We choose two neighbourhoods $U_{l,\delta}$, and $U_{\tilde{l},\tilde{\delta}}$, such that for each index $\iota \in \mathcal{I}'$ in the deformation Lemma the closure of U can be chosen to be equal to the image under π of the excluded domain with index ι of $U_{l,\delta}$. Also we choose B to be equal to the image under π of the excluded domain with index ι of $U_{\tilde{i},\tilde{\delta}}$. Taking $l > l$ for granted, this can be always attained by reducing the neighbourhood of infinity on \mathbb{P}_1 and thereby the corresponding set of indices \mathcal{I}' of excluded domains over this neighbourhood. Now we introduce the Banach space of sequences of polynomials $\alpha_{i,\iota}(\lambda)$ indexed by $i = 1, \ldots, n$ and $\iota \in \mathcal{I}'$, such that the degree of $\alpha_{i,t}$ is less than the degree d_t of $\beta_t(\lambda)$ from equation [\(12](#page-31-0)). The norm is given by

$$
\|(\alpha_{i,\iota}(\cdot))\| = \sum_{\iota \in \mathcal{I}'} \sup \left\{ \left\|\frac{\alpha_{i,\iota}}{\beta_{\iota}}\right\|_{l,\delta,\iota} \right| i = 1,\ldots,n \right\}.
$$

With the help of the deformation Lemma we can now define a map on an open subset of this Banach space into this Banach space:

$$
\left\{(\alpha_{i,\iota})_{(i,\iota)\in\{1,\ldots,n\}\times\mathcal{I}'}\mid\left\|\left(\alpha_{i,\iota}\right)\right\|\leq\epsilon\right\}\to\left\{(\alpha_{i,\iota})_{(i,\iota)\in\{1,\ldots,n\}\times\mathcal{I}'}\right\},\ \ (\alpha_{i,\iota})\mapsto(\tilde{\alpha}_{i,\iota}),
$$

such that for all $\kappa \in \mathcal{I}'$ the section \tilde{f}_{κ} of \mathcal{O}_D over the excluded domain with index κ , defined by the elementary symmetric functions $\tilde{c}_{j,\kappa} = c_{j,\kappa} + \tilde{\alpha}_{j,\kappa}/\beta_{\kappa}$ is the deformation of the section f_{κ} of $\mathcal{O}_{\mathbb{P}_1}^n$ over the same domain defined by the elementary symmetric functions

$$
c_{j,\kappa}=e_j(x_1,\ldots,x_n)+\sum_{\iota\in\mathcal{I}'\backslash\{\kappa\}}\frac{\alpha_{j,\iota}}{\beta_{\iota}}.
$$

Now Lemma [5.9](#page-32-0) gives the estimate

$$
\left\|\frac{\tilde{\alpha}_{j,\kappa}}{\beta_{\kappa}}\right\|_{l,\delta} \leq C_{\kappa}(D,\tilde{l},\tilde{\delta}) \cdot \text{constant}
$$

The constant depends only on c and x_1, \ldots, x_n . The assumption on the divisor implies that $\sum_{\kappa \in \mathcal{I}'} C_{\kappa}(D, \tilde{l}, \tilde{\delta}) < \infty$. Hence the image of this map is contained in the domain, if the neighbourhood of infinity and the corresponding set of indices \mathcal{I}' is small enough. The same argument for arbitrary $l \in \mathbb{N}$ shows that the elementary symmetric functions $\tilde{c}_{i,\kappa}$ are meromorphic functions of the form given in Example [3.9](#page-19-0). Again Lemma [5.9](#page-32-0) shows that

$$
\left\|\frac{\tilde{\alpha}_{j,\kappa}}{\beta_{\kappa}} - \frac{\tilde{\alpha}'_{j,\kappa}}{\beta_{\kappa}}\right\|_{l,\delta} \le \left\|(\alpha_{i,\iota}) - (\alpha'_{i,\iota})\right\| C_{\kappa}(D,\tilde{l},\tilde{\delta}) \cdot \text{constant}
$$

Again the constant depends only on c and x_1, \ldots, x_n . This map is a contraction, if the neighbourhood of infinity and the corresponding set of indices \mathcal{I}' is small enough. Due to Picard's fix point principle it has a unique fixed point. For such a fixed point the elementary symmetric functions $\tilde{c}_{j,k}$ do not depend on κ . Therefore they define a section of \mathcal{O}_D over some neighbourhood of infinity of the desired form. In the proof of the deformation lemma we showed that near every excluded domain this cross section can be written as $\tilde{f} = \sum_{i=1}^n \tilde{f}_{i,\kappa} u_{i,\kappa}$, where $u_{1,\kappa}, \ldots, u_{n,\kappa}$ defines an isomorphism of sheaves $\mathcal{O}_{\mathbb{P}_1}^n \simeq \pi_*(\mathcal{O}_D)$ near this excluded domain and $|\tilde{f}_i - x_i| < 2\epsilon$. Hence we can choose the values x_1, \ldots, x_n for n different cross sections u_1, \ldots, u_n of \mathcal{O}_D over some neighbourhood of infinity in a way, such that condition (ii) of Definition [5.1](#page-27-0) is satisfied. is satisfied. \Box

Proof of the implication (ii) \Rightarrow (i) of Theorem [5.5:](#page-29-0) It is quite obvious that the condition (ii) is equivalent to the assumptions of Proposition [5.10.](#page-32-0) This concludes the proof of Theorem [5.5](#page-29-0). \Box

In the case that $n = 2$ and that D is an integral divisor, the elementary symmetric functions inequation ([11](#page-31-0)) of the form given in equation [12](#page-31-0) can be calculated explicitly: Let λ_1 and λ_2 be the two base points of the two branchpoints. Furthermore let the divisor be specified by its base point λ_0 and by a choice of the branch $\sqrt{(\lambda_0 - \lambda_1)(\lambda_0 - \lambda_2)}$. Let f be the unique cross section, which takes the values x_1 and x_2 at the covering points of infinity. It is given by

$$
f^{2} - \left(x_{1} + x_{2} + \frac{x_{1} - x_{2}}{\lambda - \lambda_{0}} \sqrt{(\lambda_{0} - \lambda_{1})(\lambda_{0} - \lambda_{2})}\right) f + x_{1}x_{2} - \frac{x_{1} - x_{2}}{\lambda - \lambda_{0}} \left(\frac{x_{1} - x_{2}}{4} (2\lambda_{0} - \lambda_{1} - \lambda_{2}) + \frac{x_{1} + x_{2}}{2} \sqrt{(\lambda_{0} - \lambda_{1})(\lambda_{0} - \lambda_{2})}\right) = 0.
$$
 (13)

Now it is easy to see that the condition of Proposition [5.10](#page-32-0) is fulfilled in this case. For excluded domains, which contain only two branchpoints, the compact n-fold covering $\pi : Y_{\iota} \to \mathbb{P}_1$ decompose into one two-fold covering and $n-2$ copies of \mathbb{P}_1 . Hence we can estimate $C_i(D, l, \epsilon)$ with the help of equation (13) even in this case.

Corollary 5.11 If all the excluded domains corresponding to one $U_{l,\epsilon}$ have asymptotically no overlap, then all integral divisors of asymptotic degree $(1,1,\ldots)$ with support inside of $\pi^{-1}(\mathbb{C})$ are of Riemann-Roch type.

 \Box

The case that all domains excluded from $U_{l,\epsilon}$ have asymptotically no overlap is the generic case. Otherwise the Riemann surfaces Y_t are more complicated and may have divisors of degree equal to half the branching order, such that the corresponding direct image sheaf $\pi_*(\mathcal{O}_D)$ is not isomorphic to $\mathcal{O}_{\mathbb{P}_{\infty}}^n$.

Formula (13) shows that the coefficients of the singular parts of the elementary symmetric functions are bounded by $4 \sup\{|\lambda_0 - \lambda_1|, |\lambda_0 - \lambda_2|\}\.$ This observation was crucial in the last proof. These coefficients are bounded on the domain, where all base points of the divisor and the branchpoints are elements of a bounded subset of \mathbb{C} , iff $\pi_*(\mathcal{O}_D) \simeq \mathcal{O}_{\mathbb{P}_1}^n$ were to hold also for singular surfaces Y. But this is no more valid even in the case that $n = 2$ and that the support of D contains more than one point: If Y is the singular Riemann surface of two copies of \mathbb{P}_1 connected by one double point, and if the divisor consists of two points out of one copy of \mathbb{P}_1 with multiplicity 1 and one point out of the other copy of \mathbb{P}_1 with multiplicity -1 ,

 $\pi_*(\mathcal{O}_D)$ is not isomorphic to $\mathcal{O}_{\mathbb{P}_1}^n$. There exist similar counterexamples for $n > 2$ and D integral. These counterexamples suggest a more restrictive assumption on the location of the divisor: The statement $\pi_*(\mathcal{O}_D) \simeq \mathcal{O}_{\mathbb{P}_1}^n$ remains true, if this assumption guarantees that in the limit of singular Riemann surfaces some of the points of the divisors are deformed into the singular points. Then it can be proven that the divisors are of Riemann-Roch type.

6 The Jacobian variety

In this section we want to investigate the set of all equivalence classes of line bundles, which submit to the Riemann-Roch Theorem. The following definition is suggested from the previous section.

Definition 6.1 Let the Jacobian variety Jacobian(Y) of Y be the set of all equivalence classes in the modified sense of integral divisors of asymptotic and total degree $(1, 1, \ldots)$ in the sense of Remark [3.7](#page-17-0), which are of Riemann-Roch type.

For every divisor D of asymptotic and total degree $(1, 1, \ldots)$, which is of Riemann-Roch type there exists at least one cross section f of \mathcal{O}_D , which takes the value 1 at all covering points of infinity. Hence all such divisors are equivalent in the modified sense to some integral divisor. Moreover, if there exists only one section f of \mathcal{O}_D , which is equal to 1 at all covering points of infinity, D is equivalent in the modified sense to exactly one integral divisor. Such divisors are called non-special in the modified sense.

Lemma 6.2 For all representatives D of elements of Jacobian(Y) the following statements are equivalent:

- (i) $[D] \in Jacobian(Y)$ is non-special in the modified sense.
- (ii) $\tilde{H}^1(Y, \mathcal{O}_{D-\pi^{-1}(\infty)})=0.$
- (iii) $\pi_*({\cal O}_D)\simeq {\cal O}_{{\Bbb P}_1}^n$.

Proof: (i) \Leftrightarrow (ii): The Riemann-Roch Theorem implies

$$
\dim \tilde{H}^0(Y, \mathcal{O}_{D-\pi^{-1}(\infty)}) = \dim \tilde{H}^1(Y, \mathcal{O}_{D-\pi^{-1}(\infty)}).
$$

For all elements f of $\tilde{H}^0(Y, \mathcal{O}_{D-\pi^{-1}(\infty)})$, $1+f$ is a cross section of \mathcal{O}_D , which is equal to 1 at all covering points of infinity and all these cross sections are of the form $1 + f$, with $f \in H^0(N, \mathcal{O}_{D-\pi^{-1}(\infty)})$. Hence (i) and (ii) are equivalent.

(ii) \Leftrightarrow (iii): $\pi_*(\mathcal{O}_D)$ is isomorphic to $\mathcal{O}_{D_1} \oplus \ldots \oplus \mathcal{O}_{D_n}$, with some divisors D_1, \ldots, D_n of \mathbb{P}_1^{23} such that $\deg(D_1) + \ldots + \deg(D_n) = 0$. Then we have $\dim \tilde{H}^1(N, \mathcal{O}_{D-\pi^{-1}(\infty)}) =$ $\sum_{i=1}^n \max\{0, \deg(D_i)\}\.$ Hence is (ii) equivalent to $\deg(D_i) = 0$ for all $i = 1, \ldots, n$, which is equivalent to (iii). \square

Theorem [5.5](#page-29-0) suggests to endow $Jacobian(Y)$ with the topology induced by the infinite sums $\sum_{\iota} ||h_{l,\epsilon,\iota}(D,\cdot) - h_{l,\epsilon,\iota}(D',\cdot)||_{l,\epsilon,\iota}$. More precisely, only for large $|\iota|, h_{l,\epsilon,\iota}(D,\cdot)$ is well defined.

 $\overline{a^2}$ Thisshows the Birkhoff factorization (see e.g. [[P-S](#page-91-0), Proposition (8.11.5)]).
Hence these sums define semidefinite metrics on the set of all representatives of elements of Jacobian(Y). Each representative D decomposes into a finite collection of points of Y and sequences of divisors in the excluded domains. For the finite collection of points we use the natural topology induced by the topology of Y . For the sequences of divisors we use the topology defined by all the sums $\sum_{\iota} ||h_{l,\epsilon,\iota}(D,\cdot) - h_{l,\epsilon,\iota}(D',\cdot)||_{l,\epsilon,\iota}$ with $l \in \mathbb{N}$ and some ϵ depending on l. These together define a topology on the set of all representatives of $Jacobian(Y)$. Finally the topology of $Jacobian(Y)$ is given by the quotient topology, which is the finest topology such that the map, which maps the representatives of elements of $Jacobian(Y) \rightarrow$ elements of $Jacobian(Y)$, is continuous.

Lemma 6.3 For all integral divisors D of asymptotic degree $(1,1,\cdot)$ the following statements are equivalent:

(i) $\sum_{\iota} ||h_{l,\epsilon,\iota}(D,\cdot) - h_0||_{l,\epsilon,\iota} < \infty$ for all $l \in \mathbb{N}$ with some ϵ depending on l.

(ii) $\sum_{\iota} ||h_{l,\epsilon,\iota}(D,\cdot) - h_0||_{l,\epsilon,\iota}|_{\iota}|^k < \infty$ for any fixed l and ϵ and all $k \in \mathbb{N}$.

Proof: All entries of $h_{l,\epsilon,\iota}(D)$ are meromorphic functions on the compact n-fold covering π : $Y_{l,\epsilon,\iota} \to \mathbb{P}_1$ and at the covering points of infinity equal to the corresponding entries of h_0 . We willuse the elementary symmetric functions c_1, \ldots, c_n defined by ([11\)](#page-31-0) and of the form [\(12\)](#page-31-0). In our case the degree d of $\beta(\lambda)$ is at most $n(n-1)/2$. Now we need another lemma.

Lemma 6.4 Let $B \subset \mathbb{C} \subset \mathbb{P}_1$ be a ball containing the points $\lambda_1, \ldots, \lambda_d$, which furthermore is contained in the intersection of all balls of radius r around the points $\lambda_1, \ldots, \lambda_d$. Moreover let $A \subset \mathbb{C} \subset \mathbb{P}_1$ be a ball containing the balls with radius R around the points $\lambda_1, \ldots, \lambda_d$. Let $\|\cdot\|_{\partial B}$ and $\|\cdot\|_{\partial A}$ be the supremum norm on the boundary of B and A respectively. Finally let $\alpha(\lambda)$ be a polynomial of degree less than the degree of $\beta(\lambda) = \prod_{i=1}^{d} (\lambda - \lambda_i)$. Then we have the estimate

$$
\left\| \frac{\alpha}{\beta} \right\|_{\partial A} \le \left\| \frac{\alpha}{\beta} \right\|_{\partial B} \frac{1 - (r/R)^d}{1 - (r/R)} \frac{r}{R}
$$

.

The proof uses the same methods as the proof of Lemma [5.9](#page-32-0) \Box Conclusion of the proof of Lemma 6.3: The foregoing lemma gives an estimate

$$
C' \sum_{\iota} ||h_{\iota}(D, \cdot) - h_0||_{l', \epsilon', \iota} \leq \sum_{\iota} ||h_{\iota}(D, \cdot) - h_0||_{l, \epsilon, \iota} ||\iota||^k \leq C'' \sum_{\iota} ||h_{\iota}(D, \cdot) - h_0||_{l+k, \epsilon'', \iota}
$$

with some constants C', l', ϵ' and C'', ϵ'' depending only on l, k, ϵ and p . This lemma shows that we can use as well the sequences from (ii)in the same fashion as we used the sequences from (i) to define the topology of $Jacobian(Y)$.

Now let Isospectral(Y) be the subspace of the Fréchet space \mathcal{H}^{∞} of smooth periodic potentials \tilde{q} , such that det $(\mu \mathbf{1} - F(\lambda, \tilde{q})\mathbf{1}) = R(\lambda, \mu) = \det (\mu \mathbf{1} - F(\lambda, q))$. In Section [4](#page-19-0) we defined a map:

 $D(\cdot)$: Isospectral(Y) \rightarrow Jacobian(Y), $\tilde{q} \mapsto [D(\tilde{q})]$.

We were able to lift the shift to a flow on the image of $D : (x, [D(\tilde{q})]) \mapsto [D(T_x \tilde{q})]$. This lifting of the shift was defined by the property that the action on the corresponding line bundles is given by the tensor product with $L(x)$. In order to extend this action to the whole of *Jacobian(Y)*, let us describe the sheaves $\pi_*(\mathcal{O}_D)$ for all $[D] \in Jacobian(Y)$ by cocycles:

Let $\mathcal U$ be an open covering of X of the form

$$
\mathcal{U} = \{U\} \cup \{U_{\iota} | \iota \in \mathcal{I}\}, \text{ such that}
$$

Covering (i) U is an open neighbourhood of infinity and $\pi^{-1}(U)$ is an unbranched n-fold covering of U.

Covering (ii) U_i contains the domain excluded from U with index ι .

Covering (iii) $U_{\iota} \cap U_{\iota'} = \emptyset$ if $\iota \neq \iota'.$

Then we define the set of cocycles $C^1(\mathcal{U}, \pi_*(\mathcal{O}_Y^*))$ to be the set of holomorphic functions

 $g_{\iota}: U \cap U_{\iota} \to GL(n, \mathbb{C}), \lambda \mapsto g_{\iota}(\lambda),$ such that

Cocycle (i) for all $\iota \in \mathcal{I}$ there exists some holomorphic map

 $g_{+,i}: U_i \to GL(n,\mathbb{C}),$ such that $g_{+,i}g_i = g_i \operatorname{diagonal}(\mu_1, \ldots, \mu_n),$

where μ_i is the restriction of μ to the *i*-sheet over $U \cap U_i$.

- **Cocycle (ii)** For all $\iota \in \mathcal{I}$ det (g_{ι}) has trivial winding number around the excluded domain with index ι (the total residue of the form $\det^{-1}(g_{\iota})d \det(g_{\iota})$ on the set U_{ι} is zero).
- **Cocycle (iii)** There exists some $K > 0$, such that for all $|\iota| \geq K g_{\iota}$ extends to a holomorphic function on $U \cup X \setminus U_i$, which is equal to h_0 at $\lambda = \infty$.

Cocycle (iv) $\sum_{l} ||g_{l} - h_0||_{l,\epsilon,\iota} < \infty$ for all $l \in \mathbb{N}$ with some ϵ depending on l.

Moreover, we endow $C^1(\mathcal{U}, \pi_*(\mathcal{O}_Y^*))$ with the topology defined by the metrics

$$
d\left((g_{\iota})_{\iota\in\mathcal{I}},(\tilde{g}_{\iota})_{\iota\in\mathcal{I}}\right)=\sum_{\iota}||g_{\iota}-\tilde{g}_{\iota}||_{l,\epsilon,\iota}
$$

for all $l \in \mathbb{N}$ with some ϵ depending on l. Condition (i) ensures that the cocycles describe holomorphic vector bundles over X, which are $\pi_*(\mathcal{O})$ modules. Such vector bundles are the direct images of line bundles over Y . Due to condition (i) the asymptotic and total degrees of these line bundles are equal to $(1, 1, \ldots)$ in the sense of Remark [3.7](#page-17-0). Finally condition (iii) is analogous to the assumption (ii) on admissible divisors.

Proposition 6.5 The line bundles defined by these cocycles of $C^1(\mathcal{U}, \pi_*(\mathcal{O}_Y^*))$ are equivalent in the modified sense to some element of $Jacobian(Y)$. Moreover, the corresponding map $C^1(\mathcal{U}, \pi_*(\mathcal{O}_Y^*)) \to Jacobian(Y)$ is continuous.

Theorem 6.6 The action of the tensor product with $L(x)$ on holomorphic line bundles induces a continuous action of $\mathbb R$ on Jacobian(Y), which is denoted by

$$
\mathbb{R} \times Jacobian(Y) \to Jacobian(Y), (x, [D]) \mapsto T_x[D].
$$

of holomorphic functions

The proof²⁴ of this proposition is similar to the proof of Proposition [5.10](#page-32-0). First we need a lemma analogous to the deformation Lemma.

Lemma 6.7 Let g and \tilde{g} be analytic maps from the circle $\{\lambda \in \mathbb{C} \mid |\lambda - \lambda_0| = R\} \to GL(n, \mathbb{C}),$ such that $||g-1|| < 1$ and $||\tilde{g}-1|| < 1$, respectively. Here the norm denotes the supremum norm on the circle. Due to Lemma [5.6](#page-29-0) both elements of the Loop group admit a Birkhoff factorization $g = g_{+}g_{+}$ and $\tilde{g} = \tilde{g}_{-}\tilde{g}_{+}$, respectively. Then the following estimate holds:

$$
||g_{-} - \tilde{g}_{-}|| \le \frac{2||g - \tilde{g}||}{(1 - ||g - 1||)(1 - ||\tilde{g} - 1||)}.
$$

The proof uses the same arguments as the proof of Lemma [5.6.](#page-29-0) \Box Proof of Proposition [6.5](#page-37-0): Let $f = (f_1, \ldots, f_n)$ be a *n*-tuple of invertible holomorphic functions on U. These functions together define a section of the sheaf \mathcal{O}_Y^* over $\pi^{-1}(U)$, where the restriction of this section to the *i*-th sheet is given by f_i for all $i = 1, \ldots, n$. By abuse of notation this section is denoted by $f = (f_1, \ldots, f_n)$. Furthermore, let $(\tilde{g}_{+,i})_{i \in \mathcal{I}}$ be any sequence

$$
\tilde{g}_{+,\iota}: U_{\iota} \to GL(n,\mathbb{C}), \lambda \mapsto \tilde{g}_{+,\iota}(\lambda).
$$

Then the line bundles corresponding to two elements $(g_\iota)_{\iota \in \mathcal{I}}$ and $(\tilde{g}_\iota)_{\iota \in \mathcal{I}}$ of $C^1(\mathcal{U}, \pi_*(\mathcal{O}_Y^*))$ are equivalent in the modified sense, if

$$
\tilde{g}_{+,\iota}\tilde{g}_{\iota} = g_{\iota} \operatorname{diagonal}(f_1,\ldots,f_n) \text{ for all } \iota \in \mathcal{I}.
$$

Due to assumption (iii) on the cocycles for large $|\iota|$ the Birkhoff factorization around the open sets U_i determines \tilde{g}_i as a function depending only on g_i and $f = (f_1, \ldots, f_n)$:

$$
(h_0^{-1}\tilde{g}_{+,l}h_0)(h_0^{-1}\tilde{g}_l) = h_0^{-1}g_l \operatorname{diagonal}(f_1,\ldots,f_n).
$$

In order to prove the first statement of the proposition it suffices to show that the line bundles corresponding to the cocycles of $C^1(\mathcal{U}, \pi_*(\mathcal{O}_Y^*))$ are equivalent in the modified sense to the line bundles corresponding to some integral divisor D of asymptotic degree $(1, 1, \ldots)$ over some open neighbourhood $U_-\$ of $\lambda = \infty$ of \mathbb{P}_1 . Now we claim that to each cocycle $(g_i)_{i\in\mathcal{I}}$ there exists an invertible holomorphic function $f = (f_1, \ldots, f_n)$ on $\pi^{-1}(U)$, which is equal to 1 at all covering points of infinity, and an element $[D]$ of $Jacobian(Y)$, such that the Birkhoff factorization of $h_0^{-1}g_\iota diagonal(f_1,\ldots,f_n)$ is given by $\tilde{g}_{+,l}h_0^{-1}h_\iota(D)$ whenever $|\iota|$ is large enough. The function $h_{\iota}(D)$ was defined ahead of Theorem [5.5](#page-29-0). For this purpose we introduce the Banach space of sequences $(f_{\iota})_{\iota \in \mathcal{I}'} = (f_{1,\iota}, \ldots, f_{n,\iota})_{\iota \in \mathcal{I}'}$ of holomorphic functions

$$
f_{\iota} : \pi^{-1} (U \cup (X \setminus U_{\iota})) \to \mathbb{C},
$$

which are equal to 0 at all covering points of infinity, and the norm

$$
||(f_{\iota})_{\iota \in \mathcal{I}'}|| = \sum_{\iota \in \mathcal{I}'} ||f_{\iota}||_{l, \epsilon, \iota} < \infty \text{ with some fixed } l \text{ and } \epsilon.
$$

The set \mathcal{I}' contains the indices corresponding to all excluded domains over some neighbourhood of infinity of \mathbb{P}_1 . For all elements $(f_\iota)_{\iota \in \mathcal{I}'}$ of this Banach space, the infinite product $\prod_{\iota}(1+f_\iota)$

²⁴This proof may be passed over to Theorem [6.9.](#page-40-0)

defines a holomorphic function on $U \cap U_{l,\epsilon}$. With the help of the Birkhoff factorization we define a map of an open subset of this Banach space into the Banach space

$$
\{(f_{\iota})_{\iota \in \mathcal{I}'} || |(f_{\iota})_{\iota \in \mathcal{I}'} || < \epsilon\} \to \{(f_{\iota})_{\iota \in \mathcal{I}'} , (f_{\iota})_{\iota \in \mathcal{I}'} \mapsto (\tilde{f}_{\iota})_{\iota \in \mathcal{I}'} :
$$

Let $\tilde{g}_{+, \iota}(h_0^{-1}\tilde{g}_{\iota}) = h_0^{-1}g_{\iota}$ diagonal $\left(\prod_{\iota' \neq \iota} (1 + f_{1, \iota'}), \dots, \prod_{\iota' \neq \iota} (1 + f_{n, \iota'})\right).$

be the Birkhoff factorization around the set U_{ι} . Then \tilde{f}_{ι} is defined as

$$
\tilde{f}_{\iota} = (\tilde{f}_{1,\iota}, \dots, \tilde{f}_{n,\iota}) = \left(\frac{1}{(\tilde{g}_{\iota})_{1,1}} - 1, \dots, \frac{1}{(\tilde{g}_{\iota})_{1,n}} - 1\right)
$$

for all $\iota \in \mathcal{I}'$. Now the foregoing lemma shows that the image of this map is contained in its domain, if \mathcal{I}' is chosen small enough, and furthermore, that this map is a contraction. Hence it has an unique fixed point $(f_{fix, l})_{l\in\mathcal{I}'}$. Then there exists an integral divisor D of asymptotic degree $(1, 1, \ldots)$, such that for all $\iota \in \mathcal{I}'$ $h_{\iota}(D)$ solves the following Birkhoff factorization:

$$
\tilde{g}_{+,l}h_0^{-1}h_l(D) = g_l \prod_{\iota \in \mathcal{I}'} (1 + f_{\text{fix},\iota}).
$$

In fact, the solution $h_{\iota}(D)$ of this Birkhoff factorization diagonalizes some $\tilde{g}_{+,\iota}$ and furthermore, the first row of this matrix valued holomorphic function is equal to the first row of h_0 . Then the i-th column of this function is equal to the restriction of a meromorphic vector valued function $v(D)$ with $v(D)_1 = 1$ to the *i*- sheet of the Riemann surface Y_i over $\mathbb{P}_1 \setminus U_i$. This proves the claim. In order to prove the second statement of the proposition it suffices to show that the fixed point of the map above depends continuously on the cocycle $(g_{\iota})_{\iota \in \mathcal{I}}$. This is a consequence of the estimate of the foregoing lemma. of the estimate of the foregoing lemma.

For the proof of Theorem [6.6](#page-37-0) we again need a lemma:

Lemma 6.8 There exists some $K \geq 0$, such that for all $|\iota| \geq K$, $\ln(\mu)$ defines a holomorphic function on the excluded domain with index ι.

Proof: Obviously $\ln(\mu)$ defines a multi valued holomorphic function on $\pi^{-1}(\mathbb{C})$. In Theorem [2.6](#page-10-0) it was shown that for all $\epsilon > 0$ there exists some $c > 0$ such that

$$
|\mu_i - \exp(p_i \lambda)| < \frac{c}{|\lambda|} \exp(p_i \lambda)|
$$

for all $\lambda \in U_{0,\epsilon}$ and $i = 1, \ldots, n$. This implies the estimate

$$
\left|\ln(\mu_i) - (p_i \lambda + 2n_i \pi \sqrt{-1})\right| < \frac{c'}{|\lambda|}
$$

with some constant $c' > 0$ and some integers $n_1, \ldots, n_n \in \mathbb{Z}$. Now let us extend one branch corresponding to the integers n_i and n_j of this function from the boundary of the excluded domain with index $\iota = (i, j, k) \in \mathcal{I}$ to the interior of this excluded domain. Then the foregoing estimates on the i -th and j -th sheet imply the following condition:

$$
|n_i - n_j - k| \le \frac{2}{\epsilon} + \frac{2c'}{|\lambda|}.
$$

If |i| is large enough this implies $n_i - n_j = k$. Due to this condition $\ln(\mu)$ extends to a single valued holomorphic function on the excluded domain with index ι . valued holomorphic function on the excluded domain with index ι .

Proof of Theorem [6.6](#page-37-0): The foregoing lemma shows that the Birkhoff factorization of

$$
h_{l,\epsilon,\iota}(D,\cdot) \, diagonal(\mu_1^x,\ldots,\mu_n^x)
$$

is given for large $|\iota|$ by

$$
g_{+}(\cdot)Ad(\exp(2\pi x \sqrt{-1} \text{diagonal}(n_1,\ldots,n_n)))(h_0^{-1}h_{l,\epsilon,\iota}(D,\cdot)),
$$

with some holomorphic

 g_+ : excluded domain with index $\iota \to GL(n,\mathbb{C}), \lambda \mapsto g_+(\lambda)$

and $n_i - n_j = k$. Then Proposition [6.5](#page-37-0) proves the claim. \Box

It is quite obvious that the subset of $Jacobian(Y)$ of all equivalence classes of divisors, which are non-special in the modified sense, is an open subset of $Jacobian(Y)$. All equivalence classes of this subspace have only one integral representative. Moreover, the topology of this subspace is given directly by the topology of the set of representatives of the elements of $Jacobian(Y)$. Now let $Jacobian_0(Y) \subset Jacobian(Y)$ be the subspace $Jacobian_0(Y) =$

$$
= \{ [D] \in Jacobian(Y) | T_x[D] \text{ is non-special in the modified sense for all } x \in \mathbb{R} \}.
$$

This subset is open, too. In fact, let $[D]$ be any element of *Jacobian₀* (Y) . Due to the last theorem for all $x \in [0,1]$ there exists an open neighbourhood U_x of $[D]$ and an open interval $(x - \epsilon_x, x + \epsilon_x)$, such that for all $(x', [D']) \in (x - \epsilon_x, x + \epsilon_x) \times U_x$ the divisor $T_{x'}[D']$ is non-special in the modified sense. The open covering $\{(x - \epsilon_x, x + \epsilon_x) | x \in [0, 1]\}$ of the compact interval $[0, 1]$ has a finite subcovering. Then the intersection of the corresponding open neighbourhoods U_x is an open neighbourhood of D contained in $Jacobian_0(Y)$. Hence this subset is open.

Theorem 6.9 The map $[D(\cdot)]$: Isospectral(Y) \rightarrow Jacobian(Y) induces a homeomorphism between Isospectral(Y) and Jacobian₀ (Y) .

Proof²⁵: Theorem [4.6](#page-23-0) and Lemma [6.2](#page-35-0) shows that $D(q)$ is non-special in the modified sense for all $q \in Isospectral(Y)$ and Theorem [4.10](#page-26-0) implies the relation

$$
[D(T_xq)] = T_x[D(q)].
$$

Hence the map

$$
D(\cdot): Isospectral(Y) \to Divisors(Y), q \mapsto D(q)
$$

induces another map denoted by

$$
[D(\cdot)]: Isospectral(Y) \to Jacobian_0(Y), q \mapsto [D(q)].
$$

Due to Lemma [6.2](#page-35-0) there exists a map

$$
\mathbb{R} \times Jacobian_0(Y) \to vector \; valued \; functions \; on \; Y, (x, [D] \mapsto v(x, [D]) = \begin{pmatrix} v_1(x, [D]) \\ \vdots \\ v_n(x, [D]) \end{pmatrix},
$$

²⁵The rest of this section may be passed over. It contains the proof of this theorem.

such that

$$
\phi_{v(x,[D])}: \mathcal{O}_{\mathbb{P}_1}^n \to \pi_*(\mathcal{O}_{T_xD}), (f_1,\ldots,f_n) \mapsto \sum_i f_i v_i(x,[D])
$$

is an isomorphism and $v(x, [D])$ takes the same values as the columns of h_0 at the covering points of infinity. The proof of Proposition [5.10](#page-32-0) and the preceeding theorem show that all Taylor coefficients of $v(x, [D])$ at the covering points of infinity are continuous functions on $\mathbb{R} \times Jacobian_0(Y)$, which are periodic in x with period 1. Hence there exists a unique continuous and in x periodic potential

$$
q: \mathbb{R} \times Jacobian_0(Y) \to n \times n\text{-matrices}, (x, [D]) \mapsto q(x, [D]), \text{ such that}
$$

$$
\pi^*(h_0(p\lambda + q(x, [D]))h_0^{-1})v(x, [D]) - v(x, [D])\ln(\mu)
$$

is holomorphic on the neighbourhood of Lemma [4.7](#page-23-0) and at all covering points of infinity equal to zero. The diagonal part of $q(x, [D])$ does not depend on x and $[D]$. Now let $f(x, [D])$ be the first entry of this function:

$$
f(x, [D]) = (\pi^*(h_0(p\lambda + q(x, [D]))h_0^{-1})v(x, [D]) - v(x, [D])\ln(\mu))_1.
$$

Then the function

$$
\pi^*(h_0(p\lambda + q(x,[D]))h_0^{-1})v(x,[D]) - v(x,[D])(\ln(\mu) - f(x,[D]))
$$

extends to a meromorphic function on Y . Indeed, the function given above is a multi valued meromorphic function on Y. Then by definition of f this function is single valued and the first component is equal to zero. Now we claim that $v(x, [D])$ is differentiable with respect to x and that the derivative is equal to

$$
\frac{\partial v(x,[D])}{\partial x} = \pi^*(h_0(p\lambda + q(x,[D]))h_0^{-1})v(x,[D]) - v(x,[D])(\ln(\mu) - f(x,[D])).
$$

One way to prove this claim is to improve Proposition [6.5](#page-37-0) and Proposition [5.10](#page-32-0) and to show directly that this function is differentiable with respect to x (see footnote [27](#page-42-0)). But the proofs of these two propositions are rather extensive. There is another less tedious way to prove this claim: We define for all $x \in \mathbb{R}$ and all $\lambda \in U \setminus \{\infty\}$ with some neighbourhood U of infinity in X the function $g(x, \lambda) \in GL(n, \mathbb{C})$ such that

$$
\pi^*(g(x,\cdot))v(0,[D]) = v(0,[D])\mu^{-x}.
$$

Due to Proposition [6.5](#page-37-0) this function has a decomposition $g(x, \cdot) = g_+(x, \cdot)g_-(x, \cdot)$ into a continuous function

$$
g_+ : \mathbb{R} \times \mathbb{C} \to GL(n, \mathbb{C}), (x, \lambda) \mapsto g_+(x, \lambda),
$$

which is holomorphic with respect to λ , times a continuous function

$$
g_-: \mathbb{R} \times U \to GL(n, \mathbb{C}), (x, \lambda) \mapsto g_-(x, \lambda),
$$

which is again holomorphic with respect to λ and equal to 1 for $\lambda = \infty$, such that²⁶

$$
v(x,[D]) = \frac{\pi^*(g_-(x,\cdot))v(0,[D])}{(\pi^*(g_-(x,\cdot))v(0,[D]))_1}.
$$

²⁶The function $(\pi^*(g_-(x,\cdot))v(0,[D]))_1$ corresponds to the function $\prod_{\iota}(1+f_{\iota})$ of Proposition [6.5.](#page-37-0)

Moreover, define the $n \times n$ -matrix valued function

$$
a : \mathbb{R} \times U \to n \times n
$$
-matrices, $(x, \lambda) \mapsto a(x, \lambda)$

such that

$$
a(x, \cdot)v(x, [D]) = v(x[D])\ln(\mu).
$$

Then we saw above that this function has a decomposition

$$
a(x, \cdot) = a_{+}(x, \cdot) + a_{-}(x, \cdot), \text{ with}
$$

$$
a_{+}(x, \lambda) = h_{0}(p\lambda + q(x, [D]))h_{0}^{-1} \text{ and}
$$

$$
^{*}(a_{-}(x, \lambda))v(x, [D]) = \pi^{*}(h_{0}(p\lambda + q(x, [D]))h_{0}^{-1})v(x, [D]) - v(x, [D])\ln(\mu).
$$

Then all these functions satisfy the integral equation

$$
\int_0^x g_+(t,\lambda)a(t,\lambda)g_-(t,\lambda)dt = g_+(x,\lambda)g_-(x,\lambda) - g_+(0,\lambda)g_-(0,\lambda),
$$

or more generally²⁷

 π

$$
\int_{x}^{y} g_{+}^{-1}(x,\lambda)g_{+}(t,\lambda)a(t,\lambda)g_{-}(t,\lambda)g_{-}^{-1}(y,\lambda)dt = g_{+}^{-1}(x,\lambda)g_{+}(y,\lambda) - g_{-}(x,\lambda)g_{-}^{-1}(y,\lambda).
$$
 (14)

Now we need a

Lemma 6.10 Due to the integral equation (14), $g_{+}(\cdot, \lambda)$ and $g_{-}(\cdot, \lambda)$ obey the integral equations

$$
\int_0^x a_+(t,\lambda)g_+^{-1}(t,\lambda)dt = \mathbf{1} - g_+^{-1}(x,\lambda) \text{ and}
$$

$$
\int_0^x a_-(t,\lambda)g_-(t,\lambda)dt = g_-(x,\lambda) - \mathbf{1} \text{, respectively.}
$$

Proof: We define $G_{+,l}(x, y, \lambda) = g_+^{-1}(x, \lambda)$.

$$
\int_{x \leq t_1 \leq ... \leq t_l \leq y} Ad(g_+(t_1,\lambda))(a_+(t_1,\lambda)) \dots Ad(g_+(t_l,\lambda))(a_+(t_l,\lambda)) dt_1 \dots dt_l \cdot g_+(y,\lambda)
$$

and $G_{-,l}(x, y, \lambda) = g_{-}(x, \lambda)$.

$$
\int_{x \le t_1 \le ... \le t_l \le y} Ad(g_-^{-1}(t_1,\lambda))(a_-(t_1,\lambda)) \dots Ad(g_-^{-1}(t_l,\lambda))(a_-(t_l,\lambda)) dt_1 \dots dt_l \cdot g_-^{-1}(y,\lambda).
$$

Now we claim that for all $L \in \mathbb{N}_0$ the following equation holds:

$$
g_+^{-1}(x,\lambda)g_+(y,\lambda) - g_-(x,\lambda)g_-^{-1}(y,\lambda) =
$$

²⁷ It is easy to see that the left hand side is differentiable with respect to x at the point $x = y$ and that the derivative is equal to $a(x, \lambda)$. On the other side it is possible to improve Proposition [6.5](#page-37-0) and Proposition [5.10](#page-32-0) and to show directly that g_+ and g_- are differentiable with respect to x. Then Lemma 6.10 is an obvious consequence.

$$
= \sum_{l=1}^{L} G_{-,l}(x,y,\lambda) - \sum_{l=1}^{L} (-1)^{l} G_{+,l}(x,y,\lambda) + \sum_{l=0}^{L} (-1)^{l} \int_{x}^{y} G_{+,l}(x,t,\lambda) a(t) G_{-,L-l}(t,y,\lambda) dt.
$$

Infact, for $L = 0$ this is just the integral equation ([14\)](#page-42-0). Furthermore, this integral equation implies the equations:

$$
\sum_{l=0}^{L} (-1)^{l} \int_{x}^{y} G_{+,l}(x,t,\lambda) a(t) G_{-,L-l}(t,y,\lambda) dt =
$$
\n
$$
= \sum_{l=0}^{L+1} (-1)^{l} \int_{x}^{y} G_{+,l}(x,t,\lambda) a(t) G_{-,L+1-l}(t,y,\lambda) dt + G_{-,L+1}(x,y,\lambda) - (-1)^{L+1} G_{+,L+1}(x,y,\lambda)
$$

for all $L \in \mathbb{N}_0$. The inductive use of these equations proves the claim. The same arguments as in Section [2](#page-6-0) gives the bounds:

$$
||G_{+,l}(x, y, \lambda)|| \leq \frac{C}{l!}
$$
 and $||G_{-,l}(x, y, \lambda)|| \leq \frac{C}{l!}$

with some constant $C > 0$ depending on x, y, λ . Hence the sum

$$
\sum_{l=1}^{\infty} G_{-,l}(x, y, \lambda) - \sum_{l=1}^{\infty} (-1)^{l} G_{+,l}(x, y, \lambda)
$$

converges for any fixed x, y, λ . Moreover, the limit of the first sum is holomorphic with respect to λ for all $\lambda \in U \setminus \{\infty\}$ and equal to zero for $\lambda = \infty$. The second sum converges to an entire function with respect to λ . Then we have

$$
\sum_{l=1}^{\infty} G_{-,l}(x, y, \lambda) = 1 - g_{-}(x, \lambda) g_{-}^{-1}(y, \lambda)
$$
 for all $\lambda \in U$, and

$$
-\sum_{l=1}^{\infty} (-1)^{l} G_{+,l}(x, y, \lambda) = g_{+}^{-1}(x, \lambda) g_{+}(y, \lambda) - 1
$$
 for all $\lambda \in \mathbb{C}$, respectively.

Now let $\tilde{g}_{+}(\cdot,\lambda)$ and $\tilde{g}_{-}(\cdot,\lambda)$ be the unique solutions of the integral equations:

$$
\int_0^x \tilde{g}_+(t,\lambda) A d(g_+(t,\lambda)) (a_+(t,\lambda)) dt = 1 - \tilde{g}_+(x,\lambda)
$$
 and

$$
\int_0^x \tilde{g}_-(t,\lambda) A d(g_-^{-1}(t,\lambda)) (a_-(t,\lambda)) dt = \tilde{g}_-(x,\lambda) - 1
$$
, respectively.

These solutions can be given in terms of the infinite sums above:

$$
\tilde{g}_{+}(x,\lambda)g_{+}(x,\lambda) = \sum_{l=0}^{\infty} (-1)^{l} G_{+,l}(0,x,\lambda)
$$
 and

$$
\tilde{g}_{-}(x,\lambda)g_{-}^{-1}(x,\lambda) = \sum_{l=0}^{\infty} G_{+,l}(0,x,\lambda),
$$
 respectively.

We have seen above that both infinite sums on the right hand side converge to 1. Hence $\tilde{g}_{+}(x,\lambda)$ is equal to $g_{+}^{-1}(x,\lambda)$ and $\tilde{g}_{-}(x,\lambda)$ is equal to $g_{-}(x,\lambda)$. This proves the lemma. \Box

Conclusion of the proof of Theorem [6.9:](#page-40-0) Lemma [6.10](#page-42-0) now implies that $g_+(x, \lambda)$ and $g_-(x, \lambda)$ are differentiable with respect to x and that the derivatives are given by

$$
\frac{\partial g_+}{\partial x}(x,\lambda) = g_+(x,\lambda)a_+(x,\lambda) \text{ and } \frac{\partial g_-}{\partial x}(x,\lambda) = a_-(x,\lambda)g_-(x,\lambda), \text{ respectively.}
$$

Then we have

$$
(\pi^*(g_{-}(x,\cdot))v(0,[D]))_1 = \exp\left(\int_0^x f(t,[D])dt\right).
$$

This shows that

$$
\frac{\partial v(x,[D])}{\partial x} = \pi^*(h_0(p\lambda + q(x,[D]))h_0^{-1})v(x,[D]) - v(x,[D])(\ln(\mu) - f(x,[D])).
$$

By definition $q(x, [D])$ and $f(x, [D])$ are differentiable with respect to x up to the same order as $v(x, [D])$. Hence $v(x, [D])$, $f(x, [D])$ and $g(x, [D])$ are smooth functions with respect to x. Due to the differential equation for $g_+(x, \lambda)$ we have

$$
g_{+}(x,\lambda) = h_0 g^{-1}(x,\lambda,q(\cdot,[D]))h_0^{-1}.
$$

By definition of $g(x, \lambda)$ this implies

$$
\pi^*(F(\cdot, q(\cdot, [D])))v(0, [D]) = v(0, [D])\mu.
$$

This shows that $[D] \mapsto q(\cdot, [D])$ defines a map $Jacobian_0(Y) \rightarrow Isospectral(Y)$, such that the composition with $[D(\cdot)]$ is the identity map of *Jacobian*₀ (Y) . In Section [4](#page-19-0) we have already proven that the composition of D with this map is the identity map of $Isospectral(Y)$. All Taylor coefficients of $q(x, [D])$ with respect to x can be given in terms of the Taylor coefficients of $v(x,[D])$ with respect to λ^{-1} at the covering points of infinity (see Appendix [A\)](#page-81-0). Hence the map

$$
Jacobian_0(Y) \to Isospectral(Y), [D] \mapsto q(\cdot, [D])
$$

is continuous. Theorem [2.6](#page-10-0), the proof of the implication (i)⇒ (ii) of Theorem [5.5](#page-29-0) and Lemma [6.7](#page-38-0)
prove that the map $[D(\cdot)]$ is continuous, too. This completes the proof. \square prove that the map $[D(\cdot)]$ is continuous, too. This completes the proof. The map $q(x, [D])$ is defined even for all $(x, [D]) \in \mathbb{R} \times Jacobian(Y)$, such that $T_x[D]$ is nonspecial in the modified sense. This observation suggests that at least to all $[D] \in Jacobian(Y)$, such that $T_x[D]$ is non-special in the modified sense for all x in an open dense subset of [0, 1], there corresponds a potential with singularities. Moreover, it would be natural to establish a relation between the kind of singularity of the potential $q(\cdot, [D])$ at the point x and the index of speciality in the modified sense, which is equal to dim $\tilde{H}^1(Y, \mathcal{O}_{T_xD-\pi^{-1}(\infty)})$.

7 Darboux coordinates

Our next goal is to prove that the dynamical system of the potentials are completely integrable. In this section we take an excursion to certain Darboux coordinates.

Definition 7.1 Let $H_{\text{modified}}^0(Y, \Omega)$ be the vector space of all meromorphic differential forms, which have only poles of order at most 1 at all covering points of infinity.

Due to Theorem [3.5](#page-17-0) the sum over the residues at the covering points of infinity of all elements of $H_{\text{modified}}^0(Y, \Omega)$ is equal to zero. Hence $H_{\text{modified}}^0(Y, \Omega)$ is the space of all regular differential forms on the singular Riemann surface obtained by the identification of all covering points of infinity of Y to one multiple point (see e.g. $[Se]$).

For each $q \in \mathcal{H}^{\infty}$ let $T_q\mathcal{H}^{\infty}$ be the Fréchet space of tangent vectors δq at the point q. Due to Assumption [4.8](#page-24-0) the diagonal part of δq is equal to zero. Now let Ω_q be the map:

$$
\Omega_q: T_q\mathcal{H}^\infty \to H^0_{\text{modified}}(Y, \Omega), \delta q \mapsto \left(\frac{1}{\mu} \frac{d\mu}{dq}(\delta q)\right) d\lambda.
$$

Here we set $\frac{d\lambda}{dq} = 0$. Indeed, with this choice the expression $\frac{1}{\mu}$ $\frac{d\mu}{dq}(\delta q)$ becomes a holomorphic function on the complement of the union of all branchpoints and all covering points of infinity and due to Lemma [4.7](#page-23-0) a holomorphic function on some neighbourhood of all covering points of infinity. Moreover, it is easy to see that this function is a global section of \mathcal{O}_b and due to Assumption [4.8](#page-24-0) even a section of $\mathcal{O}_{b-\pi^{-1}(\infty)}$. This shows that $\Omega_q(\delta q)$ is an element of $H_{\text{modified}}^0(Y, \Omega)$. More generally, $\Omega_q(\delta q)$ is equal to $\left(\frac{1}{\mu}\right)$ μ $\frac{d\mu}{dq}(\delta q)\Big)\,d\lambda - \left(\frac{d\lambda}{dq}(\delta q)\right)\frac{d\mu}{\mu}$ $\frac{d\mu}{\mu}$ for arbitrary $\frac{d\lambda}{dq}$ and $\frac{d\mu}{dq}$. In fact, let $R(\lambda, \mu) = 0$ be the equation defining the Riemann surface corresponding to q. For fixed q we have

$$
\frac{\partial R}{\partial \lambda}d\lambda + \frac{\partial R}{\partial \mu}d\mu = 0
$$

and the derivatives with respect to δq obey

$$
\frac{\partial R}{\partial \lambda} \frac{d\lambda}{dq} + \frac{\partial R}{\partial \mu} \frac{d\mu}{dq} + \frac{\partial R}{\partial q} = 0.
$$

Combining these two equations we obtain

$$
\left(\frac{1}{\mu}\frac{d\mu}{dq}(\delta q)\right)d\lambda - \left(\frac{d\lambda}{dq}(\delta q)\right)\frac{d\mu}{\mu} = \left(\frac{1}{\mu}\frac{d\mu}{dq}(dq) - \frac{d\lambda}{dq}(\delta q)\frac{-\frac{\partial R}{\partial \lambda}}{\mu\frac{\partial R}{\partial \mu}}\right)d\lambda =
$$

$$
= -\frac{\frac{\partial R}{\partial q}(\delta q)}{\mu\frac{\partial R}{\partial \mu}}d\lambda = \frac{\frac{\partial R}{\partial q}(\delta q)}{\frac{\partial R}{\partial \lambda}}\frac{d\mu}{\mu}.
$$

As we mentioned before, with Assumption [4.8](#page-24-0) the space of potentials \mathcal{H}^{∞} forms a coadjoint orbit and therefore possesses a natural holomorphic symplectic structure. Let ω be the 2 form on \mathcal{H}^{∞} defined by

$$
\omega(\delta q, \delta \tilde{q}) = \sum_{i \neq j} \int_0^1 \frac{\delta q_{ij}(x) \delta \tilde{q}_{ji}(x)}{p_i - p_j} dx.
$$

With Assumption [4.8](#page-24-0) it is quite obvious that ω is a non-degenerate 2 form and extends to a holomorphic closed non-degenerate 2 form on the Hilbert space \mathcal{H}^{28}

Lemma 7.2 Let $q \in \mathcal{H}^{\infty}$ be a potential such that the corresponding Riemann surface satisfies the assumption $b_{\text{analytic}} = b_{\text{algebraic}}$. Now let $(\lambda_1, \mu_1) + \ldots + (\lambda_d, \mu_d)$ be the local part of the divisor D in some small open set of Y . After reducing this open set we can always attain $(\lambda_1,\mu_1)=\ldots=(\lambda_d,\mu_d)$. Then the functions $\sum_{i=1}^d \lambda_i^k(q) \mu_i^l(q)$ extend to holomorphic functions on some open neighbourhood of q in the Hilbert space $\mathcal{H} \supset \mathcal{H}^{\infty}$ for all $k, l \in \mathbb{N}_0$.

²⁸Those readers, which are not interested in these Darboux coordinates may pass over the rest of this section.

Proof: We know already from Theorem [2.2](#page-7-0) that $g(x, \cdot, \cdot)$ is an entire function on $\mathbb{C} \times \mathcal{H}$. Hence $v(\lambda, \mu, q)$ and $w(\lambda, \mu, q)$ defined in Lemma [4.1](#page-20-0) with normalization $v_1(\lambda, \mu, q) = 1$ and $w_1(\lambda, \mu, q) = 1$ are meromorphic functions on $\mathbb{C} \times \mathbb{C} \times \mathcal{H}$. Then there exists some $1 \leq i \leq n$, such that the local part of the divisor $D(\tilde{q})$ is given by the divisor of $1/v_i(\cdot, \cdot, \tilde{q})$ for all \tilde{q} in some open neighbourhood of q . Hence we have

$$
\sum_{i=1}^d \lambda_i^k(\tilde{q}) \mu_i^l(\tilde{q}) = \frac{1}{2\pi\sqrt{-1}} \int_{\Gamma(\tilde{q})} \lambda^k \mu^l \frac{\frac{\partial v_i}{\partial \lambda} d\lambda + \frac{\partial v_i}{\partial \mu} d\mu}{v_i},
$$

where $\Gamma(\tilde{q})$ is some loop on the Riemann surface $Y(\tilde{q})$ corresponding to the potential \tilde{q} around the local part of the divisor $D(\tilde{q})$ for all \tilde{q} in some open neighbourhood of q. Now it is easy to see that the right hand side is holomorphic on some open neighbourhood of q . In fact, Corollary [2.3](#page-8-0) gives an estimate for $||F(\lambda, q) - F(\lambda, \tilde{q})||$ in terms of $||q - \tilde{q}||$, which ensures that there exists some $\epsilon > 0$, such that the above given formula is valid for all $||q - \tilde{q}|| < \epsilon$. there exists some $\epsilon > 0$, such that the above given formula is valid for all $||q - \tilde{q}|| < \epsilon$. With the next theorem we finish our short excursion to the parametrization of the space of potentials given by the values of λ and μ at all points of the divisors. It is shown that these are almost²⁹ global coordinates and, moreover, are Darboux coordinates of the symplectic manifold. The book [\[P-T](#page-91-0)] describes from this point of view Hill's equation as a completely integrable system with action angle variables, given by these values of λ and $\ln(\mu)$ at all points of the divisor. In fact, in this case the Dirichlet isospectral sets are exactly the Lagrangian submanifolds defined by the property that the values of λ at all points of the divisor are kept fixed. The next theorem generalizes the 'basic' Theorem 2.8 and Theorem 3.5. It should be possible to carry over other parts of the beautiful analysis given in that book. After this theorem we want to return to our isospectral sets.

Theorem 7.3 Let $q \in \mathcal{H}^{\infty}$ be a potential such that the corresponding Riemann surface fulfills the assumption $b_{\text{analytic}} = b_{\text{algebraic}}$. If $\tilde{q} \in \mathcal{H}^{\infty}$ is another potential such that the values of λ and μ at all points of the divisors $D(q)$ and $D(\tilde{q})$ and the corresponding multiplicities are equal up to permutation of the points, then $q = \tilde{q}$. Moreover these coordinates are almost Darboux coordinates in the sense that

$$
\omega(\delta q, \delta \tilde{q}) = \sum_{i \in D(q)} \frac{d\lambda_i}{dq} (\delta q) \frac{d \ln(\mu_i)}{dq} (\delta \tilde{q}) - \frac{d\lambda_i}{dq} (\delta \tilde{q}) \frac{d \ln(\mu_i)}{dq} (\delta q).
$$

This means that in terms of these coordinates the symplectic form is given by

$$
\omega = \sum_i d\lambda_i \wedge d\ln(\mu_i).
$$

Proof³⁰: Let $R(\lambda, \mu, q) = 0$ and $R(\lambda, \mu, \tilde{q}) = 0$ be the two defining equations of the Riemann surfaces corresponding to q and \tilde{q} . We claim that $R(\lambda, \mu, \tilde{q}) / (\mu \frac{\partial R(\lambda, \mu, q)}{\partial \mu})$ is a section of the sheaf $\mathcal{O}_{b-D(q)}$ over $\pi^{-1}(\mathbb{C})$ of the Riemann surface $Y(q)$ corresponding to q, and moreover, that this function is bounded uniformly on $U_{0,\epsilon}$ for all $\epsilon > 0$ by $c/|\lambda|$, with some $c > 0$. In fact the divisor of the denominator is equal to b and $R(\lambda, \mu, \tilde{q})$ is assumed to have zeroes at all the

 29 In Example [10.3](#page-77-0) this coordinates fail to be one to one.

³⁰Those readers, which are not interested in this proof may jump to the next section.

points of $D(q)$. Now let $\mu_i(q)$ and $\mu_i(\tilde{q})$ be the solution of $R(\lambda, \mu, q) = 0$ and $R(\lambda, \mu, \tilde{q}) = 0$ on the *i*-th sheet of some neighbourhood of $\lambda = \infty$. Then on the *j*-sheet of this neighbourhood $R(\lambda, \mu, \tilde{q}) / (\mu \frac{\partial R(\lambda, \mu, q)}{\partial \mu})$ is equal to

$$
\frac{\mu_j(q) - \mu_j(\tilde{q})}{\mu_j(q)} \prod_{i \neq j} \frac{\mu_j(q) - \mu_i(\tilde{q})}{\mu_j(q) - \mu_i(q)} = \frac{\mu_j(q) - \mu_j(\tilde{q})}{\mu_j(q)} \prod_{i \neq j} \left(1 + \frac{\mu_i(q) - \mu_i(\tilde{q})}{\mu_j(q) - \mu_i(q)}\right).
$$

By definition of $U_{0,\epsilon}$, $\mu_i(q)$ $\mu_j(q)-\mu_i(q)$ is bounded on this set. Then Theorem [2.6](#page-10-0) shows the claim. Due to Theorem [4.3](#page-21-0) and Theorem [4.6](#page-23-0) $R(\lambda, \mu, \tilde{q}) / (\mu \frac{\partial R(\lambda, \mu, q)}{\partial \mu})$ can be written as $\sum_{i=1}^{n} f_i \frac{w_i}{w v}$ with holomorphic entire functions f_i on \mathbb{C} . Moreover, f_i are bounded near $\lambda = \infty$ by $c/|\lambda|$. Hence they are all zero, and $R(\lambda, \mu, \tilde{q})$ is zero on the Riemann surface corresponding to q. Then q and \tilde{q} correspond to the same Riemann surface and due to Section [4](#page-19-0) q and \tilde{q} are equal. Now let $q_{t,\tilde{t}}$ be the potential $q_{t,\tilde{t}} = q + t\delta q + \tilde{t}\delta \tilde{q}$. Let $v(q_{t,\tilde{t}})$ and $w(q_{t,\tilde{t}})$ be the solutions of Lemma [4.1](#page-20-0) with normalization $v_1(q_{t,\tilde{t}}) = 1 = w_1(q_{t,\tilde{t}})$. The function $(\mu_{t,\tilde{t}})^x = \exp(x \ln(\mu_{t,\tilde{t}}))$ is of course a multivalued function on $\pi^{-1}(\mathbb{C})$ of the Riemann surface corresponding to $q_{t,\tilde{t}}$. Hence $v(x, t, \tilde{t}) =$ $\pi^*(h_0g(x,\lambda, q_{t,\tilde{t}})h_0^{-1})v(q_{t,\tilde{t}})\mu_{t,\tilde{t}}^{-x}$ and $w(x,t,\tilde{t}) = \frac{1}{w(q_{t,\tilde{t}})v(q_{t,\tilde{t}})}\mu_{t,\tilde{t}}^xw(q_{t,\tilde{t}})\pi^*(h_0g^{-1}(x,\lambda,q_{t,\tilde{t}})h_0^{-1})$ are multivaluedmeromorphic solutions of ([8\)](#page-20-0) and ([9\)](#page-20-0) corresponding to the potential $T_xq_{t,\tilde{t}}$, respectively. With the choice $\frac{d\mu}{dq} = 0$ the functions $\frac{\partial v(x,t=0,0)}{\partial t}$, $\frac{\partial v(x,0,\tilde{t}=0)}{\partial \tilde{t}}$ $\frac{\partial u(x,t=0,0)}{\partial \tilde{t}}$, $\frac{\partial w(x,t=0,0)}{\partial t}$ and $\frac{\partial w(x,0,\tilde{t}=0)}{\partial \tilde{t}}$ become multivalued meromorphic functions on the Riemann surface Y corresponding to q . Let us assume that the matrix p is invertible. Otherwise the transformation $p \mapsto p+a1$ corresponds to the transformation $(\lambda, \mu) \mapsto (\lambda, \mu \exp(-a\lambda))$ without change of the Riemann surface. Let $\kappa_{t,\tilde{t}}$ be the meromorphic function $-\mu_{t,\tilde{t}}\frac{d\lambda}{d\mu_{t}}$ $\frac{d\lambda}{d\mu_{t,\tilde{t}}}$ of the Riemann surface corresponding to $q_{t,\tilde{t}}$. Theo-rem [2.6](#page-10-0) shows that at the *i*-th covering point of infinity $\kappa_{t,\tilde{t}}$ is equal to $1/p_i$. It is quite obvious that the poles of $\kappa = \kappa_{0,0}$ are the branchpoints of the covering map $\pi^{-1}(\mathbb{C}) \to \mathbb{C}$, induced by the holomorphic function μ . More precisely, the branching divisor of this covering map is given by the divisor $b - (\kappa)$. Now we need two lemmata.

Lemma 7.4 Let $P(\delta q, \delta \tilde{q})$ be the form

$$
P(\delta q, \delta \tilde{q}) = \lambda \frac{d\mu}{\mu} \int_0^1 \left(\frac{\partial (\kappa_{0, \tilde{t}=0} w(x, 0, \tilde{t}=0))}{\partial \tilde{t}} h_0 p h_0^{-1} \frac{\partial v(x, t=0, 0)}{\partial t} - \frac{\partial (\kappa_{t=0, 0} w(x, t=0, 0))}{\partial t} h_0 p h_0^{-1} \frac{\partial v(x, 0, \tilde{t}=0)}{\partial \tilde{t}} \right) dx.
$$

With the choice $\frac{d\mu}{dq} = 0$ P(δq , $\delta \tilde{q}$) becomes a meromorphic differential form on the Riemann surface Y with poles only at the branchpoints of the covering map induced by μ and the covering points of infinity. Furthermore, this form does not depend on the normalization of the solutions $v(0, t, \tilde{t})$ $v(0, t, \tilde{t})$ $v(0, t, \tilde{t})$ and $w(0, t, \tilde{t})$ of [\(8](#page-20-0)) and ([9\)](#page-20-0), respectively, whenever $w(0, t, \tilde{t})v(0, t, \tilde{t}) = 1$. The sum of residues at the covering points of infinity of $P(\delta q, \delta \tilde{q})$ is equal to $2\pi\sqrt{-1}\omega(\delta q, \delta \tilde{q})$.

Proof: The different values of $v(x, t, \tilde{t})$ and $w(x, t, \tilde{t})$ are obtained by multiplication with $\exp(2\pi kx\sqrt{-1})$ and $\exp(-2\pi kx\sqrt{-1})$ with $k \in \mathbb{Z}$, respectively. An easy calculation shows that $P(\delta q, \delta \tilde{q})$ does not depend on k and therefore is single valued. Let $f(t, t)$ be an arbitrary

smooth function with values in the meromorphic functions on the Riemann surface corresponding to $q_{t,\tilde{t}}$ and set $\tilde{v}(x, t, \tilde{t}) = f(t, \tilde{t})v(x, t, \tilde{t})$ and $\tilde{w}(x, t, \tilde{t}) = 1/f(t, \tilde{t})w(x, t, \tilde{t})$. Then we have:

$$
\int_0^1 \frac{\partial(\kappa_{0,\tilde{t}=0}\tilde{w}(x,0,\tilde{t}=0))}{\partial \tilde{t}} h_0 p h_0^{-1} \frac{\partial \tilde{v}(x,t=0,0)}{\partial t} dx -
$$
\n
$$
-\int_0^1 \frac{\partial(\kappa_{t=0,0}\tilde{w}(x,t=0,0))}{\partial t} h_0 p h_0^{-1} \frac{\partial \tilde{v}(x,0,\tilde{t}=0)}{\partial \tilde{t}} dx -
$$
\n
$$
-\int_0^1 \frac{\partial(\kappa_{0,\tilde{t}=0}\tilde{w}(x,0,\tilde{t}=0))}{\partial \tilde{t}} h_0 p h_0^{-1} \frac{\partial v(x,t=0,0)}{\partial t} dx +
$$
\n
$$
+\int_0^1 \frac{\partial(\kappa_{t=0,0}\tilde{w}(x,t=0,0))}{\partial t} h_0 p h_0^{-1} \frac{\partial v(x,0,\tilde{t}=0)}{\partial \tilde{t}} dx =
$$
\n
$$
=\frac{1}{f(0,0)} \frac{\partial f(t=0,0)}{\partial t} \frac{\partial}{\partial \tilde{t}} \int_0^1 \kappa_{0,\tilde{t}=0} w(x,0,\tilde{t}=0) h_0 p h_0^{-1} v(x,0,\tilde{t}=0) dx -
$$
\n
$$
-\frac{1}{f(0,0)} \frac{\partial f(0,\tilde{t}=0)}{\partial \tilde{t}} \frac{\partial}{\partial t} \int_0^1 \kappa_{t=0,0} w(x,t=0,0) h_0 p h_0^{-1} v(x,t=0,0) dx.
$$

The function $\frac{\partial g(x,\lambda,q)}{\partial \lambda}$ is a solution of the differential equation

$$
\left(\frac{d}{dx} + q(x) + \lambda p\right) \frac{\partial g(x, \lambda, q)}{\partial \lambda} + pg(x, \lambda, q) = 0, \ \frac{\partial g(0, \lambda, q)}{\partial \lambda} = 0.
$$

Due to Lemma [2.1](#page-6-0) $\frac{\partial g(1,\lambda,q)}{\partial \lambda}$ is given by

$$
\frac{\partial g(1,\lambda,q)}{\partial \lambda} = g(1,\lambda,q) \int_0^1 g^{-1}(x,\lambda,q) pg(x,\lambda,q) dx.
$$

Hence we have

$$
\kappa_{t,\tilde{t}} \int_0^1 w(x,t,\tilde{t}) h_0 p h_0^{-1} v(x,t,\tilde{t}) dx = -\kappa_{t,\tilde{t}} w(0,t,\tilde{t}) \pi^* \left(F^{-1}(\lambda, q_{t,\tilde{t}}) \frac{\partial F(\lambda, q_{t,\tilde{t}})}{\partial \lambda} \right) v(0,t,\tilde{t}) =
$$

=
$$
-\frac{\kappa_{t,\tilde{t}}}{\mu_{t,\tilde{t}}} \frac{d\mu_{t,\tilde{t}}}{d\lambda} = 1.
$$

This shows that $P(\delta q, \delta \tilde{q})$ does not depend on the normalization of $v(0, t, \tilde{t})$ and $w(0, t, \tilde{t})$, whenever $w(0, t, \tilde{t})v(0, t, \tilde{t}) = 1$. Therefore $P(\delta q, \delta \tilde{q})$ have poles only at the branchpoints of the covering map induced by μ and the covering points of infinity. In order to prove the last statement we claim that at all covering points of infinity $P(\delta q, \delta \tilde{q})$ has the same residues as the form

$$
\lambda d\lambda \int_0^1 \frac{\partial w(x, t=0,0)}{\partial t} h_0 p h_0^{-1} \frac{\partial v(x, 0, \tilde{t}=0)}{\partial \tilde{t}} dx - \lambda d\lambda \int_0^1 \frac{\partial w(x, 0, \tilde{t}=0)}{\partial \tilde{t}} h_0 p h_0^{-1} \frac{\partial v(x, t=0, 0)}{\partial t} dx
$$

as well with the choice $\frac{\partial \mu}{\partial q} = 0$ as with the choice $\frac{\partial \lambda}{\partial q} = 0$. The first is true because $\kappa \frac{d\mu}{\mu} = -d\lambda$ and $\frac{d\kappa_{t,\tilde{t}}}{dt}$ has a zero of order 2 at all covering points of infinity. The second is true because the

difference of $\frac{\partial v}{\partial t}$ with the choice $\frac{d\mu}{dq} = 0$ minus $\frac{\partial v}{\partial t}$ with the choice $\frac{d\lambda}{dq} = 0$ is equal to $\frac{dv}{d\lambda}$ $\frac{\partial v}{\partial t}$, where $\frac{\partial \lambda}{\partial t}$ is taken with the choice $\frac{d\mu}{dq} = 0$ and similar statements about $\frac{\partial v}{\partial \tilde{t}}$, $\frac{\partial w}{\partial t}$ and $\frac{\partial w}{\partial \tilde{t}}$. In fact, all these differences have zeroes of order al least 3 at all covering points of infinity. Hence the sum of the residues at the covering points of infinity of $P(\delta q, \delta \tilde{q})$ is equal to the sum of residues at the covering points of infinity of

$$
d\lambda \int_0^1 \frac{\partial w(x, t=0,0)}{\partial t} \pi^*(h_0 p \lambda h_0^{-1}) \frac{\partial v(x,0,\tilde{t}=0)}{\partial \tilde{t}} dx - d\lambda \int_0^1 \frac{\partial w(x,0,\tilde{t}=0)}{\partial \tilde{t}} \pi^*(h_0 p \lambda h_0^{-1}) \frac{\partial v(x,t=0,0)}{\partial t} dx.
$$

Let Σ denote the sum over all sheets of the covering map π . Then Lemma [4.5](#page-22-0) implies

$$
\sum \int_0^1 \frac{\partial w(x, t=0, 0)}{\partial t} \pi^*(h_0 p \lambda h_0^{-1}) \frac{\partial v(x, 0, \tilde{t}=0)}{\partial \tilde{t}} dx -
$$

$$
-\sum \int_0^1 \frac{\partial w(x, 0, \tilde{t}=0)}{\partial \tilde{t}} \pi^*(h_0 p \lambda h_0^{-1}) \frac{\partial v(x, t=0, 0)}{\partial t} dx =
$$

$$
= \int_0^1 tr \left(h_0^{-1} \sum \frac{\partial v(x, 0, \tilde{t}=0)}{\partial \tilde{t}} w(x, 0, 0) h_0 p \lambda h_0^{-1} \sum \frac{\partial v(x, t=0, 0)}{\partial t} w(x, 0, 0) h_0 -
$$

$$
-h_0^{-1} \sum \frac{\partial v(x, t=0, 0)}{\partial t} w(x, 0, 0) h_0 p \lambda h_0^{-1} \sum \frac{\partial v(x, 0, \tilde{t}=0)}{\partial \tilde{t}} w(x, 0, 0) h_0 \right) dx.
$$

Now the asymptotic expansions of v and w in Theorem [2.4](#page-8-0) and Theorem [2.6](#page-10-0) imply that the leading term in the asymptotic expansion of $\left(h_0^{-1}\left(\sum \frac{\partial v(x,t=0,0)}{\partial t}w(x,0,0)\right)h_0\right)$ \sum_{ij} and $\overline{\left(h_0^{-1}\left(\sum \frac{\partial v(x,0,\tilde{t}=0)}{\partial \tilde{t}} w(x,0,0)\right)h_0\right)}$ is given by $1/\lambda \frac{\delta q_{ij}}{p_j-p_i}$ and $1/\lambda \frac{\delta \tilde q_{ij}}{p_j-p_i}$ for $i \neq j$, respectively. Hence the sum of the residues at the covering points of infinity of $P(\delta q, \delta \tilde{q})d\lambda$ is equal to

$$
-2\pi\sqrt{-1}\sum_{i\neq j}\int_0^1 \frac{\delta\tilde{q}_{ij}(x)}{p_j - p_i}(p_j - p_i)\frac{\delta q_{ji}(x)}{p_i - p_j}dx = 2\pi\sqrt{-1}\sum_{i\neq j}\int_0^1 \frac{\delta q_{ij}(x)\delta\tilde{q}_{ji}(x)}{p_i - p_j}dx.
$$

This completes the proof of the lemma. \Box

Lemma 7.5 Let $Q(\delta q, \delta \tilde{q})$ be the form $Q(\delta q, \delta \tilde{q}) =$

$$
= -\frac{\kappa}{\mu} d\mu \int_0^1 \frac{\partial \lambda}{\partial t} w(x, 0, 0) h_0 p h_0^{-1} \frac{\partial v(x, 0, \tilde{t} = 0)}{\partial \tilde{t}} dx +
$$

+
$$
\frac{\kappa}{\mu} d\mu \int_0^1 \frac{\partial \lambda}{\partial \tilde{t}} w(x, 0, 0) h_0 p h_0^{-1} \frac{\partial v(x, t = 0, 0)}{\partial t} dx.
$$

With the choice $\frac{d\mu}{dq} = 0$ $Q(\delta q, \delta \tilde{q})$ is a single valued meromorphic differential form on the Riemann surface Y with poles only at the branchpoints of the covering map induced by μ and the poles of $v(q)$. If $v(q_{t,\tilde{t}})$ has a simple pole of order 1 at $(\lambda_0(t,\tilde{t}), \mu_0(t,\tilde{t}))$ and if this point is no branchpoint, the local residue of the form $Q(\delta q, \delta \tilde{q})$ at this point $(\lambda_0(t,\tilde{t}), \mu_0(t,\tilde{t}))$ is given by

$$
2\pi\sqrt{-1}\left(\frac{\partial\lambda_0}{\partial t}\frac{\partial\ln(\mu_0)}{\partial\tilde{t}}-\frac{\partial\lambda_0}{\partial\tilde{t}}\frac{\partial\ln(\mu_0)}{\partial t}\right)
$$

.

Moreover, if $v(q)$ has no poles and zeroes at the branchpoints, the total sum of all residues at the branchpoints of the covering map induced by μ of $P(\delta q, \delta \tilde{q}) + Q(\delta q, \delta \tilde{q})$ is equal to zero.

ij

Proof: If we multiply $v(x, t, \tilde{t})$ and $w(x, t, \tilde{t})$ by $\exp(2\pi kx\sqrt{-1})$ and $\exp(-2\pi kx\sqrt{-1})$, respectively, again $Q(\delta q, \delta \tilde{q})$ does not change. Hence it is single valued. The statement about the poles is obvious. If $f(t,\tilde{t})$ is locally a holomorphic function such that the divisor of $f(t,\tilde{t})$ is locally equal to $D(q_{t,\tilde{t}})$,

$$
\tilde{Q}(\delta q, \delta \tilde{q}) = -\frac{\kappa}{\mu} d\mu \int_0^1 \frac{\partial \lambda}{\partial t} \tilde{w}(x, 0, 0) h_0 p h_0^{-1} \frac{\partial \tilde{v}(x, 0, \tilde{t} = 0)}{\partial \tilde{t}} dx +
$$

$$
+ \frac{\kappa}{\mu} d\mu \int_0^1 \frac{\partial \lambda}{\partial \tilde{t}} \tilde{w}(x, 0, 0) h_0 p h_0^{-1} \frac{\partial \tilde{v}(x, t = 0, 0)}{\partial t} dx
$$

with $\tilde{v}(x, t, \tilde{t}) = f(t, \tilde{t})v(x, t, \tilde{t})$ and $\tilde{w}(x, t, \tilde{t}) = w(x, t, \tilde{t})/f(t, \tilde{t})$ is locally holomorphic. Hence the residue at $(\lambda_0(0,0), \mu_0(0,0))$ of $Q(\delta q, \delta \tilde{q})$ is equal to the residue of

$$
Q(\delta q, \delta \tilde{q}) - \tilde{Q}(\delta q, \delta \tilde{q}) = \left(\frac{1}{f(0,0)}\frac{\partial f(0,\tilde{t}=0)}{\partial \tilde{t}}\frac{\partial \lambda}{\partial t} - \frac{1}{f(0,0)}\frac{\partial f(t=0,0)}{\partial t}\frac{\partial \lambda}{\partial \tilde{t}}\right)\frac{d\mu}{\mu}.
$$

This residue is equal to

$$
2\pi\sqrt{-1}\left(\frac{\partial\lambda_0}{\partial t}\frac{\partial\ln(\mu_0)}{\partial\tilde{t}}-\frac{\partial\lambda_0}{\partial\tilde{t}}\frac{\partial\ln(\mu_0)}{\partial t}\right).
$$

To prove the last statement we first note that $P(\delta q, \delta \tilde{q}) + Q(\delta q, \delta \tilde{q})$ is equal to

$$
\frac{\partial}{\partial \tilde{t}} \lambda \kappa_{0,\tilde{t}=0} \int_0^1 w(x,0,\tilde{t}=0) h_0 p h_0^{-1} \frac{\partial v(x,t=0,\tilde{t}=0)}{\partial t} dx \frac{d\mu}{\mu} - \frac{\partial}{\partial t} \lambda \kappa_{t=0,0} \int_0^1 w(x,t=0,0) h_0 p h_0^{-1} \frac{\partial v(x,t=0,\tilde{t}=0)}{\partial \tilde{t}} dx \frac{d\mu}{\mu}.
$$

Now let $(\lambda_0(t,\tilde{t}), \mu_0(t,\tilde{t}))$ be a branchpoint of order b_0 of the covering map induced by μ such that $v(x, t, \tilde{t})$ has no pole at (λ_0, μ_0) . A local coordinate on the Riemann surface is given by $\nu^{b_0+1} = \mu - \mu_0$. Then we claim that $\frac{\kappa}{\mu} d\mu \int_0^1 w(x, t, \tilde{t}) h_0 p h_0^{-1} \frac{\partial v}{\partial t}(x, t, \tilde{t}) dx$ has a pole of order 1 at (λ_0, μ_0) , and the residue of this form is locally equal to the residue of the form

$$
\frac{1}{2} \frac{d\mu}{\mu} \frac{d}{dt} \ln \left(\frac{\partial R(\lambda, \mu, q_{t,\tilde{t}})}{\partial \lambda} \right),
$$

where again the derivative $\frac{d}{dt}$ is taken with the choice $\frac{\partial \mu}{\partial q} = 0$:

$$
\frac{d}{dt}\ln\left(\frac{\partial R(\lambda,\mu,q_{t,\tilde{t}})}{\partial\lambda}\right)=\frac{R_{\lambda t}}{R_{\lambda}}+\frac{R_{\lambda\lambda}}{R_{\lambda}}\frac{d\lambda}{dt}.
$$

Here we used the subscript to denote partial derivatives. In order to prove this claim we need some preparation. The divisor of $w(x, t, \tilde{t})$ is given by $-b$ minus the divisor of $v(x, t, \tilde{t})$. Due to the assumption $b_{\text{effective}} = b_{\text{analytic}}$, $w(x, t, t)$ has also no poles and zeroes at (λ_0, μ_0) . Now let (λ, μ) and (λ', μ) be two different points of the Riemann surface Y such that the covering map

induced by μ project them onto the same point $\mu \in \mathbb{C}$. Let v' be the value of v at (λ', μ) . Then we have

$$
\int_0^1 w(x, t, \tilde{t}) h_0 p h_0^{-1} v'(x, t, \tilde{t}) dx =
$$

$$
\frac{1}{\lambda - \lambda'} \int_0^1 w(x, t, \tilde{t}) h_0 \left(\left(\frac{d}{dx} + q(x) + p\lambda \right) - \left(\frac{d}{dx} + q(x) + p\lambda' \right) \right) h_0^{-1} v'(x, t, \tilde{t}) dx = 0.
$$

More generally, let the prime denote the value of the corresponding function at the point with α local coordinates ν' and the function without prime denotes this function at the point with local coordinates ν . We can expand the function $\int_0^1 \kappa w(x, t, \tilde{t}) h_0 p h_0^{-1} v'(x, t, \tilde{t}) dx$ in a Laurent series in ν and ν' : $\sum_{l \geq -b_0, l' \geq 0} a_{l,l'} \nu^{l} (\nu')^{l'}$. Since κ has a pole of order b_0 , we may assume $l \geq -b_0$. For $\nu =$ ν' this function is equal to $1: \sum_{l+l'=L} a_{l,l'} = \delta_{L,0}$; and for $\nu^{b_0+1} = (\nu')^{b_0+1}, \nu \neq \nu'$, this function is equal to zero. With the help of a finite Fourier transformation we obtain $a_{l,l'} = 0$, if $l + l' < 0$ and $a_{-l,l} = 1/(b_0 + 1)$ for $l = 0, 1, ..., b_0$. The form $\int_0^1 \kappa w(x, t, \tilde{t}) h_0 p h_0^{-1} dv(x, t, \tilde{t}) dx$ has a Laurent expansion of the form

$$
\sum_{l\geq -b_0,l'\geq 0} l'a_{l,l'}\nu^{l+l'-1}d\nu + \sum_{l\geq -b_0,l'\geq 0} da_{l,l'}\nu^{l+l'}.
$$

The second term vanishes and the first term has a simple pole of the form $\sum_{l=0}^{b_0} \frac{l}{b_0-l}$ b_0+1 $\frac{d\nu}{\nu}=\frac{b_0}{2}$ 2 $d\nu$ $\frac{d\nu}{\nu}$. Since $R_{\lambda}(\lambda,\mu,q_{t,\tilde{t}})$ has a pole of order b_0 at (λ_0,μ_0) , this form has the same pole as the form 1 $\frac{1}{2}d\ln(R_\lambda(\lambda,\mu,q_{t,\tilde{t}}))$. This proves the claim. If $v(q)$ has a no poles and zeroes at the branchpoints of the covering map induced by μ , the total sum of residues at the branchpoints of the form $P(\delta q, \delta \tilde{q}) + Q(\delta q, \delta \tilde{q})$ is equal to the sum of residues at the covering points of infinity of the form

$$
\frac{1}{2}\frac{d\mu}{\mu}\left(\frac{\partial\lambda}{\partial\tilde{t}}\frac{d}{dt}(\ln R_{\lambda})-\frac{\partial\lambda}{\partial t}\frac{d}{d\tilde{t}}(\ln R_{\lambda})\right),\,
$$

where all derivatives are taken according to the choice $\frac{d\mu}{dq} = 0$. At the beginning of this section we showed that $\frac{d\mu}{\mu}$ $\frac{\partial \lambda}{\partial t} = -\Omega(\delta q)$ and $\frac{d\mu}{\mu}$ $\frac{\partial \lambda}{\partial \tilde{t}} = -\Omega(\delta \tilde{q})$ are elements of $H_{\text{modified}}^0(Y, \Omega)$. Since $\frac{d}{dt}$ ln(k) and $\frac{d}{dt}$ ln(k) has zeroes of order at least 2 at all covering points of infinity, the total sum of all residues at the branchpoints of $P(\delta q, \delta \tilde{q}) + Q(\delta q, \delta \tilde{q})$ is also equal to the sum of residues at the covering points of infinity of the form

$$
\frac{1}{2}\Omega(\delta\tilde{q})\frac{d}{dt}\ln(R_{\mu}) - \frac{1}{2}\Omega(\delta q)\frac{d}{d\tilde{t}}\ln(R_{\mu}),
$$

where all derivatives are taken according to the choice $\frac{d\mu}{dq} = 0$. Now we claim that this is also true if the derivatives are taken according to the choice $\frac{d\lambda}{dq} = 0$. In fact, the difference of $\frac{d}{dt} \ln(R_\mu)$ with the choice $\frac{d\mu}{dq} = 0$ minus $\frac{d}{dt} \ln(R_\mu)$ with the choice $\frac{d\lambda}{dq} = 0$ is equal to $\frac{d}{d\lambda} \ln(R_\mu) \frac{\partial \lambda}{\partial t}$, where the derivative is taken according the choice $\frac{d\mu}{dq} = 0$. Hence the derivatives can be taken according to the choice $\frac{d\mu}{dq} = 0$ as well as to the choice $\frac{d\lambda}{dq} = 0$. Let us now take all derivatives according to the choice $\frac{d\lambda}{dq} = 0$.

$$
\frac{1}{2}\Omega(\delta\tilde{q})\frac{d}{dt}\ln(R_{\mu}) - \frac{1}{2}\Omega(\delta q)\frac{d}{d\tilde{t}}\ln(R_{\mu}) =
$$

$$
= \frac{1}{2} d\lambda \left(\frac{1}{\mu} \frac{d\mu}{d\tilde{t}} \left(\frac{R_{t\mu}}{R_{\mu}} + \frac{R_{\mu\mu}}{R_{\mu}} \frac{d\mu}{dt} \right) - \frac{1}{\mu} \frac{d\mu}{dt} \left(\frac{R_{\tilde{t}\mu}}{R_{\mu}} + \frac{R_{\mu\mu}}{R_{\mu}} \frac{d\mu}{d\tilde{t}} \right) \right) =
$$

$$
= \frac{1}{2} \Omega(\delta \tilde{q}) \frac{R_{t\mu}}{R_{\mu}} - \frac{1}{2} \Omega(\delta q) \frac{R_{\tilde{t}\mu}}{R_{\mu}}.
$$

Now let μ_i be the value of μ in the *i*-th sheet over λ . Then

$$
R(\lambda, \mu) = \prod_{i=1}^{n} (\mu - \mu_i), \ R_{\mu} = \sum_{j=1}^{n} \prod_{i \neq j} (\mu - \mu_i) \text{ and } R_{\mu t} = \sum_{j=1}^{n} \sum_{k \neq j} \frac{d\mu_k}{dt} \prod_{i \neq k, j} (\mu - \mu_i).
$$

In the *j*-th sheet $\frac{R_{\mu t}}{R_{\mu}}$ takes the value

$$
\sum_{k \neq j} \frac{\frac{d\mu_j}{dt} + \frac{d\mu_k}{dt}}{\mu_k - \mu_i}
$$

.

 $\begin{array}{c} \begin{array}{c} \begin{array}{c} \end{array}\\ \begin{array}{c} \end{array} \end{array} \end{array}$ $\frac{d}{dt} \ln(\mu)$ is bounded on some set $U_{0,\epsilon}$ by $c|\lambda|^{-1}$. In the beginning of the proof of Theorem [7.3](#page-46-0) we used the fact that $\sup\{|\mu_k|, |\mu_j|\}/|\mu_k - \mu_j|$ is bounded on all $U_{0,\epsilon}$. Hence $R_{\mu t}$ R_μ    is bounded on all $U_{0,\epsilon}$ by $c/|\lambda|$. This implies that the total sum of residues at all branchpoints of the covering map induced by μ of the form $P(\delta q, \delta \tilde{q})+Q(\delta q, \delta \tilde{q})$ converges to zero in the sense of Remark [3.7](#page-17-0), if $v(q)$ has no poles at this branchpoints. This completes the proof of Lemma [7.5](#page-49-0) \Box Completion of the proof of Theorem [7.3](#page-46-0): It is quite obvious that $Q(\delta q, \delta \tilde{q})$ has no poles at the covering points of infinity. If $D(q)$ has no poles at the branchpoints of the covering map induced by μ and furthermore is composed only of simple points, the last two lemmata show that

$$
\omega(\delta q, \delta \tilde{q}) = \sum_{i \in D(q)} \frac{d\lambda_i}{dq} (\delta q) \frac{d \ln(\mu_i)}{dq} (\delta \tilde{q}) - \frac{d\lambda_i}{dq} (\delta \tilde{q}) \frac{d \ln(\mu_i)}{dq} (\delta q),
$$

and that this sum converges. If we look more carefully at the proof, we see that it is also valid without this restriction. \Box

8 The tangent space of the Jacobian variety

It is known that in the case of the Korteweg-de Vries equation the isospectral sets are not submanifolds of the space of potentials. Hence we cannot expect that the isospectral sets are differentiable manifolds with a tangent bundle. Nevertheless we can define a subspace of the tangent space of the space of potentials, of tangent vectors along the isospectral sets. By abuse of notation this will be called the tangent space.

Definition 8.1 Let $\mathcal{L}_q \subset T_q\mathcal{H}^\infty$ be the kernel of the map $\Omega_q: T_q\mathcal{H}^\infty \to H_{\text{modified}}^0(Y, \Omega)$.

In general the map $\pi: Y \to X$ is not continuous. Nevertheless, the holomorphic functions on $\pi^{-1}(U)$ are well defined for all open sets U of X. Hence the direct image sheaf of the holomorphic functions $\pi_*(\mathcal{O}_{-\pi^{-1}(\infty)}, Y)$ is a well defined sheaf over X. Moreover, there exists a natural homomorphism $H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)}) \to H^1(X, \pi_*(\mathcal{O}_{-\pi^{-1}(\infty)})$. Indeed, let \tilde{Y} be the set Y with the unique coarsest topology containing the topology of Y and the inverse image of the topology

8 THE TANGENT SPACE OF THE JACOBIAN VARIETY 53

of X under π . Then there exists a natural inclusion $H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)}) \hookrightarrow H^1(\tilde{Y}, \mathcal{O}_{-\pi^{-1}(\infty)})$. It is quite easy to see that each covering of \tilde{Y} has a refinement, such that the first cohomology group with respect to this refinement is equal to the first cohomology group with respect to some inverse image of a covering of X. Hence $H^1(\tilde{Y}, \mathcal{O}_{-\pi^{-1}(\infty)})$ and $H^1(X, \pi_*(\mathcal{O}_{-\pi^{-1}(\infty)}))$ are isomorphic. This gives the natural inclusion $H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)}) \hookrightarrow H^1(X, \pi_*(\mathcal{O}_{-\pi^{-1}(\infty)}))$. In Section [4](#page-19-0) we already indicated that there exists a isomorphism of sheaves over X :

$$
\varepsilon_q^{-1} : \pi_*(\mathcal{O}) \to \mathcal{O}_q, f \mapsto \sum Pf,
$$

where \sum denotes the sum over all sheets of the covering map π . Now let $M_n(\mathcal{O}_{-\infty})$ denote the sheaf of rings of $n \times n$ -matrices with values in $\mathcal{O}_{-\infty}$ on X and $\mathcal{O}_{-\infty,q}$ the subsheaf of all matrices, which commute with $F(\cdot, q)$. Then there exists a natural inclusion $\mathcal{O}_{\infty,q} \hookrightarrow M_n(\mathcal{O}_{-\infty})$. Putting all this together we have the following sequence of homomorphisms:

$$
H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)}) \hookrightarrow H^1(X, \pi_*(\mathcal{O}_{-\pi^{-1}(\infty)})) \simeq H^1(X, \mathcal{O}_{-\infty, q}) \to H^1(X, M_n(\mathcal{O}_{-\infty})).
$$

Let $v(q)$ and $w(q)$ be the solutions of ([8](#page-20-0)) and ([9](#page-20-0)) with normalization $v_1(q) = 1 = w_1(q)$. Furthermore, let U_D be the complement of the support of the divisor $D(q)$ in Y.

Lemma 8.2 For each element $c \in H^1(X, \mathcal{O}_{-\infty,q})$ in the kernel of

$$
H^1(X, \mathcal{O}_{-\infty,q}) \to H^1(X, M_n(\mathcal{O}_{-\infty}))
$$

there exists a section f of \mathcal{O}_D on $\pi^{-1}(\mathbb{C})$, such that under the homomorphism

$$
H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)}) \hookrightarrow H^1(X, \pi_*(\mathcal{O}_{-\pi^{-1}(\infty)})) \simeq H^1(X, \mathcal{O}_{-\infty, q})
$$

the cocycle $[f] \in H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)})$ induced by f with respect to the covering $Y = U_D \cup \pi^{-1}(\mathbb{C})$ is mapped onto c. Moreover, there exists a unique vector valued meromorphic function δv on Y , with poles only at the poles of v and zeroes at all covering points of infinity and a section a_+ of $M_n(\mathcal{O}_{-\infty})$ on \mathbb{C} , such that

$$
vf = \pi^*(a_+)v - \delta v \text{ and } (\delta v)_1 = 0.
$$

Proof: Since c is an element of the kernel of $H^1(X, \mathcal{O}_{-\infty,q}) \to H^1(X, M_n(\mathcal{O}_{-\infty}))$ there exists a cochain $b \in C^0(\mathcal{U}, M_n(\mathcal{O}_{-\infty}))$ with some covering $\mathcal U$ of x, such that δb is a representative of $c \in C^1(\mathcal{U}, \mathcal{O}_{-\infty,q})$. For any open set U of U we define on $\pi^{-1}(U)$ the function $\delta v_U = \pi^*(b_U)v - v(\pi^*(b_U)v)_1$. If V is another element of the covering U, on $\pi^{-1}(U \cap$ V) the function $\delta v_U - \delta v_V = \pi^*(b_U - b_V)v - v(\pi^*(b_U - b_V)v)_1$. Since δb is an element $C^1(\mathcal{U}, \mathcal{O}_{-\infty,q})$, $\pi^*(b_U - b_V)v = v(\pi^*(b_U - b_V)v)_1$. Hence δv is a meromorphic function on Y with poles only at the poles of v and zeroes at all covering points of infinity. If b is an element of $C^0(\mathcal{U}, \mathcal{O}_{-\infty,q})$ the same argument shows that $\delta v = 0$. This shows that δv depends only on the element c of $H^1(X, \mathcal{O}_{-\infty,q})$. If $(\lambda_0, \mu_0) \in Y$ is a pole of v and U is an element of the covering such that $\pi^{-1}(U)$ contains this pole, then the singular part of $(\pi^*(b_U)v)_1$ does only depend on c. Indeed, if V is another element of the covering, which contains this pole, $\pi^*(b_U - b_V)v = v(\pi^*(b_U - b_V)_1$ and $(\pi^*(b_U - b_V)_1$ is holomorphic, since δb is an element of $C^1(\mathcal{U}, \mathcal{O}_{-\infty,q})$. Hence c defines a unique Mittag Leffler distribution on Y, more precisely, a globalsection of the sheaf $\mathcal{O}_D/\mathcal{O}$. Due to [[Fo](#page-90-0), Theorem 26.3] there exists a solution f of this

8 THE TANGENT SPACE OF THE JACOBIAN VARIETY 54

Mittag Leffler distribution on $\pi^{-1}(\mathbb{C})$. The function f on $\pi^{-1}(\mathbb{C})$ together with the zero function on U_D defines an element of $C^0(\{U_D, \pi^{-2}(\mathbb{C})\}, \mathcal{O}_{D-\pi^{-1}(\infty)})$. The coboundary of this element defines an element of $H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)})$ denoted by [f]. Now we claim that under the map $H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)}) \hookrightarrow H^1(X, \pi_*(\mathcal{O}_{-\pi^{-1}(\infty)})) \simeq H^1(X, \mathcal{O}_{-\infty,q})$ [f] is mapped onto c. A representative of the image of $[f]$ in $H^1(X, \pi_*(\mathcal{O}_{-\pi^{-1}(\infty)}))$ is given by the coboundary of the cochain defined by the zero section of $\pi_*(\mathcal{O}_{-\pi^{-1}(\infty)})$ on $\mathbb{P}_1\setminus \pi(supportD)$ and the section f of $\pi_*(\mathcal{O}_{D-\pi^{-1}(\infty)})$ on \mathbb{C} . Now let $\mathcal{U} \cap \{\mathbb{C}, \mathbb{P}_1 \setminus \pi(supportD)\}\$ be the covering $\{U \cap \mathbb{C}, U \cap (\mathbb{P}_1 \setminus \pi(supportD)) | U \in \mathcal{U}\}\$. Then we define the following element of $C^0(\mathcal{U} \cap {\mathbb{C}}, \mathbb{P}_1 \setminus \pi(\text{supportD})\}, \pi_*(\mathcal{O}_{-\pi^{-1}(\infty)}))$, which on $U \cap \mathbb{C}$ is equal to $f-(\pi^*(b_U)v)_1$ and on $U\cap \mathbb{P}_1\setminus \pi(supportD)$ equal to zero. The coboundary of this element is equal to the image of $[f]$ under $H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)}) \hookrightarrow H^1(X, \pi_*(\mathcal{O}_{-\pi^{-1}(\infty)}))$ minus the image of c under the isomorphism $H^1(X, \mathcal{O}_{-\infty,q}) \simeq H^1(X, \pi_*(\mathcal{O}_{-\pi^{-1}(\infty)}))$. This proves the claim. Finally we note that by construction of f the function $a_+ = \sum Pf - \sum \delta vw/ wv$ is a section of $M_n(\mathcal{O}_{-\infty})$ on \mathbb{C} , and $vf = \pi^*(a_+)v - \delta v$. The functions a_+ on \mathbb{C} and the function $a_- = -\sum \delta v w / w v$ on $\mathbb{P}_1 \setminus \pi(\text{supportD})$ define an element a of $C^0(\{\mathbb{C}, \mathbb{P}_1 \setminus \pi(\text{supportD})\}, M_n(\mathcal{O}_{-\infty}))$, such that δa is equal to the image of [f] under $H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)}) \hookrightarrow H^1(X, \pi_*(\mathcal{O}_{-\pi^{-1}(\infty)})) \simeq H^1(X, \mathcal{O}_{-\infty,q}).$ This completes the proof of the lemma. \Box

Definition 8.3 Let $H^1_{q,\text{modified}}(Y, \mathcal{O}) \subset H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)})$ be the vector space of all cocycles in the kernel of the homomorphism

$$
H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)}) \hookrightarrow H^1(X, \pi_*(\mathcal{O}_{-\pi^{-1}(\infty)})) \simeq H^1(X, \mathcal{O}_{-\infty, T_xq}) \to H^1(X, M_n(\mathcal{O}_{-\infty}))
$$

for all $x \in [0,1]^{31}$. Furthermore let $\tilde{H}^1_{q,\text{modified}}(Y, \mathcal{O}) \subset H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)})$ be the preimage under the homomorphism

$$
H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)}) \hookrightarrow H^1(X, \pi_*(\mathcal{O}_{-\pi^{-1}(\infty)})) \simeq H^1(X, \mathcal{O}_{-\infty, q})
$$

of the subspace $H^1(\mathbb{P}_1, \mathcal{O}_{-\infty,q}) \subset H^1(X, \mathcal{O}_{-\infty,q}).$

Lemma 8.4 $\tilde{H}^1_{q,\text{modified}}(Y,\mathcal{O})$ is the same for all $q \in Isospectral(Y)$. Hence we will omit the index q. Moreover, $\tilde{H}_{\text{modified}}^1(Y, \mathcal{O})$ is contained in $H^1_{q, \text{modified}}(Y, \mathcal{O})$ for all $q \in \text{Isospectral}(Y)$.

Proof: Since $H^1(\mathbb{P}_1, M_n(\mathcal{O}_{-\infty})) = 0$ we can apply the last lemma. Since \mathbb{P}_1 is compact, we can calculate the cohomology groups with respect to some finite covering. Moreover, for each open subset $U \subset \mathbb{C} \subset \mathbb{P}_1$, $H^1(U, \mathcal{O}_{-\infty,q}) \simeq H^1(\pi^{-1}(U), \mathcal{O}) = 0$ (see [\[Fo,](#page-90-0) 26.1]). Hence we can calculate the cohomology group with respect to some covering $\mathbb{P}_1 = U_+ \cup U_-,$ with $U_+ \subset \mathbb{C} \subset \mathbb{P}_1$ and some neighbourhood U_- of $\infty \in \mathbb{P}_1$. To each element $c \in H^1(\mathbb{P}_1, \mathcal{O}_{-\infty,q})$ there exists a cochain $b \in C^0(\{U_+, U_-\}, M_n(\mathcal{O}_{-\infty}))$, such that δb is a representative of c in C^1 ({ U_+, U_- } $\mathcal{O}_{-\infty,q}$). Similar arguments as in the proof of the lemma before show that the coboundary of the cochain $f \in C^0(\{\pi^{-1}(U_+), \pi^{-1}(U_-)\}, \mathcal{M})$, with $f_{\pi^{-1}(U_+)} = (\pi^*(b_{U_+})v)_1$ and $f_{\pi^{-1}(U_{-})} = (\pi^*(b_{U_{-}})v)_1$ is a representative in $C^1(\{\pi^{-1}(U_{+}), \pi^{-1}(U_{-})\}, \mathcal{O}_{-\pi^{-1}(\infty)})$ of the preimage of c under the inclusion $H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)}) \hookrightarrow H^1(\mathbb{P}_1, \mathcal{O}_{-\infty,q})$. Now let $\tilde{q} \in \text{Isospectral}(Y)$ be

³¹With the help of Proposition [6.5](#page-37-0) and Theorem [6.6](#page-37-0) one can prove that this condition is equivalent for all $x \in [0,1]$. Hence we may impose this condition only for $x = 0$. But in general $H^1_{q,\text{modified}}(Y,\mathcal{O})$ does not coincide for all $q \in Isospectral(Y)$. This vector space is the same for all q and \tilde{q} , such that $E^*(q) \otimes E(\tilde{q})$ is an element of the real part of the Picard group (Compare with Theorem [9.5\)](#page-65-0).

another potential. Then it is obvious that [f] corresponds also to an element of $H^1(\mathbb{P}_1, \mathcal{O}_{-\infty, \tilde{q}})$. This shows that $\tilde{H}^1_{q,\text{modified}}(Y,\mathcal{O})$ does not depend on q and furthermore, that $\tilde{H}^1_{\text{modified}}(Y,\mathcal{O})$ is contained in $H^1_{q,\text{modified}}(Y, \mathcal{O})$.

For compact Riemann surfaces Serre duality gives a non-degenerate pairing between the first cohomology group of the sheaf of holomorphic functions and the vector space of holomorphic differential forms. With the help of Lemma [8.2](#page-53-0) we can write down the corresponding pairing between $H^1_{q,\text{modified}}(Y,\mathcal{O})$ and $H^0_{\text{modified}}(Y,\Omega)$ as an infinite sum: Let $[f] \in H^1_{q,\text{modified}}(Y,\mathcal{O})$ be represented by a cocycle of the form defined in Lemma [8.2](#page-53-0). Then for all $\omega \in H_{\text{modified}}^1(Y, \Omega)$

$$
Res([f], \omega) = -\sum Res(f\omega),
$$

where the sum runs over all poles of f on $\pi^{-1}(\mathbb{C})$. This is equal to the sum of residues of $f\omega$ at all covering points of infinity. But in general this sum does not converge. If $[f]$ is an element of $\tilde{H}^1_{\text{modified}}(Y, \mathcal{O})$ of the form defined in Lemma [8.4](#page-54-0)

$$
Res([f], \omega) = -\sum Res\left(f_{\pi^{-1}(U_+)}\omega\right) + \sum Res\left(f_{\pi^{-1}(U_-}\omega\right),\,
$$

where the first sum runs over all poles of $f_{\pi^{-1}(U_{+})}$ on $\pi^{-1}(U_{+})$ and the second sum runs over all poles of $f_{\pi^{-1}(U_-)}$ on $\pi^{-1}(U_-)$. Since $-f_{\pi^{-1}(U_+)} + f_{\pi^{-1}(U_-)}$ is holomorphic on $\pi^{-1}(U_+ \cap U_-)$, we can omit the poles on $\pi^{-1}(U_+ \cap U_-)$. Then the first sum becomes finite and the second sum converges due to Theorem [3.5.](#page-17-0) This shows that Res is a well defined pairing between $\tilde{H}^1_{\text{modified}}(Y, \mathcal{O})$ and $H^1_{\text{modified}}(Y, \Omega)$. Now we can state the main theorem of this section:

Theorem 8.5 (i) There exists an isomorphism of vector spaces $d\Gamma_q$: $H^1_{q,\text{modified}}(Y, \mathcal{O}) \to \mathcal{L}_q$, which is uniquely determined by the property that for all $[f] \in H^1_{q,\text{modified}}(Y,\mathcal{O})$

$$
\frac{\partial v}{\partial q}(d\Gamma_q([f])) = \delta v,
$$

where δv was defined in Lemma [8.2](#page-53-0).

(ii) For all $[f] \in H^1_{q,\text{modified}}(Y,\mathcal{O})$ and all $\delta q \in T_q\mathcal{H}^\infty$ the following relation holds:

$$
\omega(d\Gamma_q([f]), \delta q) = \frac{1}{2\pi\sqrt{-1}} Res([f], \Omega_q(\delta q))
$$

and the infinite sum on the right hand converges.

- (iii) The pairing Res between $\tilde{H}_{\text{modified}}^1(Y, \mathcal{O})$ and $H_{\text{modified}}^0(Y, \Omega)$ is non-degenerate.
- (iv) \mathcal{L}_q is a maximal isotropic subspace of $T_q\mathcal{H}^\infty$ with respect to the symplectic form ω .

Proof³²: Let $[f] \in H_{q,\text{modified}}^1(Y, \mathcal{O})$ be a cocycle of the form defined in Lemma [8.2](#page-53-0) and

$$
a(x) = \sum v(T_x q) \frac{f}{w(T_x q) v(T_x q)} w(T_x q).
$$

 32 This proof may be passed over.

8 THE TANGENT SPACE OF THE JACOBIAN VARIETY 56

Due to the proof of Theorem [4.10](#page-26-0) $a(x)$ is equal to $h_0g(x, \cdot, q)h_0^{-1}a(0)h_0g^{-1}(x, \cdot, q)h_0^{-1}$. This implies that $\left[a(x), \frac{d}{dx} + h_0(p\lambda + q(x))h_0^{-1}\right] = 0$. By definition of $H_{q,\text{modified}}^1(Y, \mathcal{O})$ and due to Lemma [8.2](#page-53-0) there exists for all $x \in [0,1]$ a decomposition $a(x) = a_+(x) + a_-(x)$, such that $a_+(x)$ is an entire function from $\mathbb C$ into the $n \times n$ -matrices and $a_-(x)$ is a meromorphic function on X, which vanishes at $\lambda = \infty$. Then the commutator $[a(x), h_0(p\lambda + q(x))h_0^{-1}]$ has a decomposition of the same form and both functions $a_{+}(x)$ and $a_{-}(x)$ are smooth with respect to x. Then we define

$$
\delta q(x) = \left[h_0^{-1} a_+(x) h_0, \frac{d}{dx} + p\lambda + q(x) \right] = \left[\frac{d}{dx} + p\lambda + q(x), h_0^{-1} a_-(x) h_0 \right].
$$

This equation shows that $\delta q(x)$ is an entire function with respect to λ and is holomorphic near $\lambda = \infty$. Therefore it does not depend on λ . Moreover, it is a smooth function with respect to x. Now we claim that $\delta v = -\pi^*(a_-(0))v(q)$ satisfies the relation

$$
\pi^*(F(\cdot,q))\delta v + \pi^*\left(\frac{\partial F}{\partial q}(\cdot,q)(\delta q)\right)v = \delta v\mu.
$$

Let $\delta v(x)$ be the multivalued meromorphic function $\delta v(x) = -\pi^* (a_-(x)h_0 g(x, \cdot, q)h_0^{-1})v\mu^{-x}$. Since $a(x)$ is periodic with respect to x, we have $\delta v = \delta v(0) = \delta v(1)$. Furthermore, $\delta v(x)$ is a solution of the differential equation

$$
\pi^* \left(\frac{d}{dx} + h_0(p\lambda + q(x))h_0^{-1} \right) \delta v(x) + \delta v(x) \ln(\mu) + \pi^* \left(h_0 \delta q(x) g(x, \cdot, q) h_0^{-1} \right) v \mu^{-x} = 0.
$$

Due to Lemma [2.1](#page-6-0) $\delta v(x)$ is then equal to

$$
\delta v(x) = \pi^* \left(h_0 g(x, \cdot, q) h_0^{-1} \right) \left(\int_0^x \pi^* (h_0 g^{-1}(t, \cdot, q) \delta q(t) g(t, \cdot, q) h_0^{-1}) v + \delta v(0) \right) \mu^{-x}.
$$

For $x = 1$ we obtain the equation

$$
\delta v = \pi^*(F(\cdot, q)) \delta v \mu^{-1} + \pi^* \left(\frac{\partial F}{\partial q}(\cdot, q) (\delta q) \right) v \mu^{-1}.
$$

This proves the claim. For each $\delta q \in \mathcal{L}_q$, $\delta v = \frac{\partial v}{\partial q}(\delta q)$ is a solution of this equation. With the normalization $(\delta v)_1 = 0$ this equation has only one meromorphic solution, since there are many $\lambda \in \mathbb{C}$, such that $F(\lambda, q)$ has n different eigenvalues. This shows that δv corresponding to [f] is equal to $\frac{\partial v}{\partial q}(\delta q)$. It is quite obvious that δq is an element of \mathcal{L}_q if and only if

$$
\frac{\partial v}{\partial q}(\delta q)\mu - \pi^*(F(\cdot, q))\frac{\partial v}{\partial q}(\delta q) = \pi^*\left(\frac{\partial F(\cdot, q)}{\partial q}(\delta q)\right)v - v\frac{\partial \mu}{\partial q}(\delta q)
$$

is an element of the form $\pi^*(\delta F)v$, with some entire function $\delta F : \mathbb{C} \to n \times n$ -matrices. Hence we have a map $d\Gamma_q: H^1_{q,\text{modified}}(Y,\mathcal{O}) \to \mathcal{L}_q$. Now we claim that for each vector valued function δv , which vanishes at all covering points of infinity and furthermore satisfies the relations

$$
\delta v\mu - \pi^*(F(\cdot, q))\delta v = \pi^*(\delta F)v \text{ and } (\delta v)_1 = 0
$$

with some entire function $\delta F : \mathbb{C} \to n \times n$ -matrices, there exists one and only one element $c \in H^1(X, \mathcal{O}_{-\infty,q})$ in the kernel of $H^1(X, \mathcal{O}_{-\infty,q}) \to H^1(X, M_n(\mathcal{O}_{-\infty}))$, such that δv is equal to the corresponding meromorphic function constructed in Lemma [8.2.](#page-53-0) The relation $\delta v\mu$ − $\pi^*(F(\cdot, q))\delta v = \pi^*(\delta F)v$ is equivalent to the relation

$$
\left[\sum \frac{\delta vw}{wv}, F(\cdot, q)\right] = \delta F.
$$

To proceed further we need

Lemma 8.6 Let the following homomorphism of sheaves be an isomorphism:

$$
\mathcal{O}_{\mathbb{C}}^n \to \mathcal{O}_q, (f_1, \ldots, f_n) \mapsto \sum_{i=1}^n F^{i-1}(\cdot, q) f_i.
$$

Then for all $\lambda_0 \in \mathbb{C}$ the centralizer of $F(\lambda_0, q)$ is spanned by $\mathbb{1}, F(\lambda_0, q), \ldots, F^{n-1}(\lambda_0, q)$.

Proof: Let $R(\lambda, \mu) = \det(\mu \mathbb{1} - F(\lambda, q)) = 0$ be the eigenvalue equation and let $(\lambda_0, \mu_0) \in \mathbb{C}^2$ any solution of this equation. Then the condition implies that $R(\lambda_0, F(\lambda_0, q))(F(\lambda_0, q) - \mu_0 \mathbf{1})^{-1}$ 33 is not equal to zero. This implies the conclusion of the lemma. \Box

Due to the assumption $b_{\text{analytic}} = b_{\text{effective}} = b_{\text{algebraic}}$ the condition of the foregoing lemma is fulfilled. Then this lemma and the equation ahead of this lemma imply that in a neighbourhood of every pole of $\sum \delta vw/wv$ there exists a meromorphic section f of $\pi_*(\mathcal{O})$, such that \sum of every pole of $\sum \delta vw/ wv$ there exists a meromorphic section f of $\pi_*(\mathcal{O})$, such that $\sum Pf - \sum \delta vw/ wv$ is holomorphic on this neighbourhood. Therefore the singular part of $\sum \frac{\delta vw}{wv}$ defines a Mittag Leffler distribution on $\pi^{-1}(\mathbb{C})$. Now let f be as in Lemma [8.2](#page-53-0) a solution of this Mittag Leffler distribution on $\pi^{-1}(\mathbb{C})$. Then $vf + \delta v$ is a vector valued section of the sheaf $\mathcal{O}_{D(q)}$ over $\pi^{-1}(\mathbb{C})$. Hence there exists an entire function $a_+ : \mathbb{C} \to n \times n$ -matrices such that $v f = \pi^*(a_+)v - \delta v$. Since $(\delta v)_1 = 0$, we have $f = (\pi^*(a_+)v)_1$ and $\delta v = \pi^*(a_+)v - v(\pi^*(a_+)v)_1$. If two elements c and c' of $H^1(X, \mathcal{O}_{-\infty,q})$ correspond to the same element δv , the difference of the two corresponding 0-cochains b and b' used in the proof of Lemma [8.2](#page-53-0) is a cochain in $C^0(\mathcal{U}, \mathcal{O}_{-\infty,q})$ and $\delta(b-b') = c-c'$. Hence they are equal. This proves the claim. In particular, we have for all $\delta q \in \mathcal{L}_q$ a cocycle $[f] \in H^1(Y, \mathcal{O}_{-\pi^{-1}(\infty)})$ as in Lemma [8.2,](#page-53-0) such that $v(q)f = \pi^*(a_+)v(q) - \frac{\partial v}{\partial q}(\delta q)$, with some entire function $a_+ : \mathbb{C} \to n \times n$ -matrices. Let $v(x, q)$ be the multivalued solution $v(x,q) = \pi^*(h_0 g(x,\cdot,q) h_0^{-1}) v(q) \mu^{-x}$ of [\(8\)](#page-20-0) to the potential $T_x q$. Then we have $v(x, q)f =$

$$
= \pi^* \left(h_0 g(x, \cdot, q) h_0^{-1} \left(a_+ + h_0 \frac{\partial g(x, \cdot, q)}{\partial q} (\delta q) h_0^{-1} \right) h_0 g^{-1}(x, \cdot, q) h_0^{-1} \right) v(x, q) - \frac{\partial v(x, q)}{\partial q} (\delta q)
$$

and
$$
\pi^* (F(\cdot, T_x q)) \frac{\partial v(x, q)}{\partial q} (\delta q) - \frac{\partial v(x, q)}{\partial q} (\delta q) \mu = \pi^* \left(\frac{\partial F(T_x q)}{\partial q} (T_x \delta q) \right) v(x, q).
$$

Since the solutions of the last equations are unique up to summation of some multiple of $v(x,q), \ \frac{\partial v(x,q)}{\partial q}(\delta q)$ is equal to

$$
\frac{\partial v(T_x q)}{\partial q}(T_x \delta q) v_1(x,q) + \frac{\partial v_1(x,q)}{\partial q}(\delta q) v(x,q).
$$

³³The singularity of the function $R(\lambda_0, \mu)(\mu - \mu_0)^{-1}$ at $\mu = \mu_0$ may be removed. Hence this function should be considered to be a polynomial of degree $n-1$ with respect to μ as well as with respect to $F(\lambda_0, q)$.

Similar arguments as in Theorem [4.10](#page-26-0) show that $\frac{\partial v(x,q)}{\partial q}(\delta q)$ vanishes at all covering points of infinity. Hence we have

$$
v(T_x q) f - f_{-}(x) = \pi^*(a_{+}(x)) v(T_x q) - \frac{\partial v(T_x q)}{\partial q} (T_x \delta q), \text{ with}
$$

$$
a_{+}(x) = h_0 g(x, \cdot, q) h_0^{-1} \left(a_{+} + h_0 \frac{\partial g(x, \cdot, q)}{\partial q} (\delta q) h_0^{-1} \right) h_0 g^{-1}(x, \cdot, q) h_0^{-1} \text{ and}
$$

$$
f_{-}(x) = -\frac{\partial v_1(x, q)}{\partial q} (\delta q) \frac{v_1(T_x q)}{v_1(x, q)},
$$

which is a single valued meromorphic function on Y and vanishes at all covering points of infinity. The cocycle represented by f is of course the same as the cocycle represented by $f - f_-(x)$. Therefore the cocycle corresponding to $\frac{\partial v(T_x q)}{\partial q}(T_x q)$ does not depend on x. In particular it is an element of $H_{q,\text{modified}}^1(Y, \mathcal{O})$. This defines a map

$$
\mathcal{L}_q \to H^1_{q,\text{modified}}(Y, \mathcal{O}), \delta q \mapsto [f],
$$

such that $\delta v = \frac{\partial v}{\partial q}(\delta q)$ corresponds to $[f]$ in the sense of Lemma [8.2](#page-53-0) and this property determines a unique isomorphism $d\Gamma_q: H^1_{q,\text{modified}}(Y, \mathcal{O}) \to \mathcal{L}_q$. This proves (i). Now let $\delta q \in T_q\mathcal{H}^\infty$ and $[f] \in H_{q,\text{modified}}^1(Y, \mathcal{O})$, where $[f]$ is of the same form as in Lemma [8.2](#page-53-0). Then we define

$$
b_{-}(x) = \sum \frac{\partial v(T_x q)}{\partial q}(T_x \delta q) \frac{1}{w(T_x q)v(T_x q)} w(T_x q)
$$

and
$$
a(x) = \sum v(T_x q) \frac{f}{w(T_x q)v(T_x q)} w(T_x q) = a_{+}(x) + a_{-}(x).
$$

This implies $h_0 d\Gamma([f])(x)h_0^{-1} = \left[\frac{d}{dx} + h_0(p\lambda + q(x))h_0^{-1}, a_-(x)\right]$ and

$$
h_0\delta q(x)h_0^{-1} = \left[\frac{d}{dx} + h_0(p\lambda + q(x))h_0^{-1}, b_-(x)\right] + v(T_xq)\frac{\frac{\partial \mu}{\partial q}(\delta q)}{w(T_xq)v(T_xq)}w(T_xq),
$$

where the last derivative is taken with respect to the choice $\frac{d\lambda}{dq} = 0$. Then $2\pi\sqrt{-1}\omega(d\Gamma([f]), \delta q)$ is equal to the residue at infinity of the form

$$
d\lambda \int_0^1 tr\left(a_-(x)\left(\frac{d}{2dx} + h_0(p\lambda + q(x))h_0^{-1}\right)b_-(x) -
$$

$$
-b_-(x)\left(\frac{d}{2dx} + h_0(p\lambda + q(x)h_0^{-1}\right)a_-(x)\right)dx =
$$

$$
= d\lambda \int_0^1 tr\left(b_-(x)\left(\frac{d}{2dx} + h_0(p\lambda + q(x))h_0^{-1}\right)a_+(x) -
$$

$$
-a_+(x)\left(\frac{d}{2dx} + h_0(p\lambda + q(x)h_0^{-1}\right)b_-(x)\right)dx =
$$

8 THE TANGENT SPACE OF THE JACOBIAN VARIETY 59

$$
= -d\lambda \int_0^1 tr(a_+(x)\delta q(x))dx + d\lambda \int_0^1 tr\left(a_+(x)\sum v(T_xq)\frac{\frac{\partial \mu}{\partial q}(\delta q)}{w(T_xq)v(T_xq)}w(T_xq)\right)dx.
$$

The residue at infinity of the first summand vanishes and the residue at infinity of the second summand is equal to the residue at infinity of

$$
d\lambda \int_0^1 tr\left(a(x) \sum v(T_x q) \frac{\frac{\partial \mu}{\partial q}(\delta q)}{w(T_x q)v(T_x q)} w(T_x q)\right) dx = \sum f \Omega_q(\delta q).
$$

Therefore $\omega(d\Gamma_q([f]), \delta q)$ is equal to $\frac{1}{2\pi\sqrt{-1}} Res([f], \Omega_q(\delta q))$. This proves (ii). To each finite Mittag Leffler distribution on $\pi^{-1}(\mathbb{C})$, there exists a solution f on $\pi^{-1}(\mathbb{C})$, and the corresponding cocycle $[f]$ is of course an element of $\tilde{H}^1_{\text{modified}}(Y, \mathcal{O})$. Hence $\omega \in H^1_{\text{modified}}(Y, \Omega)$ is equal to zero, if $Res([f], \omega) = 0$ for all $[f] \in \tilde{H}^1_{\text{modified}}(Y, \mathcal{O})$. On the other hand ω is non-degenerate and therefore $[f] \in \tilde{H}_{\text{modified}}^1(Y, \mathcal{O}) \subset H_{q,\text{modified}}^1(Y, \mathcal{O})$ is equal to zero, if $\omega(d\Gamma_q([f]), \delta q) =$
 $\frac{1}{\sqrt{q}} \text{Res}([f] \Omega_{\text{total}}^1(Y, \mathcal{O}) - 0$ for all $\delta q \in T\mathcal{H}^{\infty}$. Then *Res* is a non-deconomic pairing between $\frac{1}{2\pi\sqrt{-1}}Res([f],\Omega_q(\delta q))=0$ for all $\delta q \in T_q\mathcal{H}^{\infty}$. Then Res is a non-degenerate pairing between $\tilde{H}^1_{\text{modified}}(Y, \mathcal{O})$ and $H^0_{\text{modified}}(Y, \Omega)$. This implies that $\delta q \in \mathcal{L}_q$, if and only if $\omega(\delta q, \delta \tilde{q}) = 0$ for all $\delta \tilde{q} \in \mathcal{L}_q$. This completes the proof of Theorem [8.5](#page-55-0) \Box

In the finite dimensional case the last statement of the theorem would imply that the system is completely integrable. In the infinite dimensional case this is only one possibility to define the meaning of completely integrable. But there are stronger definitions, which are more satisfactory. For example one could assume the isospectral sets to be diffeomorphic to some abelian Lie groups, where the Lie algebras are isomorphic to the tangent space. For compact Riemann surfaces the first cohomology group of the sheaf of holomorphic functions is the Lie algebra of the Picard group. If the isospectral sets were to be one connected component of the Picard group this condition would be fulfilled. In Section [4](#page-19-0) we already saw that this is not the case for Riemann surfaces of infinite genus. Nevertheless we can define some subgroup of the Picard group, which acts on all divisors of Riemann-Roch type, but this action is not transitive.

Definition 8.7 Let $\tilde{H}_{\text{modified}}^1(Y, \mathcal{O}^*)$ be the subgroup of the Picard group in the modified sense, which consists of cocycles of the form f_{+}/f_{-} on $\pi^{-1}(U_{+} \cap U_{-})$, where f_{+} is a meromorphic function on $\pi^{-1}(U_+)$ and f_- is a meromorphic function on $\pi^{-1}(U_-)$, which is equal to 1 at all covering points of infinity, and $\mathbb{P}_1 = U_+ \cup U_-$ is some open covering of \mathbb{P}_1 , such that $U_+ \subset \mathbb{C} \subset \mathbb{P}_1$ and U[−] is some open neighbourhood of infinity.

Similar arguments as in the proof of Lemma [8.4](#page-54-0) show that $\tilde{H}^1_{\text{modified}}(Y, \mathcal{O}^*)$ is the subgroup of the Picard group in the modified sense, which is mapped into $H_{\text{modified}}^1(\mathbb{P}_1, \pi_*(\mathcal{O}^*)) \simeq H_{\text{modified}}^1(\mathbb{P}_1, \mathcal{O}_q^*)$ for all $q \in \textit{Isospectral}(Y)$. Due to Proposition [6.5](#page-37-0) the connected component of the identity of $\tilde{H}^1_{\text{modified}}(Y, \mathcal{O}^*)$ acts on $Jacobian(Y)$, but neither is $Jacobian_0(Y)$ an invariant subspace under this action nor is this action transitive.

The exact sequence of sheaves

$$
0 \to \mathbb{Z} \to \mathcal{O} \to \mathcal{O}^* \to 1,
$$

induces a homomorphism

$$
\exp: \tilde{H}^1_{\text{modified}}(Y, \mathcal{O}) \to \tilde{H}^1_{\text{modified}}(Y, \mathcal{O}^*).
$$

In some sense $\tilde{H}^1_{\text{modified}}(Y, \mathcal{O})$ is the Lie algebra of $\tilde{H}^1_{\text{modified}}(Y, \mathcal{O}^*)$ and this map is the exponential map. In the next section we introduce a reality condition in order to define a real subgroup of the Picard group, which acts transitively on the real part of the isospectral sets.

9 A reality condition

In this section we will formulate a reality condition on the space of potentials. The real part of the Jacobian variety will turn out to be isomorphic to a compact abelian group. For the real part of the dynamical system we will construct action angle variables. In the simplest case, the nonlinear Schrödinger equation, there are two reality conditions: the non-focussing and the self focussing nonlinear Schrödinger equation. The non-focussing nonlinear Schrödinger equation is an extension of the real Korteweg-de Vries equation and can be treated by the methods of [[MK-T-1\]](#page-91-0). The reality condition, investigated in this section, in the simplest case turns out to be the self focussing nonlinear Schrödinger equation. In the second Appendix we give a short introduction to the relation of these two reality conditions using spectral theory.

Assumption 9.1 In this section we always assume that p_1, \ldots, p_n are imaginary.

Definition 9.2 The Fréchet space of smooth periodic potentials q, which satisfies the reality condition $q^* = -q$ is denoted by $\mathcal{H}_{\mathbb{R}}^{\infty}$. In general we will use the subscript \mathbb{R} , to denote the subsets corresponding to this reality condition.

If $q \in \mathcal{H}_{\mathbb{R}}^{\infty}$, $q(x) + p\lambda$ is an element of the Lie algebra $u(n,\mathbb{C})$, for all $\lambda \in \mathbb{R}$. It satisfies the relation

$$
(q(x) + p\lambda)^* = -(q(x) + p\overline{\lambda}).
$$

Hence $g(x, \lambda, q)$ and $F(\lambda, q)$ satisfy the relations

g

$$
g^*(x, \lambda, q) = g^{-1}(x, \bar{\lambda}, q) \text{ and } F^*(\lambda, q) = F^{-1}(\bar{\lambda}, q), \text{ respectively.}
$$

Then $R(\lambda, \mu) = \det(\mu \mathbf{1} - F(\lambda, q))$ satisfies the relation:

$$
\overline{R\left(\bar{\lambda}, \frac{1}{\bar{\mu}}\right)} = \overline{\det\left(\frac{1}{\bar{\mu}}\mathbb{1} - F(\bar{\lambda}, q)\right)} = \det\left(\frac{1}{\mu}\mathbb{1} - F^{-1}(\lambda, q)\right) =
$$

$$
= \frac{R(\lambda, \mu)}{(-\mu)^n \det(-F(\lambda, q))} = \frac{R(\lambda, \mu)}{(\mu)^n R(\lambda, 0)}.
$$

This shows that $R(\lambda, \mu) = 0$ if and only if $R(\bar{\lambda}, \bar{\mu}^{-1}) = 0$. Hence the map

$$
\theta: Y \to Y, (\lambda, \mu) \mapsto \left(\bar{\lambda}, \frac{1}{\bar{\mu}}\right)
$$

defines an anti-holomorphic involution of the Riemann surface corresponding to $q \in \mathcal{H}_{\mathbb{R}}^{\infty}$. Let us assume in this section that Y corresponds to some $q \in \mathcal{H}_{\mathbb{R}}^{\infty}$. We will see later that this implies the branching divisor of the covering map π to be invariant under the involution θ , in other

words $\theta(b_{\text{effective}}) = b_{\text{effective}}$. For every matrix valued meromorphic function or meromorphic form f let $\theta^*(f)$ denote the function or form defined by

$$
\theta^*(f)(\lambda,\mu) = f^*(\bar{\lambda}, \bar{\mu}^{-1}).
$$

By abuse of notation we will denote the natural isomorphism of the group of divisors of Y induced by θ also by θ .

Theorem 9.3 A potential $q \in \mathcal{H}^{\infty}$ satisfies the reality condition $q^* = -q$ if and only if $\theta(D(q)) = D^{t}(q)$. Moreover, every divisor D of Riemann-Roch type, which obeys the equivalence relation $D + \theta(D) \sim b$ is non-special in the modified sense. In particular

$$
Jacobian_{\mathbb{R}}(Y) = \{ [D] \in Jacobian(Y)|D + \theta(D) \sim b \text{ in the modified sense} \}
$$

is contained in Jacobian₀ (Y) , and T_x acts on this subspace.

Proof: For all $q \in \mathcal{H}^{\infty}$ we have

$$
g^*(x, \lambda, q) = g^{-1}(x, \bar{\lambda}, -q^*),
$$
 and $F^*(\lambda, q) = F^{-1}(\bar{\lambda}, -q^*).$

We assumed that $R(\lambda, \mu) = 0$ if and only if $R(\bar{\lambda}, \bar{\mu}^{-1}) = 0$. Hence μ is an eigenvalue of $F(\lambda, q)$, if and only if $\bar{\mu}^{-1}$ is an eigenvalue of $F(\bar{\lambda}, -q^*)$. If v is any meromorphic solution of

$$
\pi^*(F(\lambda, q))v = v\mu,
$$

then it is also a solution of

$$
\pi^*(F^{-1}(\lambda, q))v = v\frac{1}{\mu},
$$

and $\theta^*(v)$ is a solution of

$$
\theta^*(v)\theta^*(\pi^*(F^{-1}(\lambda, q))) = \theta^*(v)\pi^*(F^{-1^*}(\bar{\lambda}, q)) =
$$

=
$$
\theta^*(v)\pi^*(F(\lambda, -q^*)) = \theta\left(\frac{1}{\mu}\right)\theta^*(v) = \mu\theta^*(v).
$$

If $q^* = -q$, this implies $\theta^*(v) = w$ and $D^t(q) = \theta(D(q))$. Now let $D^t(q)$ be equal to $\theta(D(q))$. Since $\theta^*(\mu^x) = \mu^{-x}$, Theorem [4.10](#page-26-0) implies

$$
D^{t}(T_x q) = T_{-x} D^{t}(q) = T_{-x} \theta(D(q)) = \theta(T_x D(q)) = \theta(D(T_x)).
$$

Then we have

$$
\theta^*(v(T_xq)) = w(T_xq).
$$

Using the inverse of the map $q \mapsto D(q)$, constructed in Section [4,](#page-19-0) this implies $q^* = -q$. Now let D be any divisor of Riemann-Roch type such that

$$
D + \theta(D) - b = (f)
$$

with some meromorphic function f , which is equal to 1 at all covering points of infinity. Due to our assumption on Y , b is equal to

$$
D(q) + \theta(D(q)) + (\theta^*(v(q))v(q))
$$

with some $q \in \mathcal{H}_{\mathbb{R}}^{\infty}$. This shows that

$$
\theta(b) = b \text{ and } \theta^*(f) = f.
$$

Hence f is real on $\pi^{-1}(\mathbb{R})$. Then there exists of course some large $\lambda_0 \in \mathbb{R}$, such that D and b have no contributions at all covering points of λ_0 and the values of f at this covering points are positive. Now let g be any cross section of $\mathcal{O}_{D-\pi^{-1}(\infty)}$. By definition $\frac{\theta^*(g)gf}{\lambda-\lambda_0}$ $\frac{\lambda - \lambda_0}{\lambda - \lambda_0} d\lambda$ is a meromorphic differential form with poles only at the covering points of λ_0 . Hence the residue of this differential form is equal to the sum of the values of $2\pi\sqrt{-1}\theta^*(g)gf$ at the covering points of infinity. Due to Theorem [3.5](#page-17-0) this total residue is zero and therefore g is zero at all covering points of λ_0 . The same is of course true for all $\lambda'_0 \in [\lambda_0 - \epsilon, \lambda_0 + \epsilon]$ with some small $\epsilon > 0$. Hence g is equal to zero. We proved that

$$
H^{0}(Y, \mathcal{O}_{D-\pi^{-1}(\infty)}) \simeq H^{0}(Y, \mathcal{O}_{\theta(D)-\pi^{-1}(\infty)}) \simeq 0.
$$

The asymptotic and total degrees of D and $\theta(D)$ are equal to half the asymptotic and total degree of b. Hence they are equal to $(1, 1, \ldots)$ in the sense of Remark [3.7.](#page-17-0) Due to Serre duality this implies

$$
\tilde{H}^1(Y, \mathcal{O}_{D-\pi^{-1}(\infty)}) \simeq \tilde{H}^1(Y, \mathcal{O}_{\theta(D)-\pi^{-1}(\infty)}) \simeq 0.
$$

Due to Lemma [6.2](#page-35-0) this implies that D and $\theta(D)$ are non-special in the modified sense. As mentioned above, $\theta^*(\mu^x) = \mu^{-x}$, and therefore the action of T_x on *Jacobian*(Y) leaves the subspace $Jacobian_{\mathbb{R}}(Y)$ invariant and $Jacobian_{\mathbb{R}}(Y)$ is contained in $Jacobian_{0}(Y)$. For compact Riemann surfaces the Jacobian variety is isomorphic to the dual space of all holomorphic forms modulo the first homology group of the Riemann surface. Here the first homology group is embedded in the dual space of all holomorphic forms by integration of forms over 1-chains. On the other hand the Jacobian variety is isomorphic to the connected component of the Picard group and the dual space of the holomorphic forms is isomorphic to the Lie algebra of the Picard group. From this point of view the first homology group is isomorphic to the kernel of the exponential map from the Lie algebras of the Picard group into the connected component of the identity of the Picard group. We will see that in our case the same is true, if we restrict ourself to the real part of the Picard group. To do this let us impose the reality condition on the first homology group of our Riemann surface Y . We saw already that the real part of the Lie algebra $H_{\text{modified}}^1(Y, \mathcal{O})$ of the Picard group contains all cocycles [f], which are equal to $[-\theta^*(f)]$. The real part of the Picard group $H_{\text{modified}}^1(Y, \mathcal{O}^*)$ contains all cocycles [f], which are equal to $\frac{1}{\theta^*}$ $\left[\frac{1}{\theta^*(f)}\right]$. The involution θ induces an involution $\theta_{\#}$ of $H_1(\pi^{-1}(\mathbb{C}), \mathbb{Z})$. Hence the lattice of real periods of the Jacobian variety is given by

$$
H_{1,\mathbb{R}}(\pi^{-1}(\mathbb{C}),\mathbb{Z}) = \left\{ x \in H_1(\pi^{-1}(\mathbb{C}),\mathbb{Z}) | \theta_{\#}(x) = -x \right\}.
$$

Since the branching divisor $b_{\text{effective}}$ is equal to the zeroes of $\theta^*(v(q)))v(q)$, the support of $b_{\text{effective}}$ is contained in $\pi^{-1}(\mathbb{C} \setminus \mathbb{R})$ and $\theta(b_{\text{effective}}) = b_{\text{effective}}$. Hence we may build the Riemann surface $\pi^{-1}(\mathbb{C})$ out of n copies of \mathbb{C} by cutting and gluing along small lines, which connect the points (λ_i, μ_i) of $b_{\text{effective}}$ with the corresponding point $(\bar{\lambda}_i, \bar{\mu}_i^{-1})$ of $b_{\text{effective}}$. Hence we may choose generators of $H_{1,\mathbb{R}}(\pi^{-1}(\mathbb{C}), \mathbb{Z})$, which are in one to one correspondence with all pairs $(\lambda_{\iota}, \mu_{\iota}), (\bar{\lambda}_{\iota}, \bar{\mu}_{\iota}^{-1})$ of $b_{\text{effective}}$. Since $b_{\text{effective}}$ has asymptotic and total degree $(2, 2, \ldots)$, we may label such pairs with the index set $\mathcal I$. This shows that $H_{1,\mathbb{R}}(\pi^{-1}(\mathbb{C}),\mathbb{Z})$ is isomorphic to $\bigoplus_{\iota\in\mathcal{I}}\mathbb{Z}a_{\iota}$, where a_{ι} are

the (asymptotically uniquely defined) generators in the excluded domains with index ι . Let us now define elements $\alpha_{\iota} \in \tilde{H}^1_{\text{modified}}(Y, \mathcal{O})$, such that

$$
Res(\alpha_{\iota}, \omega) = \int_{a_{\iota}} \omega \text{ for all } \omega \in H_{\text{modified}}^{0}(Y, \Omega).
$$

Let us fix some $(\lambda_0, \mu_0) \in \pi^{-1}(\mathbb{C})$, which is no branchpoint of the covering map π . Then the function

$$
\frac{R(\lambda_0, \mu)}{(\mu - \mu_0)(\lambda - \lambda_0)R_{\mu}(\lambda_0, \mu_0)}
$$

is meromorphic on $\pi^{-1}(\mathbb{C})$ and has only one pole at (λ_0, μ_0) , such that

$$
\frac{R(\lambda_0, \mu)}{(\mu - \mu_0)(\lambda - \lambda_0)R_{\mu}(\lambda_0, \mu_0)} - \frac{1}{\lambda - \lambda_0}
$$

is holomorphic near (λ_0, μ_0) . Then the function

$$
\frac{1}{2\pi\sqrt{-1}}\int_{a_i}\frac{R(\lambda_0,\mu)}{(\mu-\mu_0)(\lambda-\lambda_0)R_{\mu}(\lambda_0,\mu_0)}d\lambda_0
$$

is holomorphic on the complement of the cycle a_t of $\pi^{-1}(\mathbb{C})$. For all $\omega \in H_{\text{modified}}^0(Y, \Omega)$ the sum of the residues at all covering points of infinity of the product of this function with ω is equal to $\int_{a_i} \omega$. Hence this function defines an element α_i of $\tilde{H}^1_{\text{modified}}(Y, \mathcal{O})$, such that for all $\omega \in H_\text{modified}^0(Y, \Omega)$

$$
Res(\alpha_{\iota}, \omega) = \int_{a_{\iota}} \omega.
$$

Since $\theta_{\#}(a_{\iota}) = -a_{\iota}$ we claim that $\theta^*(\alpha_{\iota}) = \alpha_{\iota}$. In fact, we have

$$
\theta^*(R_\mu(\lambda,\mu)) = -\frac{R_\mu(\lambda,\mu)}{\mu^{n-2}R(\lambda,0)}.
$$

Then the cocycle $\theta^*(\alpha_i)$ can be represented by the function

$$
-\frac{1}{2\pi\sqrt{-1}}\int_{a_{\iota}}\frac{\mu_0^{n-2}}{\mu^n}\frac{R(\lambda_0,0)}{R(\lambda,0)}\frac{R(\lambda_0,\mu)}{(\mu^{-1}-\mu_0^{-1})(\lambda-\lambda_0)R_{\mu}(\lambda_0,\mu_0)}d\lambda_0 =
$$

=
$$
\frac{1}{2\pi\sqrt{-1}}\int_{a_{\iota}}\left(\frac{\mu_0}{\mu}\right)^{n-1}\frac{R(\lambda_0,0)}{R(\lambda,0)}\frac{R(\lambda_0,\mu)}{(\mu-\mu_0)(\lambda-\lambda_0)R_{\mu}(\lambda_0,\mu_0)}d\lambda_0.
$$

Since

$$
\left(\frac{R(\lambda_0,0)}{R(\lambda,0)}-1\right)\frac{R(\lambda_0,\mu)}{(\mu-\mu_0)(\lambda-\lambda_0)}
$$

is a holomorphic function on $\pi^{-1}(\mathbb{C})$, the cocycle $\theta^*(\alpha_i)$ can be represented by the function

$$
\frac{1}{2\pi\sqrt{-1}}\int_{a_{\iota}}\left(\frac{\mu_0}{\mu}\right)^{n-1}\frac{R(\lambda_0,\mu)}{(\mu-\mu_0)(\lambda-\lambda_0)R_{\mu}(\lambda_0,\mu_0)}d\lambda_0.
$$

Since

$$
\left(\left(\frac{\mu_0}{\mu} \right)^{n-1} - 1 \right) \frac{R(\lambda_0, \mu)}{(\mu - \mu_0)(\lambda - \lambda_0)}
$$

is a holomorphic function on $\pi^{-1}(\mathbb{C})$, the cocycle $\theta^*(\alpha_i)$ can be represented by the function

$$
\frac{1}{2\pi\sqrt{-1}}\int_{a_{\iota}}\frac{R(\lambda_0,\mu)}{(\mu-\mu_0)(\lambda-\lambda_0)R_{\mu}(\lambda_0,\mu_0)}d\lambda_0.
$$

This shows that $\theta^*(\alpha_i)$ is equal to α_i . Since

$$
\frac{R(\lambda_0, \mu)}{(\mu - \mu_0)(\lambda - \lambda_0)R_{\mu}(\lambda_0, \mu_0)} - \frac{1}{\lambda - \lambda_0}
$$

is holomorphic near (λ_0, μ_0) , the function

$$
\exp\left(\int_{(\lambda_1,\mu_1)}^{(\lambda_2,\mu_2)} \frac{R(\lambda_0,\mu)}{(\mu-\mu_0)(\lambda-\lambda_0)R_{\mu}(\lambda_0,\mu_0)}d\lambda_0\right)
$$

is on a small open set a solution of the divisor $(\lambda_1, \mu_1) - (\lambda_2, \mu_2)$, if (λ_1, μ_1) and (λ_2, μ_2) are elements of this small open set. If we divide the path of integration into small intervals we see that the same is true for arbitrary (λ_1, μ_1) , $(\lambda_2, \mu_2) \in \pi^{-1}(\mathbb{C})$. Now let $A_{\iota}(t)$ be the line bundle over Y defined by the cocycle $\exp(2\pi\sqrt{-1}t\alpha_t)$ for all $t \in \mathbb{C}$. Then $A_t(t)$ is isomorphic in the modified sense to the trivial line bundle for all $t \in \mathbb{Z}$. For all $t \in \mathbb{C}$ it is an element of $\tilde{H}_{\text{modified}}^1(Y, \mathcal{O}^*)$ and for all $t \in \mathbb{R}$ it is even an element of the real part $\tilde{H}_{\text{modified}, \mathbb{R}}^1(Y, \mathcal{O}^*)$.

The line bundles $A_{\iota}(t)$ have another description, which has the advantage that it admits infinite tensor products. The exact sequence of sheaf homomorphisms

$$
0 \to \mathbb{Z} \to \mathcal{O} \to \mathcal{O}^* \to 1,
$$

where the homomorphism $\mathcal{O} \to \mathcal{O}^*$ is defined by $f \mapsto \exp(2\pi\sqrt{-1}f)$, induces the long exact sequence

$$
0 \to H^0(Y, \mathbb{Z}) \to H^0(Y, \mathcal{O}) \to H^0(Y, \mathcal{O}^*) \to H^1(Y, \mathbb{Z}) \to
$$

$$
\to H^1(Y, \mathcal{O}) \to H^1(Y, \mathcal{O}^*) \to H^2(Y, \mathbb{Z}) \to \dots
$$

This long exact sequence decomposes into

 $0 \to \mathbb{Z} \to \mathcal{O} \to \mathcal{O}^* \to 1$ and

$$
0 \to H^1(Y, \mathbb{Z}) \to H^1(Y, \mathcal{O}) \to H^1(Y, \mathcal{O}^*) \to H^2(Y, \mathbb{Z}) \to \dots
$$

For compact Riemann surfaces, $H^1(Y, \mathcal{O})$ is isomorphic to the Lie algebra of the Picard group $H^1(Y, \mathcal{O}^*)$. From this point of view the kernel of the exponential map of the Lie algebra of the Picard group into the Picard group is isomorphic to $H^1(Y, \mathbb{Z})$. Due to Poincaré duality this first cohomology group is isomorphic to the first homology group. Now we use a special covering of Y, in order to define the elements α_{ι} in $H_{\text{modified}}^1(Y, \mathcal{O})$, which correspond to $a_{\iota} \in H_1(\pi^{-1}(\mathbb{C}), \mathbb{Z})$. Let U₋ be the complement of all the circles $(a_i)_{i\in\mathcal{I}}$ in Y and let $(U_i)_{i\in\mathcal{I}}$ be small pairwise disjoint open tubular neighbourhoods of the circles $(a_i)_{i\in\mathcal{I}}$. The intersection $U_i \cap U_-\$ decomposes into two connected components. Now let $\alpha_i \in H_{\text{modified}}^1(Y, \mathcal{O})$ be the cocycle, which may be represented

by the element of $C^1({U_-, U_i | \iota \in \mathcal{I}}), \mathbb{Z})$, which is zero on $U_{\iota'} \cap U_-$ if $\iota' \neq \iota$ and on one component of $U_i \cap U_-\$ equal to 1 and zero on the other component. The orientation of a_i determines on which component of $U_i \cap U_-\$ the representative of α_i vanishes and on which it is equal to 1. Now $\exp(2\pi\sqrt{-1}\alpha_t)$ is of course a representative of the trivial line bundle in $H^1_{\text{modified}}(Y, \mathcal{O}^*)$. Moreover, for any sequence $(t_i)_{i\in\mathcal{I}} \in \mathbb{R}^{\mathcal{I}} \sum_{\iota\in\mathcal{I}} t_{\iota} \alpha_{\iota}$ defines an element of $H^1_{\text{modified}}(Y, \mathcal{O})$ and $\exp\left(2\pi\sqrt{-1}\sum_{\iota\in\mathcal{I}}t_{\iota}\alpha_{\iota}\right)$ defines a line bundle in $H_{\text{modified}}^1(Y,\mathcal{O}^*)$. Therefore the map

$$
(\mathbb{R}/\mathbb{Z})^{\mathcal{I}} \to H_{\text{modified}}^{1}(Y, \mathcal{O}^{*}), (t_{\iota})_{\iota \in \mathcal{I}} \mapsto \otimes_{\iota \in \mathcal{I}} A_{\iota}(t_{\iota})
$$

defines a group homomorphism from the compact abelian group $(\mathbb{R}/\mathbb{C})^{\mathcal{I}}$ into the real part of the Picard group in the modified sense.

Proposition 9.4 If the excluded domains of some $U_{l,\epsilon}$ have asymptotically no overlap, the actions of \mathbb{R}/\mathbb{Z} on Jacobian $\mathbb{R}(Y)$ defined by the tensor product with $A_{\iota}(t)$ fit together to a continuous action of the compact group $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ on Jacobian $_{\mathbb{R}}(Y)$.

Proof: We use the same arguments as in the proof of Theorem [6.6.](#page-37-0) The action of the tensor product with $A_{\iota}(t)$ does not change $g_{\iota'}$ if $\iota' \neq \iota$ and $|\iota|, |\iota'|$ are large enough. On the cocycle g_{ι} this action is given by

$$
g_{\iota}(t) = Ad\left(\exp\left(-2\pi t \sqrt{-1} \operatorname{diagonal}(n_1,\ldots,n_n)\right)\right)g_{\iota},
$$

where $n_i - n_j = 1$, and all other n's arbitrary, with $\iota = (i, j, k)$. In particular, $||g_{\iota}(t) - 1||_{l, \epsilon, \iota}$ does not depend on t. Due to Proposition [6.5](#page-37-0) the compact group $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ then acts continuously on $Jacobian_{\mathbb{R}}(Y)$.

Theorem 9.5 For all $q, \tilde{q} \in \mathcal{H}_{\mathbb{R}}^{\infty}$, $H_{q, \text{modified}}^1(Y, \mathcal{O})$ is equal to $H_{\tilde{q}, \text{modified}}^1(Y, \mathcal{O})$. Furthermore, there exists a dual basis $(\omega_i)_{i \in \mathcal{I}}$, such that for all $\omega_i \in H_{\text{modified}}^0(Y, \Omega)$ the following holds:

- (i) $\int_{a_{\iota'}} \omega_{\iota} = \delta_{\iota' \iota} = Res(\alpha_{\iota'}, \omega_{\iota}).$
- (ii) $Res([f], \omega_{\iota})$ exists for all $[f] \in H^1_{q, \text{modified}}(Y, \mathcal{O}).$
- (ii) $\omega = \sum_{\iota \in \mathcal{I}} Res(\alpha_i, \omega) \omega_{\iota}$ for all $\omega \in H_{\text{modified}}^0(Y, \Omega)$.
- (iv) $[f] = \sum_{\iota \in \mathcal{I}} Res([f], \omega_{\iota}) \alpha_{\iota}$ for all $[f] \in H^1_{q, \text{modified}}(Y, \mathcal{O}).$

Proof³⁴: For large $|\iota|$, the Riemann surface Y_{ι} decomposes into a two-fold covering of \mathbb{P}_1 and $(n-2)$ copies of \mathbb{P}_1 . Let β_{ι} and $\bar{\beta}_{\iota}$ be the values of λ at the two branchpoints of Y_{ι} . Then the two-fold covering of \mathbb{P}_1 may be described by

$$
\lambda = \frac{\beta_{\iota} - \bar{\beta}_{\iota}}{4\sqrt{-1}} \left(\kappa - \frac{1}{\kappa} \right) + \frac{\beta_{\iota} + \bar{\beta}_{\iota}}{2},
$$

where $\kappa \in \mathbb{P}_1$ gives the parameterization of the two-fold covering, which is equal to 0 and ∞ at the two covering points of infinity and equal to $\pm\sqrt{-1}$ at the two branchpoints. Now let κ_i describe the divisor $D_{\iota}(q)$. Then we claim that for all $q \in \mathcal{H}_{\mathbb{R}}^{\infty}$ the sequences $||\kappa_{\iota}|-1|$ decreases

³⁴Those readers, who are not interested in this proof may skip to Theorem [9.10.](#page-69-0)

faster than every inverse power of $|\iota|$. This is an easy consequence of the following fact: If $v(q) = \pi^*(g_{+\iota})v_{\iota}(q)$ is the Birkhoff factorization of $h(D, \cdot)$ around the excluded domain with index ι (see the proof of the implication (i)⇒ (ii) of Theorem [5.5\)](#page-29-0), $||g_{+i}-1||_{l,\epsilon,\iota}$ decreases faster than every inverse power of $|\iota|$, and the meromorphic function

$$
\theta^*(\pi^*(g_{+i})v_i(q)))\pi^*(g_{+i})v_i(q)
$$

is locally a solution of the divisor $b_{\text{effective}} - D(q) - \theta(D(q))$. In Lemma [8.2](#page-53-0) we showed that the elements of $H^1_{q,\text{modified}}(Y,\mathcal{O})$ are in one to one correspondence to Mittag Leffler distributions, or more precisely, global sections of the sheaf $\mathcal{O}_{D(q)}/\mathcal{O}$. Asymptotically such sections have the form $\frac{c_{\iota}}{\kappa - \kappa_{\iota}}$ on the Riemann surface Y.

Lemma 9.6 The sequence $(c_{\iota})_{\iota \in \mathcal{I}}$ corresponds to an element of $H^1_{q,\text{modified}}(Y, \mathcal{O})$, if and only if $\left|(\beta_{\iota}-\bar{\beta}_{\iota})c_{\iota}\right|$ decreases faster than every inverse power of $|\iota|.$

Proof: Let P_{μ} be the value of the function

$$
P\frac{d\lambda}{d\kappa} = P\frac{(\beta_{\iota} - \bar{\beta}_{\iota})(\kappa^2 + 1)}{4\sqrt{-1}\kappa^2}
$$

at the point κ_i of the divisor D. Since $\frac{d\lambda}{d\kappa}$ is zero at the branchpoints, this function is holomorphic in the excluded domain with index ι , and \underline{P}_{ι} is well defined. Let us now calculate the meromorphic function P_t on the Riemann surface Y_t corresponding to the divisor $D_{\iota}: v_{\iota}$ is given by $\begin{pmatrix} 1 \\ \frac{\kappa}{\iota+1} \end{pmatrix}$ $\kappa + \kappa_{\iota}$ $\kappa-\kappa_{\iota}$ \setminus . The transposed divisor $b_{\iota} - D_{\iota}$ admits two global sections $\left(\frac{(\kappa-\kappa_i)(\kappa-\kappa_i^{-1})}{2(\kappa^2+1)}, \frac{(\kappa-\kappa_i)(\kappa+\kappa_i^{-1})}{2(\kappa^2+1)}\right)$. Then P_ι is equal to

$$
\frac{1}{2(\kappa^2+1)}\left(\begin{array}{cc}(\kappa-\kappa_{\iota})(\kappa-\kappa_{\iota}^{-1}) & (\kappa-\kappa_{\iota})(\kappa+\kappa_{\iota}^{-1}) \\ (\kappa+\kappa_{\iota})(\kappa-\kappa_{\iota}^{-1}) & (\kappa+\kappa_{\iota})(\kappa+\kappa_{\iota}^{-1})\end{array}\right).
$$

Since $|\kappa_i|$ converges to 1 and since the restriction of P to the Riemann surface Y_i converges to P_{ι} , the sequence \parallel P_{ι} $\beta_{\iota}-\overline{\beta_{\iota}}$ \parallel is asymptotically bounded by

$$
\frac{1}{c} \le \left\| \frac{P_{\iota}}{\beta_{\iota} - \bar{\beta}_{\iota}} \right\| \le c \text{ with some } c > 1.
$$

From the proof of Theorem [8.5](#page-55-0) it follows that

$$
\delta v = \pi^* \left(\sum_{\iota} \frac{P_{\iota} c_{\iota}}{\lambda - \lambda_{\iota}} \right) v.
$$

Hence $(c_i)_{i\in\mathcal{I}}$ corresponds to an element $[f]$ of $H^1_{q,\text{modified}}(Y,\mathcal{O})$ if and only if $\sum_i \frac{P_i c_i}{\lambda - \lambda_i}$ $\frac{F_{\mu}c_{\nu}}{\lambda-\lambda_{\nu}}$ is a matrix valued meromorphic function on X . Due to Example [3.8](#page-18-0) this is equivalent to the condition that $|c_{\iota}(\beta_{\iota} - \bar{\beta}_{\iota})|$ decreases faster than every inverse power of $|\iota|$. Continuation of the proof of Theorem [9.5](#page-65-0): Let $([f_{\iota}(q)])_{\iota \in \mathcal{I}}$ denote the basis of $H^1_{q,\text{modified}}(Y,\mathcal{O})$ corresponding to the Mittag Leffler distributions, which vanish outside the excluded domain

with index ι and in this excluded domain is given by $\frac{1}{\kappa - \kappa_{\iota}}$ as used in the foregoing lemma. There exists also a dual basis of $H_{\text{modified}}^0(Y, \Omega)$: Let \underline{v}_t be the value of the function $v \frac{d\kappa}{d\lambda}(\lambda - \lambda_t)$ at the point κ_{ι} . Then we claim that the forms

$$
\omega_{\iota}(q) = \frac{w \underline{v}_{\iota}}{(\lambda - \lambda_{\iota}) w \underline{v}} d\lambda
$$

are dual to the basis $([f_{\iota}(q)])_{\iota \in \mathcal{I}}$ of $H^1_{q,\text{modified}}(Y, \mathcal{O})$. Every form $\omega \in H^0_{\text{modified}}(Y, \Omega)$ can be written as a meromorphic function times $d\lambda$. On the other hand the meromorphic functions w_1, \ldots, w_n are a basis of the meromorphic functions on Y over the meromorphic functions on X. Hence every element $\omega \in H_{\text{modified}}^0(Y, \Omega)$ can be written as a sum

$$
\omega = \sum_{i=1}^{n} \left(\frac{w_i}{wv} \pi^*(f_i) \right) d\lambda
$$
, with meromorphic functions f_i on X.

Since ω may only have poles at b, the vector valued function (f_1, \ldots, f_n) can have poles only at the base points of the divisor D. More precisely, ω must be equal to an infinite sum:

$$
\omega = \frac{1}{2\pi\sqrt{-1}} \sum_{\iota} Res([f_{\iota}(q)], \omega) \omega_{\iota}(q).
$$

On the other hand Lemma [8.2](#page-53-0) shows that

$$
[f] = \frac{1}{2\pi\sqrt{-1}} \sum_{\iota} Res([f], \omega_{\iota}(q))[f_{\iota}(q)] \text{for all } [f] \in H_{q, \text{modified}}^1(Y, \mathcal{O}).
$$

This proves the claim.

Lemma 9.7 Let $q, \tilde{q} \in Isospectral_{\mathbb{R}}(Y)$ and let the excluded domains of $U_{l,\epsilon}$ have asymptotically no overlap. Then the following estimates hold:

$$
|Res([f_{\iota}(q)], \omega_{\tilde{\iota}}(\tilde{q}))| \leq \frac{|\beta_{\iota} - \bar{\beta}_{\iota}|}{|\lambda_{\iota} - \lambda_{\tilde{\iota}}|}
$$

with some $c > 0$ for |ι| and |i| large enough and $\iota \neq \tilde{\iota}$. For $\iota = \tilde{\iota}$ we have

$$
\frac{1}{c} \leq |Res([f_{\iota}(q)], \omega_{\iota}(\tilde{q}))| \leq c.
$$

Proof: By similar arguments as before $||\underline{v}_t||$ is asymptotically bounded

$$
\frac{1}{c} \le ||\underline{v}_{\iota}|| \le c \text{ with some } c > 0.
$$

Then on the excluded domain with index ι the form $\omega_i(\tilde{q})$ becomes nearly equal to

$$
(\beta - \bar{\beta}_t) \frac{\alpha(\kappa - \tilde{\kappa}_t)(\kappa - \tilde{\kappa}_t^{-1}) + \beta(\kappa - \tilde{\kappa}_t)(\kappa - \tilde{\kappa}_t)(\kappa + \kappa_t^{-1})}{8\sqrt{-1}\kappa^2(\lambda - \lambda_{\tilde{\iota}})} d\kappa
$$

with some $\alpha, \beta \in \mathbb{C}$ such that $\frac{1}{c} \leq \sqrt{\|\alpha\|^2 + \|\beta\|^2} \leq c$. This implies

$$
|Res([f_{\iota}(q)], \omega_{\tilde{\iota}}(\tilde{q}))| \leq c \frac{|\beta_{\iota} - \bar{\beta}_{\iota}|}{|\lambda - \lambda_{\tilde{\iota}}|}
$$

if $\iota \neq \tilde{\iota}$. Finally, on the excluded domain with index ι the form $\omega_{\iota}(\tilde{q})$ becomes nearly equal to $\kappa^{-1}d\kappa$. This implies

$$
\frac{1}{c} \le |Res([f_{\iota}(q)], \omega_{\iota}(\tilde{q}))| \le c \text{ for some } c \ge 2\pi.
$$

Continuation of the proof of Theorem [9.5](#page-65-0): If $(c_i)_{i\in\mathcal{I}}$ describes an element of $H^1_{q,\text{modified}}(Y,\mathcal{O})$, the corresponding element of $H^1_{\tilde{q},\text{modified}}(Y,\mathcal{O})$ is given by

$$
\tilde{c}_t = \sum_{\iota \in \mathcal{I}} \frac{1}{2\pi\sqrt{-1}} Res([f_{\iota}(q)], \omega_{\tilde{\iota}}(\tilde{q})) c_{\iota}.
$$

Then the last two lemmata show that $H^1_{q,\text{modified}}(Y, \mathcal{O})$ and $H^1_{\tilde{q},\text{modified}}(Y, \mathcal{O})$ are equal. **Lemma 9.8** For all $\tilde{q} \in \mathcal{H}_{\mathbb{R}}^{\infty}$ and $\iota \neq \tilde{\iota}$

$$
\left| \int_{a_{\iota}} \omega_{\tilde{\iota}}(\tilde{q}) \right| \leq c \frac{|\beta_{\iota} - \bar{\beta}_{\iota}|}{|\beta_{\iota} - \bar{\beta}_{\tilde{\iota}}|}
$$

and finally, for $\iota = \tilde{\iota}$

$$
\left| \int_{a_{\iota}} \omega_{\tilde{\iota}}(\tilde{q}) \right| \geq \frac{1}{c}.
$$

The proof is similar to the proof of the lemma before. \Box

Continuation of the proof of Theorem [9.5:](#page-65-0) With this lemma it is obvious that there exists some $K \geq 0$, and a matrix $(M_{\iota\tilde{\iota}})_{\iota,\tilde{\iota}\in\mathcal{I}, |\iota|,|\tilde{\iota}| \geq K}$ such that

- (i) $\sum_{|\tilde{\iota}| \geq K} M_{\iota \tilde{\iota}} \int_{a_{\iota'}} \omega_{\tilde{\iota}}(\tilde{q}) = \delta_{\iota \iota'}$ for all $|\iota|, |\iota'| \geq K$.
- (ii) $|M_{\iota\iota}| \leq c|\beta_{\iota} \bar{\beta}_{\iota}|$ with some $c > 0$.

Lemma 9.9 Let $\omega \in H_{\text{modified}}^0(Y, \Omega)$ be an element such that $\int_{a_\iota} \omega = 0$ for all $\iota \in \mathcal{I}$. Then ω is equal to zero.

Proof³⁵: It suffices to proof the statement for a form, which satisfies the relation $\theta^*(\omega) = -\omega$. In fact, $\omega = 1/2(\omega + \theta^*(\omega)) + 1/2(\omega - \theta^*(\omega))$ and both forms $\omega - \theta^*(\omega)$ and $\sqrt{-1}(\omega + \theta^*(\omega))$ fulfil this condition. Moreover, we have

$$
\int_{a_{\iota}} \theta^*(\omega) = \overline{\int_{\theta_{\#}(a_{\iota})} \omega} = -\overline{\int_{a_{\iota}} \omega}.
$$

If $b \in H_1(\pi^{-1}(\mathbb{C}), \mathbb{Z})$, such that $\theta_{\#}(b) = b$, the assumption implies

$$
\int_b \theta^*(\omega) = \overline{\int_b \omega} = -\int_b \omega.
$$

This shows that

$$
\Re\left(\int_b \omega\right) = 0 \text{ for all } b \in H_1(\pi^{-1}(\mathbb{C}), \mathbb{Z}).
$$

³⁵Anotherproof of this lemma can be found in [[MK-T-2\]](#page-91-0), which carries over to our situation. In fact, due to our assumption all elements of $H_{\text{modified}}^0(Y, \Omega)$ can be proven to be square integrable.

The condition that $\int_{a_i} \omega = 0$ for all $\iota \in \mathcal{I}$ implies that the residue of ω at all covering points of infinity vanishes and therefore ω is holomorphic even at all covering points of infinity. Hence the function

$$
Y \to \mathbb{R}, (\lambda, \mu) \mapsto \Re \left(\int_{(\infty, \infty)}^{(\lambda, \mu)} \omega \right)
$$

is a harmonic function on Y, which vanishes at (∞, ∞) . Due to the maximum modulus Theorem
for harmonic functions (see [Col) this function is equal to zero. Hence ω is zero too. for harmonic functions (see $|Co|$) this function is equal to zero. Hence ω is zero too. Completion of the proof of Theorem [9.5:](#page-65-0) With the help of the matrices $M_{\tilde{u}}$ introduced above and the last lemma it follows that there exists a basis of holomorphic forms $(\omega_{\iota})_{\iota \in \mathcal{I}}, \omega_{\iota} \in$
 $H_{\text{loc,left}}^0(Y, \Omega)$, which fulfil the conditions (i)-(iv) of the theorem. $H_{\text{modified}}^0(Y, \Omega)$, which fulfil the conditions (i)-(iv) of the theorem.

Theorem 9.10 If the excluded domains of some $U_{l,\epsilon}$ have asymptotically no overlap, the action of $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ on any point $[D] \in Jacobian_{\mathbb{R}}(Y)$ induces a homeomorphism between $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ and Jacobian_R(Y). Moreover, the holomorphic forms $(\omega_{\iota})_{\iota \in \mathcal{I}}$ define an embedding of the real part $H_{\mathbb{R},\text{modified}}^1(Y,\mathcal{O})$ of any $H_{q,\text{modified}}^1(Y,\mathcal{O})$ onto a subspace of $\mathbb{R}^{\mathcal{I}}$, which is mapped under the exponential map onto $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$. Finally, the following diagram commutes:

Corollary 9.11 If the excluded domains of some $U_{l,\epsilon}$ have asymptotically no overlap, the group $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ acts on Isospectral_R (Y) transitively and freely. The action on any potential $q \in Isospectral_{\mathbb{R}}(Y)$ induces a homeomorphism between $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ and Isospectral_R (Y) . Moreover, there exists for any $q \in Isospectral_{\mathbb{R}}(Y)$ an embedding from $\mathcal{L}_{q,\mathbb{R}}$ onto a subspace of $\mathbb{R}^{\mathcal{I}}$, which does not depend on q, such that the flow induced by the Lie algebra element in $\mathbb{R}^{\mathcal{I}}$ corresponding to some $\delta q \in \mathcal{L}_{q,\mathbb{R}}$ is smooth in \mathcal{H}^{∞} , and the derivative at $t = 0$ is equal to δq . This action of $\mathcal{L}_{q, \mathbb{R}}$ on Isospectral_{$\mathbb{R}(Y)$} is also transitive.

In order to prove this theorem we first need a lemma:

Lemma 9.12 For any two divisors $[D], [\tilde{D}] \in Jacobian_{\mathbb{R}}(Y)$, there exists a sequence of 1-chains $(c_{\iota})_{\iota \in \mathcal{I}}$, such that

(i) c_i lies inside the excluded domain with index ι of some neighbourhood of infinity.

- (ii) $\sum_{\iota \in \mathcal{I}} \partial c_{\iota} = \tilde{D} D.$
- (iii) $\sum_{\iota \in \mathcal{I}} \int_{c_{\iota}} \omega_{\tilde{\iota}} \in \mathbb{R}$ for all $\tilde{\iota} \in \mathcal{I}$.

Proof: There exists a meromorphic function f , which is equal to 1 at all covering points of infinity such that

$$
(f) = \tilde{D} + \theta(\tilde{D}) - D - \theta(D).
$$

Asymptotically the equation $f = \frac{x}{r-1}$ $\frac{x}{x-1}$ has for all $x \in [0,1]$ in each excluded domain two solutions. This defines a sequence of 1-chains $(c_i)_{i\in\mathcal{I}}$, which fulfills condition (i) and

$$
\sum_{\iota \in \mathcal{I}} \partial c_{\iota} = \tilde{D} - D', \text{ with some } [D'] \in Jacobian_{\mathbb{R}}(Y),
$$

such that $D' + \theta(D') = D + \theta(D)$. Moreover, the arguments of the proof of Theorem [9.5](#page-65-0) show that condition (iii) is also fulfilled. It remains to prove the statement for $\tilde{D} + \theta(\tilde{D}) = D + \theta(D)$. In this case there exists a natural sequence $(c_i)_{i\in\mathcal{I}}$ of 1-chains, which fulfills condition (i)-(iii): this sequence is on Y_t asymptotically given by

$$
c_{\iota} : [0,1] \to Y_{\iota}, t \to \kappa(t) = \kappa_{\iota} \left(\frac{\tilde{\kappa}_{\iota}}{\kappa_{\iota}}\right)^{t} = \kappa_{\iota} \exp\left(t \ln\left(\frac{\tilde{\kappa}_{\iota}}{\kappa_{\iota}}\right)\right),
$$

where $\ln(\tilde{\kappa}_t/\kappa_t)$ is uniquely defined by $-\pi < \Im(\ln(\tilde{\kappa}_t/\kappa_t)) \leq \pi$. In this case c_t can be chosen to obey the relation $\theta(c_i) = -c_i$. This and again the arguments of the proof of Theorem [9.5](#page-65-0) imply condition (iii).

Proof of Theorem [9.10](#page-69-0): Let us fix some $[D] \in Jacobian_{\mathbb{R}}(Y)$. Due to Proposition [9.4](#page-65-0) the action of $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ on $[D]$ defines a continuous map $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}} \to Jacobian_{\mathbb{R}}(Y)$. On the other hand the last lemma defines a map

$$
Jacobian_{\mathbb{R}}(Y) \to \mathbb{R}^{\mathcal{I}}, [\tilde{D}] \mapsto \left(\sum_{\iota \in \mathcal{I}} \int_{c_{\iota}} \omega_{\tilde{\iota}}\right)_{\tilde{\iota} \in \mathcal{I}}, \text{ with } \sum_{\iota \in \mathcal{I}} \partial c_{\iota} = \tilde{D} - D.
$$

Now we claim that the composition of the map $Jacobian_{\mathbb{R}}(Y) \to \mathbb{R}^{\mathcal{I}}$ with the natural map $\mathbb{R}^{\mathcal{I}} \to (\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ is the inverse of the group action on $[D] : (\mathbb{R}/\mathbb{Z})^{\mathcal{I}} \to \text{Jacobian}_{\mathbb{R}}(Y)$. First we show that this map does not depend on the choice of $(c_i)_{i\in\mathcal{I}}$. If $(\tilde{c}_i)_{i\in\mathcal{I}}$ is another sequence of 1-chains, which fulfills the conditions (i)-(iii) of Lemma [9.12,](#page-69-0) then

$$
\sum_{\iota \in \mathcal{I}} \left(\int_{c_{\iota}} \omega_{\tilde{\iota}} - \int_{\tilde{c}_{\iota}} \omega_{\tilde{\iota}} \right) \in \mathbb{Z}.
$$

In fact, $\partial \sum_{\iota \in \mathcal{I}} c_{\iota} - \tilde{c}_{\iota} = 0$, and

$$
\sum_{\iota \in \mathcal{I}} \left(\int_{c_{\iota}} \omega - \int_{\tilde{c}_{\iota}} \omega + \int_{\theta(c_{\iota})} \omega - \int_{\theta(\tilde{c}_{\iota})} \omega \right) = 0.
$$

Hence $\sum_{\iota \in \mathcal{I}} c_{\iota} - \tilde{c}_{\iota}$ defines an element of $H^1(Y, \mathbb{Z})$, which lies in the eigen space with eigenvalue -1 of the involution $\theta^{\#}$. But these elements are of the form $\prod_{\iota \in \mathcal{I}} \mathbb{Z} a_{\iota}$. This proves

$$
\sum_{\iota \in \mathcal{I}} \left(\int_{c_{\iota}} \omega_{\tilde{\iota}} - \int_{\tilde{c}_{\iota}} \omega_{\tilde{\iota}} \right) \in \mathbb{Z}.
$$

Hence the map $Jacobian_{\mathbb{R}}(Y) \to \mathbb{R}^{\mathcal{I}} \to (\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ does not depend on the choice of $(c_{\iota})_{\iota \in \mathcal{I}}$. Then this map is continuous.

Secondly we prove that the map

$$
(\mathbb{R}/\mathbb{Z})^{\mathcal{I}} \to \text{Jacobian}_{\mathbb{R}}(Y) \to \mathbb{R}^{\mathcal{I}} \to (\mathbb{R}/\mathbb{Z})^{\mathcal{I}}
$$

is the identity map. For any element of $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ there exists an element of $(-1/2, 1/2]^{\mathcal{I}}$, which is mapped onto this element under the natural map

$$
(-1/2,1/2]^{\mathcal{I}} \hookrightarrow \mathbb{R}^{\mathcal{I}} \to (\mathbb{R}/\mathbb{Z})^{\mathcal{I}}.
$$

Now the multiplication with $t \in [0, 1]$ defines a continuous map from 0 to this element of $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ and hence also a continuous map $[0, 1] \rightarrow Jacobian_{\mathbb{R}}(Y)$, which is equal to $[D]$ at $t = 0$ and equal to any $[\tilde{D}]$ in the image of $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}} \to \text{Jacobian}_{\mathbb{R}}(Y)$ at $t = 1$. This defines a sequence $(c_i)_{i\in\mathcal{I}}$ of 1-chains, which fulfills condition (i)-(iii) of Lemma [9.12](#page-69-0). Lemma [9.6](#page-66-0) shows that the infinitesimal generator of this flow can be considered as an element of $H^1_{q,\text{modified}}(Y,\mathcal{O})$, and Theorem [9.10](#page-69-0) (iv) shows that the integration over all $(\omega_{\tilde{i}})_{i\in\mathcal{I}}$ along this path $(c_i)_{i\in\mathcal{I}}$ gives back the original element of $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$. Hence the map

$$
(\mathbb{R}/\mathbb{Z})^{\mathcal{I}} \to \text{Jacobian}_{\mathbb{R}}(Y) \to \mathbb{R}^{\mathcal{I}} \to (\mathbb{R}/\mathbb{Z})^{\mathcal{I}}
$$

is the identity map.

Let us finally also prove that the map

$$
Jacobian_{\mathbb{R}}(Y) \to \mathbb{R}^{\mathcal{I}} \to (\mathbb{R}/\mathbb{Z})^{\mathcal{I}} \to Jacobian_{\mathbb{R}}(Y)
$$

is the identity map³⁶. Again we want to distinguish between the two cases of the proof of Lemma [9.12.](#page-69-0) If $\tilde{D} + \theta(\tilde{D})$ is equal to $D + \theta(D)$, there exists of course a sequence $(D_n)_{n \in \mathbb{N}}$ of integral divisors of asymptotic and total degree $(1, 1, \ldots)$ such that

- (i) $D_n + \theta(D_n) = D + \theta(D)$ for all $n \in \mathbb{N}$.
- (ii) $D D_n$ is a finite divisor of degree zero.
- (iii) $([D_n])_{n\in\mathbb{N}}$ converges to $[D]$.

The divisor $D - D_n$ corresponds to an element of $\tilde{H}^1_{\mathbb{R}$, modified (Y, \mathcal{O}^*) . Moreover it corresponds to an element of the Lie algebra $\tilde{H}^1_{\mathbb{R}$, modified(Y, O). Theorem [9.5](#page-65-0) (iv) implies that for all $n \in \mathbb{N}$, $[D_n]$ is mapped onto $[D_n]$ under the map given above. Hence the same is true for $[\tilde{D}]$. In the other case there exists a continuous map

$$
[0,1] \to Jacobian_{\mathbb{R}}(Y), t \mapsto [D(t)],
$$

which is mapped onto $[D]$ for $t = 0$ and which is mapped onto $[\tilde{D}]$ for $t = 1$. The derivative of this map, which was constructed in the proof of Lemma [9.12](#page-69-0) is furthermore a map $[0, 1] \rightarrow$ $\tilde{H}^1_{\mathbb{R},\text{modified}}(Y,\mathcal{O})$. This derivative may be integrated due to Lemma [9.6](#page-66-0) to a map $[0,1] \rightarrow$ $\tilde{H}^1_{\mathbb{R},\text{modified}}(Y,\mathcal{O})$. The composition of this map $[0,1] \to \tilde{H}^1_{\mathbb{R},\text{modified}}(Y,\mathcal{O})$ with the exponential map $\tilde{H}^1_{\mathbb{R},\text{modified}}(Y,\mathcal{O}) \to \tilde{H}^1_{\mathbb{R},\text{modified}}(Y,\mathcal{O}^*)$ gives the map

 $[0,1] \to \tilde{H}_{\mathbb{R}$, modified $(Y,\mathcal{O}^*), t \mapsto \mathcal{O}$ line bundle corresponding to $D(t) - D$.

³⁶This is equivalent to Abel's Theorem for those divisors, which obey the reality condition. In general the analogous statement is more complicated. It can be proven by methods similar to those used in this section. One half can be proven with the methods of Lemma [9.12](#page-69-0) (compare with [[MK-T-2,](#page-91-0) Section 10]).
9 A REALITY CONDITION 72

Now again Theorem [9.5](#page-65-0) implies that for all $t \in [0, 1]$, $[D(t)]$ is mapped onto $[D(t)]$ under the above map. Due to Lemma [9.12](#page-69-0) the combination of these two cases cover the general case and the group action on $[D]$ defines a homeomorphism $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}} \simeq Jacobian_{\mathbb{R}}(Y)$. The rest of Theorem [9.10](#page-69-0) follows from Theorem [9.5](#page-65-0). \Box

Corollary [9.11](#page-69-0) is a direct consequence of Theorem [9.10](#page-69-0). \Box

The assumption in the last two theorems that the excluded domains have asymptotically no overlap, can be weakened. In fact, the arguments we gave can be used to prove these theorems in more general cases. But let us indicate, why this assumption must not be dropped. If $p_1(\lambda) - p_2(\lambda)$ and $p_1(\lambda) - p_3(\lambda)$ are nearly elements of $2\pi\sqrt{-1}\mathbb{Z}$, $p_2(\lambda) - p_3(\lambda)$ of course is also nearly an element of $2\pi\sqrt{-1}\mathbb{Z}$. In the case $n = 3$, the Riemann surface Y_t corresponding to the overlapping excluded domain is a three-fold covering of \mathbb{P}_1 with 6 branchpoints. In the λ -plane the cuts and branchpoints may be chosen e.g. like in figure 2, where $\{\lambda_{i,j}, \bar{\lambda}_{i,j}\}$ are the branchpoints between the *i*-th and *j*-th sheet over \mathbb{P}_1 .

If for example $\lambda_{1,2} = \lambda_{1,3}$ the Riemann surface is still non-singular. But if furthermore $\lambda_{2,3}$ becomes equal to $\lambda_{1,2} = \lambda_{1,3}$ as indicated in figure 3, the Riemann surface becomes singular.

It will be one two-fold covering of \mathbb{P}_1 and one copy of \mathbb{P}_1 connected by two ordinary double points. Such a surface is described by the algebraic equation

$$
R(\lambda, \mu) = (\mu - \lambda)(\mu^2 - 2\mu\lambda - 1) = \mu^3 - 2\mu^2\lambda + \mu(2\lambda^2 - 1) + \lambda = 0.
$$

Indeed, $\mu^2 - 2\mu\lambda - 1 = 0$ describes a two-fold covering $\lambda = \frac{\mu^2 - 1}{2\mu}$, with branchpoints at $(\lambda_1, \mu_1) = (\sqrt{-1}, \sqrt{-1}), (\lambda_2, \mu_2) = (-\sqrt{-1}, -\sqrt{-1})$ and $\frac{d\lambda}{d\mu} = \frac{2(\mu^2+1)}{\mu^2}$ $\frac{f^2+1}{\mu^2}$. The real part of the completion of the generalized Jacobian variety³⁷ of this Riemann surface turns out to decompose into two components: One non-compact three dimensional group isomorphic to $\mathbb{R} \times S^1 \times S^1$ corresponding to $b_{\text{effective}} = 3(\sqrt{-1}, \sqrt{-1}) + 3(-\sqrt{-1}, -\sqrt{-1})$ and a compact component isomorphic to S^1 corresponding to $b_{\text{effective}} = (\sqrt{-1}, \sqrt{-1}) + (\sqrt{-1}, \sqrt{-1})$. If the assumption that the excluded domains have asymptotically no overlap is dropped, this example illustrates that the perturbation of the different Riemann surfaces to each other cannot be estimated. Furthermore, in that case certain holomorphic forms are not square integrable.

10 The singular case

In this section we want to investigate, how the statements of the foregoing sections can be generalized to the singular case in the sense of Definition [4.2.](#page-21-0) Let us fix some equation $R(\lambda, \mu)$ 0, which describes the curve of eigenvalues corresponding to some potential $q \in \mathcal{H}^{\infty}$. To this curve there corresponds a unique normalization Y, which is an n-fold covering of $\mathbb{P}_1, \pi : Y \to \mathbb{P}_1$. Due to the counting lemma all zeroes of $R_u(\lambda, \mu)$ can be arranged such that asymptotically and totally to each element of $\mathcal I$ there corresponds one pair of zeroes in the excluded domain with index $i \in \mathcal{I}$. In general not all zeroes of $R_{\mu}(\lambda, \mu)$ are indeed branchpoints of the normalization $\pi: Y \to \mathbb{P}_1$. But there always exists some asymptotically and totally unique subset $\mathcal{I}_{\text{analytic}} \subset$ I , such that to each element of I_{analytic} there corresponds one pair of branch points of the covering map $\pi: Y \to \mathbb{P}_1$. In Section [4](#page-19-0) we already mentioned that the algebraic curve defined by $R(\lambda, \mu) = 0$ and the normalization Y are the extreme cases of the curves, which really

³⁷This completion of the generalized Jacobian variety is defined in the next section.

correspond to some potential $\tilde{q} \in \mathcal{H}^{\infty}$ corresponding to the curve of eigenvalues defined by the equation $R(\lambda, \mu) = 0$. Let \mathcal{O}_{μ} be the free sheaf of rings on $\mathbb{C} \subset \mathbb{P}_1$

$$
\mathcal{O}_{\mathbb{P}_1}^n \simeq \mathcal{O}_{\mu}, (f_1, \ldots, f_n) \mapsto \sum_{i=1}^n f_i \mu^{i-1}.
$$

Due to the equation $R(\lambda, \mu) = 0$, \mathcal{O}_{μ} is in fact a sheaf of rings. On the other hand $\pi_*(\mathcal{O}_Y)$ also defines a sheaf of rings on $\mathbb{C} \subset \mathbb{P}_1$. Moreover, since μ is a holomorphic function on $\pi^{-1}(\mathbb{C})$, \mathcal{O}_{μ} is a subsheaf of $\pi_*(\mathcal{O}_Y)$ and both sheaves contain $\mathcal{O}_{\mathbb{P}_1}$ as a ring subsheaf. Now let S be the set of all ring subsheaves, which contain \mathcal{O}_{μ} as a ring subsheaf. To each element of S, there corresponds a singular Riemann surface Y' with normalization $p: Y \to Y'$, such that the following diagram commutes and $\pi_*(\mathcal{O}_{Y'})$ is the corresponding subsheaf in S.

By abuse of notation we will sometimes identify Y' with the corresponding element of S . In general divisors are defined to be global sections of the sheaf $\mathcal{M}^*/\mathcal{O}^*$. Hence they define locally free submodules of the meromorphic functions. By abuse of notation we will use divisors to define finitely generated submodules of the meromorphic functions: For any singular Riemann surface Y' with normalization $p: Y \to Y'$ such that the foregoing diagram commutes let the branching divisor $b_{\text{effective}}$ be defined by the property that $\mathcal{O}_{b_{\text{effective}}}$ is the $\mathcal{O}_{Y'}$ submodule of meromorphic functions defined by:

$$
\pi'_*(\mathcal{O}_{b_{\text{effective}}}) = \left\{ g \in \pi'_*(\mathcal{M}) \, \middle| \, \sum_{\text{sheets of } \pi'} gf \in \mathcal{O}_{\mathbb{P}_1} \text{ for all } f \in \pi'_*(\mathcal{O}_{Y'}) \right\}.
$$

In general $\mathcal{O}_{b_{\text{effective}}}$ is not locally free, but the two extreme cases $\mathcal{O}_{b_{\text{analytic}}}$ as well as $\mathcal{O}_{b_{\text{algebraic}}}$ are locally free. It is quite easy to see that $\mathcal{O}_{Y'}$ can be reconstructed from the sheaf $\mathcal{O}_{b_{\text{effective}}}$:

$$
\mathcal{O}_{Y'} = \left\{ f \in p_*(\mathcal{O}_Y) | fg \in \mathcal{O}_{b_{\text{effective}} \text{ for all } g \in \mathcal{O}_{b_{\text{effective}}} \right\}.
$$

We will later give an example where S is not a countable set, but for $n = 2$ the situation is quite simple:

Proposition 10.1 If $n = 2$ locally there are only two cases

- (i) $b_{\text{analytic}} = y_0, b_{\text{algebraic}} = (2m+1)y_0, m \in \mathbb{N}$
- (ii) $b_{\text{analytic}} = 0$, $b_{\text{algebraic}} = my_1 + my_2$, with $\pi(y_1) = \pi(y_2)$.

In both cases $\mathcal{O}_{b_{\text{effective}}}$ is locally free.

In case (i) beflective is of the form beflective $=(2m'+1)y_0$, for $0 \le m' \le m$.

In case (ii) b_{effective} is of the form b_{effective} = $m'(y_1 + y_2)$, for $0 \le m' \le m$.

More generally, all finitely generated submodules $\mathcal F$ of $p_*(\mathcal{M}_Y)$ are locally free, if

$$
\{f \in p_*(\mathcal{O}_Y) | fg \in \mathcal{F} \text{ for all } g \in \mathcal{F}\} = \mathcal{O}_{Y'}.
$$

Proof: We may assume that $R(\lambda, \mu)$ has the form $\mu^2 = a(\lambda)$ with some entire function $a(\lambda)$. Hence the equation $R(\lambda, \mu) = 0$ is locally equivalent to $\mu^2 = (\lambda - \lambda_0)^k$. Now there are two cases:

- (i) $\mu^2 = (\lambda \lambda_0)^{2m+1}$. Then b_{analytic} is equal to $\pi^{-1}(\lambda_0)$ and $b_{\text{algebraic}}$ equal to $(2m+1)\pi^{-1}(\lambda_0)$. Moreover, in this case \mathcal{O}_{μ} is generated as an $\mathcal{O}_{\mathbb{P}_1}$ module by 1 and μ and $\pi_*(\mathcal{O}_Y)$ is generated as an $\mathcal{O}_{\mathbb{P}_1}$ module by 1 and $\mu(\lambda - \lambda_0)^{-m}$. For any $\mathcal{O}_{\mathbb{P}_1}$ module $\mathcal{O}_{Y'}$, which is contained in $\pi_*(\mathcal{O}_Y)$ and contains \mathcal{O}_μ , there exists an integer $0 \leq m' \leq m$ such that this module is generated by 1 and $\mu(\lambda - \lambda_0)^{-m'}$. The corresponding module $\mathcal{O}_{b_{\text{effective}}}$ is the free $\mathcal{O}_{Y'}$ module generated by $\mu(\lambda - \lambda_0)^{m-m'-1}$ and the divisor $b_{\text{effective}}$ is equal to $b_{\text{effective}} = (2m' + 1)\pi^{-1}(\lambda_0).$
- (ii) $\mu^2 = (\lambda \lambda_0)^{2m+1}$. Then $b_{\text{analytic}} = 0$ and $\pi^{-1}(\lambda_0)$ consists of two points y_1 and y_2 . b_{algebraic} is equal to $m(y_1 + y_2)$. Again \mathcal{O}_{μ} and $\pi_*(\mathcal{O}_Y)$ are generated as $\mathcal{O}_{\mathbb{P}_1}$ modules by 1, μ and 1, $\mu(\lambda - \lambda_0)^{-m}$, respectively. Any $\mathcal{O}_{\mathbb{P}_1}$ module $\mathcal{O}_{Y'}$ is again generated by 1 and $\mu(\lambda - \lambda_0)^{-m'}$, with some $0 \leq m' \leq m$. The corresponding module $\mathcal{O}_{b_{\text{effective}}}$ is the free $\mathcal{O}_{Y'}$ module generated by $\mu(\lambda - \lambda_0)^{m-m'-1}$ and the divisor $b_{\text{effective}}$ is equal to $b_{\text{effective}} = m'(y_1 + y_2).$

The same arguments show that all finitely generated submodules $\mathcal F$ are locally of the same form as $\mathcal{O}_{b_{\text{effective}}}$ with $m' \in \mathbb{Z}$ if

$$
\{f \in p_*(\mathcal{O}_Y) | fg \in \mathcal{F} \text{ for all } g \in \mathcal{F}\} = \mathcal{O}_{Y'}.
$$

 \Box

Now we explain the modification of Section [4](#page-19-0) in case of singular Riemann surfaces. In that section we investigated the dual eigen bundle of some potential q . Due to Definition [4.2](#page-21-0) the potential q completely determines the singular Riemann surface, which corresponds to this potential. In this context we will take for granted that Y' corresponds to q , or equivalently that $\varepsilon'_q : \mathcal{O}_q \to \pi'_*(\mathcal{O}_{Y'})$ is an isomorphism of sheaves. For singular Riemann surfaces the sheaves $\mathcal{O}_{D(q)}$ and $\mathcal{O}_{D^t(q)}$ have to be defined in a slightly different way: The meromorphic functions on Y and Y' coincide, hence v and w are also meromorphic functions on Y'. Now let $\mathcal{O}_{D(q)}$ and $\mathcal{O}_{D^t(q)}$ be the $\mathcal{O}_{Y'}$ submodules of $p_*(\mathcal{M}_Y)$, which are generated by v_1, \ldots, v_n and w_1, \ldots, w_n , respectively. Then Theorem [4.6](#page-23-0) is still valid. Since the singular Riemann surface Y' corresponds to q, or equivalently $\varepsilon'_q: \mathcal{O}_q \to \pi'_*(\mathcal{O}_{Y'})$ is an isomorphism, f is locally an element of $\pi'(\mathcal{O}_{Y'})$ if and only if $\sum_{\text{sheets of }\pi} fP$ is holomorphic. This implies that the entries of P generate the $\mathcal{O}_{Y'}$ module $\mathcal{O}_{b_{\text{effective}}}$. This generalizes statement (iv) of Lemma [4.5.](#page-22-0) Lemma [4.5](#page-22-0) (i)-(iii) remains valid. Due to an easy exercise the degree function extends to a unique function

deg : finitely generated submodules of $p_*(\mathcal{M}_Y) \to \mathbb{Z}$,

which obeys the two properties

(i) If $\mathcal{F} \subseteq \mathcal{F}'$ are two finitely generated submodules of $p_*(\mathcal{M}_Y)$

$$
\deg(\mathcal{F}') - \deg(\mathcal{F}) = \dim H^0(Y', \mathcal{F}'/\mathcal{F}).
$$

(ii) If D is a divisor of Y' in the correct sense, $deg(D) = deg(\mathcal{O}_D)$.

Moreover the Riemann-Roch Theorem can be generalized to this context: In fact, let $\pi': Y' \to \mathbb{P}_1$ be a singular *n*-fold covering over \mathbb{P}_1 with normalization $p: Y \to Y'$, such that the following diagram commutes.

Let F be any finitely generated $\mathcal{O}_{Y'}$ submodule of $p_*(\mathcal{M}_Y)$ and $\mathcal{O}_{b_{\text{effective}}}$ be defined as above, then the following relation holds:

$$
\dim H^{0}(Y', \mathcal{F}) - \dim H^{1}(Y', \mathcal{F}) = n + \deg(\mathcal{F}) - \frac{\deg(\mathcal{O}_{b_{\text{effective}}})}{2}.
$$

Since $\pi_*(\mathcal{O}_{b_{\text{analytic}}}) \subseteq \pi_*(\mathcal{O}_{b_{\text{effective}}}) \subseteq \pi_*(\mathcal{O}_{b_{\text{algebraic}}})$ there exists a subset $\mathcal{I}_{\text{effective}}$ of \mathcal{I} , which contains $\mathcal{I}_{\text{algebraic}}$, such that to each element of $\mathcal{I}_{\text{effective}}$ there corresponds a pair of degrees of $\mathcal{O}_{b_{\text{effective}}}$. The degree of a finitely generated submodule of $p_*(\mathcal{M}_Y)$ is a sequence of numbers indexed by $\mathcal{I}_{\text{effective}}$ in the sense of Remark [3.7.](#page-17-0) With this modification Theorem [4.6](#page-23-0) is valid also in the general case and relation [\(10](#page-23-0)) generalizes to the relation

$$
\mathcal{O}_{D(q)}\otimes\mathcal{O}_{D^t(q)}\simeq \mathcal{O}_{b_\text{effective}},
$$

where the isomorphism is given by multiplication with the function wv . The rest of Section [4](#page-19-0) can be carried over to the general case in an obvious way.

In Section [5](#page-26-0) the divisor D should be considered in the general case as a finitely generated submodule of $p_*(\mathcal{M}_Y)$. Moreover, the branching divisor has to be specified as the module $\mathcal{O}_{b_{\text{effective}}}$ defined above, and the degree of a divisor is in the general case a sequence indexed by $\mathcal{I}_{\text{effective}}$ and not by \mathcal{I} . With this modification Section [5](#page-26-0) is also true in the general case. In particular equation [\(13](#page-34-0)) is also true in the singular case $\lambda_1 = \lambda_2$, and therefore Corollary [5.11](#page-34-0) remains valid too.

In the general case Definition [6.1](#page-35-0) has to be more precise:

Definition 10.2 The Jacobian variety of a singular Riemann surface $Y' \in \mathcal{S}$ is defined to be the set of all equivalence classes of finitely generated submodules \mathcal{O}_D of $p_*(\mathcal{M}_Y)$ such that

- (i) $\{f \in p_*(\mathcal{O}_Y) | f\mathcal{O}_D \subset \mathcal{O}_D\} = \mathcal{O}_{Y'}.$
- (ii) \mathcal{O}_D is of Riemann-Roch type.
- (iii) \mathcal{O}_D is integral; this means that \mathcal{O}_D contains $\mathcal{O}_{Y'}$.
- (iv) The asymptotic and total degree of \mathcal{O}_D is equal to $(1,1,\ldots)$.

We want to emphasize that this definition is not analogous to the generalized Jacobian variety defined in [\[Se](#page-91-0)]. In fact, we do not assume that \mathcal{O}_D is locally free³⁸. We will give an example of such modules, which are not locally free. Such an integral divisor \mathcal{O}_D is called non-special in the modified sense, if there exists only one integral divisor, which is equivalent in the modified sense to \mathcal{O}_D . Lemma [6.2](#page-35-0) is valid also in the general case. Furthermore, Theorem [6.6](#page-37-0) holds even for the Jacobian variety $Jacobian(Y')$ of a singular Riemann surface. In the proof of this theorem the assumption (i) on cocycles, which correspond to line bundles on Y ′ has to be replaced in the general case by the condition

Cocycle (i)' for all $\iota \in \mathcal{I}_{\text{effective}}$ and for all holomorphic sections f of $\mathcal{O}_{Y'}$ over $\pi^{-1}(U_{\iota})$ there exists a holomorphic function $g_{+,l}: U_l \to n \times n$ -matrices, such that

 $g_{+,l}g_l = g_l \text{diagonal}(f_1, \ldots, f_n),$

where f_i is the restriction of f to the *i*-th sheet of $\pi^{-1}(U_i \cap U)$.

In general let $Isospectral(Y')$ be the subspace of all potentials, which correspond to the singular Riemann surface Y' in the sense of Definition [4.2](#page-21-0). Theorem [6.9](#page-40-0) holds in the general case.

The space of all regular forms $H_{\text{modified}}^0(Y', \Omega)$ on the singular Riemann surface Y' is equal to the vector space of all sections of $\mathcal{O}_{b_{\text{effective}}-\pi^{-1}(\infty)}$ times $d\lambda$. This is analogous to the definition of regular forms in [\[Se](#page-91-0)]. Now let q correspond to the singular Riemann surface Y' . Then the form $\frac{1}{\mu}$ $\frac{\partial \mu}{\partial q}(\delta q) d\lambda$ is an element of $H_{\text{modified}}^0(Y', \Omega)$ for all $\delta q \in T_q{\mathcal{H}}^{\infty}$. In fact, let f be a local section of $\pi'_*(\mathcal{O}_{Y'})$. Then $a_f = \sum_{\text{sheets of }\pi'} v(q) f w(q)$ is a holomorphic $n \times n$ -matrix valued function. Moreover, the following relation holds:

$$
\sum_{\text{sheets of }\pi'} \frac{1}{\mu} \frac{\partial \mu}{\partial q} (\delta q) f = tr \left(a_f F^{-1}(\cdot, q) \frac{\partial F(\cdot, \lambda)}{\partial q} (\delta q) \right).
$$

Hence the left hand side is holomorphic for all local sections of $\pi'_{*}(\mathcal{O}_{Y'})$. This shows that in general Ω_q is a map from $T_q\mathcal{H}^\infty$ into $H^0_{\text{modified}}(Y', \Omega)$.

The Darboux coordinates of singular potentials are more complicated: For a singular potential q let $D(q)$ be the sum of the divisor, which is given by the support of the sheaf $\mathcal{O}_{D(q)}/\mathcal{O}_{Y'}$ with multiplicity equal to the local dimension of this sheaf, plus the divisor, which is given by the support of the sheaf $\mathcal{O}_{b_{\text{algebraic}}}/\mathcal{O}_{b_{\text{effective}}}$ with multiplicity equal to half the local dimension of this sheaf. Then Lemma [7.2](#page-45-0) remains valid. But the first statement of Theorem [7.3](#page-46-0) is true only if $\mathcal{O}_{D(q)}$ is a locally free sheaf. In fact, in this case the proof of Theorem [7.3](#page-46-0) carries over. Now we want to give an example, in which the first statement of this theorem is false.

Example 10.3 Let $n = 3, p = diagonal(1, 0, -1)$,

$$
q = \left(\begin{array}{ccc} 0 & 0 & a \\ 0 & 0 & b \\ 0 & 0 & 0 \end{array}\right)
$$

with two constants a and b not depending on x. The corresponding spectral curve is defined by the equation

$$
R(\lambda, \mu) = (\mu - \exp(-\lambda))(\mu - 1)(\mu - \exp(\lambda)) = 0.
$$

³⁸Our definition is similar to the compactification of the generalized Jacobian variety with the help of torsion free sheaves (see[[DS\]](#page-90-0)).

It is quite obvious that the normalization Y of this singular Riemann surface is the disjoint union of three copies of \mathbb{P}_1 . Moreover, $\nu = \ln(\mu)$ is a holomorphic function on this normalization Y. It is easy to see that the singular Riemann surface Y' corresponding to q may be described as follows:

Let \tilde{Y} be the algebraic curve defined by the equation

$$
(\nu - \lambda)\nu(\nu + \lambda) = 0
$$

and $\mathbb{P}_1 \cup \mathbb{P}_1$ the corresponding normalization. Now the sheaf $\pi_*(\mathcal{O}_Y)$ can be identified with the $\mathcal{O}_{\mathbb{P}_1}$ module, which is generated by $1, \nu/\lambda, \nu^2/\lambda^2$. Then the direct image of the structure sheaf of Y' is the $\mathcal{O}_{\mathbb{P}_1}$ module generated by $1, \nu, \nu^2/\lambda$. The degree of $\mathcal{O}_{b_{\text{effective}}}$ is equal to 4. The sheaf $\mathcal{O}_{D(q)}$ is generated as an $\mathcal{O}_{Y'}$ module by

$$
1, b\frac{\nu}{\lambda} + (a+b)\frac{\nu^2}{\lambda^2}, \frac{\nu(\nu-\lambda)}{2\lambda - a - 2b}.
$$

Hence the first summand of the divisor $D(q)$ defined above is equal to

$$
(0,1)+\left(\frac{a+2b}{2}, \exp\left(\frac{a+2b}{2}\right)\right).
$$

The second summand of the divisor does not depend on a and b. This shows that the Darboux coordinates of these potentials are the same, if the value of $a + 2b$ is the same. Furthermore, the $\mathcal{O}_{Y'}$ module $\mathcal{O}_{D(q)}$ is not locally free at the singular point with $\lambda = 0$.

In the eighth section Lemma [8.2](#page-53-0) may be extended in the obvious way to the general situation. Theorem [8.5](#page-55-0) and its proof generalizes to the case in which $\mathcal{O}_{b_{\text{effective}}}$ is equal to $\mathcal{O}_{b_{\text{algebraic}}}$. The statements (ii) and (iii) of Theorem [8.5](#page-55-0) are valid for all potentials q and the corresponding possibly singular Riemann surfaces Y ′ . The other statements must be replaced by

- (iv)' Let \mathcal{L}_q^{\perp} be the subspace of $T_q\mathcal{H}^{\infty}$ of all elements δq , such that the symplectic form of δq with all elements of \mathcal{L}_q vanishes. Then \mathcal{L}_q^{\perp} is contained in \mathcal{L}_q .
- (i)' There exists a homomorphism³⁹ of vector spaces $d\Gamma_q: H^1_{q,\text{modified}}(Y',\mathcal{O}) \to \mathcal{L}_q^{\perp}$ which is uniquely determined by the property that for all $[f] \in H_{q,\text{modified}}^1(Y',\mathcal{O})$

$$
\frac{\partial v}{\partial q}(d\Gamma_q([f]) = \delta v,
$$

where δv was defined in Lemma [8.2.](#page-53-0)

 39 If $n = 2$ this homomorphism can be proven to be an isomorphism. In general this is not true, because locally there may exist additional flows (compare with Example [10.7](#page-80-0).) Moreover, in some sense the codimension of the image of the map $d\Gamma_q$ in \mathcal{L}_q is equal to the degree of $\mathcal{O}_{b_{\text{algebraic}}}$ minus the degree of $\mathcal{O}_{b_{\text{effective}}}$.

In the ninth section Theorem [9.3](#page-61-0) may be generalized in the obvious way.

Theorem 10.4 Let q satisfy the reality condition $q^* = -q$. Moreover, assume that the excluded domains of some $U_{l,\epsilon}$ of the corresponding Riemann surface Y' have asymptotically no overlap. Then this Riemann surface Y' has asymptotically no singularities; this means that the effective branching divisor is asymptotically equal to the analytic branching divisor.

Proof: Due to the assumption that the excluded domains have asymptotically no overlap, over some neighbourhood U of $\lambda = \infty$ of \mathbb{P}_1 the Riemann surface Y' can only have ordinary double points as singular points. More precisely, two points of Y , which are identified in Y' must be covering points of the same $\lambda \in \mathbb{C}$. Since the effective branching divisor $b_{\text{effective}}$ is a fixed point of θ , the value of λ at such an ordinary double point must be real. On the other hand the equation

$$
b_{\text{effective}} = D(q) + \theta(D(q)) + (\theta^*(v(q))v(q))
$$

shows that on $\pi^{-1}(\mathbb{R})$ all multiplicities of $b_{\text{effective}}$ are even. This proves that Y' has asymptotically no singular points. \Box

Due to this theorem in the rest of the ninth section the singular Riemann surface Y' may be considered as a non-singular Riemann surface over some neighbourhood U of $\lambda = \infty$ of \mathbb{P}_1 together with finitely many singularities over $\mathbb{P}_1 \setminus U$. Proposition [9.4](#page-65-0) and Theorem [9.5](#page-65-0) remain valid, if $\mathcal I$ is replaced by $\mathcal I_{\text{effective}}$. In general, all statements and proofs of the ninth section may be extended to the case $\mathcal{O}_{b_{\text{effective}}}$ being equal to $\mathcal{O}_{b_{\text{analytic}}}$. Then it is obvious that the statements of Theorem [9.10](#page-69-0) and Corollary [9.11](#page-69-0) generalize to:

Theorem 10.5 If the excluded domains of some $U_{l,\epsilon}$ have asymptotically no overlap, the action of $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ effective on any point $[D] \in Jacobian_{\mathbb{R}}(Y')$ induces a homeomorphism between $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ effective times finitely many copies of a finite dimensional abelian Lie group and Jacobian_R(Y'). Moreover, the holomorphic forms $(\omega_{\iota})_{\iota \in \mathcal{I}}$ define an embedding of the real part $H^1_{\mathbb{R}$, modified (Y, \mathcal{O}) of any $H^1_{q, \text{modified}}(Y, \mathcal{O})$ onto a subspace of $\mathbb{R}^{\mathcal{I}}$ effective, which is mapped under the exponential map onto $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ effective.

Corollary 10.6 If the excluded domains of some $U_{l,\epsilon}$ have asymptotically no overlap the group $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ effective times a finite dimensional abelian Lie group acts on each connected component of $Isospectral_{\mathbb{R}}(Y')$ transitively and freely. Moreover, the action on any potential $q \in Isospectral_{\mathbb{R}}(Y)$ induces a homeomorphism between the connected components of Isospectral_R (Y') and $(\mathbb{R}/\mathbb{Z})^{\mathcal{I}}$ effective times a finite dimensional abelian Lie group. Finally, for any $q \in Isospectral_{\mathbb{R}}(Y)$ there exists an embedding from $\mathcal{L}_{q,\mathbb{R}}^{\perp}$ onto a subspace of $\mathbb{R}^{\mathcal{I}}$ effective times the Lie algebra of the finite dimensional abelian Lie group mentioned above, and which does not depend on q, such that the flow induced by the Lie algebra element in $\mathbb{R}^{\mathcal{I}}$ effective times the finite dimensional abelian Lie algebra corresponding to some $\delta q \in \mathcal{L}^{\perp}_{q,\mathbb{R}}$ is smooth in \mathcal{H}^{∞} , and the derivative at $t = 0$ is equal to δq . This action of $\mathcal{L}_{q,\mathbb{R}}$ on Isospectral_{$\mathbb{R}(Y)$} is also transitive on the connected components.

From the point of view of Riemann surfaces the union of all the Jacobian varieties

$$
\bigcup_{Y' \in \mathcal{S}} \mathit{Jacobi}(Y')
$$

 \Box

is the completion of the Jacobian variety of the most singular Riemann surface $Y_{\text{algebraic}}$, the algebraic curve defined by the equation $R(\lambda, \mu) = 0$. In fact, in case of Riemann surfaces of finite genus this is a compactification of the generalized Jacobian variety (compare with [\[DS\]](#page-90-0)). The Jacobian variety of Riemann surfaces of infinite genus is not compact any more. Nevertheless the generalized Jacobian variety of a singular Riemann surface of infinite genus is a metrizable space and has a completion. The real part of the Jacobian variety of a Riemann surface of infinite genus is compact and the real part of the completion of the generalized Jacobian variety of a singular Riemann surface of infinite genus should be compact too. This space is homeomorphic to the subspace of all potentials \tilde{q} , such that $\det(\mu \mathbf{1} - F(\lambda, \tilde{q})) = R(\lambda, \mu)$. From the point of view of integrable systems, all Lagrangian subspaces decompose into the union of invariant subspaces under the action of $\tilde{H}_{\text{modified}}^1(Y_{\text{algebraic}}, \mathcal{O}^*)$, where $Y_{\text{algebraic}}$ is the most singular Riemann surface described by the equation $R(\lambda, \mu) = 0$:

$$
\bigcup_{Y' \in \mathcal{S}} \text{Isospectral}(Y') = \bigcup_{Y' \in \mathcal{S}} \text{Jacobian}_0(Y').
$$

Note also that the action of these Lie groups on these subspaces is given by hamiltonian flows. If this is a countable union, there can't exist further integrals of motion, which correspond to additional flows. If n is equal to 2, Proposition [10.1](#page-74-0) proves that this is the case. If we restrict our attention to the real parts of these isospectral sets, there exists a group, which acts transitively on these components. Furthermore, the real parts of the decompositions of the Lagrangian subspaces are equal to the real parts of the decompositions of the completion of the generalized Jacobian variety of $Y_{\text{algebraic}}$. Finally let us present an example, where S is not a countable set such that locally some more integrals exist. These additional integrals of course do not extend to any open set of the symplectic space.

Example 10.7 Let $n = 4^{40}$ and $p = diagonal(2, 1, -1, -2)$. We consider the constant potentials

$$
\left(\begin{array}{cccc} 0 & a & b & c \\ 0 & 0 & d & e \\ 0 & 0 & 0 & f \\ 0 & 0 & 0 & 0 \end{array}\right).
$$

The corresponding action of the finite dimensional Lie group is given by

$$
\mathbb{C}^6 \times \mathbb{C}^6 \to \mathbb{C}^6, (t_1, \dots, t_6) \times (a, b, c, d, e, f) \mapsto (\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}, \tilde{e}, \tilde{f}), \text{ with }
$$

$$
\begin{aligned}\n\tilde{a} &= \exp(t_1)a \\
\tilde{b} &= \exp(t_1 + t_2)(b + (3t_4 + 6t_6)ad) \\
\tilde{c} &= \exp(t_1 + t_2 + t_3)(c + 4t_4(ae + bf) + 4t_6(ae - bf) + (12t_4^2 + 4t_5 - 12t_6^2)adf) \\
\tilde{d} &= \exp(t_2)d \\
\tilde{e} &= \exp(t_2 + t_3)(e + (3t_4 - 6t_6)df) \\
\tilde{f} &= \exp(t_3)f.\n\end{aligned}
$$

This action decomposes into several orbits with different dimensions:

1. If $a \neq 0, d \neq 0, f \neq 0$ there is a six dimensional orbit, which is dense.

⁴⁰There exist more complicated examples for the case $n = 3$.

- 2. If $a = 0, d \neq 0, f \neq 0, b \neq 0$ or $a \neq 0, d \neq 0, f = 0, e \neq 0$ there is a five dimensional orbit. In both cases the fifth flow is trivial.
- 3. If $a \neq 0, d = 0, f \neq 0, b \neq 0, e \neq 0$ there is a one dimensional family of four dimensional orbits. Again the fifth flow is trivial and the fourth and sixth flows may be transformed into each other. The expression ae/bf is an additional integral of motion.
- 4. If $a = 0, d \neq 0, f = 0, b \neq 0, c \neq 0, e \neq 0$ the fourth, the fifth and the sixth flows are trivial. Hence there is a one dimensional family of three dimensional orbits. The value of be/dc is an additional integral of motion.
- 5. Some more orbits of lower dimensions.

Due to the third and fourth case the set S of singular curves corresponding to this spectral curve

$$
R(\lambda, \mu) = (\mu - \exp(-2\lambda))(\mu - \exp(-\lambda))(\mu - \exp(\lambda))(\mu - \exp(2\lambda))
$$

is not countable. Let us now consider the third case in some more detail. Like in Example [10.3](#page-77-0) we can define the singular Riemann surface with the help of the singular curve Y defined by the equation

$$
(\nu - 2\lambda)(\nu - \lambda)(\nu + \lambda)(\nu + 2\lambda) = 0.
$$

The normalization of this curve is the four-fold covering $\mathbb{P}_1 \cup \mathbb{P}_1 \cup \mathbb{P}_1 \cup \mathbb{P}_1$ of \mathbb{P}_1 :

The $\mathcal{O}_{\mathbb{P}_1}$ module $\pi_*(\mathcal{O}_Y)$ is generated by $1, \nu/\lambda, \nu^2/\lambda^2, \nu^3/\lambda^3$. Furthermore, the submodule $\pi'_*(\mathcal{O}_{Y'})$ is generated by

$$
1, \nu, \nu^2, \nu^3, (ae - bf)\frac{\nu^2}{\lambda} - (ae + bf)\frac{\nu^3}{\lambda^2}.
$$

There exists an additional flow on Isospectral(Y') corresponding to the additional integral of motion. Hence in this case the image of the homomorphism $d\Gamma_q: H^1_{q,\text{modified}}(Y',\mathcal{O}) \to \mathcal{L}_q^{\perp}$ is not equal to \mathcal{L}_q^{\perp} .

A Borel summability

In this Appendix we want to prove that equations([5\)](#page-9-0) and([6](#page-9-0)) establish a one to one correspondence between formal power series

$$
q(x) = \sum_{l=0}^{\infty} q_l x^l + b_0
$$
, with q_n an offdiagonal matrix for all $l \in \mathbb{N}$,

and b_0 due to Assumption [4.8](#page-24-0) a fixed diagonal matrix, and formal power series

$$
1 + \sum_{m=1}^{\infty} a_m \lambda^{-m}
$$
, with a_m an offdiagonal matrix for all $m \in \mathbb{N}$.

Furthermore, we will show that the formal power series $q(x)$ defines a analytic function in some neighbourhood of the point $x = 0$, if and only if the power series $\sum_{m=1}^{\infty} a_m \lambda^{-m} / m!$ defines an analytic function in some neighbourhood of $\lambda^{-1} = 0$. Hence the power series $\sum_{m=1}^{\infty} a_m(x)\lambda^{-m}$ of Theorem [2.4](#page-8-0) is Borel summable at some point $x \in \mathbb{R}$ if and only if $q(\cdot)$ is analytic at this point $x \in \mathbb{R}$. Moreover, the asymptotic expansion of v near infinity completely determines the potential q , if and only if q is analytic.

Now let $a_m(x) = \sum_{l=0}^{\infty} a_{m,l} x^l$ and $\sum_{l=0}^{\infty} b_{m,l} x^l$ be for all $m \in \mathbb{N}$ formal power series, such that $a_{m,l}$ are offdiagonal matrices and $b_{m,l}$ are diagonal matrices. The equation

$$
[a_{M+1}(x),p]+b_M(x) =
$$

$$
= \frac{da_M(x)}{dx} + [a_1(x), p]a_M(x) + [b_0, a_M(x)] - \sum_{m=1}^{M-1} a_m(x)b_{M-m}(x)
$$
(15)

completely determines all the formal power series $a_{M+1}(x)$ and $b_M(x)$ in terms of the power series $a_1(x), \ldots, a_M(x), b_1(x), \ldots, b_{M-1}(x)$. The inductive use of these equations determines the formal power series $a_{M+1}(x)$ and $b_M(x)$ for all $M \in \mathbb{N}$ in terms of the power series $a_1(x)$. Then equation([5\)](#page-9-0)

$$
q(x) = [a_1(x), p] + b_0
$$

shows that the formal power series

$$
1 + \sum_{m=0}^{\infty} a_m \lambda^{-m} = 1 + \sum_{m=0}^{\infty} a_m(0) \lambda^{-m}
$$

is completely determined in terms of the formal power series $q(x)$.

On the other hand the equation

$$
\left(\frac{d}{dx}\right)^{L+1} a_M(x) - \left(\frac{d}{dx}\right)^L b_M(x) =
$$
\n
$$
= [a_{M+1}^{(L)}(x), p] - [b_0, a_M^{(L)}(x)] - \sum_{l=0}^L \left(\begin{array}{c} L \\ l \end{array}\right) \left([a_1^{(l)}(x), p] a_M^{(L-l)}(x) + \sum_{M=1}^{M-1} a_m^{(l)}(x) b_{M-m}^{(L-l)} \right), \quad (16)
$$

where the superscript denotes formal derivatives with respect to x, determines the $(L + 1)$ th derivative of the power series $a_M(x)$ and the L-th derivative of the power series $b_M(x)$ in terms of derivatives of order at most L of $a_1(x), \ldots, a_{M+1}(x), b_1(x), \ldots, b_{M-1}(x)$. The inductive use of these equations determines all derivatives of the power series $a_1(x), a_2(x), \ldots$ and $b_1(x), b_2(x), \ldots$ in terms of the power series $a_1(x), a_2(x), \ldots$ In particular, all derivatives of the power series $a_1(x)$ at the point $x = 0$ are completely determined by the formal power series $1 + \sum_{m=1}^{\infty} a_m(0) \lambda^{-m}$. Hence the formal power series $q(x) = [a_1(x), p] + b_0$ is completely determined by the formal power series $1 + \sum_{m=1}^{\infty} a_m(0) \lambda^{-m}$. This shows the one to one correspondence between these two formal power series.

Now let us assume that q is a holomorphic function for all $|x| < R_x$ and bounded in norm forall $|x| \le R_x$ by some constant $C > 0$. Due to Cauchy's estimate [[Co](#page-90-0)] all coefficients are bounded:

$$
||q_l|| \leq \frac{C}{R_x^l} \Longleftrightarrow ||q^{(l)}(0)|| \leq \frac{Cl!}{R_x^l}.
$$

If we define the degree of the *l*-th derivative of q to be equal to $l + 1$, the recursion formula [\(15](#page-82-0)) shows that $a_{M+1}(x)$ and $b_M(x)$ are homogenous differential polynomials of degree $M + 1$. Now let $\alpha > 1$ be a real number greater than $1/|p_i - p_j|$ for all $n \ge i > j \ge 1$ and let the numbers $(\gamma_m)_{m \in \mathbb{N}}$ be defined inductively by

$$
\gamma_{M+1} = \alpha \left(\frac{M}{R_x} \gamma_M + \sum_{m=1}^M \gamma_m \gamma_{M+1-m} \right) \text{ and } \gamma_1 = C + ||ad(b_0)||.
$$

Then the recursion formula([15\)](#page-82-0) implies inductively that

$$
||a_{M+1}(0)|| \le \gamma_{M+1} \text{ and } ||b_M(0)|| \le \gamma_{M+1}.
$$

It is easy to see that

$$
\gamma_{M+1} \leq M!(C+\|ad(b_0)\|)\left(\left(C+\|ad(b_0)\|+\frac{1}{R_x}\right)\alpha\right)^M.
$$

This implies that for all

$$
|\lambda^{-1}| < \frac{R_x}{\alpha(R_x(C + \|ad(b_0)\|) + 1)}
$$

the power series $1 + \sum_{m=1}^{\infty} a_m(0) \lambda^{-m} / m!$ defines a holomorphic function.

Now let the real positive numbers $\alpha_M^{(L)}$ and $\beta_M^{(L)}$ satisfy the recursion relations

$$
\|\alpha_M^{(L+1)}\| = \|ad(p)\|\alpha_{M+1}^{(L)}\| + \|ad(b_0)\|\alpha_M^{(L)}\| + \beta_M^{(L)}\tag{17}
$$

$$
\beta_M^{(L)} = \sum_{l=0}^L \binom{L}{l} \left(\|ad(p)\| \alpha_1^{(l)} \alpha_M^{(L-l)} + \sum_{m=1}^{M-1} \alpha_m^{(l)} \beta_{M-m}^{(L-l)} \right). \tag{18}
$$

Then the inductive use of equation([16\)](#page-82-0) implies that the following estimates hold

$$
||a_M^{(L)}(0)|| \le \alpha_M^L
$$
 and $||b_M^{(L-1)}(0)|| \le \beta_M^{(L-1)}$ for all $M, L \in \mathbb{N}$,

whenever $||a_M(0)|| \le \alpha_M^{(0)}$ for all $M \in \mathbb{N}$. If $1 + \sum_{m=1}^{\infty} a_m(0)\lambda^{-m}$ is Borel summable, we may assume

$$
||a_m(0)|| \le \frac{C(m-1)!}{R_{\lambda}^{m-1}}
$$

with some $R_{\lambda} > 0$ and $C > 0$. Hence we set

$$
\alpha_m^{(0)} = \frac{C(m-1)!}{R_{\lambda}^{m-1}}.
$$

Now we claim that due to the recursion relations([17\)](#page-83-0) and [\(18](#page-83-0)) the following estimates hold:

$$
\alpha_M^{(L+1)} \le \frac{C(M+L)!}{R_{\lambda}^{M-1}} \left(\frac{\|ad(p)\|}{R_{\lambda}} + \frac{\|ad(b_0)\|}{C(2\|ad(p)\| + R_{\lambda})} + 1 \right)^{L+1} \left(C\left(\|ad(p)\| + \frac{R_{\lambda}}{2} \right) \right)^{M+L}
$$

$$
\beta_M^{(L)} \le \frac{C(M+L)!}{R_{\lambda}^{M-1}} \left(\frac{\|ad(p)\|}{R_{\lambda}} + \frac{\|ad(b_0)\|}{C(2\|ad(p)\| + R_{\lambda})} + 1 \right)^L \left(C\left(\|ad(p)\| + \frac{R_{\lambda}}{2} \right) \right)^{M+L}
$$

for all $M \in \mathbb{N}, L \in \mathbb{N}_0$. For the proof of the claim we first note that for all $L \in \mathbb{N}_0$, all $0 \leq l \leq L$ and all $M \in \mathbb{N} \setminus \{1\}$

$$
\binom{L}{l} \sum_{m=1}^{M-1} (m+l-1)!(M+L-m-l)! \le \frac{(L-l+1)(M+L)!}{(L+1)(L+2)}.
$$

For $M = 2$ this is obvious, and for $M > 2$ it follows by induction in M. This implies that

$$
\sum_{l=0}^{L} \binom{L}{l} \sum_{m=1}^{M-1} (m+l-1)!(M+L-m-l)! \le \frac{(M+L)!}{2}.
$$

Furthermore, the following estimate is obvious:

$$
\sum_{l=0}^{L} \binom{L}{l} l!(M+L-l-1)! \le (M+L)!.
$$

With these estimates the claim is an easy calculation. This claim directly shows that $q(x) =$ $[a_1(x), p] + b_0$ defines a holomorphic function on some neighbourhood of $x = 0 \in \mathbb{C}$, if the formal power series $1 + \sum_{m=1}^{\infty} a_m(0) \lambda^{-m}$ is Borel summable. We want to remark that this result cannot be proven with the well known theorem of Watson[[W-W\]](#page-91-0), which gives an explicit formula to reconstruct a holomorphic function out of his asymptotic expansion, if the asymptotic expansion is Borel summable. But there is another way to prove it. In fact, the method of Segal and Wilson [\[S-W](#page-91-0)] to produce solutions of the Korteweg-de Vries equation with the help of the Birkhoff factorization can be generalized to this situation. In [\[H-S-S\]](#page-90-0) there was given a modification of the usual Birkhoff factorization (see e.g.[[P-S\]](#page-91-0)) in order to cover more solutions of the Korteweg-de Vries and nonlinear Schrödinger equations. It is possible to go further in this direction:

- Let $L^-GL(n,\mathbb{C})$ be the group of all formal power series

$$
g_{-}(\lambda) = 1 + \sum_{m=1}^{\infty} a_m \lambda^{-m}
$$
, with $n \times n$ -matrices a_m for all $m \in \mathbb{N}$,

such that $\sum_{m=1}^{\infty} a_m \lambda^{-m} / m!$ defines a holomorphic entire function in λ^{-1} .

- Let $L^+GL(n,\mathbb{C})$ be the group of holomorphic entire functions

$$
g_+ : \mathbb{C} \to GL(n, \mathbb{C}), \lambda \mapsto g_{+(\lambda)}
$$
 of type 1,

i.e. g_{+} is asymptotically bounded by

$$
||g_+(\lambda)|| \le \exp(\alpha|\lambda|)
$$
 with some $\alpha > 0$.

Then the product of elements of $L^-GL(n,\mathbb{C})$ with elements of $L^+GL(n,\mathbb{C})$ is well defined. Moreover, let $LGL(n,\mathbb{C})$ be the group of invertible elements

$$
g(\lambda) = \sum_{m \in \mathbb{Z}} a_m \lambda^m,
$$

such that $1 + \sum_{m=-1}^{\infty} a_m \lambda^m$ is an element of $L^- GL(n, \mathbb{C})$ and $\sum_{m=0}^{\infty} a_m \lambda^m$ defines an entire function $\mathbb{C} \to n \times n$ -matrices of type 1. Then the Birkhoff factorization ([[P-S](#page-91-0), Theorem 8.1.2]) can be carried over to this modification of the loop group of $GL(n,\mathbb{C})$. With the help of this Birkhoff factorization one can reconstruct the potential $q(\cdot)$ out of the asymptotic expansion $1 + \sum_{m=1}^{\infty} a_m(0) \lambda^{-m}$ $1 + \sum_{m=1}^{\infty} a_m(0) \lambda^{-m}$ $1 + \sum_{m=1}^{\infty} a_m(0) \lambda^{-m}$ defined in Theorem [2.4](#page-8-0) (see [[H-S-S](#page-90-0)]).

In [\[Sch](#page-91-0)] it is shown that all higher flows of the Korteweg-de Vries and nonlinear Schrödinger equations correspond to the series of hamilton functions, given by the asymptotic expansion of ln(μ) in terms of λ^{-1} of Theorem [2.6](#page-10-0). Hence the statement of this Appendix is related to a statement [\[MK-T-1](#page-91-0), Theorem 10.1] of McKean and Trubowitz, which gives a condition for the spanning of the 'tangent' space of the isospectral sets in the sense of Section [8](#page-52-0) by all the local flows.

B Another reality condition

In this Appendix we indicate, how to treat other reality conditions in a fashion similar to that of Section [9](#page-60-0). In the case of the nonlinear Schrödinger equation two reality conditions are known: the so called non-focussing and the self focussing nonlinear Schrödinger equation. We will see that these two reality conditions are related to the two covering maps induced by λ and μ , respectively. In fact, let us now consider the reality condition corresponding to the covering map induced by μ . For the nonlinear Schrödinger equation this corresponds to the non-focussing case. The methods of [\[MK-T-1](#page-91-0)] can be carried over to this case, but we will give a different approach.

For this purpose we assume that the matrix p is invertible. Otherwise the transformation $p \mapsto pa + 1$ corresponds to the transformation $(\lambda, \mu) \mapsto (\lambda, \mu \exp(-a\lambda))$ without change of the Riemann surface, as mentioned in the proof of Theorem [7.3](#page-46-0). In addition we assume that p_1, \ldots, p_n are imaginary numbers. Then the Lax equation can be written as an eigenvalue equation

$$
\left(-\frac{d}{dx}p^{-1} - q(x)p^{-1}\right)\phi(x) = \lambda\phi(x)
$$
\n(19)

for a vector valued function $\phi(x) =$ $\sqrt{ }$ $\overline{}$ $\phi_1(x)$. . $\phi_n(x)$ \setminus . With the domain given as the set of absolutely $\overline{}$

continuous differentiable functions, such that $\phi(1) = \mu \phi(0)$, the operator

$$
L(\mu) = -\left(\frac{d}{dx}p^{-1} + q(x)p^{-1}\right)
$$

extendsto a closed unbounded operator (see e.g. [[R-S\]](#page-91-0)) of the Hilbert space $L^2([0,1],\mathbb{C}^n)$ of square integrable functions with the scalar product

$$
\left\langle \phi, \tilde{\phi} \right\rangle = \int_0^1 \phi^*(x) \tilde{\phi}(x) dx.
$$

Moreover, if $|\mu| = 1$ and $(q(x)p^{-1})^* = q(x)p^{-1}$ this operator is essentially self-adjoint. Hence we introduce the reality condition

$$
pq + q^*p = 0.
$$

The solution of the eigenvalue equation is given by

$$
\phi(x) = pg(x, \lambda, q)p^{-1}\phi(0).
$$

The boundary condition requires $\phi(0)$ to be an eigen vector of $pg(1, \lambda, q)p^{-1}$ with eigenvalue μ . Hence the equation

$$
R(\lambda, \mu) = \det(\mu \mathbf{1} - g(1, \lambda, q)) = 0
$$

describes the eigenvalues μ of $g(1, \lambda, q)$ and $F(\lambda, q)$ depending on λ , as well as the eigenvalues λ of the unbounded operator $L(\mu)$ depending on μ . From the second point of view $R(\lambda, \mu) = 0$ describes an infinite-fold covering over $\mu \in \mathbb{C} \setminus \{0\}$, such that all covering points of some $\mu \in \mathbb{C} \setminus \{0\}$ fit together to form the spectrum of $L(\mu)$. The reality condition implies

$$
L^*(\mu) = L\left(\frac{1}{\bar{\mu}}\right) \text{ for all } \mu \in \mathbb{C} \setminus \{0\},\
$$

since $L(\mu)$ is an unbounded operator valued holomorphic function. We leave aside the analytic aspects of such functions and let us just transform the reality condition into a condition on $g(x, \lambda, q)$. Let $U(p)$ be the Lie subgroup of $GL(n, \mathbb{C})$ which fixes the hermitian form defined by p:

$$
U(p) = \{ g \in GL(n, \mathbb{C}) | g^*pg = p \}.
$$

The corresponding Lie algebra is given by

$$
u(p) = \{a \in gl(n, \mathbb{C})|pa + a^*p = 0\}.
$$

Then the reality condition is equivalent to the condition that $q(x) + p\lambda$ is an element of $u(p)$ for all real λ . This implies that $q(x, \lambda, q)$ is an element of $U(p)$ for all real λ . More generally, the following relation holds:

$$
g^*(x, \bar{\lambda}, q)pg(x, \lambda, q) = p \text{ for all } \lambda \in \mathbb{C}.
$$

Hence we have

$$
\overline{R\left(\bar{\lambda}, \frac{1}{\bar{\mu}}\right)} = \overline{\det\left(\frac{1}{\bar{\mu}}\mathbb{1} - g(\bar{\lambda})\right)} = \det\left(\frac{1}{\mu}\mathbb{1} - pg^{-1}(\lambda)p^{-1}\right) =
$$

$$
= \det\left(\frac{1}{\mu}\mathbb{1} - g^{-1}(\lambda)\right) = \frac{R(\lambda, \mu)}{(-\mu)^n \det(g(\lambda))} = \frac{R(\lambda, \mu)}{(\mu)^n R(\lambda, 0)}.
$$

Again we have an antilinear involution of the Riemann surface Y :

$$
\theta: Y \to Y, (\lambda, \mu) \mapsto \left(\bar{\lambda}, \frac{1}{\bar{\mu}}\right).
$$

Let us now find a condition on the divisors, which is equivalent to the reality condition. For this purpose we carry over equation [\(10](#page-23-0)) to the case of the covering map induced by μ . The solution of the eigenvalue equation with quasi-periodic boundary conditions is given by

$$
\phi(x) = \pi^*(pg(x,\cdot,q)h_0^{-1})v.
$$

For the transposed eigenvalue equation

$$
\frac{d}{dx}\psi(x)p^{-1} - q(x)p^{-1}\psi(x) = \lambda\psi(x)
$$
\n(20)

for a vector valued function $\psi(x) = (\psi_1(x), \dots, \psi_n(x))$ with boundary condition $\psi(1) = \frac{1}{\mu}\psi(0)$ we have the solution

$$
\psi(x) = w\pi^*(h_0g^{-1}(x,\cdot,q)).
$$

Hence the analogous operator P of Lemma [4.5](#page-22-0) is given by

$$
P: L^{2}([0,1], \mathbb{C}^{n}) \to L^{2}([0,1], \mathbb{C}^{n}), \chi \mapsto P\chi = \phi \frac{\int_{0}^{1} \psi(x)\chi(x)dx}{\int_{0}^{1} \psi(x)\phi(x)dx}.
$$

Similar to Lemma [4.5](#page-22-0) we have:

Lemma B.1 (i) $P^2 = P$.

- (ii) $L(\mu)P = PL(\mu) = \lambda P$.
- (iii) $\sum_{\text{sheets of the covering map induced by } \mu} P = 1$.
- (iv) The divisor of P is equal to $-b_{\mu}$, the branching divisor of the covering map induced by μ .

We only want to indicate the proof of this lemma. The first two statements may be verified directly. If $L(\mu)$ is normal, the statement (iii) is a consequence of the spectral decomposition of $L(\mu)$. Since both sides are meromorphic functions, this implies (iii). In the proof of Lemma [7.4](#page-47-0) we used the fact that

$$
\int_0^1 \psi(x)\phi(x)dx = \int_0^1 w\pi^*(h_0g^{-1}(x,\lambda,q)pg(x,\lambda,q)h_0^{-1})vdx =
$$

= $-wF^{-1}(\lambda,q)\frac{\partial F(\lambda,q)}{\partial \lambda}v = -\frac{1}{\mu}\frac{d\mu}{d\lambda}wv.$

The divisor of the meromorphic function $\mu^{-1} \frac{d\mu}{d\lambda}$ is equal to the branching divisor of the covering map induced by μ minus the branching divisor of the covering map induced by λ . A similar argument to that in Lemma [4.5](#page-22-0) now shows (iv). \Box

For all matrix valued meromorphic functions f on Y, let $\theta^*(f)$ again be the function $\theta^*(f)$ = $(f \circ \theta)^*$. As we saw above, the reality condition implies

$$
F^{-1^*}(\bar{\lambda}, q) = h_0 p h_0^{-1} F(\lambda, q) h_0 p^{-1} h_0^{-1}.
$$

Then we have

$$
\theta^*(v)h_0ph_0^{-1}\pi^*(F(\cdot,q)) = \theta^*(v)\theta^*(\pi^*(F^{-1}(\cdot,q)))h_0ph_0^{-1} = \mu\theta^*(v)h_0ph_0^{-1}.
$$

Thusthe reality condition transforms to the relation: $w = \theta^*(v)h_0ph_0^{-1}$ is a solution of ([9\)](#page-20-0), ifand only if v is a solution of (8) (8) . Of course this is not compatible with the normalization $v_1 = 1 = w_1$. Hence we choose another normalization: For every potential q let $\phi(x)$ and $\psi(x)$ be the unique solutions of [\(19](#page-85-0)) with boundary condition $\phi(1) = \phi(0)\mu$ and [\(20](#page-87-0)) with boundary condition $\psi(0) = \mu^{-1}\psi(1)$, such that $\sum_{i=1}^{n} \phi_i(0) = \sqrt{n} = \sum_{i=1}^{n} \psi_i(0)$, respectively. Due to this normalization ϕ and ψ can be considered as $L^2([0,1],\mathbb{C}^n)$ valued meromorphic functions on Y with no zeroes. If v and w are the solutions of [\(8](#page-20-0)) and [\(9](#page-20-0)) with normalization $v_1 = 1 = w_1$, respectively, ϕ and ψ are given by

$$
\phi(x) = \pi^*(pg(x, \cdot, q)h_0^{-1})v \frac{1}{(h_0 ph_0^{-1} v)_1}
$$

$$
\psi(x) = w\pi^*(h_0 g^{-1}(x, \cdot, q)).
$$

Hence the reality condition transforms to the condition

$$
\phi(x) = \theta^*(\psi(x)) = \theta^*(w\pi^*(h_0g^{-1}(x,\cdot,q))).
$$

It is obvious that the divisor of ψ is equal to $-D^t(q)$ and the divisor of ϕ is equal to $-D(q)$ – $((h_0ph_0^{-1}v)_1)$. This function takes the values $p_1\sqrt{n}, \ldots, p_n\sqrt{n}$ at the covering points of infinity. Therefore it does not define an equivalence relation in the modified sense. As a direct consequence of Lemma [B.1](#page-87-0) we have

$$
(\phi) + (\psi) + \left(\int_0^1 \psi(x)\phi(x)dx\right) = b_\mu.
$$
\n(21)

With the normalization given above the function $\int_0^1 \psi(x) \phi(x) dx$ is equal to

$$
-\frac{1}{\mu}\frac{d\mu}{d\lambda}\frac{wv}{(wh_0ph_0^{-1}v)_1}.
$$

This function takes the same value at all covering points of infinity. Hence it defines an equivalence relation in the modified sense. Now we can state a theorem analogous to Theorem [9.3](#page-61-0):

Theorem B.2 In general the set of fixed points of the involution θ decomposes into several connected components⁴¹. If a meromorphic function f gives an equivalence relation in the

⁴¹In the hyperelliptic case $n = 2$ it is more convenient to use the antilinear involution $(\lambda, \mu) \rightarrow (\bar{\lambda}, \bar{\mu})$. The number of the connected components of the set of fixed points of both involutions is equal to the genus of theRiemann surface plus one. Such Riemann surfaces are called M-curves (see e.g. [Kr-2] [Kr-2] [Kr-2]). Then the reality condition simplifies to the condition, that the divisor is invariant under this involution.

REFERENCES 89

modified sense between $D + \theta(D)$ and b_{μ} , then on each connected component of the set of fixed points of the involution θ this function is either non-negative or non-positive. Hence the real part of the Jacobian variety

$$
Jacobian_{\mathbb{R}}(Y) = \{ [D] \in Jacobian(Y)|D + \theta(D) \sim b_{\mu} \text{ in the modified sense } \}
$$

decomposes into several connected components being characterized by the sign of f on each connected component of the set of fixed points of the involution θ . Now let Jacobian_{R,0}(Y) be the connected component of Jacobian_R(Y) corresponding to the case that f is non-negative on the whole set of fixed points of the involution θ . Then Jacobian_{R,0}(Y) is contained in Jacobian₀(Y), and T_x acts on this subspace. Moreover, a potential satisfies the reality condition $pq + q^*p = 0$ if and only if the inverse of the restriction of the divisor of ϕ to $\pi^{-1}(\mathbb{C})$ with the normalization given above is an element of Jacobian_{R,0} (Y) .

The proof is quite similar to the proof of Theorem [9.3.](#page-61-0) We only indicate the modifications. If ψ is equal to $\theta^*(\phi)$ the function $\int_0^1 \psi(x)\phi(x)dx$ is positive on the set of fixed points of θ , and the inverse of the restriction of the divisor of ϕ to $\pi^{-1}(\mathbb{C})$ is an element of *Jacobian*_{R,0} (Y) . On the other hand let f be a function, such that (f) gives an equivalence relation in the modified sense between $D + \theta(D)$ and b_{μ} . Then $\theta^*(f)/f$ is a holomorphic function of Y, which is equal to 1 at all covering points of infinity. Hence $\theta^*(f)$ is equal to f and f is real valued on the set of fixed points of the involution θ . Moreover, f can only have zeroes of even order on this set. Then f is either non-negative or non-positive on each connected component of this set. Now let μ_0 be any value with $|\mu_0| = 1$ and let g be any cross section of $\mathcal{O}_{D-\pi^{-1}(\infty)}$. Then the total residue of the form $\theta^*(g)gf(\mu-\mu_0)^{-1}d\mu$ converges to zero. This again shows that D is non-special in the modified sense, if D is an element of $Jacobi\pi_{\mathbb{R},0}(Y)$. The rest of the proof is the same as the proof of Theorem [9.3.](#page-61-0) \Box

Finally we want to mention that Theorem [9.3](#page-61-0) and Theorem [B.2](#page-88-0) are related to the following fact: Let $LG, L-G, L^+G$ be the subgroups of $LGL(n, \mathbb{C}), L^-GL(n, \mathbb{C}), L^+GL(n, \mathbb{C})$ of elements, which obey the relation

$$
g^*(\bar{\lambda}) = g^{-1}(\lambda)
$$
 for all λ , with $|\lambda| = 1$.

Then the Birkhoff factorization defines a diffeomorphism onto the whole of LG:

$$
L^-G \times L^+G \to LG, (g_-,g_+) \mapsto g_-g_+.
$$

Acknowledgements. The first part of this work was performed while the author was visiting the Forschungsinstitut für Mathematik (ETH, Zürich). He would like to thank J. Moser for his hospitality. Discussions with H. Knörrer and E. Trubowitz have been very stimulating and helpful. Furthermore, the author thanks F. J. Archer, P. G. Grinevich, J. Mund and R. Schrader for comments and corrections.

References

[A-H-H] Adams M. R., Harnad J., Hurtubise J.: Darboux coordinates and Liouville-Arnold integration in Loop algebras. Commun. Math. Phys. 155, 385-413 (1993).

REFERENCES 90

- [A-vM] Adler M., van Moerbeke P.: Completely integrable systems, Euclidian Lie algebras and curves. Adv. Math. 38, 267-317 (1980); Lineraization of Hamiltonian systems, Jacobi varieties and representation theory. Adv. Math. 38, 318-379 (1980).
- [B-S] Beals R., Sattinger D. H.: On the complete integrability of completely integrable systems. Commun. Math. Phy. 138, 409-436 (1991).
- [Bo] Bourgain J.: Fourier transform restriction phenomena for certain lattice subsets and applications to non-linear evolution equations. Preliminary version.
- [Co] Conway J. B.: Functions of one complex variable. Berlin, Heidelberg, New York: Springer 1973.
- [D-K-N] Dubrovin B. A., Krichever I. M., Novikov S. P.: Integrable systems I. In: Arnold V. I., Novikov S. P. (eds.) Dynamical Systems IV. Encyclopaedia of Mathematical Sciences vol.4, pp. 173-280. Springer,(1990).
- [DS] D'Souza C.: Compactification of generalized Jacobians. Pro. Indian Acad. Sci. 88, 419-457 (1979).
- [F-T] Faddeev L. D., Takhtajan L. A.: Hamiltonian methods in the theory of solitons. Berlin, Heidelberg, New York: Springer 1987.
- [F-K-T] Feldman J., Knörrer H., Trubowitz E.: Riemann surfaces of infinite genus. Preliminary version.
- [F-K] Fordy A. P., Kulish P. P.: Nonlinear Schrödinger equations and simple Lie algebras: Commun. Math. Phys. 89, 427-443 (1983).
- [Fo] Forster O.: Lectures on Riemann surfaces. Berlin, Heidelberg, New York: Springer 1981.
- [Fr] Frenkel I. B.: Orbital theory for affine Lie algebras. Invent. Math. 77, 301-352 (1984).
- [Gu] Gunning R. C.: Lectures on vector bundles over Riemann surfaces. Princeton, New Jersey: Princeton University Press 1967.
- [H-S-S] Haak G., Schmidt M., Schrader R.: Group theoretic formulation of the Segal-Wilson approach to integrable systems with applications: Rev. Math. Phys. 4, 451-499 (1992).
- [Ho] Hochstadt H.: On the asymptotic spectrum of Hill's equation. Archiv d. Mathematik 14, 34-38 (1963).
- [I-M] Its A. R., Matveev V. B.: Hill's operator with finitely many gaps. Functional Anal. Apll. 9, 65-66 (1975).
- [Kr-1] Krichever I. M.: Methods of algebraic geometry in the theory of non-linear equations. Russian Math. Surveys 32 (6), 185-213 (1977).
- [Kr-2] Krichever I. M.: Spectral theory of two-dimensional periodic operators and its applications. Russian Math. Surveys 44, 145-225 (1989).
- [MK-T-1] McKean H., Trubowitz E.: Hill's operator and hyperelliptic function theory in the presence of infinitely many branchpoints. Commun. Pure Appl. Math. 29, 143-226 (1976).
- [MK-T-2] McKean H., Trubowitz E.: Hill's surfaces and their theta functions. American Math. Soc. 84 (6), 1052-1085 (1978).
- [M-W] Magnus W., Winkler S.: Hill's equation. New York, London, Sydney: Interscience Publishers 1966.
- [P-S] Pressley A., Segal G.: Loop groups: Oxford: Clarendon Press 1986.
- [P-T] Pöschel J., Trubowitz E.: Inverse spectral theory. Orlando, Florida: Academic Press 1987.
- [R-S] Reed M., Simon B.: Functional analysis. San Diego, California: Academic Press 1980.
- [R-S-T] Reyman A. G., Semenov-Tian-Shansky M. A.: Reduction of Hamiltonian systems, affine Lie algebras and Lax equations I and II. Inventiones math. 54, 81-100 (1979); Inventiones math. 63, 423-432 (1981).
- [Sch] Schmidt M. U.: Solitonen und Schleifengruppen. Berlin: Thesis 1990.
- [S-W] Segal G., Wilson G.: Loop groups and equations of KdV type. Publ. Math. I. H. E. S. 61, 5-65 (1985).
- [Se] Serre J. P.: Algebraic groups and class fields. Berlin, Heidelberg, New York: Springer 1988.
- [W-W] Whittaker E. T., Watson G. N.: A course of modern analysis. Cambridge: University Press 1927 (ed. 4).