# Solvability of the  $G_2$  Integrable System

Marcos Rosenbaum<sup>1</sup>, Alexander Turbiner<sup>2,3</sup> and Antonio Capella

Instituto de Ciencias Nucleares, UNAM, Apartado Postal 70-543, 04510 Mexico D.F., Mexico

# Abstract

It is shown that the 3-body trigonometric  $G_2$  integrable system is exactlysolvable. If the configuration space is parametrized by certain symmetric functions of the coordinates then, for arbitrary values of the coupling constants, the Hamiltonian can be expressed as a quadratic polynomial in the generators of some Lie algebra of differential operators in a finite-dimensional representation. Four infinite families of eigenstates, represented by polynomials, and the corresponding eigenvalues are described explicitly.

Intern.Journ.Mod.Phys. A13, 3885-3904 (1998)

<sup>1</sup>mrosen@xochitl.nuclecu.unam.mx

<sup>2</sup> turbiner@xochitl.nuclecu.unam.mx

<sup>3</sup>On leave of absence from the Institute for Theoretical and Experimental Physics, Moscow 117259, Russia.

#### 1. INTRODUCTION

<span id="page-1-0"></span>Consider the Hamiltonian describing a three-body system with pairwise and three-body interactions

$$
\mathcal{H}_{G_2}^{(r)} = \frac{1}{2} \sum_{k=1}^3 \left[ -\frac{\partial^2}{\partial x_k^2} + \omega^2 x_k^2 \right] + g \sum_{k(1.1)
$$

where g,  $g_1$  are the coupling constants. This model, now known as the rational  $G_2$  model, was originally proposed by J. Wolfes[[1](#page-17-0)] who studied its bound states and proved its exact solvability. The scattering problem for the same model was explored in [\[2\]](#page-17-0). It is worth mentioning that in the limit  $g_1 \to 0$ , the model (1.1) becomes the celebrated Calogero model[[3](#page-17-0)].

Using the Hamiltonian reduction method [\[4\]](#page-17-0) (see also the review [\[5](#page-17-0)]) Olshanetsky and Perelomov have shown that the model (1.1) is nothing but a particular (rational) case of a more general model related to the root system of the exceptional group  $G_2$ . They called this model the  $G_2$  model and proved its complete integrability. However, the question of whether even a trigonometric version of the general  $G_2$  model

$$
\mathcal{H}_{G_2} = -\frac{1}{2} \sum_{k=1}^3 \frac{\partial^2}{\partial x_k^2} + \frac{g}{4} \sum_{k(1.2)
$$

is exactly-solvable remained open for many years. Recently, Quesne [\[6\]](#page-17-0) showed its exact solvability for the particular case  $g = 0$ . In the present paper we demonstrate the exact solvability of (1.2) for the general case  $g, g_1 \neq 0$ . We use a notion of exact solvability based on the existence of a flag of functional spaces with an explicit basis preserved by the Hamiltonian. A constructive criterion for exact-solvability consists on checking whether the flag is related to finite-dimensional representation spaces of a Lie algebra of differential operators [\[7\]](#page-17-0). If this criterion is fulfilled, then the corresponding algebra is called the hidden algebra of the system studied, and therefore the Hamiltonian can be written in terms of the generators of this algebra.

In[[8](#page-17-0)] it was shown that the eigenfunctions of the N-body Calogero and Sutherland mod-els[[3](#page-17-0), [9](#page-17-0)] form the flag of finite-dimensional representation spaces of the algebra  $gl(N)$ , realized by first order differential operators. The corresponding Hamiltonians were rewritten as quadratic polynomials in the generators of the Borel subalgebra of  $gl(N)$ , and the coupling constants appear only in the coefficients of these polynomials. Recently, it was shown that this statement holds for all the ABCD Olshanetsky-Perelomov integrable systems (and also for their SUSY generalizations which turn out to be associated to the hiddenalgebra  $gl(N|N-1)$ , see [[10\]](#page-17-0)).

Here we show that for the 3-body Calogero-Sutherland models, there exists one more hidden algebra that we shall call  $g^{(2)}$  (see Appendix B), besides the above-mentioned hidden algebra  $gl(3)$ . This property is maintained for the general rational  $G_2$  model (1.1) and also for the degenerate  $(g = 0)$  trigonometric  $G_2$  model. However, for the general trigonometric  $G_2$  case only the hidden algebra  $g^{(2)}$  remains, while  $gl(3)$  is no longer a hidden algebra.

<span id="page-2-0"></span>The paper is organized as follows. In Section 2 the algebraic and, the  $gl(3)$  and  $g^{(2)}$ Lie-algebraic forms of the 3-body Calogero and Sutherland models are given. Section 3 is devoted to the demonstration of a phenomenon, which we call 'complementarity', as a consequence of which the 3-body Calogero and Sutherland models are shown to be equivalentto degenerate  $G_2$  models corresponding to [\(1.1](#page-1-0)) and ([1.2\)](#page-1-0) with  $g = 0$ . The special coordinates leading to the algebraic and the  $g^{(2)}$  Lie-algebraic forms of a general  $G_2$  model are used in Section 4, in other to demonstrate the exact solvability of this model. A realization of the algebra  $ql(3)$  in terms of first order differential operators, acting on the plane, is given in Appendix A. Appendix B is devoted to a description of the infinite-dimensional algebra of differential operators  $g^{(2)}$ , admitting finite-dimensional representations in terms of polynomials in two variables. Finally, in Appendix C the explicit formulas for the first several eigenfunctions of the general  $G_2$  model are presented.

### 2. Algebraic and Lie algebraic forms of the 3-body Calogero and Sutherland models

In this section we represent the Hamiltonians of the 3-body Calogero and Sutherland models in an algebraic form, by making use of two different sets of translationally invariant, permutationally-symmetric coordinates. The procedure leads to two different Lie-algebraic forms: (i) in terms of the generators of the  $gl(3)$  algebra [\[8\]](#page-17-0) and (ii) in terms of some generators of the infinite-dimensional algebra  $g^{(2)}$  (see Appendix B).

### 2.1. Calogero model. The Hamiltonian of the 3-body Calogero model is defined by

$$
\mathcal{H}_{\text{Cal}} = \frac{1}{2} \sum_{i=1}^{3} \left[ -\frac{\partial^2}{\partial x_i^2} + \omega^2 x_i^2 \right] + g \sum_{i < j}^{3} \frac{1}{(x_i - x_j)^2} \,,\tag{2.1}
$$

where  $g = v(v - 1) > -\frac{1}{4}$  is the coupling constant and  $\omega$  is the harmonic oscillator frequency. The ground state eigenfunction is given by

$$
\Psi_0^{(c)}(x) = \Delta^{\nu}(x)e^{-\omega \frac{X^2}{2}}, \qquad (2.2)
$$

where  $\Delta(x) = \prod_{i < j} |x_i - x_j|$  is the Vandermonde determinant and  $X^2 = \sum_i x_i^2$ . It was shown by Calogero that the eigenfunctions for this model can be expressed as

$$
\Psi(x) = \Psi_0^{(c)}(x) P_c(x) , \qquad (2.3)
$$

where  $P_c(x)$  is a polynomial symmetric under permutations of any two  $x_i$ 's. The operator having these polynomials as eigenfunctions can be obtained by performing on (2.1) the gauge rotation

$$
h_{\text{Cal}} = -2(\Psi_0^{(c)}(x))^{-1} \mathcal{H}_{\text{Cal}} \Psi_0^{(c)}(x) . \qquad (2.4)
$$

In order to study the internal dynamics of the system we introduce the center-of-mass coordinate  $Y = \sum_{j=1}^{3} x_j$  and the translation-invariant Perelomov relative coordinates [\[11](#page-17-0)]:

$$
y_i = x_i - \frac{1}{3}Y, \quad i = 1, 2, 3 \quad , \tag{2.5}
$$

<span id="page-3-0"></span>which obey the constraint  $y_1 + y_2 + y_3 = 0$ . To incorporate the permutation symmetry and the translational invariance we consider the coordinates[[8](#page-17-0)]:

$$
\tau_2 = -y_1^2 - y_2^2 - y_1 y_2 , \qquad (2.6)
$$

$$
\tau_3 = -y_1 y_2 (y_1 + y_2) \,. \tag{2.7}
$$

In terms of these coordinates we obtain, after extracting the center-of-mass motion, the first algebraic form for the Hamiltonian([2.4\)](#page-2-0):

$$
h_{\text{Cal}} = -2\tau_2 \partial_{\tau_2 \tau_2}^2 - 6\tau_3 \partial_{\tau_2 \tau_3}^2 + \frac{2}{3} \tau_2^2 \partial_{\tau_3 \tau_3}^2 - [4\omega \tau_2 + 2(1+3\nu)]\partial_{\tau_2} - 6\omega \tau_3 \partial_{\tau_3} \; . \tag{2.8}
$$

It is worth emphasizing that for a fixed coupling constant  $g$  there are 2 different solutions for  $\nu$ , hence the operator (2.8) has two different sets of polynomial eigenfunctions. In fact, as shown in  $[8]$ , the operator  $(2.8)$  can be rewritten in a Lie-algebraic form in terms of the gl(3)-algebra generators (see (A.1) where  $n = 0$  and  $x, y$  replaced by  $\tau_2, \tau_3$ , respectively). The corresponding expression is

$$
h_{\text{Cal}} = -2J_{2,2}^0 J_2^- - 6J_{3,3}^0 J_2^- + \frac{2}{3} J_{2,3}^0 J_{2,3}^0 - 4\omega J_{2,2}^0 - 2(1+3\nu)J_2^- - 6\omega J_{3,3}^0 \,. \tag{2.9}
$$

The polynomial eigenfunctions of  $(2.8)$ – $(2.9)$  have the form given in  $(A.2)$  (see also Fig.1 in Appendix A). Further note that the operator (2.8) is invariant under  $\tau_3 \to -\tau_3$ , resulting fromthe reflection symmetry  $(x \to -x)$  of the original Hamiltonian ([2.1\)](#page-2-0). We incorporate this symmetry by introducing the new coordinates

$$
\lambda_1 = \tau_2
$$
,  $\lambda_2 = \tau_3^2$ , (2.10)

and then the operator (2.8) takes another algebraic form given by

$$
h_{\text{Cal}} = -2\lambda_1 \partial_{\lambda_1 \lambda_1}^2 - 12\lambda_2 \partial_{\lambda_1 \lambda_2}^2 + \frac{8}{3} \lambda_1^2 \lambda_2 \partial_{\lambda_2 \lambda_2}^2
$$
\n
$$
- [4\omega \lambda_1 + 2(1+3\nu)] \partial_{\lambda_1} - (12\omega \lambda_2 - \frac{4}{3} \lambda_1^2) \partial_{\lambda_2}.
$$
\n
$$
(2.11)
$$

(cf.(2.8)). Now  $h_{\text{Cal}}$  can be rewritten in terms of the generators (B.2) of the algebra  $gl_2 \Join$  $R^3 \subset g^{(2)}$  (see Appendix B with  $n = 0$  and  $x, y$  are replaced by  $\lambda_1, \lambda_2$ , respectively) only, rather than in terms of  $gl(3)$ . The corresponding *Lie-algebraic* form for the Hamiltonian is then given by

$$
h_{\text{Cal}} = -2L^2L^1 - 12L^3L^1 + \frac{8}{3}L^7L^3 - 2(1+3\nu)L^1
$$
  

$$
-4\omega L^2 - 12\omega L^3 - \frac{4}{3}L^7.
$$
 (2.12)

The polynomial eigenfunctions of  $(2.11)$ – $(2.12)$  have the form  $(B.1)$  (see also Fig.2 in Appendix B).

Both Lie-algebraic forms (2.9), (2.12) do not contain the positive-grading generators  $J_2^+, J_3^+$  and  $L^4$ , respectively. This ensures the preservation of the corresponding flags of polynomial spaces (A.2) and (B.1), and the exact solvability of the Calogero model following the criterion formulated in the Introduction.

<span id="page-4-0"></span>2.2. Sutherland model. The Hamiltonian for the 3-body Sutherland model is defined by

$$
\mathcal{H}_{\text{Suth}} = -\frac{1}{2} \sum_{k=1}^{3} \frac{\partial^2}{\partial x_k^2} + \frac{g}{4} \sum_{k
$$

where  $g = \nu(\nu - 1) > -\frac{1}{4}$  $\frac{1}{4}$  is the coupling constant. The ground state of this Hamiltonian is

$$
\Psi_0^{\text{(Suth)}}(x) = (\Delta^{(trig)}(x))^{\nu} , \qquad (2.14)
$$

where  $\Delta^{(trig)}(x) = \prod_{i is the trigonometric analog of the Vandermonde$ determinant. Sutherland showed that for this model any eigenfunction can be written as

$$
\Psi(x) = \Psi_0^{\text{(Suth)}}(x) P_s(e^{ix}) \;, \tag{2.15}
$$

where  $P_s(e^{ix})$  is a polynomial, symmetric under permutations of any two  $x_i$ 's. These polynomials are the so called Jack-Sutherland polynomials. As for the Calogero case, the operator which has these polynomials as eigenfunctions can be obtained by performing on (2.13) a gauge transformation

$$
h_{\text{Suth}} = -2(\Psi_0^{\text{(Suth)}}(x))^{-1} \mathcal{H}_{\text{Suth}} \Psi_0^{\text{(Suth)}}(x) . \tag{2.16}
$$

To exhibit the internal structure of this system, we introduce the translation-invariant, permutation-symmetric, periodic coordinates[[8](#page-17-0)]

$$
\eta_2 = \frac{1}{\alpha^2} [\cos(\alpha y_1) + \cos(\alpha y_2) + \cos(\alpha (y_1 + y_2)) - 3], \qquad (2.17)
$$

$$
\eta_3 = \frac{2}{\alpha^3} [\sin(\alpha y_1) + \sin(\alpha y_2) - \sin(\alpha (y_1 + y_2))]. \tag{2.18}
$$

In the limit  $\alpha \to 0$ , these coordinates become (2.6)–(2.7). After extracting the center-ofmass motion, the Hamiltonian  $(2.16)$  in these coordinates takes the *algebraic* form

$$
h_{\text{Suth}} = -(2\eta_2 + \frac{\alpha^2}{2}\eta_2^2 - \frac{\alpha^4}{24}\eta_3^2)\partial_{\eta_2\eta_2}^2 - (6 + \frac{4\alpha^2}{3}\eta_2)\eta_3\partial_{\eta_2\eta_3}^2 + (\frac{2}{3}\eta_2^2 - \frac{\alpha^2}{2}\eta_3^2)\partial_{\eta_3\eta_3}^2 - [2(1+3\nu) + 2(\nu + \frac{1}{3})\alpha^2\eta_2]\partial_{\eta_2} - 2(\nu + \frac{1}{3})\alpha^2\eta_3\partial_{\eta_3}.
$$
\n(2.19)

It is worth emphasizing that similarly to the Calogero case, there are 2 different solutions for  $\nu$  for a fixed coupling constant g. Consequently the operator (2.19) has two different setsof polynomial eigenfunctions. As shown in  $[8]$  $[8]$  $[8]$ , the operator  $(2.19)$  can be rewritten in a Lie-algebraic form in terms of the  $gl(3)$ -algebra generators (see (A.1) with  $n = 0$  and  $x, y$  are replaced by  $\eta_2, \eta_3$ , respectively). This procedure yields

$$
h_{\text{Suth}} = -2J_{2,2}^{0}J_{2}^{-} - 6J_{3,3}^{0}J_{2}^{-} + \frac{2}{3}J_{2,3}^{0}J_{2,3}^{0} - 2(1+3\nu)J_{2}^{-} + \frac{\alpha^{4}}{24}J_{3,2}^{0}J_{3,2}^{0}
$$
\n
$$
-\alpha^{2}\left[\frac{1}{2}J_{2,2}^{0}J_{2,2}^{0} + \frac{4}{3}J_{2,2}^{0}J_{3,3}^{0} + \frac{1}{2}J_{3,3}^{0}J_{3,3}^{0} + 2(\nu + \frac{1}{12})(J_{2,2}^{0} + J_{3,3}^{0})\right].
$$
\n(2.20)

The polynomial eigenfunctions of  $(2.19)$ – $(2.20)$  have the form  $(A.2)$  (see also Fig.1 in Appendix A).

<span id="page-5-0"></span>Also, as in the Calogero case, the Sutherland Hamiltonian possesses a reflection symmetry  $(x \to -x)$ . We can incorporate this symmetry in a new set of variables by defining

$$
\sigma_1 = \eta_2 \;, \qquad \qquad \sigma_2 = \eta_3^2 \;.
$$
 (2.21)

In these variables the operator [\(2.16](#page-4-0)) becomes

$$
h_{\text{Suth}} = -(2\sigma_1 + \frac{\alpha^2}{2}\sigma_1^2 - \frac{\alpha^4}{24}\sigma_2)\partial_{\sigma_1\sigma_1}^2 - (12 + \frac{8\alpha^2}{3}\sigma_1)\sigma_2\partial_{\sigma_1\sigma_2}^2 + (\frac{8}{3}\sigma_1^2\sigma_2 - 2\alpha^2\sigma_2^2)\partial_{\sigma_2\sigma_2}^2 - [2(1 + 3\nu) + 2(\nu + \frac{1}{3})\alpha^2\sigma_1]\partial_{\sigma_1} + \left[\frac{4}{3}\sigma_1^2 - (\frac{7}{3} + 4\nu)\alpha^2\sigma_2\right]\partial_{\sigma_2},
$$
\n(2.22)

which is another *algebraic* form for the Sutherland Hamiltonian (cf.(2.19)). The operator  $(2.22)$  can be represented in terms of the generators of the algebra  $g^{(2)}$  (see (B.2), (B.4) with  $n = 0$  and x, y are replaced by  $\sigma_1, \sigma_2$ , respectively):

$$
h_{\text{Suth}} = -2L^2L^1 - 12L^3L^1 + \frac{8}{3}L^7L^3 - \alpha^2(\frac{L^2L^2}{2} + \frac{8L^3L^2}{3} + 2L^3L^3) + \frac{\alpha^4}{24}T - 2(1+3\nu)L^1 - \frac{4}{3}L^7 - 2(\nu + \frac{1}{12})\alpha^2L^2 - 4(\nu + \frac{1}{12})\alpha^2L^3,
$$
\n(2.23)

which is the  $g^{(2)}$  Lie-algebraic form of the Sutherland Hamiltonian. The polynomial eigenfunctions of  $(2.22)$  or  $(2.23)$  have the form  $(B.1)$  (see also Fig.2 in Appendix B). The operator $(2.23)$  contains now the generator T, given by  $(B.4)$  $(B.4)$ , which does not belong to  $gl_2 \ltimes R^3$ . Note that this generator T is not of a positive-grading.

Since both Lie-algebraic forms([2.20\)](#page-4-0), (2.23) do not contain the positive-grading generators  $J_2^+, J_3^+$  and  $L^4$ , hence the corresponding flags of the polynomial spaces (A.2) and (B.1) are preserved. This demonstrates the exact solvability of the Sutherland model following the criterion formulated in the Introduction. Observe that if we put  $\omega = 0$  and  $\alpha = 0$ , the algebraic and the Lie-algebraic forms of the Sutherland model reduce to the corresponding forms of the Calogero model.

#### 3. Complementarity

Let us consider a quantum-mechanical 3-body system. There are two systems of translation-invariant relative coordinates  $y$  and  $\tilde{y}$  related by:

$$
\tilde{y}_1 = y_1 - y_2 ,\n\tilde{y}_2 = y_2 - y_3 ,\n\tilde{y}_3 = y_3 - y_1 .
$$
\n(3.1)

Let the Hamiltonian for this 3-body system be given by

$$
\mathcal{H} = -A\frac{\partial^2}{\partial Y^2} - B\left(\frac{\partial^2}{\partial y_1^2} + \frac{\partial^2}{\partial y_2^2} - \frac{\partial^2}{\partial y_1 \partial y_2}\right) + V_1(y) + V_2(\tilde{y}),\tag{3.2}
$$

where

$$
V_1(y) = \sum_{i=1}^3 V_1(y_i) , V_2(\tilde{y}) = \sum_{i=1}^3 V_2(\tilde{y}_i) .
$$

<span id="page-6-0"></span>Here Y is the center-of-mass coordinate and  $A, B$  are some constants. Suppose the coordinates  $y$  in (3.2) are the Perelomov coordinates [\(2.5](#page-2-0)), then one can easily see that  $A = 3, B = \frac{2}{3}$  $\frac{2}{3}$  and the coordinates  $\tilde{y}$  are nothing but the Jacobi coordinates

$$
y_i = x_i - x_{i+1} \tag{3.3}
$$

On the other hand, if we take the y's as the Jacobi coordinates, then in [\(3.2\)](#page-5-0)  $A = 3, B = 2$ and the  $\tilde{y}$ 's become the Perelomov coordinates. So beginning from some Hamiltonian with a definite choice of relative coordinates, we arrive, after the change of variables  $(3.1)$  $(3.1)$ , at a Hamiltonian where the potentials  $V_1$  and  $V_2$  are interchanged. Of course, the spectra remain the same under such an operation, which makes it possible to connect the Hamiltonians with different coupling constants for two- and three-body interactions. Thus, we can put in correspondence two distinct physical problems by regarding them as related by some kind of 'complementarity' (equivalence). It is important to note that this peculiar feature appears only for 3-body problems.

In order to illustrate the above discussed complementarity, let us consider the Hamil-tonian [\(2.22\)](#page-5-0).Introduce the new coordinates (by replacing  $y \to \tilde{y}$  in ([2.21\)](#page-5-0))

$$
\tilde{\sigma}_1 := \sigma_1(\tilde{y}) = \frac{1}{\alpha^2} \left[ \cos(\alpha(y_1 - y_2)) + \cos(\alpha(y_2 - y_3)) + \cos(\alpha(y_3 - y_1)) - 3 \right], (3.4)
$$

$$
\tilde{\sigma}_2 := \sigma_2(\tilde{y}) = \frac{4}{\alpha^6} \left[ \sin(\alpha(y_1 - y_2)) + \sin(\alpha(y_2 - y_3)) + \sin(\alpha(y_3 - y_1)) \right]^2.
$$
 (3.5)

Itcan be verified that these coordinates are related to the original  $\sigma$  coordinates ([2.21\)](#page-5-0) by the following non-trivial algebraic relations

$$
\tilde{\sigma}_1 = 3\sigma_1 + \frac{1}{2}\alpha^2\sigma_1^2 + \frac{1}{8}\alpha^4\sigma_2 , \qquad (3.6)
$$

$$
\tilde{\sigma}_2 = -27\sigma_2 - 4\sigma_1^3 - 9\alpha^2 \sigma_1 \sigma_2 - \alpha^2 \sigma_1^4 - \frac{1}{2} \alpha^4 \sigma_1^2 \sigma_2 - \frac{1}{16} \alpha^6 \sigma_2^2 \,. \tag{3.7}
$$

Adding back the center-of-mass coordinate to the Hamiltonian [\(2.22\)](#page-5-0) with  $\sigma_1, \sigma_2$  replaced by  $\tilde{\sigma}_1, \tilde{\sigma}_2$ , followed by the gauge transformation

$$
\mathcal{H}_{3B} = \Psi_0^{(3B)}(x) h_{\text{Suth}}(\tilde{\sigma}_1, \tilde{\sigma}_2) (\Psi_0^{(3B)}(x))^{-1} , \qquad (3.8)
$$

where the gauge factor is defined by

$$
\Psi_0^{(3B)}(x) = (\Delta_1^{(trig)}(x))^{\nu} , \qquad (3.9)
$$

with  $\Delta_1^{(trig)}(x) = \prod_{i \le j}^3 i, j \ne k \vert \sin \frac{1}{2}(x_i + x_j - 2x_k) \vert$ , we finally arrive at the 3-body Hamiltonian with the 3-body interaction studied by Quesne[[6](#page-17-0)]

$$
\mathcal{H}_{3B} = -\frac{1}{2} \sum_{k=1}^{3} \frac{\partial^2}{\partial x_k^2} + \frac{g_1}{4} \sum_{\substack{k < l \\ k, l \neq m}}^{3} \frac{1}{\sin^2(\frac{1}{2}(x_k + x_l - 2x_m))} \tag{3.10}
$$

Note that now the coupling constant  $g_1$  is given as

$$
g_1=3\nu(\nu-1)\ .
$$

Thus, using the 'complementarity', we have been able to relate the 3-body Sutherland model with two-body interactions and the coupling constant  $g = \nu(\nu - 1)$  to a three-body

<span id="page-7-0"></span>model with 3-body interactions and the coupling constant  $g_1 = 3\nu(\nu - 1)$ . Although the solvability of the last problem was already proved in [\[6](#page-17-0)] by finding the hidden algebra  $gl(3)$ , the complementarity gives an immediate explanation for the solvability and a very short and transparent way of proving it. It also clarifies why the polynomial eigenfunctions of this model are given by Jack-Sutherland polynomials as noted in [\[6](#page-17-0)].

## 4. SOLVABILITY OF THE  $G_2$  INTEGRABLE SYSTEM

Making use of the previously introduced relative coordinates, we shall derive in this section the algebraic and Lie-algebraic forms of the rational and trigonometric  $G_2$  models which lead to polynomial eigenfunctions. This fact together with the explicit calculation of the eigenvalues, exhibit the exact solvability of the model.

4.1. The rational  $G_2$  model. As pointed out in the Introduction, this model was solved exactly for the bound states by J. Wolfes [\[1\]](#page-17-0), who used the procedure of separation of variables. In this section the algebraic and Lie-algebraic nature of the solvability of this model will be shown.

We begin by recalling the Hamiltonian for the rational  $G_2$  model

$$
\mathcal{H}_{G_2}^{(r)} = \frac{1}{2} \sum_{k=1}^3 \left[ -\frac{\partial^2}{\partial x_k^2} + \omega^2 x_k^2 \right] + g \sum_{k(4.1)
$$

where  $g = \nu(\nu - 1) > -\frac{1}{4}$  $\frac{1}{4}$  and  $g_1 = 3\mu(\mu - 1) > -\frac{3}{4}$  $\frac{3}{4}$  are the coupling constants associated with the 2-body and 3-body interactions, respectively. In this case we have two coupling constants and two different solutions for each value of  $\nu$  and  $\mu$ , which implies an existence of four families of solutions. The ground-state eigenfunction is given by

$$
\Psi_0^{(r)}(x) = (\Delta^{(r)}(x))^{\nu} (\Delta_1^{(r)}(x))^{\mu} e^{-\frac{1}{2}\omega \sum x_i^2}, \qquad (4.2)
$$

where  $\Delta^{(r)}(x) = \prod_{i \leq j}^{3} |x_i - x_j|$  and  $\Delta_1^{(r)}(x) = \prod_{i \leq j; i, j \neq k} |x_i + x_j - 2x_k|$ . Wolfes showed explicitly that any solution to (4.1) can be written in factorizable form as

$$
\Psi(x) = \Psi_0^r(x) P_{G_2}^{(r)}(x) , \qquad (4.3)
$$

where  $P_{G_2}^{(r)}$  $G_2^{(r)}(x)$  is a polynomial symmetric under permutations of any two of the  $x_i$ 's. The operator having these polynomials as eigenfunctions can be obtained by gauge rotating  $(4.1):$ 

$$
h_{\mathcal{G}_2}^{(r)} = -2(\Psi_0^{(r)}(x))^{-1} \mathcal{H}_{\mathcal{G}_2}^{(r)} \Psi_0^{(r)}(x) . \qquad (4.4)
$$

It then follows that in terms of the coordinates  $\tau$ , given in [\(2.6](#page-3-0))–([2.7\)](#page-3-0), the operator (4.4) takes the form

$$
h_{\mathcal{G}_2}^{(r)} = -2\tau_2 \partial_{\tau_2 \tau_2}^2 - 6\tau_3 \partial_{\tau_2 \tau_3}^2 + \frac{2}{3} \tau_2^2 \partial_{\tau_3 \tau_3}^2 - \{4\omega \tau_2 + 2[1 + 3(\mu + \nu)]\} \partial_{\tau_2} - 6\omega \tau_3 \partial_{\tau_3} , \tag{4.5}
$$

afterdropping out the center-of-mass dependence (cf.  $(2.8)$  $(2.8)$ ). This operator is the first algebraic form of the rational  $G_2$  model. It can be immediately rewritten in terms of the <span id="page-8-0"></span>generators of the algebra  $gl(3)$  (see (A.1) where  $n = 0$  and  $x, y$  are replaced by  $\tau_2, \tau_3$ , respectively)

$$
h_{\mathcal{G}_2}^{(r)} = -2J_{2,2}^0 J_2^- - 6J_{3,3}^0 J_2^- + \frac{2}{3}J_{2,3}^0 J_{2,3}^0 - 4\omega J_{2,2}^0 - 2[1 + 3(\mu + \nu)]J_2^- - 6\omega J_{3,3}^0 ,\tag{4.6}
$$

(cf.([2.9\)](#page-3-0)). Eq. (4.6) is the  $gl(3)$  Lie-algebraic form of the rational  $G_2$  model.

The model  $h_{G_2}^{(r)}$  $G_2^{(r)}$  admits another set of algebraic and Lie-algebraic forms. They can easilybe obtained if the operator ([4.4\)](#page-7-0) is written in terms of the coordinates  $\lambda_1, \lambda_2$  given by [\(2.10\)](#page-3-0). Thus,

$$
h_{\mathcal{G}_2}^{(r)} = -2\lambda_1 \partial_{\lambda_1 \lambda_1}^2 - 12\lambda_2 \partial_{\lambda_1 \lambda_2}^2 + \frac{8}{3} \lambda_1^2 \lambda_2 \partial_{\lambda_2 \lambda_2}^2 - \left\{ 4\omega \lambda_1 + 2[1 + 3(\mu + \nu)] \right\} \partial_{\lambda_1} - (12\omega \lambda_2 - \frac{4}{3} \lambda_1^2) \partial_{\lambda_2} ,
$$
\n(4.7)

 $(cf.(2.11))$  $(cf.(2.11))$  $(cf.(2.11))$ . This is the second *algebraic* form of the rational  $G_2$  model, which also admits a representation in terms of the generators of the algebra  $g^{(2)}$  given in (B.2), where  $n = 0$ and x, y are replaced by  $\lambda_1, \lambda_2$ , respectively, as

$$
h_{\mathcal{G}_2}^{(r)} = -2L^2L^1 - 12L^3L^1 + \frac{8}{3}L^7L^3 - 2[1 + 3(\mu + \nu)]L^1 - 4\omega L^2 - 12\omega L^3 - \frac{4}{3}L^7
$$
\n(4.8)

(cf.([2.12\)](#page-3-0)). Eq. (4.8) is the  $g^{(2)}$  Lie-algebraic form of the rational  $G_2$  model. The operator (4.8) depends on the generators of the  $gl_2 \ltimes R^3$ -algebra only. This implies that (4.8) possesses two invariant subspaces,  $W_n$  and  $\tilde{W}_n$  (see (B.1), (B.3)). Thus, it leads to the conclusion that there exists a family of eigenfunctions depending only on the variable  $\lambda_1$ . This, in fact, is already known both for this model [\[1\]](#page-17-0) as well as for the general many-body Calogero model[[3](#page-17-0)]. This fact was used in[[12\]](#page-17-0) to construct quasi-exactly-solvable manybody problems which generalize the Calogero model. We should emphasize that the same property allows one to construct also quasi-exactly-solvable generalizations of the rational  $G_2$  model. Such generalizations will be presented elsewhere.

In conclusion, one can state that the rational  $G_2$  model, as it is for the case of the Calogero and Sutherland models, admits two different algebraic and also two different Liealgebraic forms and, eventually, is characterized by two different hidden algebras. Since the Lie-algebraic forms (4.6), (4.8) contain no positive-grading generators, four infinite families of polynomial eigenfunctions of([4.5](#page-7-0)) and (4.7) occur according to the different values of  $\nu, \mu$ .

4.2. The trigonometric  $G_2$  model. The Hamiltonian for the trigonometric  $G_2$  model has the form

$$
\mathcal{H}_{G_2} = -\frac{1}{2} \sum_{k=1}^3 \frac{\partial^2}{\partial x_k^2} + \frac{g}{4} \sum_{k(4.9)
$$

where  $g = \nu(\nu - 1) > -\frac{1}{4}$  $\frac{1}{4}$  and  $g_1 = 3\mu(\mu - 1) > -\frac{3}{4}$  $\frac{3}{4}$  are the coupling constants associated with the 2-body and 3-body interactions, respectively. Again, as in the previous case,

<span id="page-9-0"></span>there are four families of solutions depending on the permitted values for  $\nu$  and  $\mu$ . The ground-state eigenfunction is given by

$$
\Psi_0^{(t)}(x) = (\Delta^{(trig)}(x))^{\nu} (\Delta_1^{(trig)}(x))^{\mu} , \qquad (4.10)
$$

where  $\Delta^{(trig)}(x)$ ,  $\Delta_1^{(trig)}$  $1^{(trig)}(x)$  are the trigonometric analogies of the Vandermonde determinant and are defined by

$$
\Delta^{(trig)}(x) = \prod_{i < j}^{3} |\sin \frac{1}{2}(x_i - x_j)|,
$$
\n
$$
\Delta_1^{(trig)}(x) = \prod_{\substack{k < l \\ k, l \neq m}}^{3} |\sin \frac{1}{2}(x_i + x_j - 2x_k)|.
$$

Guided by what occurs with the Calogero, Sutherland, rational and degenerate trigonometric  $G_2$  models, let us check whether there exist families of eigenfunctions of  $(4.9)$  of the type  $\Psi(x) = \Psi_0^{(t)}(x) P_{(G_2)}(x)$ , where the  $P_{(G_2)}$  are polynomials. If this is the case, there is a chance that following the conjectures made in  $[7]$  the trigonometric  $G_2$  model also possesses a hidden algebraic structure. For this purpose we make the gauge transformation of([4.9\)](#page-8-0) with the ground state eigenfunction (4.10) as the gauge factor. We get

$$
h_{\mathcal{G}_2} = -\frac{2}{3} (\Psi_0^{(t)}(x))^{-1} \mathcal{H}_{\mathcal{G}_2} (\Psi_0^{(t)}(x)) . \tag{4.11}
$$

Rewriting (4.11) in terms of the new coordinates  $\tilde{\sigma}$  given in [\(3.4\)](#page-6-0)–([3.5](#page-6-0)) we arrive, after some calculations accompanied by some miraculous cancellations, at an operator with coefficients which are surprisingly polynomials. The resulting expression is

$$
h_{G_2} = (2\tilde{\sigma}_1 + \frac{\alpha^2}{2}\tilde{\sigma}_1^2 - \frac{\alpha^4}{24}\tilde{\sigma}_2)\partial_{\tilde{\sigma}_1\tilde{\sigma}_1}^2 + (12 + \frac{8\alpha^2}{3}\tilde{\sigma}_1)\tilde{\sigma}_2\partial_{\tilde{\sigma}_1\tilde{\sigma}_2}^2
$$
\n
$$
-(\frac{8}{3}\tilde{\sigma}_1^2\tilde{\sigma}_2 - 2\alpha^2\tilde{\sigma}_2^2)\partial_{\tilde{\sigma}_2\tilde{\sigma}_2}^2 + \{2[1 + 3(\mu + 2\nu)] + \frac{2}{3}(1 + 3\mu + 4\nu)\alpha^2\tilde{\sigma}_1\}\partial_{\tilde{\sigma}_1}
$$
\n
$$
-\left\{\frac{4}{3}(1 + 4\nu)\tilde{\sigma}_1^2 - [\frac{7}{3} + 4(\mu + \nu)]\alpha^2\tilde{\sigma}_2\right\}\partial_{\tilde{\sigma}_2},
$$
\n(4.12)

(cf.([2.11\)](#page-3-0),([2.22\)](#page-5-0), ([4.7](#page-8-0))). Eq. (4.12) is the *algebraic* form of the trigonometric  $G_2$  model, and can be written in terms of the generators of the algebra  $g^{(2)}$  containing the generators of its subalgebra  $gl_2 \times R^3$ , plus the extra generator T. The explicit expression is given by

$$
h_{G_2} = 2L^2L^1 + 12L^3L^1 - \frac{8}{3}L^7L^3 + \alpha^2(\frac{L^2L^2}{2} + \frac{8L^3L^2}{3} + 2L^3L^3) - \frac{\alpha^4}{24}T \quad (4.13)
$$
  
+2[1 + 3(\mu + 2\nu)]L^1 + \frac{4}{3}(1 - 4\nu)L^7  
+ (2\mu + \frac{8}{3}\nu + \frac{1}{6})\alpha^2L^2 + \left[\frac{1}{3} + 4(\mu + \nu)\right]\alpha^2L^3 ,

where in the generators (B.2),(B.4) the parameter  $n = 0$  and x, y are replaced by  $\tilde{\sigma}_1$ ,  $\tilde{\sigma}_2$ , re-spectively (cf.([2.23](#page-5-0)),[\(4.8](#page-8-0))). This is the Lie-algebraic form of the trigonometric  $G_2$  model. However, unlike all previously discussed examples including the rational  $G_2$  model, there does not exist anymore a  $gl(3)$  Lie-algebraic representation for the trigonometric  $G_2$  model.

<span id="page-10-0"></span>A partial explanation for this fact is related to the non-existence of the algebraic representation of  $\sigma_{1,2}$  in terms of  $\tilde{\sigma}_{1,2}$ , similar to the representation [\(3.6\)](#page-6-0)–([3.7\)](#page-6-0).

The Lie-algebraic form  $(4.13)$  contains no positive-grading generators such as  $L<sup>4</sup>$  in the algebra  $g^{(2)}$  and, hence, the flag of polynomial spaces (B.1) is preserved. This demonstrates the exact solvability of the trigonometric  $G_2$  model, once again following the criterion formulated in the Introduction. Note that there are four infinite families of polynomial eigenfunctions labeled by the different values of  $\nu$ ,  $\mu$ . These polynomials are generalizations of the Jack-Sutherland polynomials.

The next question is to find the spectrum of the Hamiltonian [\(4.9\)](#page-8-0). First, let us note thatthe operator ([4.12\)](#page-9-0) has a block triangular form in the  $\tilde{\sigma}$  variables. There is a simple trick allowing us to reduce this operator to pure triangular form, based on the introduction of the new variables

$$
\rho_1 = \tilde{\sigma}_1 , \qquad \qquad \rho_2 = \tilde{\sigma}_2 + \frac{4}{\alpha^2} \tilde{\sigma}_1^2 . \qquad (4.14)
$$

It is worth noticing that if  $\alpha \to 0$  this change of variables becomes singular, reflecting the non-existence of bound states for the rational Calogero and  $G_2$  models in the absence of the harmonic oscillator term in potential. This coordinate transformation has the very attractive property of leaving the space (B.1) invariant.

In these new coordinates the Hamiltonian([4.12\)](#page-9-0) takes the form

$$
h_{G_2} = (2\rho_1 + \frac{2}{3}\alpha^2\rho_1^2 - \frac{\alpha^4}{24}\rho_2)\partial_{\rho_1\rho_1}^2 + (12\rho_2 + 2\alpha^2\rho_1\rho_2 - \frac{16}{\alpha^2}\rho_1^2)\partial_{\rho_1\rho_2}^2 + (2\alpha^2\rho_2^2 + \frac{96}{\alpha^2}\rho_1\rho_2 - \frac{256}{\alpha^4}\rho_1^3)\partial_{\rho_2\rho_2}^2 + [2(1+3\mu+6\nu) + \frac{2}{3}(1+3\mu+4\nu)\alpha^2\rho_1]\partial_{\rho_1} + \{2(1+2\mu+2\nu)\alpha^2\rho_2 + \frac{16}{\alpha^2}(2+3\mu+6\nu)\rho_1\}\partial_{\rho_2},
$$
\n(4.15)

and it is easy to check that it is indeed a triangular operator. Evidently, this operator can be rewritten in terms of the  $g^{(2)}$ -generators as

$$
h_{G_2} = 2L^2L^1 + 12L^1L^3 - \frac{\alpha^4}{24}T + \frac{2}{3}\alpha^2L^2L^2 + 2\alpha^2L^2L^3 + 2\alpha^2L^3L^3 \qquad (4.16)
$$
  

$$
-\frac{16}{\alpha^2}L^7L^1 + \frac{96}{\alpha^2}L^3L^6 - \frac{256}{\alpha^2}L^6L^7 + 2(1 + 3\mu + 6\nu)L^1
$$
  

$$
+(2\mu + \frac{8}{3}\nu)\alpha^2L^2 + 4(\mu + \nu)\alpha^2L^3 + \frac{16}{\alpha^2}(2 + 3\mu + 6\nu)L^6,
$$

where in the generators (B.2),(B.4) the parameter  $n = 0$  and x, y are replaced by  $\rho_1, \rho_2$ , respectively.

Using either one of the representations  $(4.15)$  or  $(4.16)$  the energy levels of the Hamiltonian  $H_{G_2}$  can be easily found and are given by

$$
\mathcal{E}_{n,m} = \left[ \frac{2}{3}(n-2m-1)(n+m+1) + 2n\mu + \frac{4}{3}(2n-m)\nu + \frac{2}{3} \right] \alpha^2 ,\qquad (4.17)
$$

where  $n, m$  are quantum numbers with

$$
n = 0, 1, 2, 3, ...
$$
,  $0 \le m \le \left[\frac{n}{2}\right]$ .

The explicit expressions for the first several eigenfunctions of [\(4.15](#page-10-0)) are presented in Appendix C.

## 5. Conclusion

We have found that the general trigonometric  $G_2$  integrable model with two arbitrary coupling constants is exactly-solvable. This model is characterized by a certain hidden algebra, which is an infinite-dimensional Lie algebra of the vector fields. It is quite amazing that for all degenerations of the general  $G_2$  model and also generalizations of the rational cases obtained by adding the harmonic oscillator interaction, there exists an alternative hidden algebra  $gl(3)$ , which disappears only in the general case of two non-vanishing coupling constants. For completeness of the presentation, the explicit forms for some of the eigenvalues of the general trigonometric  $G_2$  integrable model are given.

It is important to emphasize that the algebraic forms for the Hamiltonians of the Calogero, Sutherland, rational and trigonometric  $G_2$  integrable models  $(2.8)$ ,  $(2.11)$ ,  $(2.19)$ ,  $(2.22)$ ,  $(4.5)$ ,  $(4.7)$ ,  $(4.12)$ ,  $(4.15)$  are not contained in the list of algebraic forms of the Schroedinger operators possessing hidden algebraic structures, given in the papers [\[13](#page-17-0), [14, 15](#page-17-0)].

#### APPENDIX A. REPRESENTATION OF THE ALGEBRA  $gl(3)$

The algebra  $gl(3)$  has a realization in terms of first order differential operators acting on the  $(x, y)$ -plane which are given by

$$
J_2^- = \partial_x \quad (-1,0) , \qquad J_{2,2}^0 = x \partial_x \quad (0,0) , \qquad J_{3,2}^0 = y \partial_x \quad (-1,+1) ,
$$
  
\n
$$
J_3^- = \partial_y \quad (0,-1) , \qquad J_{3,3}^0 = y \partial_y \quad (0,0) , \qquad J_{2,3}^0 = x \partial_y \quad (+1,-1) ,
$$
  
\n
$$
J^0 = n - x \partial_x - y \partial_y \quad (0,0) , \qquad J_2^+ = x J^0 \quad (+1,0) , \qquad J_3^+ = y J^0 \quad (+1,0) ,
$$
  
\n(A.1)

where  $n \in R$ . In this realization a grading  $(\alpha, \beta)$  can assigned to the generators  $(A.1)$  by observing that  $Jx^py^q \propto x^{p+\alpha}y^{q+\beta}$ . If n is a non-negative integer, the representation (A.1) becomes finite-dimensional and the representation space

$$
V_n = (x^p y^q | 0 \le (p+q) \le n),
$$
\n(A.2)

is characterized by the Newton diagram (see Fig.1).



Fig.1 The Newton diagram illustrating the finite-dimensional representation space (A.2) of the algebra  $gl(3)$ .

The algebra (A.1) acts on the space (A.2) irreducibly.

## APPENDIX B. REPRESENTATION OF THE ALGEBRA  $g^{(2)}$

We define the algebra  $g^{(2)}$  as an infinite-dimensional algebra of differential operators on the  $(x, y)$ -plane possessing a finite-dimensional representation which acts irreducibly on the space of inhomogeneous polynomials

$$
W_n = \langle x^p y^q | 0 \le (p + 2q) \le n \rangle,
$$
\n(B.1)

which is diagrammatically represented by Fig.2.

<span id="page-13-0"></span>

Fig.2 The Newton diagram illustrating the finite-dimensional representation space (B.1) of the algebras  $g^{(2)}$  and  $gl_2 \ltimes R^3$ .

The algebra  $g^{(2)}$  contains the subalgebra  $gl_2 \ltimes R^3$  of first order differential operators

$$
L^{1} = \partial_{x} \qquad (-1,0) , \qquad L^{2} = x\partial_{x} - \frac{n}{3} \qquad (0,0) ,
$$
  
\n
$$
L^{3} = y\partial_{y} - \frac{n}{6} \qquad (0,0) , \qquad L^{4} = x^{2}\partial_{x} + 2xy\partial_{y} - nx \quad (+1,0) ,
$$
  
\n
$$
L^{5} = \partial_{y} \qquad (0,-1) , \qquad L^{6} = x\partial_{y} \qquad (+1,-1) ,
$$
  
\n
$$
L^{7} = x^{2}\partial_{y} \qquad (+2,-1) , \qquad (B.2)
$$

where  $($ ,  $)$  denotes the grading (defined in the similar way as in Appendix A) and n is a non-negative integer. The  $gl_2 \ltimes R^3$ -algebra, (B.2) is characterized by the commutation relations

$$
\begin{array}{llll}\n[L^1, L^2] = L^1, & [L^1, L^3] = 0, & [L^1, L^4] = 2(L^2 + L^3), \\
[L^2, L^3] = 0, & [L^2, L^4] = L^4, & [L^3, L^4] = 0, \\
[L^{5+k}, L^1] = -kL^{5+k-1}, & [L^{5+k}, L^2] = kL^{5+k}, & [L^{5+k}, L^3] = 0, \\
[L^{5+k}, L^4] = (2 - k)L^{5+k+1}, & [L^{5+k}, L^{5+m}] = 0,\n\end{array}
$$

and the generator  $L^4$  is the only positive-root generator of the algebra  $gl_2 \ltimes R^3$ . If this generator is omitted the remaining generators form the Borel subalgebra of  $gl_2 \ltimes R^3$ . The algebra  $gl_2 \ltimes R^3$  acts on (B.1) reducibly possessing an invariant subspace <sup>4</sup>

$$
\tilde{W}_n = \langle x^p | 0 \le p \le n \rangle . \tag{B.3}
$$

In order to obtain the Lie-algebraic forms of Calogero and rational  $G_2$ -integrable models it is sufficient to use the  $gl_2 \times R^3$  generators only representing their Hamiltonians. However, the Lie-algebraic forms of the Sutherland and trigonometric  $G_2$ -integrable models require more generators than these the  $gl_2 \ltimes R^3$  algebra can provide. One of such extra generators leaving  $W_n$  invariant has the form

$$
T = y \partial_{xx}^2 \t{,} \t(B.4)
$$

and is characterized by the grading  $(-2, +1)$ . Clearly it does not belong to the universal enveloping algebra of the  $g\bar{l}_2 \ltimes R^3$ -algebra. The generator (B.4) does act on the space (B.3). Thus, the space (B.3) is no longer a common invariant subspace with respect to

<sup>&</sup>lt;sup>4</sup>We are grateful to R. Bautista for fruitful discussion on the  $gl_2 \ltimes R^3$ -algebra

the action of the operators  $(B.2)$ ,  $(B.4)$ . Consequently, the operators  $(B.2)$ ,  $(B.4)$  act on (B.1) irreducibly and one can state that the  $g^{(2)}$ -algebra coincides with the algebra of all polynomials in the generators  $(B.2)$ ,  $(B.3)$ . The proof is based on Burnside theorem (see discussion in [\[7](#page-17-0)]).

Due to the fact that the Hamiltonians we study are represented by second order differential operators, it would be instructive to classify all second order differential operators acting on  $W_n$ , or, equivalently, these having  $W_n$  as the invariant subspace. It is clear that some of these operators are given by quadratic elements of the universal enveloping algebra of  $gl_2 \ltimes R^3$  being quadratic polynomials in the generators (B.2). In order to find the remaining ones we should consider all possible the commutators, double-commutators etc between the operator T and the generators of the algebra  $gl_2 \ltimes R^3$ .

Firstly, let us check the commutation relations of T with the  $gl_2 \ltimes R^3$ -generators:

$$
[L^{1}, T] = 0 , \t [L^{2}, T] = -2T , \t [L^{3}, T] = T ,
$$
  
\n
$$
[L^{4}, T] = -2xy\partial_{xx}^{2} - 4y^{2}\partial_{xy}^{2} + 2(n - 1)y\partial_{x} \equiv T^{1} \t (-1, +1) ,
$$
  
\n
$$
[L^{5+k}, T] = x^{k}\partial_{xx}^{2} - k(k - 1)x^{k-2}y\partial_{y} - 2kx^{k-1}y\partial_{xy}^{2} \t (B.5)
$$
  
\n
$$
= \begin{cases} L^{1}L^{1} , & k = 0 \\ L^{2}L^{1} - 2L^{1}L^{3} , & k = 1 \\ L^{2}L^{2} - 4L^{2}L^{3} - L^{2} \\ - (2 + \frac{4}{3}n)L^{3} - (\frac{2}{3} + \frac{n}{9})n , & k = 2 . \end{cases}
$$

We find that except for the commutator  $[L^4, T] \equiv T^1$  all other commutators belong to the universal enveloping algebra of  $gl_2 \ltimes R^3$ . Now among the double commutators we should consider only those involving the commutators of  $T^1$  with the  $gl_2 \ltimes R^3$ -generators, namely

$$
[L^1, T^1] = -2T , \t [L^5, T^1] = -2L^1(L^2 + 4L^3) ,
$$
  
\n
$$
[L^2, T^1] = -T^1 , \t [L^6, T^1] = -2L^4L^1 + 4L^3L^3 - 2L^2
$$
  
\n
$$
[L^3, T^1] = T^1 , \t -2(1 + \frac{n}{3})L^3 - n(1 + \frac{2}{3}n) ,
$$
  
\n
$$
[T, T^1] = 0 , \t [L^7, T^1] = -2(L^2 - 2L^3)L^4 + 2L^4
$$
  
\n
$$
[L^4, T^1] = 2x^2y\partial_{xx}^2 + 8xy^2\partial_{xy}^2 + 8y^3\partial_{yy}^2
$$
  
\n
$$
+ 4(n - 1)xy\partial_x + (12 - 8n)y^2\partial_y + 2n(n - 1)y
$$
  
\n
$$
\equiv T^2 (0, +1) .
$$
  
\n(B.6)

Hence, the only new operator that appears is  $T^2$ . Repeating the above procedure with  $T^2$ we get

$$
[L1, T2] = -2T1 ,\n[L3, T2] = T2 ,\n[L4, T2] = 0 ,\n[L5, T2] = -2L1L4 + 24L3L3 + 12L2L3 - 4(1 + n)L3 - \frac{2}{3}n ,\n[L6, T2] = 2L2L4 + 8L3L4 ,\n[L7, T2] = 2L4L4 ,\n[L7, T2] = 2L4L4 , (B.7)
$$

As seen there are no new operators found. Therefore the three generators  $T, T^1, T^2$  exhaust all generators which can be represented by second order differential operators and which do not belong to the universal enveloping algebra of the  $gl_2\ltimes R^3$  algebra. We should emphasize the surprising fact that these new generators  $T, T^1, T^2$  form an abelian subalgebra of  $g^{(2)}$ (see  $(B.6), (B.7)$ ).

APPENDIX C. EIGENFUNCTIONS OF THE  $G_2$  trigonometric model

We present here the explicit expressions for the first six eigenfunctions and eigenvalues ofthe general trigonometric  $G_2$  model for the Hamiltonian ([4.15\)](#page-10-0).

$$
\begin{array}{rcl} \mathcal{E}_{1,0} & = & \left( \frac{2}{3} + 2\mu + \frac{8}{3}\nu \right) \alpha^2 \ , \\ \mathcal{E}_{2,0} & = & \rho_1 + \frac{1+3\mu+6\nu}{(1+3\mu+4\nu)\alpha^2} \ ; \\ \mathcal{E}_{2,0} & = & \left( \frac{8}{3} + 4\mu + \frac{16}{3}\nu \right) \alpha^2 \ , \\ \mathcal{E}_{2,1} & = & \left( -2+4\mu+4\nu \right) \alpha^2 \ , \\ \mathcal{E}_{2,1} & = & \left( -2+4\mu+4\nu \right) \alpha^2 \ , \\ \mathcal{E}_{2,1} & = & \left( -2+4\mu+4\nu \right) \alpha^2 \ , \\ \mathcal{E}_{2,1} & = & \left( -2+4\mu+4\nu \right) \alpha^2 \ , \\ \mathcal{E}_{2,1} & = & \left( -2+4\mu+4\nu \right) \alpha^2 \ , \\ \mathcal{E}_{2,1} & = & \rho_1^2 + \frac{\alpha^2}{48}\rho_2 + \frac{3(2+\beta\mu+6\nu)}{2(-4+3\mu+2\nu)} \rho_1 + \frac{13(2+3\mu+6\nu)(1+3\mu+6\nu)}{2(-1+2\mu+2\nu)(-4+2\nu+3\mu)\alpha^4} \ ; \\ \mathcal{E}_{3,0} & = & \left( 6+6\mu+8\nu \right) \alpha^2 \ , \\ \mathcal{W}_{3,0} & = & \rho_1^3 - \frac{3\alpha^2}{16(1+\nu)} \rho_1 \rho_2 + \frac{9(12\nu^2 + (6\mu+12)\nu+5+3\mu)}{2(-1+2\mu+2\nu)(-4+2\nu+3\mu)\alpha^2} \ , \\ \mathcal{E}_{3,1} & = & \left( 6+6\mu+8\nu \right) \alpha^2 \ , \\ \mathcal{W}_{3,0} & = & \rho_1^3 - \frac{3\alpha^2}{16(1+\nu)(6+3\mu+6\nu)(2+2\nu+\mu)(6+4\mu+3\mu)\alpha^2} \ , \\ \mathcal{E}_{3,1} & = & \left( \frac{9}{4} + 6\mu + \frac{20}{3}\nu \right) \alpha^2 \ , \\ \mathcal{E}_{3,1
$$

#### **REFERENCES**

- <span id="page-17-0"></span>[1] J. Wolfes, "On the three-body linear problem with three-body interaction", J. Math. Phys. 15 (1974) 1420-1424.
- [2] F. Calogero and C. Marchioro, "Exact solution of a one-dimensional three-body scattering problem with two-body and/or three-body inverse-square potential", J. Math. Phys. 15 (1974) 1425–1430.
- [3] F. Calogero, "Solution of a three-body problem in one dimension", J. Math. Phys. 10 (1969) 2191–2196;

F. Calogero, "Ground state of a one-dimensional N-body problem", J. Math. Phys. 10 (1969) 2197–2200;

F. Calogero, "Solution of the one-dimensional N-body problem with quadratic and/or inversely quadratic pair potentials", J. Math. Phys. 12 (1971) 419–436.

- [4] M. A. Olshanetsky and A. M. Perelomov, "Quantum completely integrable systems connected with semi-simple Lie algebras", Lett. Math. Phys. 2 (1977) 7-13.
- [5] M. A. Olshanetsky and A. M. Perelomov, "Quantum integrable systems related to Lie algebras," Phys. Rep. 94 (1983) 313.
- [6] C. Quesne, "An exactly solvable three particle problem with three body interaction", hep-th/9612173.
- C. Quesne, "Three-body Generalizations of the Sutherland Problem", hep-th/9706067.
- [7] A. V. Turbiner, "Lie algebras and linear operators with invariant subspace", in Lie algebras, cohomologies and new findings in quantum mechanics (N. Kamran and P. J. Olver, eds.), AMS, vol. 160, pp. 263–310, 1994; funct-an/9301001 "Lie-algebras and Quasi-exactly-solvable Differential Equations", in CRC Handbook of Lie Group Analysis of Differential Equations, Vol.3: New Trends in Theoretical Developments and Computational Methods, Chapter 12, CRC Press (N. Ibragimov, ed.), pp. 331-366, 1995; hep-th/9409068.
- [8] W. Rühl and A. Turbiner, "Exact solvability of the Calogero and Sutherland models", Mod. Phys. Lett.  $A10$  (1995) 2213-2222, hep-th/9506105.
- [9] B. Sutherland, "Exact results for a quantum many-body problem in one dimension", Phys. Rev. A4 (1971) 2019–2021; B. Sutherland, "Exact results for a quantum many-body problem in one dimension, II", Phys. Rev. A5 (1972) 1372–1376.
- [10] L. Brink, A. Turbiner and N. Wyllard, "Hidden Algebras of the (super) Calogero and Sutherland models", Journ. Math. Phys.39 (1998) 1285-1315, hep-th/9705219
- [11] A. M. Perelomov, "Algebraic approach to the solution of a one-dimensional model of N interacting particles", Teor. Mat. Fiz. 6 (1971) 364 (in Russian); English translation: Sov. Phys. – Theor. and Math. Phys. 6 (1971) 263.
- [12] A. Minzoni, M. Rosenbaum and A. Turbiner, "Quasi-Exactly-Solvable Many-Body Problems", Mod. Phys. Lett. A11 (1996) 1977-1984, hep-th/9606092.
- [13] M.A. Shifman and A.V. Turbiner, "Quantal problems with partial algebraization of the spectrum", Comm. Math. Phys. 126 (1989) 347-365.
- [14] A. González-Lopéz, N. Kamran and P.J. Olver, "New quasi-exactly-solvable Hamiltonians in two dimensions", Comm.Math.Phys., 159 (1994) 503-537.
- [15] R. Milson, "On the Construction of Quasi-Exactly-Solvable Schroedinger Operators on Homogeneous Spaces", Journ. Math. Phys. 36 (1995) 6004–6027.