

The nondynamical r-matrix structure of the elliptic Calogero-Moser Model

Bo-yu Hou^b and Wen-li Yang^{a,b *†}

^a CCAST (World Laboratory), P.O.Box 8730 ,Beijing 100080, China

^b Institute of Modern Physics, Northwest University, Xian 710069, China [‡]

October 29, 2018

Abstract

1

In this paper, we construct a new Lax operator for the elliptic Calogero-Moser model with $N=2$. The nondynamical r-matrix structure of this Lax operator is also studied . The relation between our Lax operator and the Lax operator given by Krichever is also obtained.

Mathematics Subject Classification : 70F10 , 70H33 , 81U10.

1 Introduction

The elliptic Calogero-Moser (CM) model[1-4] is the system of N one-dimensional particles interacting by two-particle potential of the elliptic type. It is well-known that the CM model is completely integrable[1-8]. The Lax operator of this system , which is the most effective way to construct the complete set of integrals of motion, was found by Krichever[6] . The classical r-matrix structure of the Lax operator given by Krichever for the CM model was obtained by Sklyanin[7]. This r-matrix is a natural generalization of the matrix found by Avan et al [9] for the trigonometric potential. There exists a specific feature that the r-matrix for these model turns out to be of dynamical type (i.e it depends on the dynamical variables)

*e-mail :wlyang@nwu.edu.cn

†Fax :0086-029-8303511

‡Mailing address

¹This paper was given as a lecture in the international meeting “ New developments in quantum field theories” in Urumqi,China,10-19 August,1996

and satisfy a generalized Yang-Baxter equation[7-9]. Very recently development was connected with the geometrical interpretation of the dynamical r-matrix in terms of the Hamiltonian reduction[10].

The difficulty presented by the dynamic aspect of the r-matrix is : **I.** the Poisson algebra of a model ,whose structural constants are given by a dynamical r-matrix, is generally speaking no longer closed; **II.** To solve the quantization problem is still an open problem. So far ,only for one particular example—the spin generalization of the CM model—a proper algebraic setting (the Gervais-Neveu-Felder equation) was found [11] which also allows to quantize the model. To overcome the above difficulties may be that whether another Lax operator for the CM model which has a numerical r-matrix structure could be found. In this paper, we construct a new Lax operator for the CM model with $N=2$. In this new Lax representation , we find its r-matrix being of numerical type , which satisfies the classical Yang-Baxter equation. Further, the relation between the old one (given by Krichever) and ours is also obtained.

2 Review of the CM model

The Calogero-Moser model is the system of N one-dimensional particles interactin by the two-particle potential

$$\begin{aligned} V(q_{ij}) &= Q(q_{ij}) \quad , \quad q_{ij} = q_i - q_j \quad , \quad i, j = 1, \dots, N \quad , \\ Q(q) - Q(u) &= E(u, q)E(u, -q) \end{aligned} \quad (1)$$

and $E(u, q)$ is an elliptic function which is defined in Eq.(3). The identity Eq.(3a) is used in the second equation. In this paper, we restrict ourselves to the case $N=2$.

In terms of the canonical variables $p_i \quad , \quad q_i \quad (i=1,2)$

$$\{p_i, p_j\} = 0 \quad , \quad \{q_i, q_j\} = 0 \quad , \quad \{p_i, q_j\} = \delta_{i,j} \quad ,$$

the Hamiltonian of the system is expressed as

$$H = \sum_{i=1}^2 p_i^2 + \sum_{i \neq j} V(q_{ij}). \quad (2)$$

The Hamiltonian Eq.(2) with the potential Eq.(1) is known to be completely integrable. The most effective way to show its integrability is to construct the Lax representation for the system (namely, to find the Lax operator). One Lax representation for the CM model was first found by Krichever[6]

$$\tilde{L}(u) = \begin{pmatrix} p_1 & E(u, q_{12}) \\ E(u, q_{21}) & p_2 \end{pmatrix} \quad , \quad E(u, q) = \frac{\sigma(u+q)}{\sigma(u)\sigma(q)} \quad , \quad (3)$$

where the elliptic function $\sigma(u)$ is defined in Eq.(6). The Hamiltonian in Eq.(2) can be written as

$$H = tr(\tilde{L}^2(u)) + V(u)$$

where $V(u)$ does not depend upon the dynamical variables and the identity

$$E(u, q_{12})E(u, q_{21}) = V(q_{12}) - V(u) \quad (3a)$$

is used. The motion equation can be written in the following form [6]

$$\frac{d}{dt} \tilde{L}(u) = \{\tilde{L}(u), H\} = [\tilde{L}(u), \tilde{M}(u)]$$

The Poisson bracket of the Lax operator $\tilde{L}(u)$ can be described in terms of r-matrix form[7]

$$\{\tilde{L}_1(u), \tilde{L}_2(v)\} = [\tilde{r}_{12}(u, v), \tilde{L}_1(u)] - [\tilde{r}_{21}(v, u), \tilde{L}_2(v)] \quad (4)$$

$$\tilde{r}_{12}(u, v) = \begin{pmatrix} a & d_{12} & & \\ d_{21} & & c_{12} & \\ & & c_{21} & d_{12} \\ & & & d_{21} & a \end{pmatrix}$$

$$a = -\xi(u-v) - \xi(v) \quad , \quad \xi(u) = \partial_u \{\ln \sigma(u)\}$$

$$c_{ij} = -E(u-v, q_{ij}) \quad d_{ij} = -\frac{1}{2}E(v, q_{ij})$$

It can be seen that the classical r-matrix $\tilde{r}(u, v)$ is a dynamical r-matrix (i.e the matrix element of $\tilde{r}(u, v)$ depends upon the dynamical variables q_i). The Poisson bracket Eq.(4) leads to the evolution integrals $tr(\tilde{L}^n(u))$ of the motion. Sklyanin also shown that the dynamical r-matrix $\tilde{r}(u, v)$ defined by Eq.(4) satisfies the generalized Yang-Baxter equation[7]

$$[R^{(123)}, \tilde{L}^{(1)}] + [R^{(231)}, \tilde{L}^{(2)}] + [R^{(312)}, \tilde{L}^{(3)}] = 0 \quad (5)$$

where

$$R^{(123)} \equiv \tilde{r}_{(123)} - \{\tilde{r}_{13}, \tilde{L}^{(2)}\} + \{\tilde{r}_{12}, \tilde{L}^{(3)}\},$$

and

$$\tilde{r}_{(123)} = [\tilde{r}_{12}, \tilde{r}_{13}] + [\tilde{r}_{12}, \tilde{r}_{23}] - [\tilde{r}_{13}, \tilde{r}_{32}]$$

Due to the r-matrix $\tilde{r}(u, v)$ depending on the dynamical variables, the Poisson bracket of $\tilde{L}(u)$ is no longer closed. The quantum version of Eq.(4) and the generalized Yang-Baxter equation is still not found except the spin generalization of the CM model in which the Gervais-Neveu-Felder equation was found[11].

3 The new Lax representation for CM model

The Lax representation $\tilde{L}(u)$ of the CM model given by Krichever in Eq.(3) and its classical r-matrix $\tilde{r}(u, v)$ given by Sklyanin in Eq.(4) leads to some difficulty[7] in the investigation of the CM model : the Poisson algebra of the Lax operator is no longer closed and the quantum version of Eq.(4) is still not found. This motivate us to find a new Lax representation of the CM model. Fortunately, we find the new Lax representation of the CM model, in which the classical r-matrix is numerical one (This kind Lax representation does not always exist for generic system with dynamical r-matrix). In this section, we construct this new lax operator (in contrast to the Lax operator given by Krichever, we call the Lax operator found by us as the new Lax operator).

First,let us define some elliptic functions

$$\begin{aligned}
\theta^{(j)}(u) &= \theta \left[\begin{array}{c} \frac{1}{2} - \frac{j}{2} \\ \frac{1}{2} \end{array} \right] (u, 2\tau) \\
\sigma(u) &= \theta \left[\begin{array}{c} \frac{1}{2} \\ \frac{1}{2} \end{array} \right] (u, \tau) \\
\theta \left[\begin{array}{c} a \\ b \end{array} \right] (u, \tau) &= \sum_{m=-\infty}^{\infty} \exp\{i\pi[(m+a)^2\tau + 2(m+a)(z+b)]\} \\
\theta^{(j)}(u) &= \partial_u \{\theta^{(j)}(u)\}
\end{aligned} \tag{6}$$

where τ is a complex number with $Im(\tau) > 0$. We find that there exist another Lax representation for the CM model and denote it by $L(u)$

$$L(u) = \begin{pmatrix} L_{11}(u) & L_{12}(u) \\ L_{21}(u) & L_{22}(u) \end{pmatrix}, \tag{7}$$

where

$$\begin{aligned}
L_{11}(u) &= \{\theta_1(u - 2\bar{q}_1)\theta_2(u - 2\bar{q}_2)p_1 - \theta_1(u - 2\bar{q}_2)\theta_2(u - 2\bar{q}_2)E(u, q_{21}) \\
&\quad + \theta_1(u - 2\bar{q}_1)\theta_2(u - 2\bar{q}_1)E(u, q_{12}) - \theta_1(u - 2\bar{q}_2)\theta_2(u - 2\bar{q}_1)p_2\} \Delta^{-1} \\
L_{12}(u) &= \{-\theta_1(u - 2\bar{q}_1)\theta_1(u - 2\bar{q}_2)p_1 + \theta_1(u - 2\bar{q}_2)\theta_1(u - 2\bar{q}_2)E(u, q_{21}) \\
&\quad - \theta_1(u - 2\bar{q}_1)\theta_1(u - 2\bar{q}_1)E(u, q_{12}) + \theta_1(u - 2\bar{q}_2)\theta_1(u - 2\bar{q}_1)p_2\} \Delta^{-1} \\
L_{21}(u) &= \{\theta_2(u - 2\bar{q}_1)\theta_2(u - 2\bar{q}_2)p_1 - \theta_2(u - 2\bar{q}_2)\theta_2(u - 2\bar{q}_2)E(u, q_{21}) \\
&\quad + \theta_2(u - 2\bar{q}_1)\theta_2(u - 2\bar{q}_1)E(u, q_{12}) - \theta_2(u - 2\bar{q}_2)\theta_2(u - 2\bar{q}_1)p_2\} \Delta^{-1} \\
L_{22}(u) &= \{-\theta_2(u - 2\bar{q}_1)\theta_1(u - 2\bar{q}_2)p_1 + \theta_2(u - 2\bar{q}_2)\theta_1(u - 2\bar{q}_2)E(u, q_{21}) \\
&\quad - \theta_2(u - 2\bar{q}_1)\theta_1(u - 2\bar{q}_1)E(u, q_{12}) + \theta_2(u - 2\bar{q}_2)\theta_1(u - 2\bar{q}_1)p_2\} \Delta^{-1} \\
\Delta &= \sigma(q_1 - q_2)\sigma(u - q_1 - q_2)
\end{aligned}$$

and $\bar{q}_1 = \frac{q_1 - q_2}{2} = -\bar{q}_2$. The Hamiltonian Eq.(2) is equal to

$$H = \text{tr}(\tilde{L}^2(u)) + V(u) = \text{tr}(L^2(u)) + V(u)$$

The motion equation can also be described in the commutator form

$$\frac{d}{dt}L(u) = \{L(u), H\} = [L(u), M(u)]$$

The very ‘‘good’’ properties of our Lax operator defined in Eq.(7) is that the basic Poisson bracket of the Lax operator $L(u)$ can be written in the usual Poisson-Lie bracket form with a purely numerical r-matrix (i.e. nondynamical r-matrix). Namely,

$$\{L_1(u), L_2(v)\} = [r_{12}(u-v), L_1(u) + L_2(v)] \quad (8)$$

and the numerical r-matrix $r(u)$ reads as

$$r(u) = \begin{pmatrix} a(u) & & & d(u) \\ & b(u) & c(u) & \\ & c(u) & b(u) & \\ d(u) & & & a(u) \end{pmatrix}$$

and

$$\begin{aligned} a(u) &= \frac{\theta^{(0)}(u)}{\theta^{(0)}(u)} - \frac{\sigma'(u)}{\sigma(u)} \quad , \quad b(u) = \frac{\theta^{(1)}(u)}{\theta^{(1)}(u)} - \frac{\sigma'(u)}{\sigma(u)} \\ c(u) &= \frac{\theta^{(0)}(0)\theta^{(1)}(u)}{\theta^{(0)}(u)\theta^{(1)}(0)} \quad , \quad d(u) = \frac{\theta^{(0)}(0)\theta^{(0)}(u)}{\theta^{(1)}(u)\theta^{(1)}(0)} \end{aligned}$$

where $a(u), b(u), c(u), d(u)$ are all independent upon dynamical variable. The numerical r-matrix $r(u)$ defined in Eq.(8) satisfies the classical Yang-Baxter equation

$$[r_{12}(u-v), r_{13}(u-\eta)] + [r_{12}(u-v), r_{23}(v-\eta)] + [r_{13}(u-\eta), r_{23}(v-\eta)] = 0 \quad (9)$$

and enjoys in the antisymmetric properties

$$r_{12}(u) = -r_{21}(-u) \quad (10)$$

The standard Poisson-Lie bracket Eq.(8) of the Lax operator $L(u)$ and the numerical r-matrix $r(u)$ enjoying in the classical Yang-Baxter equation Eq.(9) and antisymmetry Eq.(10), make it possible to construct the quantum theory of the CM model.

4 The relation between two Lax representation

In fact, the new Lax representation given by us in Eq.(7) can be obtained from the old one given by Krichever in Eq.(3) through a dynamical gauge transformation as follows

$$L(u) = g(u) \tilde{L}(u) g^{-1}(u) \quad (11)$$

where

$$g(u) = \begin{pmatrix} \theta_1(u - 2\bar{q}_1) & -\theta_1(u - 2\bar{q}_2) \\ -\theta_2(u - 2\bar{q}_1) & \theta_2(u - 2\bar{q}_2) \end{pmatrix}$$

Due to the transformation Eq.(11) being dependent up the dynamical variable q_i , the classical r-matrix could not be transferred by a similarity transformation as that of the Lax operator in Eq.(11). Fortunately, through a straightforward calculation, we find that the numerical r-matrix $r(u - v)$ can be obtained from the dynamical one $\tilde{r}(u, v)$ as follows

$$r_{12}(u - v) = g_1(u)g_2(v) \tilde{r}_{12}(u, v)g_1^{-1}(u)g_2^{-1}(v) + g_2(v)\{g_1(u), \tilde{L}_2(v)\}g_1^{-1}(u)g_2^{-1}(v)$$

up to some matrix which does commute with $L_1(u) + L_2(v)$.

Summary

The numerical r-matrix $r(u)$ Eq.(6) could provide a mean to construct a separation of variables for the CM model in the same manner as that in the case of the integrable magnetic chains[13]. Moreover, the r-matrix $r(u)$ also make it possible that one can construct the dressing transformation for the CM model. The dressing group of this system would be an analogue to the classical limit of Sklyanin algebra[12,14]. A possible way to approach quantization of the CM model would be to look for a quantum version of the Poisson-Lie bracket Eq.(8). The most possible candidate is the elliptic Sklyanin algebra[12,14] which satisfies

$$R_{12}(u - v)T_1(u)T_2(v) = T_2(v)T_1(u)R_{12}(u - v) \quad (12)$$

where $R_{12}(u - v)$ is the eight-vertex Baxter's R-matrix and satisfies the quantum Yang-Baxter equation

$$R_{12}(u - v)R_{13}(u - \eta)R_{23}(v - \eta) = R_{23}(v - \eta)R_{13}(u - \eta)R_{12}(u - v) \quad (13)$$

and the unitary condition

$$R_{12}(u)R_{21}(-u) = 1 \quad (14)$$

The classical numerical r-matrix $r(u)$ have the relation with the quantum one $R(u)$ Eq.(12) as follows

$$R(u) = 1 + wr(u) + o(w^2) \quad , \quad \text{when the crossing parameter } w \longrightarrow 0 \quad (15)$$

In this paper, we only consider the special case of $N = 2$ for the CM model. However, the results can be generalized to the generic case of $2 \leq N$. We will present the further results in the further paper.

References

- [1] F.Calogero , Lett. Nuovo. Cim. **13** (1975) 411; Lett. Nuovo. Cim. **16** (1976) 77.
- [2] J.Moser, Adv. Math. **16** (1975) 1.
- [3] S.N.M.Ruijsenaars and H.Schneider, Ann. Phys. **Vol.170** (1986) 370.
- [4] S.N.M.Ruijsenaars, Comm. Math. Phys. **Vol.115** (1988) 127.
- [5] M.A.Olshanetsky and A.M.Perelomov , Phys. Rep. **71** (1981) 313.
- [6] I.M.Krichever, Func. Annal. Appl. **14** (1980) 282.
- [7] E.K.Sklyanin, *Dynamical r-matrices for the elliptic Calogero-Moser model*, hep-th/9308060 (1993).
- [8] H.W.Braden and T.Suzaki, Lett. Math. Phys. **Vol.30** (1994) 147.
- [9] J.Avan and M.Talon, Phys. Lett. **B 303** (1993) 33.
- [10] Yuri B. Suris, *Why are the rational and hyperbolic Ruijsenaars-Schneider hierarchies governed by the the same R-operators as the Calogero-Moser ones*, hep-th/9602160 (1996).
- [11] J.Avan, O.Babelon and E.Billey, *The Gervais-Neveu-Felder equation and the quantum Calogero-Moser systems*, hep-th/9505091 (1995).
- [12] B.Y.Hou, K.J.Shi and Z.X.Yang, J. Phys. **A 26**(1993) 4951.
- [13] E.K.Sklyanin, Comm.Math.Phys. **Vol.150**(1992) 181.
- [14] E.K.Sklyanin, Funct. Anal. Appl. **16**(1982) 263 ; Funct. Anal. Appl. **17**(1983) 320.