

A BILINEAR APPROACH TO DISCRETE MIURA TRANSFORMATIONS

N. JOSHI

*Dept. of Pure Mathematics
University of Adelaide
Adelaide 5005 Australia*

A. RAMANI

*CPT, Ecole Polytechnique
CNRS, UPR 14
91128 Palaiseau, France*

B. GRAMMATICOS

*GMPIB, Université Paris VII
Tour 24-14, 5^e étage
75251 Paris, France*

Abstract

We present a systematic approach to the construction of Miura transformations for discrete Painlevé equations. Our method is based on the bilinear formalism and we start with the expression of the nonlinear discrete equation in terms of τ -functions. Elimination of τ -functions from the resulting system leads to another nonlinear equation, which is a “modified” version of the original equation. The procedure therefore yields Miura transformations. In this letter, we illustrate this approach by reproducing previously known Miura transformations and constructing new ones.

The transformations named after Miura relate the solutions of a given nonlinear equation to those of some other, also nonlinear equation. The original, most famous, transformation of this type is the one relating the solutions of the KdV (Korteweg-deVries) and modified-KdV equations [1]. When the transformation relates the solutions of the *same* equation, corresponding to different values of the parameters of the equation, we call them auto-Bäcklund transformations. Schlesinger transformations are a particular kind of auto-Bäcklund. In the case of the Painlevé equations they correspond to special changes of parameters: integer or half-integer shifts of the monodromy exponents.

Miura transformations are of particular interest since they reveal the various interrelations between nonlinear systems. They are often the first step towards the construction of the auto-Bäcklund transformations and Lax pairs. In the domain of the (continuous) Painlevé equations, the systematic derivation and study of Miura, auto-Bäcklund and Schlesinger transformations has been a subject of intense activity [2]. It is worth pointing out that the Schlesinger transformations of the sixth Painlevé equation (P_{VI}) have only recently been established [3].

As far as the discrete Painlevé equations (d- \mathbb{P}) are concerned, the question of how to systematically derive interrelationships is far from settled. This is not only due to the fact that the d- \mathbb{P} 's have only recently made their appearance [4] but also because there are so many of them. In a series of works we have obtained the Miura transformations for several d- \mathbb{P} 's [5,6,7,8]. The approach we used was based on the observation that all the Miura transformations (in perfect analogy to the continuous case) are rational expressions involving the ratio of “Riccati forms”:

$$y = \frac{\alpha\bar{x}x + \beta\bar{x} + \gamma x + \delta}{\epsilon\bar{x}x + \zeta\bar{x} + \eta x + \theta} \quad (1)$$

where $x = x(n)$, $\bar{x} = x(n+1)$, $\underline{x} = x(n-1)$ and similarly for y . Postulating such a form (1), one starts with x satisfying a given d- \mathbb{P} and obtains another d- \mathbb{P} for y , by determining the coefficients $\alpha, \beta, \dots, \theta$. The procedure is quite cumbersome, in particular if one works with the full generality of (1). In practice, a good deal of intuition is needed in the derivation of the Miura transformations of various d- \mathbb{P} 's.

In this letter we shall present a new, systematic, approach to the construction of Miura transformations of discrete Painlevé equations. It is based on the bilinear formalism. As we have shown in our previous works, the τ -functions entering the bilinear transformations are defined by the singularity structure of the discrete equation. In particular, they are determined by the singularity pattern contained in an orbit that becomes nearly singular [9]. Moreover, the number of τ -functions corresponds to the number of types of singularity that can occur. The τ -functions are known to play a major role in the Miura, auto-Bäcklund and Schlesinger transformations of the d- \mathbb{P} 's and, in particular, in their self-dual description [10]. For the derivation of the Miura transformations, as we shall see below, we will need only their discrete logarithmic derivatives.

Let us start with an example based on a well-known (and well-studied) discrete Painlevé equation, the “standard” d-P_{II}:

$$\bar{x} + \underline{x} = \frac{zx + a}{1 - x^2}, \quad (2)$$

where $z = \alpha n + \beta$. The bilinearisation of (2) was given in [9,11]. We have argued there that, due to the singularity structure of (2) i.e. $\{\pm 1, \infty, \mp 1\}$, two τ -functions, F and G are needed and we proposed the transformation:

$$x = 1 - \frac{\bar{F}G}{FG} = -1 + \frac{F\bar{G}}{FG}. \quad (3)$$

As we hinted above and since our aim is not the bilinearisation of d-P_{II}, we can simplify our calculation by introducing the auxiliary variables $u = F/\bar{F}$, $v = \underline{G}/G$. Thus we have

$$x = 1 - \frac{v}{u} = -1 + \frac{\underline{u}}{\bar{v}}. \quad (4)$$

The important “trick” in the derivation of the “modified” equation is to eliminate one of the two τ -functions or, equivalently, one of the u, v appearing in the expression (4), for instance u . (Note that one has to up- and/or down-shift both (2) and (4) in order to have enough equations for this elimination.) The net result is a *homogeneous* 5-point equation for v . Given the homogeneity one can integrate it by putting $v/\bar{v} = w$. The resulting equation involves only products of $w\bar{w}$ (up- or down-shifted). We can integrate the equation once more by putting $w\bar{w} = \omega$. We then obtain for ω the three-point equation:

$$(\bar{\omega} + \omega - \bar{z})(\omega + \underline{\omega} - z) = \frac{(2\omega - a - z)(2\omega + a - \bar{z})}{\omega}. \quad (5)$$

This is a discretization [12] of equation XXXIV of the Painlevé-Gambier list [13] which is well known to be related to P_{II} by a Miura transformation. The standard form of d-P₃₄ can be recovered by putting $\omega = y + z/2 + \alpha/4$, where $\alpha = z - \underline{z}$. We then find:

$$(\bar{y} + y)(y + \underline{y}) = \frac{4y^2 - (a - \alpha/2)^2}{y + z/2 + \alpha/4}. \quad (6)$$

This equation can be regarded as a modified version of dP_{II} if we can explicitly construct the Miura transformation relating the two. We do this by retracing the above sequence of calculations. From the definitions of ω , w , and v we have:

$$y = \frac{v}{\bar{v}} - \frac{z}{2} - \frac{\alpha}{4} \quad (7)$$

Using (4) we find $v = (1 - x)u$ and $\bar{v} = \underline{u}/(1 + \bar{x})$. This leads to

$$y = (1 + \bar{x})(1 - x) - \frac{z}{2} - \frac{\alpha}{4}, \quad (8)$$

i.e. the well-known Miura transformation of d-P_{II} (see eq (9a) of [5]). Using (8) and the d-P_{II} equation (2) we can construct in a straightforward way the inverse of the Miura transformation.

Another most interesting equation is the alternate d-P_{II}, studied in detail in [14]:

$$\frac{z}{1+x\bar{x}} + \frac{\underline{z}}{1+x\underline{x}} = \frac{1}{x} - x + \underline{z} + \mu \quad (9)$$

This equation has two singularity patterns $\{0, \infty\}$ and $\{\infty, 0\}$ which suggest the following expression in terms of τ -functions:

$$x = \frac{F\underline{G}}{\underline{F}G}. \quad (10)$$

As in the previous case, we work with the discrete logarithmic derivatives of the τ -functions: $u = \underline{F}/F$, $v = \underline{G}/G$. In [14] we have shown that for the bilinearization of (9) we must introduce a separation through a first bilinear condition:

$$\overline{F}\underline{G} + \underline{F}\overline{G} = zFG. \quad (11)$$

This equation will also be our starting point here. Rewriting (11) in terms of u , v , we have:

$$\frac{v}{\underline{u}} + \frac{u}{\overline{v}} - z = 0 \quad (12)$$

Using (12) and (9) we can eliminate u . The result is a 4-point homogeneous equation in v which we can integrate once through $v/\overline{v} = y$. We obtain thus

$$(\overline{y} - \underline{y})^2 y^4 + 2z^2(\overline{y} + \underline{y})y^2(1 - y) + z^2\left(4y^3 + y^2(z^2 - \mu^2 - 4) - 2yz^2 + z^2\right) = 0 \quad (13)$$

which is precisely the equation obtained in [14]. We remark that (13) is a three-point correspondence and not a simple mapping. As we have remarked in [14], this indicates that the actual form of the modified equation is a 4-point one and equation (13) is its integral. Again, the Miura transformation

$$y = \frac{zx}{1+x\bar{x}}$$

follows from the above calculations.

Up to now we have used our method on equations for which the result was already known, having been obtained by a different approach. It is interesting to see whether we can use it to obtain new results corresponding to equations that have not yet been studied. Two examples will be presented in what follows.

We start with a q -P_{II} equation which was first identified in [15]:

$$(\overline{xx} - 1)(x\underline{x} - 1) = \frac{z\underline{z}}{a^2} \frac{x}{x - az} \quad (14)$$

where $z = z_0\lambda^n$ (and $\underline{z} = z/\lambda$, $\bar{z} = z\lambda$). The singularity structure of (14) is easily obtained. Two patterns exist, namely $\{az, \infty, 0\}$ and $\{0, \infty, az\}$, a fact that suggests the introduction of two τ -functions F and G . We are thus led to the ansatz:

$$x = \frac{\overline{FG}}{FG} = az + \frac{F\overline{G}}{FG} \quad (15)$$

Using the τ -functions we can rewrite (14) in the form:

$$(\overline{\overline{FG}} - F\overline{G})(\overline{FG} - \underline{FG}) = \frac{z\bar{z}}{a^2}\overline{FGFG} \quad (16)$$

which suggests immediately the separation:

$$\overline{\overline{FG}} - F\overline{G} = \frac{z}{a}\overline{FG}. \quad (17)$$

As in the previous case, we introduce $u = \overline{F}/F$ and $v = G/\underline{G}$ and rewrite equations (15) and (17) as:

$$\frac{u}{v} = az + \frac{\bar{v}}{\underline{u}} \quad (18a)$$

$$\frac{\bar{u}}{v} = \frac{z}{a} + \frac{\bar{v}}{u} \quad (18b)$$

Next, we eliminate v and introduce $y = \bar{u}/u$. We obtain for the latter the equation:

$$(y\bar{y} - 1)(y\underline{y} - 1) = \frac{z\bar{z}}{y}(y - a^2)(y - \frac{1}{a^2}) \quad (19)$$

Equation (19) is indeed a q -form of equation P₃₄ as was shown in [15].

Our final example will be again based on a q -P_{II} equation:

$$\overline{\underline{x}} = az \frac{x + z}{x(x - 1)} \quad (20)$$

which was identified in [15]. The singularity patterns of (20) are $\{1, \infty, 0\}$ and $\{0, \infty, 1\}$ suggesting the following ansatz in terms of two τ -functions F , G :

$$x = \frac{\overline{FG}}{FG} = 1 + \frac{F\overline{G}}{FG}. \quad (21)$$

As previously, we introduce $v = \underline{G}/G$ and $u = F/\overline{F}$. Expressing (20) and (21) in terms of u , v , we eliminate v . Next we introduce the nonlinear variable:

$$y = \frac{u}{\bar{u}} \quad (22)$$

and obtain the equation

$$(y\underline{y} - az^2)(y\bar{y} - a\lambda^2z^2) = -\lambda za(y - az)(y - \lambda z). \quad (23)$$

Equation (23) can be brought into canonical form if we put $y = Yz\sqrt{\lambda a}$. We thus obtain

$$(Y\underline{Y} - 1)(Y\overline{Y} - 1) = -\frac{1}{z}(Y - c)(Y - \frac{1}{c}). \quad (24)$$

where $c = \sqrt{a/\lambda}$. The explicit construction of the Miura transformation is straightforward. From (22), we find:

$$y = \overline{x}(x - 1) \quad (25a)$$

and using (20) or (24) we obtain the second half of the Miura:

$$x = \frac{y\underline{y} - az^2}{az - y}. \quad (25b)$$

It is interesting here to actually use these transformations in order to construct the Schlesinger transformation of the q - P_{II} (20). We start by introducing another Miura, through the elimination of u rather than v . We find:

$$\psi = \frac{v}{\overline{v}} = \underline{x}(x - 1) \quad (26)$$

and the canonical equation is obtained for Ψ where $\psi = \Psi z\sqrt{a/\lambda}$:

$$(\Psi\overline{\Psi} - 1)(\Psi\underline{\Psi} - 1) = -\frac{1}{z}(\Psi - \gamma)(\Psi - \frac{1}{\gamma}) \quad (27)$$

where $\gamma = \sqrt{\lambda a} = \lambda c$. Here is how one constructs the auto-Bäcklund: we start from x associated to a given a . We construct ψ (and Ψ) related to $\gamma = \sqrt{\lambda a}$. Then since Y and Ψ obey the same equation for different parameter we can consider that Ψ is indeed a Y provided we change the parameter of (24) to $\tilde{c} = \gamma = \sqrt{\tilde{a}/\lambda}$ which means that $\tilde{a} = \lambda^2 a$. From $Y = \Psi$ we can easily compute the relation between y and ψ : $y = \lambda^2 \psi$ (because Y is computed at $\tilde{a} = \lambda^2 a$). Combining now (26) and (25b) we find:

$$\tilde{x} = \frac{az(\lambda\underline{x}(x - 1) - z)}{x(az - \underline{x}(x - 1))} = \frac{z(\lambda a(x + z) - x\overline{x})}{x(x(\overline{x} - 1) - z)} \quad (28)$$

the two expressions in the r.h.s. of (28) being perfectly equivalent because x satisfies (20). In an analogous way we can obtain the Schlesinger:

$$\tilde{x} = \frac{az(\overline{x}(x - 1)/\lambda - z)}{x(az - \overline{x}(x - 1))} = \frac{z(a(x + z)/\lambda - x\underline{x})}{x(x(\underline{x} - 1) - z)} \quad (29)$$

where \underline{x} is a solution of (20) for $a = a/\lambda^2$.

Let us summarize our findings here. In this letter we have adressed the problem of the systematic construction of the Miura transformations for discrete Painlevé equations and obtained a new approach. Our method is based on the bilinear formalism and the construction of the modified equation proceeds through the elimination of one of the τ -functions. The method can be applied to both difference- and q - discrete equations.

Its usefulness lies in the fact that using the Miura transformations we can proceed to the construction of the auto-Bäcklund and Schlesingers in a straightforward way. In a future publication we expect to apply this approach to the long list of d - and q - \mathbb{P} 's that have not yet been examined.

ACKNOWLEDGEMENT

The research reported in this letter was partially supported by the Australian Research Council.

REFERENCES

- [1] R. M. Miura, J. Math. Phys. 9 (1968) 1202.
- [2] A.S. Fokas and M.J. Ablowitz, J. Math. Phys. 23 (1982) 2033,
A.S. Fokas, U. Mugan and M.J. Ablowitz, Physica D 30 (1988) 247,
A.S. Fokas, and X. Zhou, Comm. Math. Phys. 144 (1992) 601.
A.P. Bassom, P.A. Clarkson, and A.C. Hicks, Stud. Appl. Math. 95 (1995) 1.
A.E. Milne, P.A. Clarkson, and A.P. Bassom, Stud. Appl. Math. 98 (1997) 139.
- [3] U. Mugan and A. Sakka, J. Math. Phys. 36 (1995) 1284.
- [4] A. Ramani, B. Grammaticos, J. Hietarinta, Phys. Rev. Lett. 67 (1991) 1829.
- [5] A. Ramani, B. Grammaticos, Jour. Phys. A 25 (1992) L633
- [6] M. Jimbo, H. Sakai, A. Ramani, B. Grammaticos, Phys. Lett. A 217 (1996) 111
- [7] K. M. Tamizhmani, B. Grammaticos, A. Ramani, Lett. Math. Phys. 29 (1993) 49
- [8] K.M. Tamizhmani, A. Ramani, B. Grammaticos, Y. Ohta, Lett. Math. Phys. 38 (1996) 289.
- [9] A. Ramani, B. Grammaticos, J. Satsuma, Jour. Phys. A 28 (1995) 4655.
- [10] A. Ramani, Y. Ohta, J. Satsuma and B. Grammaticos, *Self-duality and Schlesinger chains for the asymmetric d - P_{II} and q - P_{III} equations*, to appear in Comm. Math. Phys.
- [11] J. Satsuma, K. Kajiwara, B. Grammaticos, J. Hietarinta, A. Ramani, Jour. Phys. A 28 (1995) 3541.
- [12] A. S. Fokas, B. Grammaticos, A. Ramani, J. of Math. Anal. and Appl. 180 (1993) 342.
- [13] E.L. Ince, *Ordinary differential equations*, Dover, New York, 1956.
- [14] F. Nijhoff, J. Satsuma, K. Kajiwara, B. Grammaticos, A. Ramani, Inv. Probl. 12 (1996) 697.
- [15] A. Ramani, B. Grammaticos, Physica A 228 (1996) 160.