

TREE CLASS E_7 : DIFFERENTIALS OF THE BIMODULE RESOLUTION

Since Q_i are direct sums, their elements can be concerned as column vectors, hence differentials can be described as matrixes (which are being multiplied by column vectors from the right). Now let us describe the matrixes of differentials componentwisely.

Remark 1. Numeration of lines and columns always starts with zero.

Notation.

- (1) Denote by $w_{i \rightarrow j}$ the way that starts in i th vertex and ends in j th.
- (2) Fot j th column of differential matrix let j_2 be the aliquot of division of j by s .

Description of the d_0
 $d_0 : Q_1 \rightarrow Q_0$ – is an $(7s \times 8s)$ matrix.

If $0 \leq j < 2s$, then

$$(d_0)_{ij} = \begin{cases} w_{7(j+m) \rightarrow 7(j+m)+1+3f(j_2,1)} \otimes e_{7j}, & i = j_s; \\ -e_{7(j+m)+1+3f(j_2,1)} \otimes w_{7j \rightarrow 7j+1+3f(j_2,1)}, & i = j + s + 2sf(j_2,1); \\ 0, & \text{otherwise.} \end{cases}$$

If $2s \leq j < 5s$, then

$$(d_0)_{ij} = \begin{cases} w_{7(j+m)+j_2-1 \rightarrow 7(j+m)+j_2+2f(j_2,4)} \otimes e_{7j+j_2-1}, & i = j - s; \\ -e_{7(j+m)+j_2+2f(j_2,4)} \otimes w_{7j+j_2-1 \rightarrow 7j+j_2+2f(j_2,4)}, & i = j + 2sf(j_2,4); \\ 0, & \text{otherwise.} \end{cases}$$

If $5s \leq j < 7s$, then

$$(d_0)_{ij} = \begin{cases} w_{7(j+m)+j_2-1 \rightarrow 7(j+m)+j_2} \otimes e_{7j+j_2-1}, & i = j - s; \\ -e_{7(j+m)+j_2} \otimes w_{7j+j_2-1 \rightarrow 7j+j_2}, & i = j; \\ 0, & \text{otherwise.} \end{cases}$$

If $7s \leq j < 8s$, then

$$(d_0)_{ij} = \begin{cases} -e_{7(j+m+1)} \otimes w_{7j+6 \rightarrow 7(j+1)}, & i = (j+1)s; \\ w_{7(j+m)+6 \rightarrow 7(j+m+1)} \otimes e_{7j+6}, & i = j - s; \\ 0, & \text{otherwise.} \end{cases}$$

Description of the d_1
 $d_1 : Q_2 \rightarrow Q_1$ – is an $(8s \times 7s)$ matrix.

If $0 \leq j < s$, then

$$(d_1)_{ij} = \begin{cases} w_{7(j+m)+1+j_1+2f(j_1,3) \rightarrow 7(j+m+1)-1} \otimes w_{7j \rightarrow 7j+j_1}, \\ \quad i = j + s(j_1 + 1 - f(j_1, 0)), \quad 0 \leq j_1 < 4; \\ -w_{7(j+m+1)-3+j_1 \rightarrow 7(j+m+1)-1} \otimes w_{7j \rightarrow 7j+j_1+3(1-f(j_1,0))}, \\ \quad i = j + s(j_1 + 4 - 3f(j_1, 0)), \quad 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_1)_{ij} = \begin{cases} w_{7(j+m+1)+1+j_1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+1 \rightarrow 7(j+1)+j_1}, \\ \quad i = (j+1)_s + 2sf(j_1, 1), \quad 0 \leq j_1 < 2; \\ w_{7(j+m)+2+j_1+2f(j_1, 2, 3) \rightarrow 7(j+m+1)+2} \otimes w_{7j+1 \rightarrow 7j+1+j_1+2f(j_1, 3)}, \\ \quad i = j + s(1 + j_1 + 2f(j_1, 3)), \quad 0 \leq j_1 < 4; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_1)_{ij} = \begin{cases} w_{7(j+m+1)+1+j_1 \rightarrow 7(j+m+1)+3} \otimes w_{7j+2 \rightarrow 7(j+1)+j_1}, \\ \quad i = (j+1)_s + s(j_1 + 1 - f(j_1, 0)), \quad 0 \leq j_1 < 3; \\ w_{7(j+m)+3+j_1+2f(j_1, 1, 2) \rightarrow 7(j+m+1)+3} \otimes w_{7j+2 \rightarrow 7j+2+j_1+2f(j_1, 2)}, \\ \quad i = j + s(1 + j_1 + 2f(j_1, 2)), \quad 0 \leq j_1 < 3; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_1)_{ij} = \begin{cases} e_{7(j+m+1)+4} \otimes w_{7j+3 \rightarrow 7(j+1)}, \quad i = s + (j+1)_s; \\ w_{7(j+m)+6 \rightarrow 7(j+m+1)+4} \otimes e_{7j+3}, \quad i = j + s; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+4} \otimes w_{7j+3 \rightarrow 7j+6}, \quad i = j + 4s; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_1)_{ij} = \begin{cases} w_{7(j+m+1)+4+j_1 \rightarrow 7(j+m+1)+5} \otimes w_{7j+4 \rightarrow 7(j+1)+4f(j_1, 1)}, \\ \quad i = (j+1)_s + s(1 + 4f(j_1, 1)), \quad 0 \leq j_1 < 2; \\ w_{7(j+m)+5+j_1 \rightarrow 7(j+m+1)+5} \otimes w_{7j+4 \rightarrow 7j+4+j_1}, \quad i = j + s(j_1 + 1), \quad 0 \leq j_1 < 3; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_1)_{ij} = \begin{cases} e_{7(j+m+1)+1} \otimes w_{7j+5 \rightarrow 7(j+1)}, \quad i = (j+1)_s; \\ w_{7(j+m)+6 \rightarrow 7(j+m+1)+1} \otimes e_{7j+5}, \quad i = j + s; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+1} \otimes w_{7j+5 \rightarrow 7j+6}, \quad i = j + 2s; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_1)_{ij} = \begin{cases} w_{7(j+m+1)+1+j_1+2f(j_1, 3, 4) \rightarrow 7(j+m+2)} \otimes w_{7j+6 \rightarrow 7(j+1)+j_1+2f(j_1, 4)}, \\ \quad i = (j+1)_s + s(j_1 + f(j_1, 1, 4) + 2f(j_1, 4)), \quad 0 \leq j_1 < 5; \\ w_{7(j+m+1) \rightarrow 7(j+m+2)} \otimes e_{7j+6}, \quad i = j + s; \\ 0, \quad \text{otherwise.} \end{cases}$$

Description of the d_2
 $d_2 : Q_3 \rightarrow Q_2$ – is an $(7s \times 9s)$ matrix.

If $0 \leq j < s$, then

$$(d_2)_{ij} = \begin{cases} w_{7(j+m-1)+6 \rightarrow 7(j+m)+2} \otimes e_{7j}, & i = j; \\ -e_{7(j+m)+2} \otimes w_{7j \rightarrow 7j+1}, & i = j + s; \\ w_{7(j+m)+1 \rightarrow 7(j+m)+2} \otimes w_{7j \rightarrow 7j+5}, & i = j + 5s; \\ 0, & \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_2)_{ij} = \begin{cases} w_{7(j+m-1)+6 \rightarrow 7(j+m)+5} \otimes e_{7j}, & i = j_s; \\ -w_{7(j+m)+4 \rightarrow 7(j+m)+5} \otimes w_{7j \rightarrow 7j+3}, & i = j + 2s; \\ e_{7(j+m)+5} \otimes w_{7j \rightarrow 7j+4}, & i = j + 3s; \\ 0, & \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_2)_{ij} = \begin{cases} w_{7(j+m)+2 \rightarrow 7(j+m)+3} \otimes e_{7j+1}, & i = j - s; \\ -e_{7(j+m)+3} \otimes w_{7j+1 \rightarrow 7j+2}, & i = j; \\ 0, & \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_2)_{ij} = \begin{cases} -e_{7(j+m)+6} \otimes w_{7j+2 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ w_{7(j+m)+3 \rightarrow 7(j+m)+6} \otimes e_{7j+2}, & i = j - s; \\ -w_{7(j+m)+4 \rightarrow 7(j+m)+6} \otimes w_{7j+2 \rightarrow 7j+3}, & i = j; \\ 0, & \text{otherwise.} \end{cases}$$

If $4s \leq j < 6s$, then

$$(d_2)_{ij} = \begin{cases} w_{7(j+m)+6 \rightarrow 7(j+m+1)-f(j_2,5)} \otimes w_{7j+j_2-1 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ w_{7(j+m)+j_2 \rightarrow 7(j+m+1)-f(j_2,5)} \otimes e_{7j+j_2-1}, & i = j - s; \\ -w_{7(j+m+1)-6f(j_2,5) \rightarrow 7(j+m+1)-f(j_2,5)} \otimes w_{7j+j_2-1 \rightarrow 7j+10-j_2}, & i = j + 2sf(j_2, 4); \\ 0, & \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_2)_{ij} = \begin{cases} w_{7(j+m)+1 \rightarrow 7(j+m+1)} \otimes e_{7j+5}, & i = j - s; \\ -e_{7(j+m+1)} \otimes w_{7j+5 \rightarrow 7j+6}, & i = j; \\ 0, & \text{otherwise.} \end{cases}$$

If $7s \leq j < 8s$, then

$$(d_2)_{ij} = \begin{cases} -w_{7(j+m)+6 \rightarrow 7(j+m+1)+1} \otimes w_{7j+6 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ -e_{7(j+m+1)+1} \otimes w_{7j+6 \rightarrow 7(j+1)+5}, & i = 5s + (j+1)_s; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+1} \otimes e_{7j+6}, & i = j - s; \\ 0, & \text{otherwise.} \end{cases}$$

If $8s \leq j < 9s$, then

$$(d_2)_{ij} = \begin{cases} -e_{7(j+m+1)+4} \otimes w_{7j+6 \rightarrow 7(j+1)+3}, & i = 3s + (j+1)_s; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+4} \otimes e_{7j+6}, & i = j - 2s; \\ 0, & \text{otherwise.} \end{cases}$$

Description of the d_3
 $d_3 : Q_4 \rightarrow Q_3$ – is an $(9s \times 10s)$ matrix.

If $0 \leq j < s$, then

$$(d_3)_{ij} = \begin{cases} (1 - 2f(j_1, 1))w_{7(j+m)+2+3f(j_1, 1)+f(j_1, 2)+4f(j_1, 3, 4) \rightarrow 7(j+m)+6} \otimes w_{7j \rightarrow 7j+j_1-1+f(j_1, 0)+f(j_1, 4)}, & i = j + s(j_1 + f(j_1, 4)), 0 \leq j_1 < 5; \\ 0, & \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_3)_{ij} = \begin{cases} (1 - 2f(j_1, 2, 3))w_{7(j+m+1)-2f(j_1, 0)-f(j_1, 2) \rightarrow 7(j+m+1)} \otimes w_{7j \rightarrow 7j+j_1+2-2f(j_1, 0)}, & i = j + s(j_1 + 2 - 2f(j_1, 0)), 0 \leq j_1 < 4; \\ 0, & \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_3)_{ij} = \begin{cases} w_{7(j+m+1)+2+j_1 \rightarrow 7(j+m+1)+3} \otimes w_{7j+1 \rightarrow 7(j+1)+j_1}, & i = (j+1)_s + 2sj_1, 0 \leq j_1 < 2; \\ w_{7(j+m)+3+3j_1 \rightarrow 7(j+m+1)+3} \otimes w_{7j+1 \rightarrow 7j+1+j_1}, & i = j + sj_1, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_3)_{ij} = \begin{cases} w_{7(j+m)+6+j_1 \rightarrow 7(j+m+1)+4} \otimes w_{7j+2 \rightarrow 7j+2+j_1}, & i = j + sj_1, 0 \leq j_1 < 2; \\ e_{7(j+m+1)+4} \otimes w_{7j+2 \rightarrow 7j+6}, & i = j + 5s; \\ 0, & \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_3)_{ij} = \begin{cases} w_{7(j+m+1)+j_1 \rightarrow 7(j+m+1)+1} \otimes w_{7j+3 \rightarrow 7j+3+3j_1}, & i = j + 3j_1s, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_3)_{ij} = \begin{cases} -e_{7(j+m+1)+5} \otimes w_{7j+3 \rightarrow 7(j+1)}, & i = s + (j+1)_s; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+5} \otimes e_{7j+3}, & i = j - s; \\ w_{7(j+m+1)+4 \rightarrow 7(j+m+1)+5} \otimes w_{7j+3 \rightarrow 7j+6}, & i = j + 3s; \\ 0, & \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_3)_{ij} = \begin{cases} w_{7(j+m)+6+j_1 \rightarrow 7(j+m+1)+1} \otimes w_{7j+4 \rightarrow 7j+4+j_1}, & i = j + s(j_1 - 1), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $7s \leq j < 8s$, then

$$(d_3)_{ij} = \begin{cases} e_{7(j+m+1)+2} \otimes w_{7j+5 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+2} \otimes e_{7j+5}, & i = j-s; \\ w_{7(j+m+1)+1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+5 \rightarrow 7j+6}, & i = j; \\ 0, & \text{otherwise.} \end{cases}$$

If $8s \leq j < 9s$, then

$$(d_3)_{ij} = \begin{cases} w_{7(j+m+1) \rightarrow 7(j+m+1)+4} \otimes e_{7j+5}, & i = j-2s; \\ e_{7(j+m+1)+4} \otimes w_{7j+5 \rightarrow 7j+6}, & i = j; \\ 0, & \text{otherwise.} \end{cases}$$

If $9s \leq j < 10s$, then

$$(d_3)_{ij} = \begin{cases} w_{7(j+m+1)+2+j_1+2f(j_1,2) \rightarrow 7(j+m+1)+6} \otimes w_{7j+6 \rightarrow 7(j+1)+j_1}, & i = (j+1)_s + s(j_1 + f(j_1, 1, 2)), 0 \leq j_1 < 3; \\ w_{7(j+m+1)+1 \rightarrow 7(j+m+1)+6} \otimes e_{7j+6}, & i = j-2s; \\ -w_{7(j+m+1)+4 \rightarrow 7(j+m+1)+6} \otimes e_{7j+6}, & i = j-s; \\ 0, & \text{otherwise.} \end{cases}$$

Description of the d_4
 $d_4 : Q_5 \rightarrow Q_4$ – is an $(10s \times 10s)$ matrix.

If $0 \leq j < s$, then

$$(d_4)_{ij} = \begin{cases} w_{7(j+m-1)+6+j_1 \rightarrow 7(j+m)+4} \otimes e_{7j}, & i = j + sj_1, 0 \leq j_1 < 2; \\ -e_{7(j+m)+4} \otimes w_{7j \rightarrow 7j+2}, & i = j + 3s; \\ e_{7(j+m)+4} \otimes w_{7j \rightarrow 7j+5}, & i = j + 8s; \\ 0, & \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_4)_{ij} = \begin{cases} w_{7(j+m-1)+6 \rightarrow 7(j+m)+3} \otimes e_{7j}, & i = js; \\ -e_{7(j+m)+3} \otimes w_{7j \rightarrow 7j+1}, & i = j+s; \\ (-1)^{j_1+1} w_{7(j+m)+1+j_1 \rightarrow 7(j+m)+3} \otimes w_{7j \rightarrow 7j+4+j_1}, & i = j + (5 + j_1)s, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_4)_{ij} = \begin{cases} (-1)^{j_1} w_{7(j+m)+f(j_1,1,2) \rightarrow 7(j+m)+1} \otimes w_{7j \rightarrow 7j+2+j_1-2f(j_1,0)}, & i = j + s(2j_1 - f(j_1, 0)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_4)_{ij} = \begin{cases} -e_{7(j+m)+6} \otimes w_{7j+1 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ (-1)^{j_1} w_{7(j+m)+3+j_1 \rightarrow 7(j+m)+6} \otimes w_{7j+1 \rightarrow 7j+1+j_1}, & i = j + s(j_1 - 1 + f(j_1, 2)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_4)_{ij} = \begin{cases} -e_{7(j+m+1)} \otimes w_{7j+2 \rightarrow 7(j+1)}, & i = s + (j+1)_s; \\ (-1)^{j_1} w_{7(j+m)+4+j_1 \rightarrow 7(j+m+1)} \otimes w_{7j+2 \rightarrow 7j+2+j_1}, & i = j + s(2j_1 - 1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_4)_{ij} = \begin{cases} e_{7(j+m)+6} \otimes w_{7j+3 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ (2f(j_1, 0) - 1) w_{7(j+m)+4+j_1-3f(j_1, 0) \rightarrow 7(j+m)+6} \otimes w_{7j+3 \rightarrow 7j+3+3f(j_1, 2)}, & i = j + s(j_1 - 1 + 3f(j_1, 2)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_4)_{ij} = \begin{cases} w_{7(j+m)+6+j_1 \rightarrow 7(j+m+1)} \otimes w_{7j+4 \rightarrow 7(j+1)}, & i = (j+1)_s + sj_1, 0 \leq j_1 < 2; \\ (1 - 2f(j_1, 1)) w_{7(j+m)+1+j_1 \rightarrow 7(j+m+1)} \otimes w_{7j+4 \rightarrow 7j+4+j_1}, & i = j + sj_1, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $7s \leq j < 8s$, then

$$(d_4)_{ij} = \begin{cases} w_{7(j+m)+2 \rightarrow 7(j+m)+6} \otimes e_{7j+5}, & i = j; \\ -w_{7(j+m)+4+2j_1 \rightarrow 7(j+m)+6} \otimes w_{7j+5 \rightarrow 7j+5+j_1}, & i = j + s(j_1 + 1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $8s \leq j < 9s$, then

$$(d_4)_{ij} = \begin{cases} (2f(j_1, 2) - 1) w_{7(j+m)+6+j_1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+6 \rightarrow 7(j+1)+3f(j_1, 2)+5f(j_1, 3)}, & i = (j+1)_s + s(j_1 + 2f(j_1, 2) + 4f(j_1, 3)), 0 \leq j_1 < 4; \\ w_{7(j+m)+6 \rightarrow 7(j+m+1)+2} \otimes e_{7j+6}, & i = j + s; \\ 0, & \text{otherwise.} \end{cases}$$

If $9s \leq j < 10s$, then

$$(d_4)_{ij} = \begin{cases} (-1)^{j_1+1} w_{7(j+m+1)+4+j_1 \rightarrow 7(j+m+1)+5} \otimes w_{7j+6 \rightarrow 7(j+1)+2+j_1}, & i = (3 + 2j_1)s + (j+1)_s, 0 \leq j_1 < 2; \\ w_{7(j+m)+6 \rightarrow 7(j+m+1)+5} \otimes e_{7j+6}, & i = j; \\ 0, & \text{otherwise.} \end{cases}$$

Description of the d_5
 $d_5 : Q_6 \rightarrow Q_5$ – is an $(10s \times 12s)$ matrix.

If $0 \leq j < s$, then

$$(d_5)_{ij} = \begin{cases} (2f(j_1, 0) - 1) w_{7(j+m)+4-j_1-f(j_1, 2) \rightarrow 7(j+m)+6} \otimes e_{7j}, & i = j + j_1 s, 0 \leq j_1 < 3; \\ (2f(j_1, 2) - 1) e_{7(j+m)+6} \otimes w_{7j \rightarrow 7j+1+2j_1}, & i = j + (3 + 2j_1)s, 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_5)_{ij} = \begin{cases} w_{7(j+m)+3 \rightarrow 7(j+m+1)} \otimes e_{7j}, & i = j; \\ w_{7(j+m)+6+f(j_1,1,2) \rightarrow 7(j+m+1)} \otimes w_{7j \rightarrow 7j+1+j_1+f(j_1,2)}, & i = j + (2 + j_1 + f(j_1, 2))s, 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_5)_{ij} = \begin{cases} e_{7(j+m+1)+4} \otimes w_{7j+1 \rightarrow 7(j+1)}, & i = (j+1)s; \\ w_{7(j+m)+6+j_1 \rightarrow 7(j+m+1)+4} \otimes w_{7j+1 \rightarrow 7j+1+j_1}, & i = j + s(j_1 + 1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_5)_{ij} = \begin{cases} e_{7(j+m+1)+1} \otimes w_{7j+2 \rightarrow 7(j+1)}, & i = 2s + (j+1)s; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+1} \otimes e_{7j+2}, & i = j + s; \\ 0, & \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_5)_{ij} = \begin{cases} w_{7(j+m+1)+4 \rightarrow 7(j+m+1)+5} \otimes w_{7j+2 \rightarrow 7(j+1)}, & i = (j+1)s; \\ (2f(j_1, 0) - 1)w_{7(j+m+1+f(j_1,2))-j_1 \rightarrow 7(j+m+1)+5} \otimes w_{7j+2 \rightarrow 7j+2+j_1+2f(j_1,2)}, & i = j + s(j_1 + 3f(j_1, 2)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_5)_{ij} = \begin{cases} w_{7(j+m+1)+1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+3 \rightarrow 7(j+1)}, & i = 2s + (j+1)s; \\ w_{7(j+m)+6+3j_1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+3 \rightarrow 7j+3+3j_1}, & i = j + 3sj_1, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_5)_{ij} = \begin{cases} w_{7(j+m+1)-j_1+4f(j_1,2) \rightarrow 7(j+m+1)+2} \otimes w_{7j+4 \rightarrow 7j+4+j_1}, & i = j + sj_1, 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $7s \leq j < 8s$, then

$$(d_5)_{ij} = \begin{cases} -e_{7(j+m+1)+4} \otimes w_{7j+4 \rightarrow 7(j+1)}, & i = (j+1)s; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+4} \otimes e_{7j+4}, & i = j - s; \\ 0, & \text{otherwise.} \end{cases}$$

If $8s \leq j < 9s$, then

$$(d_5)_{ij} = \begin{cases} w_{7(j+m+1)+3-2j_1 \rightarrow 7(j+m+1)+3} \otimes w_{7j+5 \rightarrow 7(j+1)}, & i = s(j_1 + 1) + (j+1)s, 0 \leq j_1 < 2; \\ w_{7(j+m)+6+3j_1 \rightarrow 7(j+m+1)+3} \otimes w_{7j+5 \rightarrow 7j+5+j_1}, & i = j + (j_1 - 1)s, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $9s \leq j < 10s$, then

$$(d_5)_{ij} = \begin{cases} w_{7(j+m)+6 \rightarrow 7(j+m+1)+5} \otimes e_{7j+5}, & i = j - 2s; \\ e_{7(j+m+1)+5} \otimes w_{7j+5 \rightarrow 7j+6}, & i = j; \\ 0, & \text{otherwise.} \end{cases}$$

If $10s \leq j < 11s$, then

$$(d_5)_{ij} = \begin{cases} w_{7(j+m+1+f(j_1,2))+3-2j_1 \rightarrow 7(j+m+1)+6} \otimes w_{7j+6 \rightarrow 7(j+1)+f(j_1,2)}, & i = s(j_1 + 1) + (j + 1)_s, 0 \leq j_1 < 3; \\ (-1)^{j_1} w_{7(j+m+1)+2+3j_1 \rightarrow 7(j+m+1)+6} \otimes e_{7j+6}, & i = j + s(j_1 - 2), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $11s \leq j < 12s$, then

$$(d_5)_{ij} = \begin{cases} e_{7(j+m+2)} \otimes w_{7j+6 \rightarrow 7(j+1)+2}, & i = 4s + (j + 1)_s; \\ w_{7(j+m+1)+5 \rightarrow 7(j+m+2)} \otimes e_{7j+6}, & i = j - 2s; \\ 0, & \text{otherwise.} \end{cases}$$

Description of the d_6
 $d_6 : Q_7 \rightarrow Q_6$ – is an $(12s \times 12s)$ matrix.

If $0 \leq j < s$, then

$$(d_6)_{ij} = \begin{cases} w_{7(j+m-1)+6 \rightarrow 7(j+m)+2} \otimes e_{7j}, & i = j; \\ w_{7(j+m) \rightarrow 7(j+m)+2} \otimes e_{7j}, & i = j + s; \\ (-1)^{j_1+1} w_{7(j+m)+2-f(j_1,0) \rightarrow 7(j+m)+2} \otimes w_{7j \rightarrow 7j+2+j_1}, & i = j + s(j_1 + 4 - f(j_1, 0)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_6)_{ij} = \begin{cases} w_{7(j+m-1)+6 \rightarrow 7(j+m)+5} \otimes e_{7j}, & i = j_s; \\ (2f(j_1, 0) - 1) w_{7(j+m)+5-f(j_1,0) \rightarrow 7(j+m)+5} \otimes w_{7j \rightarrow 7j+1+j_1+2f(j_1,2)}, & i = j + s(1 + 2j_1 + 3f(j_1, 2)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_6)_{ij} = \begin{cases} (2f(j_1, 0) - 1) w_{7(j+m)+4(1-f(j_1,0)) \rightarrow 7(j+m)+4} \otimes w_{7j \rightarrow 7j+j_1+2f(j_1,2)}, & i = j + s(j_1 - 1 + 4f(j_1, 2)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_6)_{ij} = \begin{cases} -w_{7(j+m)+6+j_1 \rightarrow 7(j+m+1)} \otimes w_{7j+1 \rightarrow 7(j+1)}, & i = (j + 1)_s + sj_1, 0 \leq j_1 < 2; \\ (-1)^{j_1} w_{7(j+m)+4-3j_1 \rightarrow 7(j+m+1)} \otimes w_{7j+1 \rightarrow 7j+1+j_1}, & i = j + s(j_1 - 1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_6)_{ij} = \begin{cases} e_{7(j+m)+6} \otimes w_{7j+2 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ w_{7(j+m)+1 \rightarrow 7(j+m)+6} \otimes e_{7j+2}, & i = j - s; \\ (2f(j_1, 2) - 1)w_{7(j+m)+5-3f(j_1, 1)+f(j_1, 2) \rightarrow 7(j+m)+6} \otimes w_{7j+2 \rightarrow 7j+2+j_1+2f(j_1, 2)}, \\ & i = j + s(j_1 + 4f(j_1, 2)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_6)_{ij} = \begin{cases} e_{7(j+m+1)} \otimes w_{7j+3 \rightarrow 7(j+1)}, & i = s + (j+1)_s; \\ (2f(j_1, 0) - 1)w_{7(j+m)+5+j_1-3f(j_1, 0) \rightarrow 7(j+m+1)} \otimes w_{7j+3 \rightarrow 7j+6-3f(j_1, 0)}, \\ & i = j + s(j_1 + 4 - 4f(j_1, 0)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_6)_{ij} = \begin{cases} -e_{7(j+m)+6} \otimes w_{7j+4 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ (2f(j_1, 0) - 1)w_{7(j+m)+2+2j_1-3f(j_1, 2) \rightarrow 7(j+m)+6} \otimes w_{7j+4 \rightarrow 7j+4+f(j_1, 2)}, \\ & i = j + sj_1, 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $7s \leq j < 8s$, then

$$(d_6)_{ij} = \begin{cases} (2f(j_1, 0) - 1)w_{7(j+m)+4+j_1-f(j_1, 0) \rightarrow 7(j+m)+6} \otimes w_{7j+5 \rightarrow 7j+5+f(j_1, 2)}, \\ & i = j + s(j_1 + 1), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $8s \leq j < 9s$, then

$$(d_6)_{ij} = \begin{cases} (-1)^{j_1} w_{7(j+m)+5+2j_1 \rightarrow 7(j+m+1)} \otimes w_{7j+5 \rightarrow 7j+5+j_1}, & i = j + s(2j_1 + 1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $9s \leq j < 10s$, then

$$(d_6)_{ij} = \begin{cases} -e_{7(j+m+1)+4} \otimes w_{7j+6 \rightarrow 7(j+1)+1}, & i = 2s + (j+1)_s; \\ w_{7(j+m)+6+j_1 \rightarrow 7(j+m+1)+4} \otimes e_{7j+6}, & i = j + s(j_1 + 1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $10s \leq j < 11s$, then

$$(d_6)_{ij} = \begin{cases} (1 - 2f(j_1, 2))w_{7(j+m)+6+3j_1-2f(j_1, 2) \rightarrow 7(j+m+1)+3} \otimes w_{7j+6 \rightarrow 7(j+1)+3j_1-f(j_1, 2)}, \\ & i = (j+1)_s + s(5j_1 - 2f(j_1, 2)), 0 \leq j_1 < 3; \\ w_{7(j+m)+6 \rightarrow 7(j+m+1)+3} \otimes e_{7j+6}, & i = j; \\ 0, & \text{otherwise.} \end{cases}$$

If $11s \leq j < 12s$, then

$$(d_6)_{ij} = \begin{cases} -e_{7(j+m+1)+1} \otimes w_{7j+6 \rightarrow 7(j+1)+2}, & i = 3s + (j+1)_s; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+1} \otimes e_{7j+6}, & i = j; \\ 0, & \text{otherwise.} \end{cases}$$

Description of the d_7
 $d_7 : Q_8 \rightarrow Q_7$ – is an $(12s \times 13s)$ matrix.

If $0 \leq j < s$, then

$$(d_7)_{ij} = \begin{cases} (1 - 2f(j_1, 1, 2))w_{7(j+m)+2+j_1+2f(j_1,1)+f(j_1,3)-f(j_1,5)\rightarrow 7(j+m)+6} \otimes \\ \quad w_{7j\rightarrow 7j+2f(j_1,3)+4f(j_1,4)+5f(j_1,5)}, \\ \quad i = j + s(j_1 + f(j_1, 3) + 2f(j_1, 4, 5)), \quad 0 \leq j_1 < 6; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_7)_{ij} = \begin{cases} (1 - 2f(j_1, 1, 3))w_{7(j+m+1)-2f(j_1,0)-f(j_1,2)\rightarrow 7(j+m+1)} \otimes w_{7j\rightarrow 7j+j_1+f(j_1,4)}, \\ \quad i = j + s(1 + j_1 - f(j_1, 0) + 2f(j_1, 4)), \quad 0 \leq j_1 < 5; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_7)_{ij} = \begin{cases} w_{7(j+m+1)-f(j_1,1)+f(j_1,3)\rightarrow 7(j+m+1)+1} \otimes w_{7j+1\rightarrow 7j+1+j_1+2f(j_1,3)}, \\ \quad i = j + s(1 + j_1 + 5f(j_1, 3)), \quad 0 \leq j_1 < 4; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_7)_{ij} = \begin{cases} w_{7(j+m+1)+5-j_1\rightarrow 7(j+m+1)+5} \otimes w_{7j+1\rightarrow 7(j+1)}, \quad i = s(j_1 + 1) + (j + 1)_s, \quad 0 \leq j_1 < 2; \\ w_{7(j+m+1)\rightarrow 7(j+m+1)+5} \otimes e_{7j+1}, \quad i = j; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_7)_{ij} = \begin{cases} -e_{7(j+m+1)+2} \otimes w_{7j+2\rightarrow 7(j+1)}, \quad i = (j + 1)_s; \\ w_{7(j+m)+6+j_1\rightarrow 7(j+m+1)+2} \otimes w_{7j+2\rightarrow 7j+2+j_1+2f(j_1,2)}, \\ \quad i = j + s(j_1 + 5f(j_1, 2)), \quad 0 \leq j_1 < 3; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_7)_{ij} = \begin{cases} -w_{7(j+m+1)+2\rightarrow 7(j+m+1)+3} \otimes w_{7j+3\rightarrow 7(j+1)}, \quad i = (j + 1)_s; \\ w_{7(j+m+1)+3j_1-5f(j_1,2)\rightarrow 7(j+m+1)+3} \otimes w_{7j+3\rightarrow 7j+6-3f(j_1,0)}, \\ \quad i = j + s(4 + j_1 - 4f(j_1, 0)), \quad 0 \leq j_1 < 3; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_7)_{ij} = \begin{cases} -e_{7(j+m+1)+4} \otimes w_{7j+3\rightarrow 7(j+1)}, \quad i = 2s + (j + 1)_s; \\ w_{7(j+m+1)+4j_1\rightarrow 7(j+m+1)+4} \otimes w_{7j+3\rightarrow 7j+3+3j_1}, \quad i = j + s(4j_1 - 1), \quad 0 \leq j_1 < 2; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $7s \leq j < 8s$, then

$$(d_7)_{ij} = \begin{cases} w_{7(j+m)+6+4f(j_1,2)\rightarrow 7(j+m+1)+3} \otimes w_{7j+4\rightarrow 7j+4+j_1}, \\ \quad i = j + s(j_1 - 1 + 2f(j_1, 2)), \quad 0 \leq j_1 < 3; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $8s \leq j < 9s$, then

$$(d_7)_{ij} = \begin{cases} e_{7(j+m+1)+5} \otimes w_{7j+4\rightarrow 7(j+1)}, \quad i = s + (j+1)_s; \\ w_{7(j+m)+6\rightarrow 7(j+m+1)+5} \otimes e_{7j+4}, \quad i = j - 2s; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $9s \leq j < 10s$, then

$$(d_7)_{ij} = \begin{cases} w_{7(j+m)+6+j_1+3f(j_1,2)\rightarrow 7(j+m+1)+4} \otimes w_{7j+5\rightarrow 7j+5+f(j_1,2)}, \\ \quad i = j + s(j_1 - 2), \quad 0 \leq j_1 < 3; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $10s \leq j < 11s$, then

$$(d_7)_{ij} = \begin{cases} w_{7(j+m+1)+j_1\rightarrow 7(j+m+1)+1} \otimes w_{7j+5\rightarrow 7j+5+j_1}, \quad i = j + s(3j_1 - 2), \quad 0 \leq j_1 < 2; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $11s \leq j < 12s$, then

$$(d_7)_{ij} = \begin{cases} (-1)^{j_1} w_{7(j+m+1)+2+2j_1\rightarrow 7(j+m+1)+6} \otimes w_{7j+6\rightarrow 7(j+1)+4f(j_1,2)}, \\ \quad i = (j+1)_s + 2s(j_1 + f(j_1, 2)), \quad 0 \leq j_1 < 3; \\ (2f(j_1, 0) - 1) w_{7(j+m+1)+4-j_1-f(j_1,2)\rightarrow 7(j+m+1)+6} \otimes e_{7j+6}, \\ \quad i = j + s(j_1 - 2), \quad 0 \leq j_1 < 3; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $12s \leq j < 13s$, then

$$(d_7)_{ij} = \begin{cases} (2f(j_1, 1, 2) - 1) w_{7(j+m+1)+2+2j_1+f(j_1,2)-2f(j_1,3)\rightarrow 7(j+m+2)} \otimes w_{7j+6\rightarrow 7(j+1)+f(j_1,2)+4f(j_1,3)}, \\ \quad i = (j+1)_s + s(1 + j_1 - f(j_1, 0) + 2f(j_1, 3)), \quad 0 \leq j_1 < 4; \\ w_{7(j+m+1)+3\rightarrow 7(j+m+2)} \otimes e_{7j+6}, \quad i = j - 2s; \\ 0, \quad \text{otherwise.} \end{cases}$$

Description of the d_8
 $d_8 : Q_9 \rightarrow Q_8$ – is an $(13s \times 12s)$ matrix.

If $0 \leq j < s$, then

$$(d_8)_{ij} = \begin{cases} w_{7(j+m-1)+6+j_1\rightarrow 7(j+m)+5} \otimes e_{7j}, \quad i = j + sj_1, \quad 0 \leq j_1 < 2; \\ (1 - 2f(j_1, 2, 3)) w_{7(j+m)+4+f(j_1,0)+f(j_1,2)\rightarrow 7(j+m)+5} \otimes w_{7j\rightarrow 7j+2+j_1-f(j_1,0)}, \\ \quad i = j + s(3 + 3j_1 - f(j_1, 2) - 3f(j_1, 3)), \quad 0 \leq j_1 < 4; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_8)_{ij} = \begin{cases} w_{7(j+m-1)+6 \rightarrow 7(j+m)+3} \otimes e_{7j}, & i = j_s; \\ (-1)^{j_1+1} w_{7(j+m)+3-f(j_1,0) \rightarrow 7(j+m)+3} \otimes w_{7j \rightarrow 7j+2+j_1}, & i = j + s(3 + j_1 + f(j_1, 2)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_8)_{ij} = \begin{cases} (1 - 2f(j_1, 2)) w_{7(j+m)+1-f(j_1,0) \rightarrow 7(j+m)+1} \otimes w_{7j \rightarrow 7j+j_1+3f(j_1,2)}, & i = j + s(j_1 - 1 + 7f(j_1, 2)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_8)_{ij} = \begin{cases} -e_{7(j+m)+6} \otimes w_{7j+1 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ (2f(j_1, 0) - 1) w_{7(j+m)+1+4f(j_1,1)+f(j_1,2) \rightarrow 7(j+m)+6} \otimes w_{7j+1 \rightarrow 7j+1+f(j_1,2)}, & i = j + s(j_1 - 1), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_8)_{ij} = \begin{cases} w_{7(j+m)+6+j_1 \rightarrow 7(j+m+1)} \otimes w_{7j+2 \rightarrow 7(j+1)}, & i = (j+1)_s + sj_1, 0 \leq j_1 < 2; \\ (-1)^{j_1} w_{7(j+m)+2+j_1+3f(j_1,2) \rightarrow 7(j+m+1)} \otimes w_{7j+2 \rightarrow 7j+2+j_1+2f(j_1,2)}, & i = j + s(j_1 + 6f(j_1, 2)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_8)_{ij} = \begin{cases} (-1)^{j_1} w_{7(j+m)+3+j_1+f(j_1,2) \rightarrow 7(j+m)+6} \otimes w_{7j+3 \rightarrow 7j+3+3f(j_1,2)}, & i = j + s(j_1 + 4f(j_1, 2)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_8)_{ij} = \begin{cases} -e_{7(j+m)+6} \otimes w_{7j+4 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ (1 - 2f(j_1, 1, 2)) w_{7(j+m)+2+j_1+f(j_1,0)+2f(j_1,1)-4f(j_1,3) \rightarrow 7(j+m)+6} \otimes w_{7j+4 \rightarrow 7j+4+f(j_1,2,3)+2f(j_1,4)}, & i = j + s(j_1 + 1), 0 \leq j_1 < 5; \\ 0, & \text{otherwise.} \end{cases}$$

If $7s \leq j < 8s$, then

$$(d_8)_{ij} = \begin{cases} -e_{7(j+m+1)} \otimes w_{7j+4 \rightarrow 7(j+1)}, & i = s + (j+1)_s; \\ w_{7(j+m)+5 \rightarrow 7(j+m+1)} \otimes e_{7j+4}, & i = j + s; \\ 0, & \text{otherwise.} \end{cases}$$

If $8s \leq j < 9s$, then

$$(d_8)_{ij} = \begin{cases} (2f(j_1, 0) - 1) w_{7(j+m)+4+j_1-4f(j_1,1) \rightarrow 7(j+m+1)} \otimes w_{7j+5 \rightarrow 7j+5+f(j_1,2,3)}, & i = j + s(j_1 + 1), 0 \leq j_1 < 4; \\ 0, & \text{otherwise.} \end{cases}$$

If $9s \leq j < 10s$, then

$$(d_8)_{ij} = \begin{cases} (-1)^{j_1+1} w_{7(j+m+1)+1+j_1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+6 \rightarrow 7(j+1)+1+j_1}, \\ \quad i = (j+1)_s + 2s(j_1+1), \quad 0 \leq j_1 < 2; \\ w_{7(j+m)+6+j_1 \rightarrow 7(j+m+1)+2} \otimes e_{7j+6}, \quad i = j + s(2+j_1), \quad 0 \leq j_1 < 2; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $10s \leq j < 11s$, then

$$(d_8)_{ij} = \begin{cases} -e_{7(j+m+1)+5} \otimes w_{7j+6 \rightarrow 7(j+1)+4}, \quad i = 8s + (j+1)_s; \\ w_{7(j+m)+6 \rightarrow 7(j+m+1)+5} \otimes e_{7j+6}, \quad i = j + s; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $11s \leq j < 12s$, then

$$(d_8)_{ij} = \begin{cases} (1 - 2f(j_1, 3))w_{7(j+m)+6+j_1+3f(j_1, 2)+2f(j_1, 3) \rightarrow 7(j+m+1)+4} \otimes w_{7j+6 \rightarrow 7(j+1)+3f(j_1, 2)+5f(j_1, 3)}, \\ \quad i = (j+1)_s + s(j_1 + 4f(j_1, 2) + 6f(j_1, 3)), \quad 0 \leq j_1 < 4; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+4} \otimes e_{7j+6}, \quad i = j + s; \\ 0, \quad \text{otherwise.} \end{cases}$$

Description of the d_9
 $d_9 : Q_{10} \rightarrow Q_9$ – is an $(12s \times 12s)$ matrix.

If $0 \leq j < s$, then

$$(d_9)_{ij} = \begin{cases} (2f(j_1, 0) - 1)w_{7(j+m)+5-2j_1 \rightarrow 7(j+m)+6} \otimes e_{7j}, \quad i = j + sj_1, \quad 0 \leq j_1 < 3; \\ (1 - 2f(j_1, 2))e_{7(j+m)+6} \otimes w_{7j \rightarrow 7j+2+j_1-f(j_1, 0)}, \quad i = j + s(4 + j_1 - f(j_1, 0)), \quad 0 \leq j_1 < 3; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_9)_{ij} = \begin{cases} w_{7(j+m+1)-4f(j_1, 0)-f(j_1, 2) \rightarrow 7(j+m+1)} \otimes w_{7j \rightarrow 7j+2j_1-2f(j_1, 3)-3f(j_1, 4)}, \\ \quad i = j + s(3 + j_1 - 3f(j_1, 0) - f(j_1, 1)), \quad 0 \leq j_1 < 5; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_9)_{ij} = \begin{cases} -w_{7(j+m+1)+1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+1 \rightarrow 7(j+1)}, \quad i = 2s + (j+1)_s; \\ w_{7(j+m)+6+f(j_1, 1) \rightarrow 7(j+m+1)+2} \otimes w_{7j+1 \rightarrow 7j+1+j_1}, \quad i = j + s(j_1 + 1), \quad 0 \leq j_1 < 3; \\ -e_{7(j+m+1)+2} \otimes w_{7j+1 \rightarrow 7j+6}, \quad i = j + 7s; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_9)_{ij} = \begin{cases} -w_{7(j+m+1)+3-2j_1 \rightarrow 7(j+m+1)+3} \otimes w_{7j+2 \rightarrow 7(j+1)}, \quad i = (j+1)_s + s(j_1 + 1), \quad 0 \leq j_1 < 2; \\ w_{7(j+m+1)-j_1 \rightarrow 7(j+m+1)+3} \otimes w_{7j+2 \rightarrow 7j+2+j_1}, \quad i = j + s(j_1 + 1), \quad 0 \leq j_1 < 2; \\ -w_{7(j+m+1)+2 \rightarrow 7(j+m+1)+3} \otimes w_{7j+2 \rightarrow 7j+6}, \quad i = j + 6s; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_9)_{ij} = \begin{cases} w_{7(j+m+1) \rightarrow 7(j+m+1)+4} \otimes e_{7j+2}, & i = j; \\ -e_{7(j+m+1)+4} \otimes w_{7j+2 \rightarrow 7j+6}, & i = j + 7s; \\ 0, & \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_9)_{ij} = \begin{cases} w_{7(j+m)+6 \rightarrow 7(j+m+1)+5} \otimes e_{7j+3}, & i = j; \\ -e_{7(j+m+1)+5} \otimes w_{7j+3 \rightarrow 7j+6}, & i = j + 5s; \\ 0, & \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_9)_{ij} = \begin{cases} w_{7(j+m+1)-f(j_1,0)+4f(j_1,3) \rightarrow 7(j+m+1)+4} \otimes w_{7j+4 \rightarrow 7j+3+j_1+f(j_1,0)}, & i = j + s(j_1 + 2f(j_1, 3)), 0 \leq j_1 < 4; \\ 0, & \text{otherwise.} \end{cases}$$

If $7s \leq j < 8s$, then

$$(d_9)_{ij} = \begin{cases} e_{7(j+m+1)+1} \otimes w_{7j+4 \rightarrow 7(j+1)}, & i = 2s + (j+1)_s; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+1} \otimes e_{7j+4}, & i = j; \\ 0, & \text{otherwise.} \end{cases}$$

If $8s \leq j < 9s$, then

$$(d_9)_{ij} = \begin{cases} w_{7(j+m+1)+2j_1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+5 \rightarrow 7j+5+j_1}, & i = j + sj_1, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $9s \leq j < 10s$, then

$$(d_9)_{ij} = \begin{cases} -e_{7(j+m+1)+5} \otimes w_{7j+5 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ w_{7(j+m+1)+6-j_1-6f(j_1,0) \rightarrow 7(j+m+1)+5} \otimes w_{7j+5 \rightarrow 7j+6-f(j_1,0)}, & i = j + s(j_1 - f(j_1, 0)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $10s \leq j < 11s$, then

$$(d_9)_{ij} = \begin{cases} w_{7(j+m+1)+5+j_1 \rightarrow 7(j+m+1)+6} \otimes w_{7j+6 \rightarrow 7(j+1)+j_1}, & i = (j+1)_s + 3sj_1, 0 \leq j_1 < 2; \\ (2f(j_1, 0) - 1)w_{7(j+m+1)+6-j_1-4f(j_1,0) \rightarrow 7(j+m+1)+6} \otimes e_{7j+6}, & i = j + s(j_1 - 1), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $11s \leq j < 12s$, then

$$(d_9)_{ij} = \begin{cases} e_{7(j+m+2)} \otimes w_{7j+6 \rightarrow 7(j+1)+4}, & i = 7s + (j+1)_s; \\ w_{7(j+m+1)+5 \rightarrow 7(j+m+2)} \otimes e_{7j+6}, & i = j - s; \\ 0, & \text{otherwise.} \end{cases}$$

Description of the d_{10}
 $d_{10} : Q_{11} \rightarrow Q_{10}$ – is an $(12s \times 10s)$ matrix.

If $0 \leq j < s$, then

$$(d_{10})_{ij} = \begin{cases} w_{7(j+m-1)+6+j_1 \rightarrow 7(j+m)+2} \otimes e_{7j}, & i = j + sj_1, 0 \leq j_1 < 2; \\ -w_{7(j+m)+2-f(j_1,1) \rightarrow 7(j+m)+2} \otimes w_{7j \rightarrow 7j+3+j_1-2f(j_1,0)}, & i = j + s(j_1 + 6 - 4f(j_1, 0)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_{10})_{ij} = \begin{cases} (2f(j_1, 0) - 1)w_{7(j+m)+4-4f(j_1,0) \rightarrow 7(j+m)+4} \otimes w_{7j \rightarrow 7j+2j_1}, & i = j + s(2j_1 + 1 - f(j_1, 0)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_{10})_{ij} = \begin{cases} -e_{7(j+m+1)} \otimes w_{7j+1 \rightarrow 7(j+1)}, & i = s + (j+1)_s; \\ (-1)^{j_1} w_{7(j+m)+2+j_1 \rightarrow 7(j+m+1)} \otimes w_{7j+1 \rightarrow 7j+1+j_1}, & i = j + sj_1, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_{10})_{ij} = \begin{cases} -e_{7(j+m)+6} \otimes w_{7j+2 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ (1 - 2f(j_1, 1, 2))w_{7(j+m)+3+j_1 \rightarrow 7(j+m)+6} \otimes w_{7j+2 \rightarrow 7j+2+f(j_1,2)+4f(j_1,3)}, & i = j + s(j_1 + 4f(j_1, 3)), 0 \leq j_1 < 4; \\ 0, & \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_{10})_{ij} = \begin{cases} w_{7(j+m)+5+2j_1 \rightarrow 7(j+m+1)} \otimes w_{7j+3 \rightarrow 7j+3+3j_1}, & i = j + s(1 + 6j_1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_{10})_{ij} = \begin{cases} -w_{7(j+m)+6+j_1 \rightarrow 7(j+m+1)} \otimes w_{7j+4 \rightarrow 7(j+1)}, & i = (j+1)_s + sj_1, 0 \leq j_1 < 2; \\ (1 - 2f(j_1, 1, 2))w_{7(j+m)+2j_1+1+3f(j_1,0)-2f(j_1,1) \rightarrow 7(j+m+1)} \otimes w_{7j+4 \rightarrow 7j+3+j_1+f(j_1,0)}, & i = j + s(2j_1 + f(j_1, 0)), 0 \leq j_1 < 4; \\ 0, & \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_{10})_{ij} = \begin{cases} (2f(j_1, 0) - 1)w_{7(j+m)+4+j_1-2f(j_1,0) \rightarrow 7(j+m)+6} \otimes w_{7j+5 \rightarrow 7j+5+f(j_1,2)}, & i = j + s(j_1 + 2), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $7s \leq j < 8s$, then

$$(d_{10})_{ij} = \begin{cases} (1 - 2f(j_1, 0, 1))w_{7(j+m)+6+j_1+3f(j_1,2,3)+2f(j_1,4) \rightarrow 7(j+m+1)+5} \otimes w_{7j+6 \rightarrow 7(j+1)+j_1-f(j_1,1)+f(j_1,4)}, & i = (j+1)_s + s(j_1 + 2f(j_1, 2, 3) + 5f(j_1, 4)), 0 \leq j_1 < 5; \\ w_{7(j+m)+6+j_1 \rightarrow 7(j+m+1)+5} \otimes e_{7j+6}, & i = j + s(3 + j_1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $8s \leq j < 9s$, then

$$(d_{10})_{ij} = \begin{cases} (-1)^{j_1+1} w_{7(j+m+1)+2+j_1 \rightarrow 7(j+m+1)+3} \otimes w_{7j+6 \rightarrow 7(j+1)+1+j_1}, \\ \quad i = (j+1)_s + s(j_1+2), \quad 0 \leq j_1 < 2; \\ w_{7(j+m)+6 \rightarrow 7(j+m+1)+3} \otimes e_{7j+6}, \quad i = j+2s; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $9s \leq j < 10s$, then

$$(d_{10})_{ij} = \begin{cases} -e_{7(j+m+1)+1} \otimes w_{7j+6 \rightarrow 7(j+1)+4}, \quad i = 7s + (j+1)_s; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+1} \otimes e_{7j+6}, \quad i = j+2s; \\ 0, \quad \text{otherwise.} \end{cases}$$

Description of the d_{11}
 $d_{11} : Q_{12} \rightarrow Q_{11}$ – is an $(10s \times 10s)$ matrix.

If $0 \leq j < s$, then

$$(d_{11})_{ij} = \begin{cases} (1 - 2f(j_1, 1) - 2f(j_1, 5))w_{7(j+m)+3+j_1-f(j_1,0)+2f(j_1,2)-f(j_1,5)-3f(j_1,6) \rightarrow 7(j+m+1)} \otimes \\ \quad w_{7j \rightarrow 7j+j_1-1+f(j_1,0)}, \\ \quad i = j + sj_1, \quad 0 \leq j_1 < 7; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_{11})_{ij} = \begin{cases} w_{7(j+m+1)+2 \rightarrow 7(j+m+1)+3} \otimes w_{7j+1 \rightarrow 7(j+1)}, \quad i = (j+1)_s; \\ (1 - 2f(j_1, 2))w_{7(j+m+1)-j_1+5f(j_1,2) \rightarrow 7(j+m+1)+3} \otimes w_{7j+1 \rightarrow 7j+1+j_1+3f(j_1,2)}, \\ \quad i = j + s(j_1+1+4f(j_1,2)), \quad 0 \leq j_1 < 3; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_{11})_{ij} = \begin{cases} e_{7(j+m+1)+4} \otimes w_{7j+1 \rightarrow 7(j+1)}, \quad i = s + (j+1)_s; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+4} \otimes e_{7j+1}, \quad i = j; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_{11})_{ij} = \begin{cases} -w_{7(j+m+1)+4 \rightarrow 7(j+m+1)+5} \otimes w_{7j+2 \rightarrow 7(j+1)}, \quad i = s + (j+1)_s; \\ (1 - 2f(j_1, 2))w_{7(j+m)+6+j_1+4f(j_1,2) \rightarrow 7(j+m+1)+5} \otimes w_{7j+2 \rightarrow 7j+2+j_1+2f(j_1,2)}, \\ \quad i = j + s(j_1+2f(j_1,2)), \quad 0 \leq j_1 < 3; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_{11})_{ij} = \begin{cases} (-1)^{j_1} w_{7(j+m+1)+j_1 \rightarrow 7(j+m+1)+1} \otimes w_{7j+3 \rightarrow 7j+3+3j_1}, \quad i = j + 5sj_1, \quad 0 \leq j_1 < 2; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_{11})_{ij} = \begin{cases} e_{7(j+m+1)+2} \otimes w_{7j+4 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ (-1)^{j_1} w_{7(j+m+1)+j_1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+4 \rightarrow 7j+4+2j_1}, & i = j + 4sj_1, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_{11})_{ij} = \begin{cases} (2f(j_1, 0) - 1) w_{7(j+m+1)-j_1+7f(j_1, 2) \rightarrow 7(j+m+1)+5} \otimes w_{7j+4 \rightarrow 7j+4+j_1}, & i = j + s(j_1 - 1), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $7s \leq j < 8s$, then

$$(d_{11})_{ij} = \begin{cases} w_{7(j+m)+6+4j_1 \rightarrow 7(j+m+1)+3} \otimes w_{7j+5 \rightarrow 7j+5+j_1}, & i = j + s(2j_1 - 1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $8s \leq j < 9s$, then

$$(d_{11})_{ij} = \begin{cases} w_{7(j+m+1)+6-4f(j_1, 0) \rightarrow 7(j+m+1)+6} \otimes w_{7j+6 \rightarrow 7(j+1)+2j_1+f(j_1, 2)}, & i = (j+1)_s + 3sj_1, 0 \leq j_1 < 3; \\ (2f(j_1, 0) - 1) w_{7(j+m+1)+5-2j_1 \rightarrow 7(j+m+1)+6} \otimes e_{7j+6}, & i = j + s(j_1 - 1), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $9s \leq j < 10s$, then

$$(d_{11})_{ij} = \begin{cases} e_{7(j+m+2)} \otimes w_{7j+6 \rightarrow 7(j+1)+1}, & i = 2s + (j+1)_s; \\ w_{7(j+m+1)+3 \rightarrow 7(j+m+2)} \otimes e_{7j+6}, & i = j - s; \\ 0, & \text{otherwise.} \end{cases}$$

Description of the d_{12}

$d_{12} : Q_{13} \rightarrow Q_{12}$ – is an $(10s \times 9s)$ matrix.

If $0 \leq j < s$, then

$$(d_{12})_{ij} = \begin{cases} (2f(j_1, 0) + 2f(j_1, 3) - 1) w_{7(j+m)+j_1-1+f(j_1, 0)+3f(j_1, 1) \rightarrow 7(j+m)+3} \otimes w_{7j \rightarrow 7j+j_1+1-f(j_1, 0, 1)}, & i = j + s(2j_1 - 1 + f(j_1, 0) + f(j_1, 2)), 0 \leq j_1 < 5; \\ 0, & \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_{12})_{ij} = \begin{cases} w_{7(j+m) \rightarrow 7(j+m)+5} \otimes e_{7j}, & i = j_s; \\ (2f(j_1, 2) - 1) w_{7(j+m)+5-f(j_1, 0) \rightarrow 7(j+m)+5} \otimes w_{7j \rightarrow 7j+2j_1+f(j_1, 0)}, & i = j + s(j_1 + 1 + 2f(j_1, 2)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_{12})_{ij} = \begin{cases} (2f(j_1, 0) + 2f(j_1, 3) - 1) w_{7(j+m)+3+j_1-5f(j_1, 3)-f(j_1, 4) \rightarrow 7(j+m)+6} \otimes \\ w_{7j+1 \rightarrow 7j+j_1+f(j_1, 0)+2f(j_1, 4)}, & i = j + s(j_1 - 1 + 3f(j_1, 4)), 0 \leq j_1 < 5; \\ 0, & \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_{12})_{ij} = \begin{cases} -e_{7(j+m+1)} \otimes w_{7j+2 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ (1 - 2f(j_1, 1))w_{7(j+m)+4+j_1+f(j_1, 0)-4f(j_1, 1) \rightarrow 7(j+m+1)} \otimes w_{7j+2 \rightarrow 7j+6-4f(j_1, 0)-3f(j_1, 1)}, \\ & i = j + s(j_1 + 3f(j_1, 2, 3)), 0 \leq j_1 < 4; \\ 0, & \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_{12})_{ij} = \begin{cases} (-1)^{j_1} w_{7(j+m+1)+j_1 \rightarrow 7(j+m+1)+1} \otimes w_{7j+3 \rightarrow 7(j+1)+3j_1}, \\ & i = (j+1)_s + 4sj_1, 0 \leq j_1 < 2; \\ (2f(j_1, 0) - 1)w_{7(j+m)+5+j_1-4f(j_1, 0) \rightarrow 7(j+m+1)+1} \otimes w_{7j+3 \rightarrow 7j+6-3f(j_1, 0)}, \\ & i = j + s(3 + j_1 - 3f(j_1, 0)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_{12})_{ij} = \begin{cases} (2f(j_1, 0) - 1)w_{7(j+m)+2+3f(j_1, 1)+f(j_1, 2)+4f(j_1, 3) \rightarrow 7(j+m)+6} \otimes w_{7j+4 \rightarrow 7j+3+j_1+f(j_1, 0)}, \\ & i = j + sj_1, 0 \leq j_1 < 4; \\ 0, & \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_{12})_{ij} = \begin{cases} (-1)^{j_1} w_{7(j+m)+3+4j_1 \rightarrow 7(j+m+1)} \otimes w_{7j+5 \rightarrow 7j+5+j_1}, & i = j + s(2j_1 + 1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $7s \leq j < 8s$, then

$$(d_{12})_{ij} = \begin{cases} (2f(j_1, 1) - 1)w_{7(j+m+1)+j_1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+6 \rightarrow 7(j+1)+2+j_1-2f(j_1, 0)}, \\ & i = (j+1)_s + s(3 + j_1 - 3f(j_1, 0)), 0 \leq j_1 < 3; \\ w_{7(j+m)+6+j_1 \rightarrow 7(j+m+1)+2} \otimes e_{7j+6}, & i = j + s(j_1 + 1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $8s \leq j < 9s$, then

$$(d_{12})_{ij} = \begin{cases} -e_{7(j+m+1)+4} \otimes w_{7j+6 \rightarrow 7(j+1)+1}, & i = 2s + (j+1)_s; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+4} \otimes e_{7j+6}, & i = j + s; \\ 0, & \text{otherwise.} \end{cases}$$

Description of the d_{13}
 $d_{13} : Q_{14} \rightarrow Q_{13}$ – is an $(9s \times 7s)$ matrix.

If $0 \leq j < s$, then

$$(d_{13})_{ij} = \begin{cases} (-1)^{j_1} w_{7(j+m)+6-3f(j_1, 0)-f(j_1, 1) \rightarrow 7(j+m)+6} \otimes w_{7j \rightarrow 7j+j_1-1+f(j_1, 0)+2f(j_1, 3)}, \\ & i = j + s(j_1 + 2f(j_1, 3)), 0 \leq j_1 < 4; \\ 0, & \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_{13})_{ij} = \begin{cases} e_{7(j+m+1)+5} \otimes w_{7j+1 \rightarrow 7(j+1)}, & i = s + (j+1)_s; \\ (1 - 2f(j_1, 2))w_{7(j+m)+6+j_1+3f(j_1, 2) \rightarrow 7(j+m+1)+5} \otimes w_{7j+1 \rightarrow 7j+1+j_1+3f(j_1, 2)}, \\ & i = j + s(j_1 + 1 + 4f(j_1, 2)), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_{13})_{ij} = \begin{cases} w_{7(j+m+1)+j_1 \rightarrow 7(j+m+1)+1} \otimes w_{7j+2 \rightarrow 7j+2+j_1}, & i = j + s(j_1 + 1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_{13})_{ij} = \begin{cases} w_{7(j+m+1)+1+j_1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+3 \rightarrow 7j+3+3j_1}, & i = j + s(3j_1 + 1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_{13})_{ij} = \begin{cases} e_{7(j+m+1)+3} \otimes w_{7j+4 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ w_{7(j+m)+6+j_1+f(j_1, 2) \rightarrow 7(j+m+1)+3} \otimes w_{7j+4 \rightarrow 7j+4+j_1}, & i = j + s(j_1 + 1), 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_{13})_{ij} = \begin{cases} w_{7(j+m+1)+4j_1 \rightarrow 7(j+m+1)+4} \otimes w_{7j+5 \rightarrow 7j+5+j_1}, & i = j + s(2j_1 + 1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_{13})_{ij} = \begin{cases} w_{7(j+m+2)-4f(j_1, 0)-f(j_1, 1) \rightarrow 7(j+m+2)} \otimes w_{7j+6 \rightarrow 7(j+1)+j_1+2f(j_1, 3)}, \\ & i = (j+1)_s + s(j_1 + 1 - f(j_1, 0) + 2f(j_1, 3)), 0 \leq j_1 < 4; \\ (-1)^{j_1} w_{7(j+m+1)+2+2j_1 \rightarrow 7(j+m+2)} \otimes e_{7j+6}, & i = j + s(j_1 + 1), 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

Description of the d_{14}
 $d_{14} : Q_{15} \rightarrow Q_{14}$ – is an $(7s \times 8s)$ matrix.

If $0 \leq j < s$, then

$$(d_{14})_{ij} = \begin{cases} -e_{7(j+m)+6} \otimes w_{7j \rightarrow 7(j+1)}, & i = (j+1)_s; \\ (-1)^{j_1} w_{7(j+m-1)+6+j_1+5f(j_1, 1) \rightarrow 7(j+m)+6} \otimes w_{7j \rightarrow 7j+j_1}, \\ & i = j + sj_1, 0 \leq j_1 < 6; \\ 0, & \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_{14})_{ij} = \begin{cases} w_{7(j+m)+6 \rightarrow 7(j+m+1)} \otimes w_{7j+1 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ (-1)^{j_1} w_{7(j+m)+j_1+5f(j_1, 0) \rightarrow 7(j+m+1)} \otimes w_{7j+1 \rightarrow 7j+1+j_1}, \\ & i = j + sj_1, 0 \leq j_1 < 3; \\ -e_{7(j+m+1)} \otimes w_{7j+1 \rightarrow 7j+6}, & i = j + 5s; \\ 0, & \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_{14})_{ij} = \begin{cases} -w_{7(j+m)+6+2j_1 \rightarrow 7(j+m+1)+1} \otimes w_{7j+2 \rightarrow 7(j+1)+2j_1}, & i = (j+1)_s + 2sj_1, 0 \leq j_1 < 2; \\ (-1)^{j_1} w_{7(j+m)+1+j_1 \rightarrow 7(j+m+1)+1} \otimes w_{7j+2 \rightarrow 7j+2+j_1}, & i = j + sj_1, 0 \leq j_1 < 2; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+1} \otimes w_{7j+2 \rightarrow 7j+6}, & i = j + 4s; \\ 0, & \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_{14})_{ij} = \begin{cases} (-1)^{j_1} w_{7(j+m+1)+1+j_1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+3 \rightarrow 7(j+1)+2+j_1}, & i = (j+1)_s + s(2+j_1), 0 \leq j_1 < 2; \\ w_{7(j+m)+6 \rightarrow 7(j+m+1)+2} \otimes w_{7j+3 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ (-1)^{j_1} w_{7(j+m)+2+5j_1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+3 \rightarrow 7j+3+3j_1}, & i = j + 3sj_1, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_{14})_{ij} = \begin{cases} (2f(j_1, 0) - 1) w_{7(j+m)+3+j_1+2f(j_1, 2) \rightarrow 7(j+m+1)} \otimes w_{7j+4 \rightarrow 7j+4+j_1}, & i = j + sj_1, 0 \leq j_1 < 3; \\ 0, & \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_{14})_{ij} = \begin{cases} -e_{7(j+m+1)+4} \otimes w_{7j+5 \rightarrow 7(j+1)+5}, & i = 5s + (j+1)_s; \\ w_{7(j+m)+4+3j_1 \rightarrow 7(j+m+1)+4} \otimes w_{7j+5 \rightarrow 7j+5+j_1}, & i = j + sj_1, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_{14})_{ij} = \begin{cases} -w_{7(j+m)+6 \rightarrow 7(j+m+1)+3} \otimes w_{7j+6 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ (-1)^{j_1+1} w_{7(j+m+1)+1+j_1 \rightarrow 7(j+m+1)+3} \otimes w_{7j+6 \rightarrow 7(j+1)+2+j_1}, & i = (j+1)_s + s(2+j_1), 0 \leq j_1 < 3; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+3} \otimes e_{7j+6}, & i = j; \\ 0, & \text{otherwise.} \end{cases}$$

If $7s \leq j < 8s$, then

$$(d_{14})_{ij} = \begin{cases} -w_{7(j+m+1)+5-j_1 \rightarrow 7(j+m+1)+5} \otimes w_{7j+6 \rightarrow 7(j+1)+1+4j_1}, & i = (j+1)_s + s(1+4j_1), 0 \leq j_1 < 2; \\ w_{7(j+m+1) \rightarrow 7(j+m+1)+5} \otimes e_{7j+6}, & i = j - s; \\ 0, & \text{otherwise.} \end{cases}$$

Description of the d_{15}
 $d_{15} : Q_{16} \rightarrow Q_{15}$ – is an $(8s \times 7s)$ matrix.

If $0 \leq j < s$, then

$$(d_{15})_{ij} = \begin{cases} w_{7(j+m)+6+j_1 \rightarrow 7(j+m+1)} \otimes w_{7j \rightarrow 7j+j_1}, & i = j + sj_1, 0 \leq j_1 < 2; \\ -e_{7(j+m+1)} \otimes w_{7j \rightarrow 7j+4}, & i = j + 4s; \\ 0, & \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_{15})_{ij} = \begin{cases} w_{7(j+m+1)+j_1 \rightarrow 7(j+m+1)+1} \otimes w_{7j+1 \rightarrow 7j+1+j_1}, & i = j + sj_1, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_{15})_{ij} = \begin{cases} w_{7(j+m+1)+1+j_1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+2 \rightarrow 7j+2+j_1}, & i = j + sj_1, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_{15})_{ij} = \begin{cases} w_{7(j+m+1)+2+j_1 \rightarrow 7(j+m+1)+3} \otimes w_{7j+3 \rightarrow 7j+3+3j_1}, & i = j + 3sj_1, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_{15})_{ij} = \begin{cases} w_{7(j+m+1)+4+j_1 \rightarrow 7(j+m+1)+4} \otimes w_{7j+4 \rightarrow 7j+4+j_1}, & i = j + sj_1, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_{15})_{ij} = \begin{cases} (-1)^{j_1} w_{7(j+m+1)+4+j_1 \rightarrow 7(j+m+1)+5} \otimes w_{7j+5 \rightarrow 7j+5+j_1}, & i = j + 2sj_1, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_{15})_{ij} = \begin{cases} e_{7(j+m+1)+6} \otimes w_{7j+6 \rightarrow 7(j+1)}, & i = (j+1)_s; \\ (-1)^{j_1} w_{7(j+m+1)+3+2j_1 \rightarrow 7(j+m+1)+6} \otimes e_{7j+6}, & i = j + sj_1, 0 \leq j_1 < 2; \\ 0, & \text{otherwise.} \end{cases}$$

Description of the d_{16}
 $d_{16} : Q_{17} \rightarrow Q_{16}$ – is an $(7s \times 7s)$ matrix.

If $0 \leq j < s$, then

$$(d_{16})_{ij} = \begin{cases} -e_{7(j+m+1)} \otimes w_{7j \rightarrow 7(j+1)}, & i = (j+1)_s; \\ (-1)^{j_1} w_{7(j+m)+j_1 \rightarrow 7(j+m+1)} \otimes w_{7j \rightarrow 7j+j_1}, & i = j + sj_1, 0 \leq j_1 < 7; \\ 0, & \text{otherwise.} \end{cases}$$

If $s \leq j < 2s$, then

$$(d_{16})_{ij} = \begin{cases} (-1)^{j_1} w_{7(j+m+1)+j_1 \rightarrow 7(j+m+1)+1} \otimes w_{7j+1 \rightarrow 7(j+1)+j_1}, \\ \quad i = (j+1)_s + sj_1, 0 \leq j_1 < 2; \\ (-1)^{j_1} w_{7(j+m)+1+j_1 \rightarrow 7(j+m+1)+1} \otimes w_{7j+1 \rightarrow 7j+1+j_1}, & i = j + sj_1, 0 \leq j_1 < 3; \\ -w_{7(j+m)+6 \rightarrow 7(j+m+1)+1} \otimes w_{7j+1 \rightarrow 7j+6}, & i = j + 5s; \\ 0, & \text{otherwise.} \end{cases}$$

If $2s \leq j < 3s$, then

$$(d_{16})_{ij} = \begin{cases} (-1)^{j_1+1} w_{7(j+m+1)+j_1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+2 \rightarrow 7(j+1)+j_1}, \\ \quad i = (j+1)_s + sj_1, \quad 0 \leq j_1 < 3; \\ (-1)^{j_1} w_{7(j+m)+2+j_1 \rightarrow 7(j+m+1)+2} \otimes w_{7j+2 \rightarrow 7j+2+j_1}, \quad i = j + sj_1, \quad 0 \leq j_1 < 2; \\ w_{7(j+m)+6 \rightarrow 7(j+m+1)+2} \otimes w_{7j+2 \rightarrow 7j+6}, \quad i = j + 4s; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $3s \leq j < 4s$, then

$$(d_{16})_{ij} = \begin{cases} (-1)^{j_1} w_{7(j+m+1)+j_1 \rightarrow 7(j+m+1)+3} \otimes w_{7j+3 \rightarrow 7(j+1)+j_1}, \\ \quad i = (j+1)_s + sj_1, \quad 0 \leq j_1 < 4; \\ (-1)^{j_1} w_{7(j+m)+3+3j_1 \rightarrow 7(j+m+1)+3} \otimes w_{7j+3 \rightarrow 7j+3+3j_1}, \quad i = j + 3sj_1, \quad 0 \leq j_1 < 2; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $4s \leq j < 5s$, then

$$(d_{16})_{ij} = \begin{cases} -w_{7(j+m+1)+4j_1 \rightarrow 7(j+m+1)+4} \otimes w_{7j+4 \rightarrow 7(j+1)+4j_1}, \quad i = (j+1)_s + 4sj_1, \quad 0 \leq j_1 < 2; \\ (-1)^{j_1} w_{7(j+m)+4+j_1 \rightarrow 7(j+m+1)+4} \otimes w_{7j+4 \rightarrow 7j+4+j_1}, \quad i = j + sj_1, \quad 0 \leq j_1 < 3; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $5s \leq j < 6s$, then

$$(d_{16})_{ij} = \begin{cases} w_{7(j+m+1) \rightarrow 7(j+m+1)+5} \otimes w_{7j+5 \rightarrow 7(j+1)}, \quad i = (j+1)_s; \\ (-1)^{j_1} w_{7(j+m+1)+4+j_1 \rightarrow 7(j+m+1)+5} \otimes w_{7j+5 \rightarrow 7(j+1)+4+j_1}, \\ \quad i = (j+1)_s + s(4+j_1), \quad 0 \leq j_1 < 2; \\ (-1)^{j_1} w_{7(j+m)+5+j_1 \rightarrow 7(j+m+1)+5} \otimes w_{7j+5 \rightarrow 7j+5+j_1}, \quad i = j + sj_1, \quad 0 \leq j_1 < 2; \\ 0, \quad \text{otherwise.} \end{cases}$$

If $6s \leq j < 7s$, then

$$(d_{16})_{ij} = \begin{cases} (-1)^{j_1+1} w_{7(j+m+1)+j_1 \rightarrow 7(j+m+1)+6} \otimes w_{7j+6 \rightarrow 7(j+1)+j_1}, \\ \quad i = (j+1)_s + sj_1, \quad 0 \leq j_1 < 7; \\ w_{7(j+m)+6 \rightarrow 7(j+m+1)+6} \otimes e_{7j+6}, \quad i = j; \\ 0, \quad \text{otherwise.} \end{cases}$$