Noninvertible Gauge Symmetry in (2+1)d Topological Orders: A String-Net Model Realization

Yu Zhao^{*a*} Yidun Wan^{1*a,b*}

 ^a State Key Laboratory of Surface Physics, Center for Astronomy and Astrophysics, Department of Physics, Center for Field Theory and Particle Physics, and Institute for Nanoelectronic devices and Quantum Computing, Fudan University, 2005 Songhu Road, Shanghai 200433, China
 ^b Shanghai Research Center for Quantum Sciences, 99 Xiupu Road, Shanghai 201315, China

E-mail: yuzhao20@fudan.edu.cn, ydwan@fudan.edu.cn

ABSTRACT: We develop a systematic framework for understanding symmetries in topological phases in 2 + 1 dimensions using the string-net model, encompassing both gauge symmetries that preserve anyon species and global symmetries permuting anyon species, including both invertible symmetries describable by groups and noninvertible symmetries described by categories. As an archetypal example, we reveal the first noninvertible categorical gauge symmetry of topological orders in 2 + 1 dimensions: the Fibonacci gauge symmetry of the doubled Fibonacci topological order, described by the Fibonacci fusion 2-category. Our approach involves two steps: first, establishing duality between different string-net models with Morita equivalent input fusion categories that describe the same topological order; and second, constructing symmetry transformations within the same string-net model when the dual models have isomorphic input fusion categories, achieved by composing duality maps with isomorphisms of degrees of freedom between the dual models.

¹Corresponding author

Contents

1	Introduction	2
2	Sketch of Our Approach	3
3	Fibonacci String-Net Model	4
4	Duality Map and Enlarging the Hilbert Space	5
5	Symmetry Transformation of the Doubled Fibonacci Topological Order	7
6	Fibonacci Categorical Gauge Symmetry	10
\mathbf{A}	Review of the Extended String-net Model	12
	A.1 Topological Features	14
	A.2 Excited States	15
в	Frobenius Algebras and Bimodules	17
	B.1 Frobenius Algebra	17
	B.2 Bimodules over a Frobenius Algebra	18
	B.3 The Bimodule Fusion Category over a Frobenius Algebra	19
С	General Constructions of Dualities and Symmetry Transformations in	
	the Extended String-Net Model	21
	C.1 Enlarging the Hilbert Space	22
	C.2 Duality	23
	C.3 Symmetry Transformation	24
	C.4 Braiding of Bimodules	24
D	The \mathbb{Z}_2 Global Symmetry transformation of the \mathbb{Z}_2 Toric Code String-Net	
	Model	25
	D.1 The toric code string-net model	25
	D.2 The global symmetry transformation of the toric code topological order	26
	D.3 Criterion for Distinguishing Gauge Symmetries from Global Symmetries	28
\mathbf{E}	Frobenius Algebra of Fibonacci Fusion Categories and Its Simple Bimod-	
	ules	2 8

1 Introduction

Symmetry is a central concept in modern physics, traditionally described by groups. Topological orders in 2 + 1 dimensions, whose low-energy effective descriptions are topological gauge field theories, are phases of matter beyond the Landau-Ginzburg paradigm, implying novel symmetry structures. Nevertheless, gauge symmetries¹ in topological orders are not completely clear. In the Dijkgraaf-Witten theory[1–4] or their lattice Hamiltonian model—the (twisted) quantum double model of topological orders[5–13], the gauge structure, i.e., gauge group and gauge transformation, has been well understood. More generally, such as in the Turaev-Viro topological field theory[2, 14–17] or their lattice Hamiltonian model—the string-net model[18–29], the gauge symmetry is often not known. There, if a gauge symmetry exists, it is likely not describable by any group but by categories[30–40], invoking the concept of noninvertible symmetry in physics. Noninvertible symmetry has been extensively explored in 1+1 dimensions[46–50] but remain largely an open problem in 2+1 dimensions, despite studies in certain (2 + 1)d systems[51–53] and attempts in applying noninvertible symmetry to QFT and M theory[54–58].

This article tackles this problem by systematically formulating the symmetry transformations of (2+1)-dimensional topological orders as operators of the string-net model. We show that such symmetry transformations can be either gauge symmetries that preserve anyon species or global symmetries permuting anyon species and can be invertible (describable by groups) or noninvertible (described by a fusion 2-categories). As a key result, our construction reveals the first noninvertible gauge symmetry of topological orders in 2 + 1dimensions:

The noninvertible gauge symmetry (to be called the Fibonacci gauge symmetry) of the doubled Fibonacci topological order described by the Fibonacci fusion 2-category.

This is an archetypal example illustrating our general construction, which is detailed in Appendix C. In the main text, we shall provide a brief overview of our approach and expound on the Fibonacci gauge symmetry. Besides, we also provide a clear criterion (see Appendix D.3) determining whether a symmetry transformation generates a global or a gauge symmetry of the topological order. An example of global symmetry so constructed is shown in Appendix D.

Our construction can help realize anyon condensations [31, 59–63] in the string-net model. Fluxon and simple-current condensations—special types of anyon condensation—have been realized in string-net model[29, 64], but not general non-Abelian anyon condensation. As to be reported elsewhere, we can apply our symmetry transformations to the string-net model and then perform all types of anyon condensations in familiar forms in Refs. [29, 64] in the transformed model.

¹A historical misnomer; it should be more appropriately called gauge invariance or gauge redundancy because it is mathematical redundancy where different states in the Hilbert space describe the same physical state.

2 Sketch of Our Approach

As the low-energy effective theories of topological orders, topological quantum field theories (TQFTs) are believed to be gauge theories. For the Dijkgraaf-Witten TQFT, its gauge symmetry and transformation are manifest and clear. For more general topological orders, whose effective theory is the Turaev-Viro TQFT, however, its gauge symmetry and transformations are in general obscure.

To reveal the gauge structure of topological orders in general, we first note that topological orders are primarily observed in strongly correlated electron systems [65-72], where anyons are collective excitations of the electrons, the fundamental degrees of freedom (d.o.f) comprising the physical Hilbert space, where anyons are represented as excited states. While an Abelian anyon is represented by one excited state up to phase, a non-Abelian anyon is represented by a multi-dimensional subspace of the Hilbert space $[73-78]^2$. That is, a physical anyon may have an internal space [26, 29, 80]. Nevertheless, TQFT fails to capture such internal spaces of anyons because anyons are elementary and indecomposable ingredients in TQFT. In contrast, in the string-net model—a Hamiltonian extension of the Turaev-Viro TQFT—of topological orders, anyon excitations are concrete excited states of the model. Anyons are represented by orthogonal subspaces (one- or multi-dimensional) in the Hilbert space. We show that in the string-net model, it is possible to transform the fundamental d.o.f. while preserving topological invariance and the Hamiltonian. Such a transformation would induce a linear transformation of the Hilbert space of the model. When this linear transformation preserves the subspaces representing the anyons, it is qualified as a gauge transformation of the model. The model and thus the topological order it describes has a corresponding gauge symmetry. When this transformation can turn the anyons to one another, it is global.

We construct such symmetry transformations in two steps. Firstly, we establish a duality between different string-net models describing the same topological order. While the fundamental d.o.f.s in a usual gauge theory take value in a group, those in the Turaev-Viro TQFT or a string-net model are simple objects in a fusion category—the input of the model. Our duality maps two string-net models whose input fusion categories are Morita equivalent to each other[81] and classifies all string-net models describing the same topological order. This duality generalizes the electromagnetic duality[11, 13, 82] in topological orders and topological field theories.

In the second step, symmetry transformations can be constructed when two dual stringnet models have isomorphic input fusion categories. In such a case, the duality in the first step can be composed with an isomorphism to form a symmetry transformation on the same string-net model. Such a symmetry transformation can be global or a gauge depending on whether it permutes anyon species. Such a symmetry, when it is noninvertible, is described by a fusion 2-category. We now construct the gauge symmetry transformation of the doubled Fibonacci topological order to illustrate our approach. We shall first briefly review the Fibonacci string-net model.

²This is why non-Abelian anyons can support topological quantum computation[79].



Figure 1: Part of the string-net model lattice. A tail (wavy line) is attached to an arbitrary edge of every plaquette.

3 Fibonacci String-Net Model

We take the form of the string-net model defined in [26] because its Hilbert space encompasses the full anyon spectra of the topological orders. The model is defined on a 2-dimensional trivalent lattice, e.g., that in Fig. 1. Each plaquette has a tail attached to any of its edges³. Each edge or tail carries a label—fundamental d.o.f—taking value in the simple objects of the model's input fusion category \mathscr{F} . The Hilbert space is spanned by all possible assignments of the labels, such that the labels on three edges (tails) meeting at any vertex satisfy the fusion rules of \mathscr{F} .

Let's solve our example: the doubled Fibonacci topological order with the string-net model. The input Fibonacci fusion category Fibo has two simple objects 1 and τ , the basic d.o.f. on edges/tails. The Hilbert space \mathcal{H}_{Fibo} is spanned by all possible assignments of 1 and τ to all edges/tails, dictated by the nonzero fusion rule $\delta_{ijk} = 1$ for three edges/tails i, j, k meeting at any vertex:

$$\delta_{111} = \delta_{1\tau\tau} = \delta_{\tau1\tau} = \delta_{\tau\tau1} = \delta_{\tau\tau\tau} = 1.$$

The Hamiltonian is a sum of commuting projectors:

$$H_{\mathsf{Fibo}} := -\sum_{\mathrm{Plaquettes} \ P} Q_P \ ,$$

where operators Q_P are detailed in Appendix A. The ground states are common eigenstates of all Q_P operators with +1 eigenvalues. An excited state $|\psi\rangle$ is another common eigenstate that satisfies $Q_P |\psi\rangle = 0$ for one or more plaquettes P, each said to hold an anyon. A ground state has a trivial anyon in each plaquette. The doubled Fibonacci topological order has four anyon species:

$$1\overline{1}, \quad \tau\overline{1}, \quad 1\overline{\tau}, \quad \tau\overline{\tau},$$

 $^{^{3}}$ The original string-net model in Ref. [18], which has no such tails, cannot fully describe charge excitations. These added tails carry the charges of anyons, thus enlarging the Hilbert space to encompass the complete anyon spectrum.

where $1\overline{1}$ is the trivial anyon.

It is the kairos to invoke the concept of anyon's *internal space*. Unlike Abelian anyons, a non-Abelian anyon has a nontrivial internal space because multiple non-Abelian anyons occupy a multi-dimensional Hilbert space in TQFT[73–78]. Nevertheless, the internal space of a single non-Abelian anyon is generally hidden in the language of TQFT. The string-net model is however able to manifest such internal spaces because an excited state is labeled by both the anyon species J in each plaquette and the anyon's internal charge p—the d.o.f. on the tail where the anyon resides. A non-Abelian anyon carries more than one charge type and is thus represented on a certain multi-dimensional Hilbert subspace of excited states of the model. The gauge symmetry transformation to be constructed may mix the internal charges p while preserving the anyon species J of a non-Abelian anyon. The Fibonacci string-net model has five allowed pairs (J, p):

$$(1\bar{1},1),$$
 $(1\bar{\tau},\tau),$ $(\tau\bar{1},\tau),$ $(\tau\bar{\tau},1),$ $(\tau\bar{\tau},\tau).$

An anyon $\tau \bar{\tau}$ has two possible charges 1 and τ in the model. Anyons $\tau \bar{1}$ and $1\bar{\tau}$ are both non-Abelian but seem to carry only one charge τ . Yet, the gauge structure of the doubled Fibonacci topological order will urge us to enlarge the model's Hilbert space, such that anyons $\tau \bar{1}$ and $1\bar{\tau}$ will each carry two different charges.

4 Duality Map and Enlarging the Hilbert Space

A fusion category \mathscr{F} has *Frobenius algebra* objects. It is a theorem[81] that the bimodules a special class of representations of a given Frobenius algebra \mathcal{A} in \mathscr{F} —form another fusion category $\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$ that is *categorically Morita equivalent* to \mathscr{F} . This fact implies that two string-net models with input fusion categories $\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$ and \mathscr{F} are equivalent in describing the same topological order. We now explicitly establish this equivalence by a duality map \mathcal{D} between these two models.

Fusion category Fibo has a nontrivial Frobenius algebra

$$\mathcal{A} := \{ \alpha 1 + \beta \tau \mid \alpha, \beta \in \mathbb{C}, \ \tau^2 = 1 + \phi^{-\frac{3}{4}} \tau \},$$

$$(4.1)$$

where simple objects 1 and τ are regarded as the two basis elements of \mathcal{A} . \mathcal{A} has two simple (viz irreducible) bimodules: The trivial bimodule $M_1 = (P_1, V_1)$, with representation space

$$V_1 := \{ \alpha 1 + \beta \tau | \alpha, \beta \in \mathbb{C} \},\$$

and the nontrivial bimodule $M_{\tau} = (P_{\tau}, V_{\tau})$ with the representation space

$$V_{\tau} := \{ \alpha 1 + \beta \tau_1 + \gamma \tau_2 | \alpha, \beta, \gamma \in \mathbb{C} \}.$$

Here, τ_1 and τ_2 are both the simple object τ , but regarded as different basis elements in V_{τ} because they are acted on by \mathcal{A} differently⁴. The indices 1 and 2 of τ in M_{τ} are the

⁴This is analogous to the scenario where an irreducible representation of a group can appear more than once in a certain reducible representation of the group. When physics kicks in, the different occurrences of the same irreducible representation are distinguishable.

multiplicity labels. Functions $P_i : \mathcal{A}^2 \times V_i \times \{1, \tau\} \times V_i \to \mathbb{C}$ represent pairs $(a, b) \in \mathcal{A}^2$ as rank-3 tensors on the representation spaces V_i , where $i = 1, \tau$. The components of P_i are listed in Appendix E. These tensors indicate that a and b act sequentially on $x \in V_i$, transforming it to z with the coefficient $\sum_{y} [P_i]_{xyz}^{ab}$. The intermediate object y varies in $\{1, \tau\}$ to satisfy the fusion rule $\delta_{axy} = \delta_{byz} = 1$.

The string-net model with the input fusion category $\mathsf{Bimod}_{\mathsf{Fibo}}(\mathcal{A})$ is the *dual Fibological string-net model*. The d.o.f. on dual model's edges/tails are simple objects in $\mathsf{Bimod}_{\mathsf{Fibo}}(\mathcal{A})$ —the simple bimodules M_1 and M_{τ} . By the definition of bimodules, we construct a duality map \mathcal{D} that embeds the d.o.f. of the dual model as superpositions of those in the original model:

$$\mathcal{D} \left| M_i \right| := \sum_{a,b,y=1,\tau} \sum_{x,z \in L_{M_i}} [P_i]^{ab}_{xyz} = \begin{bmatrix} z \\ y \\ x \end{bmatrix}, \quad (4.2)$$

where $i = 1, \tau, L_{M_1} = \{1, \tau\}$, and $L_{M_{\tau}} = \{1, \tau_1, \tau_2\}$. A black line can be an edge or tail. The red lines are auxiliary and will be annihilated by topological moves, resulting in a unitary transformation between the two models, which can be understood plaquette by plaquette:

where $I_k, E_k, M \in \{M_1, M_\tau\}$, and " \cdots " omits the expansion coefficients detailed in Appendix C. After the topological moves, the d.o.f. τ on any edge will cease to have any multiplicity index, while that on any tail will keep a multiplicity index if it belongs to L_{M_τ} . Therefore, to justify this duality, we have to enlarge the Hilbert space $\mathcal{H}_{\mathsf{Fibo}}$ of the original Fibonacci string-net model to \mathcal{H}^* by distinguishing τ_1 and τ_2 on each tail but not on the edges. This enlargement is physically sound: The tail carries an anyon's internal charge that reflects the action of \mathcal{A} , which can only be told when different occurrences of τ in the bimodules of \mathcal{A} are distinguished by multiplicity indices. In contrast, the d.o.f. on edges are pertaining to ground states because any path along edges has to be a closed loop. At any vertex along such a loop, fusion rules are met; they treat τ_1 and τ_2 the same⁵.

Due to orthonormality, the actual enlargement is done by embedding $\mathcal{H}_{\mathsf{Fibo}}$ in \mathcal{H}^* as

$$\begin{cases} \tau \\ \end{array} \implies \left(\frac{1}{2\phi} + \frac{\sqrt{\phi}}{2}\right) \\ \end{cases}^{\tau_1} + \left(\frac{1}{2\phi} - \frac{\sqrt{\phi}}{2}\right) \\ \end{cases}^{\tau_2}$$
(4.4)

 $^{{}^{5}}$ As an analogy: It makes no sense to question the electric charge in a closed electric flux loop because the Gauss law (analogous to fusion rules) is met everywhere along the loop. Only when the loop is cut open to be a path, one can ask about the charges at the ends of the path where the Gauss law is broken.

for each *tail*, as shown in Fig. 2a. The original Hilbert space $\mathcal{H}_{\mathsf{Fibo}}$ of the string-net model is a subspace of \mathcal{H}^* .

5 Symmetry Transformation of the Doubled Fibonacci Topological Order

We can further construct a symmetry transformation of the doubled Fibonacci phase based on the duality \mathcal{D} defined in Eqs. (4.2) and (4.3) because their input fusion categories are isomorphic:

$$\mathcal{F}_{\mathcal{A}} : \mathsf{Fibo} \to \mathsf{Bimod}_{\mathsf{Fibo}}(\mathcal{A}), \quad 1 \mapsto M_1, \quad \tau \mapsto M_{\tau}.$$
 (5.1)

This isomorphism $\mathcal{F}_{\mathcal{A}}$ induces an isomorphic map $\varphi_{\mathcal{A}}$ between the Fibonacci string-net model and its dual model, and thus a *unitary transformation* \mathcal{G} of the Fibonacci string-net model in the enlarged Hilbert space \mathcal{H}^* :

$$\mathcal{G} := \mathcal{D} \circ \varphi_{\mathcal{A}} , \quad \varphi_{\mathcal{A}} \quad \left| i \right| := \left| \begin{array}{c} M_i \\ M_i \end{array}, \quad i = 1, \tau. \end{array} \right.$$
(5.2)

Here, the line can be either an edge or a tail. That is, \mathcal{G} transforms the local d.o.f. 1 and τ on edges (tails) as

$$\begin{vmatrix} i & \Longrightarrow & \sum_{a,b,y=1,\tau;x,z\in L_{M_i}} [P_i]_{xyz}^{ab} & a = \begin{vmatrix} z & b \\ y & b \\ x & x \end{vmatrix}, \quad i = 1, \tau.$$
(5.3)

Unitary transformation \mathcal{G} (5.2) does not preserve the original Hilbert space $\mathcal{H}_{\mathsf{Fibo}}$ of the string-net model but rather rotates it within the enlarged space \mathcal{H}^* . Nevertheless, the symmetry transformation does not alter the observables of the quantum system; we can only measure the anyons' charges in the basis comprising the original d.o.f.s 1 and τ on tails. Consequently, after the transformation \mathcal{G} , we must project the transformed states in \mathcal{H}^* back into $\mathcal{H}_{\mathsf{Fibo}}$ using a projector \mathcal{P} . Hence, in the doubled Fibonacci topological order, the symmetry transformation cannot simply be \mathcal{G} . Instead, it should be

$$\tilde{\mathcal{G}} := \mathcal{P}\mathcal{G}\mathcal{P}.$$
 (5.4)

As discussed in the Appendix D.3, transformation \mathcal{G} preserves anyon species but acts nontrivially on each anyon's enlarged internal space spanned by $\{1, \tau_1, \tau_2\}$ on the tail where the anyon resides. Different anyon species experience distinct actions, which blockdiagonalizes \mathcal{G} (5.2) in \mathcal{H}^* :

$$\mathcal{G} = \prod_{\text{Plaquettes}} \sum_{P \text{ Anyons } J} \mathcal{G}_P^J \Pi_P^J, \tag{5.5}$$

where Π_P^J is measures whether plaquette P has an anyon J (Π_P^J 's components are detailed in Appendix A), and \mathfrak{G}_P^J acts nontrivially only on the Hilbert subspace spanned by excited states with same anyon species in all plaquettes and varying charges of anyon J on the tail in plaquette P.

Consequently, transformation $\tilde{\mathcal{G}}$, projected from \mathcal{G} , is a gauge symmetry transformation that preserves anyon species and can also be block-diagonalized along with \mathcal{G} :

$$\tilde{\mathcal{G}} = \prod_{\text{Plaquettes}} \sum_{P \text{ Anyons } J} \tilde{\mathcal{G}}_P^J \Pi_P^J, \quad \tilde{\mathcal{G}}_P^J := \mathcal{P} \mathcal{G}_P^J \mathcal{P}.$$
(5.6)

1. The trivial anyon $1\overline{1}$ has only charge 1 that is invariant under the symmetry transformation:

$$\mathfrak{G}_{P}^{1\bar{1}} = \tilde{\mathfrak{G}}_{P}^{1\bar{1}} = \mathbb{1} , \qquad \mathfrak{G}_{P}^{1\bar{1}} \stackrel{\slashed{scalar}}{\underset{slashed{scalar}}{\overset{slashed{scalar}}}{\overset{slashed{scalar}}{\overset{slashed{scalar}}}{\overset{slashed{scalar}}{\overset{slashed{scalar}}}{\overset{slashed{scalar}}{\overset{slashed{scalar}}}{\overset{slashed{scalar}}{\overset{slashed{scalar}}}}{\overset{slashed{scalar}}}{\overset{slashed{scalar}}}{\overset{slashed{scalar}}}{\overset{slashed{scalar}}}{\overset{slashed{scalar}}}{\overset{slashed{scalar}}}{\overset{slashed{scalar}}}{\overset{slashed{scalar}}}{\overset{slashed{scalar}}}{\overset{slashed{scalar}}}{\overset{slashed{scalar}}}{$$

The ground states of the topological order are gauge invariant.

2. Anyon $\tau \bar{1}$ has only charge τ now embedded in the enlarged space spanned by $\{\tau_1, \tau_2\}$ as a superposition (4.4). The transformation is expressed in this enlarged space:

$$\begin{split} \mathfrak{G}_{P}^{\tau\bar{1}} &= \begin{pmatrix} \frac{\phi\sqrt{D}-1}{2\phi\sqrt{\phi}} & \frac{\sqrt{D}+\phi^{2}}{2\phi^{2}} \\ -\frac{\sqrt{D}+\phi^{2}}{2\phi\sqrt{\phi}} & \frac{\phi\sqrt{D}-1}{2\phi\sqrt{\phi}} \end{pmatrix}, \end{split}$$
(5.8)
$$\tilde{\mathfrak{G}}_{P}^{\tau\bar{1}} &= \begin{pmatrix} \frac{(\phi\sqrt{D}-1)(\sqrt{\phi}+1)}{4\phi^{2}} & \frac{1-\phi\sqrt{D}}{4\phi^{2}\sqrt{\phi}} \\ \frac{1-\phi\sqrt{D}}{4\phi^{2}\sqrt{\phi}} & \frac{(\phi\sqrt{D}-1)(\sqrt{\phi}-1)}{4\phi^{2}} \end{pmatrix}, \end{cases}$$
$$\mathfrak{G}_{P}^{\tau\bar{1}} &\stackrel{\stackrel{}{\gtrless}}{} \tau &= \stackrel{\stackrel{}{\$} \tau_{\tau\bar{1}}}{=} \frac{4\sqrt{5}-1}{2\sqrt{\phi}} \stackrel{\stackrel{}{\$} \tau_{1}}{=} -\frac{4\sqrt{5}+1}{2\sqrt{\phi}} \stackrel{\stackrel{}{\$} \tau_{2}}{=} , \end{cases}$$

Here, $D = \phi^2 + 1$. These matrices are 2×2 because anyon $\tau \bar{1}$ does not have charge 1. Unitary transformation \mathcal{G} (5.2) rotates the physical charge τ of anyon $\tau \bar{1}$ out of $\mathcal{H}_{\mathsf{Fibo}}$, necessitating the application of the projected transformation $\tilde{\mathcal{G}}_P^{\tau \bar{1}}$ to revert to $\mathcal{H}_{\mathsf{Fibo}}$ (Fig. 2b). Since $\det(\tilde{\mathcal{G}}_P^{\tau \bar{1}}) = 0$, matrix $\tilde{\mathcal{G}}_P^{\tau \bar{1}}$ is not invertible. Hence, symmetry transformation $\tilde{\mathcal{G}}$ (5.4) represents a noninvertible symmetry.

3. Anyon $1\bar{\tau}$ also has merely charge τ . Unitary transformation \mathcal{G} (5.2) rotates the physical charge τ (4.4) of the $1\bar{\tau}$ anyon also within the $\{\tau_1, \tau_2\}$ space:

$$\mathcal{G}_{P}^{1\bar{\tau}} = \begin{pmatrix} -\frac{\phi\sqrt{D}+1}{2\phi\sqrt{\phi}} & \frac{\phi\sqrt{\phi}-\sqrt[4]{5}}{2\phi\sqrt{\phi}} \\ \frac{4\sqrt{5}-\phi\sqrt{\phi}}{2\phi\sqrt{\phi}} & -\frac{\phi\sqrt{D}+1}{2\phi\sqrt{\phi}} \end{pmatrix},\tag{5.9}$$

as depicted in Fig. 2b. Matrix $\tilde{\mathcal{G}}_P^{1\bar{\tau}}$ is noninvertible.

4. Anyon $\tau \bar{\tau}$ has two gauge charges 1 and τ in the string-net model. Transformations \mathcal{G} are expressed in the enlarged 3-dimensional space spanned by 1, τ_1 , τ_2 :

$$\begin{split} \mathbf{g}_{P}^{\tau\bar{\tau}} &= \begin{pmatrix} \frac{1}{\phi^{2}} & \frac{\phi^{2}\sqrt{5}-\sqrt{\phi}}{2\phi^{4}} \sqrt{5} - \frac{\phi^{2}\sqrt{5}+\sqrt{\phi}}{2\phi^{4}} \sqrt{5} \\ \frac{\sqrt{5}+\phi\sqrt{D}}{2\phi^{2}} & \frac{1-\phi^{2}\sqrt{5\phi}}{2\phi^{4}} & \frac{\phi\sqrt{5}-2\sqrt{\phi}}{2\phi^{3}} \\ \frac{\sqrt{5}-\phi\sqrt{D}}{2\phi^{2}} & -\frac{\phi\sqrt{5}+2\sqrt{\phi}}{2\phi^{2}} & -\frac{1+\phi^{2}\sqrt{5\phi}}{2\phi^{2}\sqrt{\phi}} \\ \frac{\sqrt{5}+\phi\sqrt{D}}{2\phi^{2}} & -\frac{1+\sqrt{\phi}}{2\phi^{2}\sqrt{\phi}} & \frac{1}{2\phi^{3}\sqrt{\phi}} \\ \frac{\sqrt{5}-\phi\sqrt{D}}{2\phi^{2}} & -\frac{1+\sqrt{\phi}}{2\phi^{3}} & \frac{1}{2\phi^{3}\sqrt{\phi}} \\ \frac{\sqrt{5}-\phi\sqrt{D}}{2\phi^{2}} & \frac{1}{2\phi^{3}\sqrt{\phi}} & \frac{1-\sqrt{\phi}}{2\phi^{3}} \\ \frac{\sqrt{5}-\phi\sqrt{D}}{2\phi^{2}} & \frac{1}{2\phi^{3}\sqrt{\phi}} & \frac{1-\sqrt{\phi}}{2\phi^{3}} \\ \end{pmatrix}, \\ \mathbf{g}_{P}^{\tau\bar{\tau}} &= \mathbf{g}_{P}^{\tau\bar{\tau}} &= \frac{\sqrt{\phi^{2}+1}}{\phi^{2}} \quad \mathbf{g}_{1}^{\tau} & - \frac{\sqrt{\phi}}{\phi^{2}} \quad \mathbf{g}_{\tau}^{\tau} \\ \mathbf{g}_{P}^{\tau\bar{\tau}} &= \frac{\sqrt{\phi^{2}+1}}{\phi^{2}} \quad \mathbf{g}_{1}^{\tau} & - \frac{\sqrt{\phi}}{\phi^{3}} \quad \mathbf{g}_{\tau}^{\tau} \\ \mathbf{g}_{P}^{\tau\bar{\tau}} &= \frac{\sqrt{\phi^{2}+1}}{\phi^{2}} \quad \mathbf{g}_{1}^{\tau} & - \frac{\sqrt{\phi}}{\phi^{3}} \quad \mathbf{g}_{\tau}^{\tau} \\ \mathbf{g}_{P}^{\tau\bar{\tau}} &= \frac{\sqrt{\phi^{2}+1}}{\phi^{2}} \quad \mathbf{g}_{1}^{\tau} & - \frac{\sqrt{\phi}}{\phi^{3}} \quad \mathbf{g}_{\tau}^{\tau} \\ \mathbf{g}_{P}^{\tau\bar{\tau}} &= \frac{\sqrt{\phi^{2}+1}}{\phi^{2}} \quad \mathbf{g}_{1}^{\tau} & - \frac{\sqrt{\phi}}{\phi^{3}} \quad \mathbf{g}_{\tau}^{\tau} \\ \mathbf{g}_{P}^{\tau\bar{\tau}} &= \frac{\sqrt{\phi^{2}+1}}{\phi^{2}} \quad \mathbf{g}_{1}^{\tau} \\ \mathbf{g}_{P}^{\tau\bar{\tau}} \\ \mathbf{g}_{P}^{\tau\bar{\tau}} &= \frac{\sqrt{\phi^{2}+1}}{\phi^{2}} \quad \mathbf{g}_{1}^{\tau} \\ \mathbf{g}_{P}^{\tau\bar{\tau}} \\ \mathbf{g}_{P$$

See Fig. 2c. Although these transformations affect apparently only the d.o.f. on the tail in P, an excited state is transformed into superpositions of *excited states* with the same anyon species but different charges in plaquette P, leading to varying d.o.f. on plaquette P's edges.

Note that while the symmetry transformations are obtained in the string-net model, the symmetry is model-independent and intrinsic to the doubled Fibonacci topological order. We now reveal the symmetry structure.



Figure 2: (a) Physical d.o.f.s (depicted by black vectors) $1, \tau$ embedded in the space $\{1, \tau_1, \tau_2\}$. (b) Charges τ of anyon species $\tau \overline{1}, 1\overline{\tau}$ after transformations \mathcal{G} (depicted by blue vectors) and $\tilde{\mathcal{G}}$ (by orange vectors). (c) Charges $1, \tau$ of anyon $\tau\overline{\tau}$ after transformations \mathcal{G} (depicted by blue vectors) and $\tilde{\mathcal{G}}$ (by orange vectors).

6 Fibonacci Categorical Gauge Symmetry

We now show that the symmetry of the doubled Fibonacci topological order is a *categorical* gauge symmetry characterized by the Fibonacci fusion 2-category.

Recall that in a usual gauge theory with gauge group G, a gauge field configuration $g \in G$ is transformed to $g' = hgh^{-1} \in G$ by a gauge transformation characterized by $h \in G$. The gauge group G is both the gauge field's configuration space and the space of the gauge transformations. Now the question in our case is: What is the symmetry structure of the doubled Fibonacci topological order analogous to the gauge group G together with its symmetry transformation in a usual gauge theory?

Looking back to how \mathcal{G} transforms the fundamental d.o.f. as in Eq. (5.3), the two sides of (5.3) seem different in lattice structures, although the red lines on the RHS are auxiliary. But the two sides do have the same lattice structure even before annihilating the red lines for both mathematical and physical reasons. Mathematically, in Fibo, the simple objects 1 and τ are simple bimodules over the trivial Frobenius algebra $\mathcal{A}_0 = \mathbb{C}[1]$, so

Fibo =
$$\mathsf{Bimod}_{\mathsf{Fibo}}(\mathcal{A}_0)$$
.

Thus, an edge/tail labeled by simple object $i \in \{1, \tau\}$ in the original model must also carry two red lines:

$$1 \underbrace{|}_{i}^{i} \xrightarrow{1} \sum_{a,b,y=1,\tau} \sum_{x,z \in L_{M_{i}}} [P_{\tau}]_{xyz}^{ab} a_{x} \underbrace{|}_{x}^{b}; \qquad (6.1)$$

however, since these two red lines are labeled by the trivial object 1, they are omitted for simplicity. Physically, the simple objects of the input fusion category of a string-net model are the pure charges, defined concerning the trivial flux characterized by the Frobenius algebras. When Fibo is the input fusion category, the trivial flux is characterized by $\mathcal{A}_0 = \mathbb{C}[1]$. When $\mathsf{Bimod}_{\mathsf{Fibo}}(\mathcal{A})$ is the input fusion category, the trivial flux is characterized by \mathcal{A} (4.1). Hence, the RHS of (6.1), where red lines carry elements of \mathcal{A} , defines precisely how pure charges M_1 and M_{τ} appear in the string-net model with input fusion category $\mathsf{Bimod}_{\mathsf{Fibo}}(\mathcal{A})$. In conclusion, the lattice structures before and after transformation $\mathcal{G}(5.2)$ are the same. On the other hand, \mathcal{G} also retains the input fusion category Fibo because

 $\mathsf{Bimod}_{\mathsf{Fibo}}(\mathcal{A}) \subset \mathsf{Fibo}, \qquad \mathsf{Bimod}_{\mathsf{Fibo}}(\mathcal{A}) \cong \mathsf{Fibo}.$

It only transforms the model's d.o.f.s on edges and tails from simple objects $1, \tau$ to composite objects M_1, M_{τ} in Fibo.

Inspired by the above discussion, we find that a fusion 2-category, the **Fibonacci** fusion 2-category, exists to describe this gauge symmetry coherently. This 2-category consists of the following three ingredients⁶:

- 1. Objects are bimodule categories over Frobenius algebras in Fibo. These bimodule categories are subcategories of and isomorphic to Fibo. Two such objects are $Bimod_{Fibo}(\mathcal{A}_0) = Fibo$ and $Bimod_{Fibo}(\mathcal{A}) \subset Fibo$.
- 2. One-morphisms are isomorphism functors between objects. Two such 1-morphisms are the identity functor and $\mathcal{F}_{\mathcal{A}}$ defined in Eq. (5.1):

$$\operatorname{id}(1) = 1$$
, $\operatorname{id}(\tau) = \tau$, $\mathcal{F}_{\mathcal{A}}(1) = M_1$, $\mathcal{F}_{\mathcal{A}}(\tau) = M_{\tau}$.

3. Two-morphisms are natural transformations between 1-morphisms. We are particularly interested in the 2-morphism—the *composition* of 1-morphisms. Consider, for example, the composition of two 1-morphisms \$\mathcal{F}_{\mathcal{A}}\$ and \$\mathcal{F}_{\mathcal{B}}\$, where \$\mathcal{A}\$ and \$\mathcal{B}\$ are Frobenius algebras in Fibo. Since the bimodule category Bimod_{Fibo}(\$\mathcal{A}\$) is isomorphic to Fibo via the functor \$\mathcal{F}_{\mathcal{A}}\$, the object \$\mathcal{F}_{\mathcal{A}}\$(\$\mathcal{B}\$) becomes a Frobenius algebra in Bimod_{Fibo}(\$\mathcal{A}\$). This allows us to define the bimodule category over \$\mathcal{F}_{\mathcal{A}}\$(\$\mathcal{B}\$) within Bimod_{Fibo}(\$\mathcal{A}\$):

$$\mathsf{Bimod}_{\mathsf{Bimod}_{\mathsf{Fibo}}(\mathcal{A})}(\mathcal{F}_{\mathcal{A}}(\mathcal{B})).$$

This category is a subcategory of and is isomorphic to $\mathsf{Bimod}_{\mathsf{Fibo}}(\mathcal{A})$. Hence, it is a subcategory of Fibo and is also isomorphic to Fibo. In other words, fusion category $\mathsf{Bimod}_{\mathsf{Bimod}_{\mathsf{Fibo}}(\mathcal{A})}(\mathcal{F}_{\mathcal{A}}(\mathcal{B}))$ is an object in the Fibonacci fusion 2-category. The isomorphism from Fibo to $\mathsf{Bimod}_{\mathsf{Bimod}_{\mathsf{Fibo}}(\mathcal{A})}(\mathcal{F}_{\mathcal{A}}(\mathcal{B}))$ is interpreted as the composition of two functors:

$$\mathcal{F}_{\mathcal{F}_{\mathcal{A}}(\mathcal{B})} = \mathcal{F}_{\mathcal{B}} \circ \mathcal{F}_{\mathcal{A}}.$$

Unitary transformation $\mathcal{G}(5.2)$ is a representation of 1-morphism $\mathcal{F}_{\mathcal{A}}$ over the enlarged Hilbert space \mathcal{H}^* :

$$\rho(\mathrm{id}) = \mathbb{1}, \qquad \rho(\mathcal{F}_{\mathcal{A}}) = \mathcal{G}.$$

Composing gauge transformations here differs from composing transformations in a usual gauge theory. Each time transformation \mathcal{G} is applied, it is necessary to first enlarge the

⁶For a general definition and properties of fusion 2-categories, refer to Ref. [83].

Hilbert space. On the other hand, the symmetry transformation is the projection of the composition of \mathcal{G} back into $\mathcal{H}_{\mathsf{Fibo}}$ by projector \mathcal{P} , such as $\tilde{\mathcal{G}}$ (5.4).

Now, to answer the question raised at the beginning of this section, the Fibonacci fusion 2-category is analogous to a gauge field's *phase space* comprising the gauge field's configuration space and its conjugate momenta. In a usual (in particular a lattice) gauge theory, the gauge field's configuration space is the gauge group, which per se is the representation space of the gauge transformations—the adjoint representation of the gauge group. Consider gauge group U(1) as a simple example, the gauge transformation $d/d\theta$ is the conjugate momentum of $\exp(i\theta) \in U(1)$. In the gauge theory of the doubled Fibonacci topological order, the gauge field's configuration space is Fibo, and the gauge transformations are 1-morphisms of the Fibonacci fusion 2-category. Mathematically, the Fibonacci fusion 2-category serves as the "adjoint representation" of Fibo, with 1-morphisms acting on the representation space—Fibo, as all bimodule categories are subcategories of Fibo.

We refer to the gauge symmetry of the doubled Fibonacci topological order and the Fibonacci Turaev-Viro TQFT as the *Fibonacci 2-categorical gauge symmetry*. Each bimodule category in the Fibonacci fusion 2-category is a gauge choice, preserving anyon species. The possible internal charges of an anyon are the two simple objects in the chosen bimodule category. Since we chose the original input fusion category Fibo as our gauge, after the symmetry transformation, we must project the transformed states back into the original Hilbert space \mathcal{H}_{Fibo} , such that anyons' charges are measured in the basis comprising the original d.o.f. 1 and τ .

Acknowledgments

We thank Davide Gaiotto, Lukas Mueller, Ling-Yan Hung, Hongguang Liu, Yuting Hu, Chenjie Wang and Yinan Wang for inspiring and helpful discussions. YW is supported by the General Program of Science and Technology of Shanghai No. 21ZR1406700, and the Shanghai Municipal Science and Technology Major Project (Grant No. 2019SHZDZX01). The authors are grateful for the hospitality of the Perimeter Institute during his visit, where the main part of this work is done. This research was supported in part by the Perimeter Institute for Theoretical Physics. Research at Perimeter Institute is supported by the Government of Canada through the Department of Innovation, Science and Economic Development and by the Province of Ontario through the Ministry of Research, Innovation and Science.

A Review of the Extended String-net Model

In this section, we briefly review the string-net model defined in Ref. [26]. The string-net model is an exactly solvable Hamiltonian model defined on a 2-dimensional lattice. An example lattice is depicted in Fig. 1. All vertices are trivalent. Within each plaquette of the lattice, a tail is attached to an arbitrary edge of the plaquette, pointing inward. We will later demonstrate that different choices of the edge to which the tail is attached are

equivalent. In general cases, each edge and tail is oriented, and different choices of directions are equivalent. Nevertheless, for the case of the Fibonacci string-net model presented in the main body, different direction choices are the same, so we omit the directions of edges and tails in the main body.

The input data of the string-net model is a fusion category \mathscr{F} , described by a finite set $L_{\mathscr{F}}$, whose elements are called *simple objects*, equipped with three functions $N: L^3_{\mathscr{F}} \to \mathbb{N}$, $d: L_{\mathscr{F}} \to \mathbb{R}$, and $G: L^6_{\mathscr{F}} \to \mathbb{C}$. The function N sets the *fusion rules* of the simple objects, satisfying

$$\sum_{e \in L_{\mathscr{F}}} N^e_{ab} N^d_{ec} = \sum_{e \in L_{\mathscr{F}}} N^d_{ae} N^e_{bc}, \qquad \qquad N^c_{ab} = N^{b^*}_{c^*a}$$

There exists a special simple object $0 \in L_{\mathscr{F}}$, called the *trivial object*, such that for any $a, b \in L_{\mathscr{F}}$,

$$N_{0a}^b = N_{0b}^a = \delta_{ab},$$

where δ is the Kronecker symbol. For each $a \in L_{\mathscr{F}}$, there exists a unique simple object $a^* \in L_{\mathscr{F}}$, called the *opposite object* of a, such that

$$N^0_{ab} = N^0_{ba} = \delta_{ba^*}$$

We only consider the case where for any $a, b, c \in L_{\mathscr{F}}, N_{ab}^{c} = 0$ or 1. In this case, we define

$$\delta_{abc} = N_{ab}^{c^*} \in \{0, 1\}.$$

The basic configuration of the string-net model is established by labeling each edge and tail with a simple object in $L_{\mathscr{F}}$, subject to the constraint on all vertices that $\delta_{ijk} = 1$ for the three incident edges or tails meeting at this vertex, all pointing toward the vertex and respectively counterclockwise labeled by $i, j, k \in L_{\mathscr{F}}$. We can reverse the direction of any edge or tail and simultaneously conjugate its label as $j \to j^*$, which keeps the configuration invariant. The Hilbert space \mathcal{H} of the model is spanned by all possible configurations of these labels on the edges and tails.

The function d returns the quantum dimensions of the simple objects in $L_{\mathscr{F}}$. It is the largest eigenvalues of the fusion matrix and forms the 1-dimensional representation of the fusion rule.

$$d_a d_b = \sum_{c \in L_{\mathscr{F}}} N_{ab}^c d_c.$$

In particular, $d_1 = 1$, and for any $a \in L_{\mathscr{F}}, d_a = d_{a^*} \neq 0$.

The function G defines the 6j-symbols of the fusion algebra. It satisfies

$$\sum_{n} d_{n} G_{v^{*}u^{*}a}^{pqn} G_{j^{*}i^{*}b}^{ivn} G_{q^{*}p^{*}c}^{ijn} = G_{i^{*}pu^{*}}^{abc} G_{vq^{*}j}^{c^{*}b^{*}a^{*}}, \qquad \sum_{n} d_{n} G_{kln}^{ijp} G_{l^{*}k^{*}n}^{j^{*}i^{*}q} = \frac{\delta_{pq^{*}}}{d_{p}} \delta_{ijp} \delta_{klq},$$
$$G_{kln}^{ijm} = G_{ijn^{*}}^{klm^{*}} = G_{lkn^{*}}^{jim} = G_{nk^{*}l^{*}}^{mij} = \alpha_{m} \alpha_{n} \overline{G_{l^{*}k^{*}n^{*}}^{j^{*}i^{*}m^{*}}},$$

where $\alpha_a = \operatorname{sgn}(a)$.

The Hamiltonian of the string-net model reads

$$H := -\sum_{\text{Plaquettes } P} Q_P, \qquad Q_P := \frac{1}{D} \sum_{s \in L_{\mathscr{F}}} Q_P^s, \qquad D := \sum_{a \in L_{\mathscr{F}}} d_a^2, \qquad (A.1)$$

where the plaquette operator Q_P^s acts on edges surrounding plaquette P and has the following matrix elements on a hexagonal plaquette⁷:

$$Q_P^s \xrightarrow[e_6]{i_1, \dots, p}_{i_6, i_6, i_6} e_4 := \delta_{p,0} \sum_{j_k \in L_\mathscr{F}} \prod_{k=1}^6 \left(\sqrt{d_{i_k} d_{j_k}} \ G_{sj_{k+1}^* j_k}^{e_k i_k i_{k+1}^*} \right) \xrightarrow[e_6]{i_1, \dots, p}_{i_6, i_6} e_4$$

It turns out that

$$(Q_P^s)^{\dagger} = Q_P^{s^*}, \qquad Q_P^r Q_P^s = \sum_{t \in L_{\mathscr{F}}} N_{rs}^t Q_P^t, \qquad Q_P^2 = Q_P, \qquad Q_{P_1} Q_{P_2} = Q_{P_2} Q_{P_1}$$

The summands Q_P in Hamiltonian H are commuting projectors, so the Hamiltonian is exactly solvable. The ground-state subspace \mathcal{H}_0 of the system is the projection

$$\mathcal{H}_0 = \left[\prod_{\text{Plaquettes } P} Q_P\right] \mathcal{H}.$$
 (A.2)

If the lattice has the sphere topology, the model has a unique ground state $|\Phi\rangle$ up to scalars.

A.1 Topological Features

We briefly review the topological nature of the ground-state subspace of the string-net model defined in Ref. [26]. Any two lattices with the same topology can be transformed into each other by so-called *Pachner moves*. There are unitary linear maps between the Hilbert spaces of two string-net models with the same input fusion category on different lattices associated with these moves, denoted as \mathcal{T} . The ground states are invariant under such linear transformations. There are three kinds of elementary Pachner moves, whose

⁷We only show the actions of Q_P operator on a hexagonal plaquette. The matrix elements of Q_P operators on other types of plaquettes are defined similarly.

corresponding linear transformations are:

$$\mathfrak{T} \stackrel{a}{\longrightarrow} \stackrel{c}{\longrightarrow} \stackrel{c}{\longrightarrow} \stackrel{c}{\longrightarrow} = \sum_{n \in L_{\mathscr{F}}} \sqrt{d_m d_n} G_{kln}^{ijm} \stackrel{a}{\longrightarrow} \stackrel{n}{\longrightarrow} \stackrel{$$

Here we use red " \times " to mark the plaquettes to contract. Any other Pachner moves and their corresponding linear transformations of Hilbert spaces are compositions of these three elementary moves. Given initial and final lattices, there are multiple ways to compose these elementary moves, but different ways result in the same transformation matrices on the ground-state Hilbert space.

We have also noted that different selections of the edge to which the tail is attached are equivalent. These variations lead to distinct lattice configurations and, consequently, different Hilbert spaces for the lattice model. The equivalence of states in such Hilbert spaces is established by the following linear transformation \mathcal{T}' :

The states where tails attach to other edges can be obtained recursively in this manner.

For convenience, in certain instances, we will temporarily incorporate auxiliary states with multiple tails within a single plaquette. These states, despite having multiple tails in one plaquette, are all equivalent to states within the Hilbert space:



A.2 Excited States

An excited state $|\varphi\rangle$ of the string-net model is an eigenstate of the Hamiltonian such that $Q_P|\varphi\rangle = 0$ at some plaquettes P. In such a state, there are anyons in these plaquettes P.

We also refer to the ground states as trivial excited states, in which there are only *trivial anyons* in all plaquettes. We assume the sphere topology, in which the model has a unique ground state; nevertheless, the results in this section apply to other topologies.

We start with the simplest excited states with a pair of anyons in two *adjacent* plaquettes with a common edge E. This state can be generated by ribbon operator $W_E^{J;pq}$:

$$W_E^{J;pq} \stackrel{i_2}{\underset{e_n \quad i_n}{\overset{j}{\underset{i_n}{\underset{k \in L_{\mathscr{F}}}{\sqrt{\frac{d_k}{d_j}}}}}} := \sum_{\substack{k \in L_{\mathscr{F}}}{\sqrt{\frac{d_k}{d_j}}} \overline{z_{pqj}^{J;k}} \stackrel{e_1 \quad i_2}{\underset{p^* \quad \dots \quad q}{\underset{e_n \quad i_n}{\underset{i_n}{\underset{i_n}{\underset{k \in L_{\mathscr{F}}}{\sqrt{\frac{d_k}{d_j}}}}}}, \quad (A.6)$$

where j is the label on edge E, and \bar{z} is the complex conjugate. Here, the four-indexed tensor $z_{pqj}^{J;k}$ is called the *half-braiding tensor*, defined by the following equation:

$$\frac{\delta_{jt} N_{rs}^t}{d_t} z_{pqt}^{J;w} = \sum_{u,l,v \in L_{\mathscr{F}}} z_{lqr}^{J;v} z_{pls}^{J;u} \cdot d_u d_v G_{p^*wu^*}^{r^*s^*t} G_{qw^*v}^{srj^*} G_{rv^*w}^{s^*ul^*}.$$

The label J, called the *anyon species*, labels different minimal solutions of the z tensor that cannot be the sum of any other nonzero solutions. The ribbon operator $W_E^{J;pq}$ creates in the two adjacent plaquettes a pair of anyons J^* and J with charges p^* and q. An anyon species J may have multiple possible charges p, causing multiple excited states of the string-net model to represent the same anyon. Categorically, anyon species J are labeled by simple objects in the *center* of the input fusion category \mathscr{F} , a modular tensor category whose categorical data record all topological properties of the topological order that the string-net model describes:

$$J \in L_{\mathcal{Z}(\mathscr{F})}$$
.

States with two quasiparticles in two non-adjacent plaquettes are generated by ribbon operators along longer paths. These ribbon operators result from concatenating shorter ribbon operators. For example, to create two quasiparticles J^* and J with charges p_0^* and p_n in two non-adjacent plaquettes P_0 and P_n , we can choose a sequence of plaquettes (P_0, P_1, \dots, P_n) , where P_i and P_{i+1} are adjacent plaquettes with their common edge E_i . The ribbon operator $W_{P_0P_n}^{J;p_0p_n}$ is

$$W_{P_0P_n}^{J;p_0p_n} := \left[\sum_{p_1p_2\cdots p_{n-1}\in L_{\mathscr{F}}}\prod_{k=1}^{n-1} \left(d_{p_k}B_{P_k}W_{E_k}^{J;p_kp_{k+1}}\right)\right]W_{E_0}^{J;p_0p_1}.$$

Different choices of plaquette paths (P_0, P_1, \dots, P_n) give the same operator $W_{P_0P_n}^{J;p_0p_n}$ if these sequences can deform continuously from one to another. Following the same method, we can also define the creation operator of three or more anyons.

At the end of this section, we define the measurement operator Π_P^J measuring whether there is an anyon J excited in plaquette P:

$$\Pi_{P}^{J} \xrightarrow[i_{1}]{i_{1}} \underset{e_{6}}{\overset{i_{1}}{\underset{i_{5}}}}{\overset{i_{1}}{\underset{e_{5}}}} \overset{e_{3}}{\underset{e_{4}}{\underset{i_{5}}}} := \sum_{s,t \in L_{\mathscr{F}}} \frac{d_{s}d_{t}}{d_{p}} z_{pps}^{J;t} \xrightarrow[i_{1}]{i_{1}} \underset{i_{7}}{\overset{p}{\underset{p}{\underset{i_{5}}}}}{\overset{i_{1}}{\underset{i_{6}}}} ... (A.7)$$

The set of measurement operators are orthonormal and complete:

$$\Pi_P^J \Pi_P^K = \delta_{JK} \Pi_P^J, \qquad \sum_{J \in L_{\mathcal{Z}(\mathscr{F})}} \Pi_P^J = \mathbb{1}.$$

B Frobenius Algebras and Bimodules

It is a mathematical theorem [81] that two fusion categories \mathscr{F} and \mathscr{F}' have isomorphic centers if and only if they are *categorically Morita equivalent*. That is, two string-net models with categorically Morita equivalent input fusion categories describe the same topological order. Category theory also tells that if a fusion category \mathscr{F}' is categorically Morita equivalent to \mathscr{F} , there must be a *Frobenius algebra* \mathcal{A} in \mathscr{F} , such that \mathscr{F}' is isomorphic to the *bimodule category* over \mathcal{A} in \mathscr{F} :

$$\mathscr{F}' \cong \mathsf{Bimod}_{\mathscr{F}}(\mathcal{A}).$$
 (B.1)

Therefore, different string-net models describing the same topological order are classified by all Frobenius algebras \mathcal{A} in a particular input fusion category \mathscr{F} . Such equivalent models have bimodule categories $\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$ as their input fusion categories. We can establish the duality maps between these equivalent models. In this section, we briefly review the definition of Frobenius algebras in a given fusion category and their bimodules and leave the duality maps for the next section.

B.1 Frobenius Algebra

A Frobenius algebra \mathcal{A} in a fusion category \mathscr{F} is characterized by a pair of functions (n, f). Function $n: L_{\mathscr{F}} \to \mathbb{N}$ returns the *multiplicity* n_a of $a \in L_{\mathscr{F}}$ appearing in the Frobenius algebra \mathcal{A} , satisfying $n_a = n_{a^*}$. The simple objects of \mathcal{A} are labeled by pairs a_{α} , where $a \in L_{\mathscr{F}}$ satisfies $n_a > 0$, and $\alpha = 1, 2, \ldots, n_a$ is the *multiplicity index*. We denote the set of all simple objects in \mathcal{A} as $L_{\mathcal{A}}$.

The algebraic multiplication of \mathcal{A} is given by a function $f: L^3_{\mathcal{A}} \to \mathbb{C}$, satisfying:

$$\sum_{t_{\tau}\in L_{\mathcal{A}}} f_{r_{\rho}s_{\sigma}t_{\tau}} f_{a_{\alpha}b_{\beta}c_{\tau}^{*}} G_{abc}^{rst} \sqrt{d_{c}d_{t}} = \sum_{\gamma=1}^{n_{c}} f_{a_{\alpha}c_{\gamma}s_{\sigma}} f_{r_{\rho}c_{\gamma}^{*}b_{\beta}} ,$$

$$\sum_{a_{\alpha}b_{\beta}\in L_{\mathcal{A}}} f_{a_{\alpha}b_{\beta}c_{\gamma}} f_{b_{\beta}^{*}a_{\alpha}^{*}c_{\gamma}^{*}} \sqrt{d_{a}d_{b}} = d_{\mathcal{A}}\sqrt{d_{c}}, \qquad f_{a_{\alpha}b_{\beta}c_{\gamma}} = f_{b_{\beta}c_{\gamma}a_{\alpha}}, \qquad f_{0a_{\alpha}b_{\beta}} = \delta_{ab^{*}}\delta_{\alpha\beta},$$
(B.2)

where

$$d_{\mathcal{A}} := \sum_{a \in L_{\mathscr{F}}} n_a d_a \tag{B.3}$$

is the quantum dimension of the Frobenius algebra \mathcal{A} . This definition aligns with the one in the main body, where a Frobenius algebra \mathcal{A} is expressed as a vector space spanned by simple objects, and the algebraic multiplicity rule is given by function f:

$$\mathcal{A} = \mathbb{C}[L_{\mathcal{A}}], \qquad a_{\alpha}b_{\beta} = \sum_{c_{\gamma} \in L_{\mathcal{A}}} f_{a_{\alpha}b_{\beta}c_{\gamma}^{*}} c_{\gamma} \in \mathbb{C}[L_{\mathcal{A}}].$$

For convenience, in a lattice model, we use red edges or tails to indicate that this edge or tail is labeled by a simple object in Frobenius algebra \mathcal{A} , and a red dot on a vertex to represent a coefficient f multiplied to this state.

$$\begin{array}{c|c} & a_{\alpha} \\ & b_{\beta} \\ \hline & c_{\gamma} \end{array} \end{array} \right\rangle \qquad := \qquad f_{a_{\alpha}b_{\beta}c_{\gamma}} \\ & b_{\beta} \\ \hline & c_{\gamma} \end{array} \right\rangle. \tag{B.4}$$

We also use dashed red edges or tails to represent that we are summing over all states with labels on this edge in $L_{\mathcal{A}}$. The definition (B.2) of Frobenius algebra \mathcal{A} can then be illustrated graphically by the Pachner moves.



B.2 Bimodules over a Frobenius Algebra

A bimodule M over a Frobenius algebra \mathcal{A} in a fusion category \mathscr{F} is characterized by a pair of functions (n^M, P_M) . The function $n^M : L_{\mathscr{F}} \to \mathbb{N}$ returns the *multiplicity* n_a^M of $a \in L_{\mathscr{F}}$ appearing in bimodule M, satisfying $n_a^M = n_{a^*}^M$. The simple objects of M are labeled by pairs a_i , where $a \in L_{\mathscr{F}}$ satisfies $n_a^M > 0$, and $i = 1, 2, \ldots, n_a^M$ labels the multiplicity index. We denote the set of all simple objects in bimodule M as L_M .

The action of Frobenius algebra \mathcal{A} on bimodule M is characterized by function P_M : $L^2_{\mathcal{A}} \times L_M \times L_{\mathscr{F}} \times L_M \to \mathbb{C}$, satisfying the following defining equations:

$$\sum_{uv \in L_{\mathscr{F}}} \sum_{y_{\upsilon} \in L_{M}} [P_{M}]_{x_{\chi}uy_{\upsilon}}^{a_{\alpha}r_{\rho}} [P_{M}]_{y_{\upsilon}vz_{\zeta}}^{b_{\beta}s_{\sigma}} G_{urw}^{v^{*}by} G_{axc}^{w^{*}bu} G_{wrt^{*}}^{sz^{*}\upsilon} \sqrt{d_{u}d_{\upsilon}d_{w}d_{w}d_{y}d_{c}d_{t}}$$

$$= \sum_{\gamma=1}^{n_{c}} \sum_{\tau=1}^{n_{t}} P_{x_{\chi}wz_{\zeta}}^{c_{\gamma}t_{\tau}} f_{a_{\alpha}c_{\gamma}^{*}b_{\beta}} f_{r_{\rho}s_{\sigma}t_{\tau}},$$

$$[P_{M}]_{x_{\chi}yz_{\zeta}}^{00} = \delta_{xy}\delta_{yz}\delta_{\chi\upsilon}\delta_{\upsilon\zeta}, \qquad [P_{M}]_{x_{\chi}yz_{\zeta}}^{a_{\alpha}b_{\beta}} = [P_{M}]_{z_{\zeta}^{*}y^{*}x_{\chi}^{*}}^{b_{\beta}a_{\alpha}}.$$
(B.5)

This definition aligns with the one in the main body, where a bimodule M is expressed as a vector space spanned by simple objects $M = \mathbb{C}[L_M]$. A pair of Frobenius algebra elements $(a_{\alpha}, b_{\beta}) \in \mathbb{C}[L_A]^2$ is represented as a three-index tensor P_M on the bimodule space $\mathbb{C}[L_M]$.

For convenience, in a lattice model, we use a blue line to indicate that this line is labeled by a simple object in bimodule M and a wavy blue line to represent summing over all states with labels on this edge in $L_{\mathscr{F}}$ with coefficient P_M :

$$a_{\alpha} \xrightarrow{z_{\zeta}} M \xrightarrow{b_{\beta}} \dots = \sum_{y \in L_{\mathscr{F}}} [P_M]^{a_{\alpha}b_{\beta}}_{x_{\chi}yz_{\zeta}} \qquad a_{\alpha} \xrightarrow{z_{\zeta}} b_{\beta} \xrightarrow{z_{\zeta}} \dots \quad (B.6)$$

The definition (B.5) of bimodule M can then be depicted graphically by Pachner moves:



B.3 The Bimodule Fusion Category over a Frobenius Algebra

The set of all bimodules over a given Frobenius algebra \mathcal{A} in a fusion category \mathscr{F} forms a fusion category, denoted as $\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$. In this section, we briefly introduce the categorical data of the fusion category $\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$.

1. A bimodule M in $Bimod_{\mathscr{F}}(\mathcal{A})$ is *simple* if it cannot be written as a direct sum of two other bimodules. That is, we cannot find two bimodules M_1 and M_2 such that:

$$n_{a}^{M} = n_{a}^{M_{1}} + n_{a}^{M_{2}}, \quad [P_{M}]_{x_{\chi}yz_{\zeta}}^{a_{\alpha}b_{\beta}} = \begin{cases} [P_{M_{1}}]_{x_{\chi}yz_{\zeta}}^{a_{\alpha}b_{\beta}}, & (\chi \leq n_{x}^{M_{1}}, \zeta \leq n_{z}^{M_{1}}), \\ [P_{M_{2}}]_{x_{(\chi-n_{x}^{M_{1}})}^{a_{\alpha}b_{\beta}}yz_{(\zeta-n_{z}^{M_{2}})}, & (\chi > n_{x}^{M_{1}}, \zeta > n_{z}^{M_{1}}), \\ 0. & (\text{otherwise}). \end{cases}$$

2. The quantum dimension of a bimodule M in $\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$ is

$$d_M = \frac{1}{d_{\mathcal{A}}} \sum_{a \in L_{\mathscr{F}}} n_a^M d_a.$$
(B.7)

3. For any three bimodules M_1 , M_2 , and M_3 , we can represent their fusion rules in terms of their simple objects. Define the matrix $[\Delta_{M_1M_2M_3}]$ that represents how the

basis elements in the bimodule spaces are connected when the three bimodules fuse:

$$\begin{split} [\Delta_{M_1M_2M_3}]^{x_{\chi}y_{\upsilon}z_{\zeta}}_{r_{\rho}s_{\sigma}t_{\tau}} := & \frac{1}{d_{\mathcal{A}}^3} \sum_{a_{\alpha}b_{\beta}c_{\gamma} \in L_{\mathcal{A}}} \sum_{p \in L_{\mathscr{F}}} \sum_{u_{\rho} \in L_{M_1}} \sum_{v_{\sigma} \in L_{M_2}} \sum_{w_{\lambda} \in L_{M_3}} [P_1]^{b_{\beta}c_{\gamma}^*}_{x_{\chi}ur_{\rho}} [P_2]^{c_{\gamma}a_{\alpha}^*}_{y_{\upsilon}vs_{\sigma}} \times \\ & [P_3]^{a_{\alpha}b_{\beta}^*}_{z_{\zeta}wt_{\tau}} G^{bxu^*}_{c^*r^*p} G^{swp}_{br^*t^*} G^{pvz}_{aw^*s^*} G^{xyz}_{vpc} \sqrt{d_ud_vd_wd_ad_bd_cd_rd_sd_t} d_p . \end{split}$$

$$(B.8)$$

This definition can be depicted graphically:



The fusion rule of three bimodules M_1, M_2, M_3 is

$$\delta_{M_1 M_2 M_3} = \text{Tr}[\Delta_{M_1 M_2 M_3}]. \tag{B.9}$$

4. The Frobenius algebra \mathcal{A} itself is the trivial bimodule M_0 over \mathcal{A} :

$$L_{M_0} = L_{\mathcal{A}}, \qquad [P_{M_0}]_{x_{\chi}yz_{\zeta}}^{a_{\alpha}b_{\beta}} = \sum_{\nu=1}^{n_y} f_{a_{\alpha}x_{\chi}y_{\nu}^*} f_{y_{\nu}b_{\beta}z_{\zeta}^*}.$$
 (B.10)

Given a bimodule M, its opposite bimodule M^* is

$$L_{M^*} = L_M, \qquad [P_{M^*}]_{x_{\chi}yz_{\zeta}}^{a_{\alpha}b_{\beta}} = ([P_M]_{x_{\chi}yz_{\zeta}}^{a_{\alpha}b_{\beta}})^*.$$
(B.11)

5. The bimodule conditions induces that matrix $\Delta_{M_1M_2M_3}$ is a projector:

$$\Delta_{M_1M_2M_3}^2 = \Delta_{M_1M_2M_3}.$$

If $\delta_{M_1M_2M_3} \neq 0$, we can find the normalized eigenvectors $\mathcal{V}_{M_1M_2M_3}^{r_\rho s_\sigma t_\tau} \in \mathbb{C}$, such that

$$\sum_{r_{\rho}\in L_{M_{1}}}\sum_{s_{\sigma}\in L_{M_{2}}}\sum_{t_{\tau}\in L_{M_{3}}} [\Delta_{M_{1}M_{2}M_{3}}]_{r_{\rho}s_{\sigma}t_{\tau}}^{x_{\chi}y_{\upsilon}z_{\zeta}} \mathcal{V}_{M_{1}M_{2}M_{3}}^{r_{\rho}s_{\sigma}t_{\tau}} = \mathcal{V}_{M_{1}M_{2}M_{3}}^{x_{y}y_{j}z_{k}},$$

$$\sum_{x_{\chi}\in L_{M_{1}}}\sum_{y_{\upsilon}\in L_{M_{2}}}\sum_{z_{\zeta}\in L_{M_{3}}} |\mathcal{V}_{M_{1}M_{2}M_{3}}^{x_{\chi}y_{\upsilon}z_{\zeta}}|^{2} \sqrt{d_{x}d_{y}d_{z}} = d_{\mathcal{A}}^{2} \sqrt{d_{M_{1}}d_{M_{2}}d_{M_{3}}}.$$
(B.12)

For convenience, in a lattice model, we use blue lines labeled by a bimodule M to represent summing over all states with labels in L_M on this line. Additionally, blue dots represent a state multiplied by a coefficient \mathcal{V} :

$$\begin{vmatrix} M_1 \\ M_2 \\ M_3 \end{vmatrix} \qquad := \sum_{x_{\chi} \in L_{M_1}} \sum_{y_{\upsilon} \in L_{M_2}} \sum_{z_{\zeta} \in L_{M_3}} \mathcal{V}_{M_1 M_2 M_3}^{x_{\chi} y_{\upsilon} z_{\zeta}} \end{vmatrix} _{y_{\upsilon}} \qquad (B.13)$$

Such a state is invariant under $\mathcal{D}_{M_1M_2M_3}$ matrix. The 6j-symbol of $\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$ category is

$$G_{M_{3}M_{4}M'}^{M_{1}M_{2}M} = \frac{1}{d_{\mathcal{A}}^{3}\sqrt{d_{M_{1}}d_{M_{2}}d_{M_{3}}d_{M_{4}}d_{M}d_{M'}}} \mathcal{T} \xrightarrow{M_{1}}_{M_{2}} \times M_{3} \times M' \quad (B.14)$$

C General Constructions of Dualities and Symmetry Transformations in the Extended String-Net Model

Given a fusion category \mathscr{F} and a Frobenius algebra $\mathcal{A} \in \mathscr{F}$, two string-net models with \mathscr{F} and $\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$ as the input data describe the same topological order. Categorically, $\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$ is defined by an injective functor

$$\mathcal{D}: \mathsf{Bimod}_{\mathscr{F}}(\mathcal{A}) \to \mathscr{F}, \qquad M \mapsto \bigoplus_{a \in L_{\mathscr{F}}} n_a^M a, \tag{C.1}$$

and for any morphisms $\phi_{M_1M_2}^{M_3} \in \mathsf{Bimod}_{\mathscr{F}}(\mathcal{A}) : M_1 \otimes M_2 \to M_3 \text{ and } \varphi_{xy}^z \in \mathscr{F} : x \otimes y \to z,$

$$\mathcal{D}(\phi_{M_1M_2}^{M_3}) = \bigoplus_{z_{\zeta} \in L_{M_3}} \bigg[\bigoplus_{x_{\chi} \in L_{M_1}} \bigoplus_{y_{\upsilon} \in L_{M_2}} \mathcal{V}_{M_1M_2M_3^*}^{x_{\chi}y_{\upsilon}z_{\zeta}^*} \varphi_{x_{\chi}y_{\upsilon}}^{z_{\zeta}} \bigg],$$

where x_{χ} , y_{v} , and z_{ζ} are respectively the χ -th x object, v-th y, and ζ -th z in the direct sum $\mathcal{D}(M)$.

Such a functor \mathcal{D} induces a duality between the two models with \mathcal{F} and $\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$ as the input data:

$$M \implies \sum_{a_{\alpha}, b_{\beta} \in L_{\mathcal{A}}} \sum_{x_{\chi}, z_{\zeta} \in L_{M}} a_{\alpha} \stackrel{z_{\zeta}}{\longrightarrow} b_{\beta} .$$
(C.2)

This duality induces a unitary morphism between the Hilbert spaces $\mathcal{H}_{\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})}$ and $\mathcal{H}_{\mathscr{F}}$ of these two models. Such a unitary linear transformation can be understood plaquette by plaquette:

$$\begin{array}{c} & \stackrel{N_2}{\longrightarrow} & \stackrel{N_2}{\longrightarrow} & \stackrel{N_3}{\longrightarrow} & \stackrel{M_4}{\longrightarrow} & \stackrel{N_3}{\longrightarrow} & \stackrel{M_4}{\longrightarrow} & \stackrel{N_3}{\longrightarrow} & \stackrel{M_4}{\longrightarrow} & \stackrel{N_3}{\longrightarrow} & \stackrel{N_4}{\longrightarrow} & \stackrel{N_2}{\longrightarrow} & \stackrel{N_2}{\longrightarrow} & \stackrel{N_2}{\longrightarrow} & \stackrel{N_3}{\longrightarrow} & \stackrel{N_3}{\longrightarrow$$



Note that the black edges and tails labeled by $M_i, N_i \in \mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$ represent basis states in the dual model, where $\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$ is the input fusion category and M_i, N_i are simple objects. In contrast, the blue edges and tails labeled by $M_i, N_i \in \mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$ represent superposition states in the original model with \mathscr{F} as the input fusion category, where the superpositions are defined in Eqs. (B.6) and (B.13).

This understanding differs slightly from the unitary duality transformation in the sense that, after the above transformation (C.3) is applied to all plaquettes, the resulting basis state $|\psi\rangle$ satisfies

$$\langle \psi | \psi \rangle = d_{\mathcal{A}}^{g-2}, \tag{C.4}$$

where g is the genus of the surface on which the lattice is embedded. By applying the duality map and normalizing the resulting basis states, we obtain a unitary morphism between the two string-net models.

After the topological moves in Eq. (C.3), the degree of freedom on any edge will cease to have any multiplicity index of simple objects in bimodules, while that on any tail will still have a multiplicity index. Therefore, to make sense of this duality and make it unitary, we are urged to enlarge the Hilbert space of the original Fibonacci string-net model on each tail but not on the edges, such that two simple objects $a_{\alpha}, a_{\beta} \in L_M$ with different multiplicity indices $\alpha \neq \beta$ are distinguishable on tails.

C.1 Enlarging the Hilbert Space

In the enlarged Hilbert space, each tail carries a degree of freedom labeled by a pair a_{α} , where

$$a \in L_{\mathscr{F}}, \qquad \alpha = 1, 2, \cdots, N_a^{\mathcal{A}}, \qquad N_a^{\mathcal{A}} = \max_{M \in L_{\operatorname{Bimod}}_{\mathscr{F}}(\mathcal{A})} \{n_a^M\},$$
 (C.5)

where $L_{\mathsf{Bimod}_{\mathcal{F}}(\mathcal{A})}$ is the set of all simple bimodules over Frobenius algebra \mathcal{A} . But the basic degrees of freedom on edges remain to take value varying the simple objects of the input fusion category \mathscr{F} . The Hilbert space on the tail is spanned by all enlarged degrees of

freedom on tails and original degrees of freedom on edges, subject to the fusion rules on all vertices.

For any bimodule M, its simple object $x_{\chi}^M \in L_M$ corresponds to a superposition state $|x_{\chi}^M\rangle$ in the local Hilbert space of a tail:

$$\left|x_{\chi}^{M}\right\rangle := \sum_{i=1}^{N_{x}^{\mathcal{A}}} A_{\chi,i}^{x,M} |x_{i}\rangle.$$
(C.6)

All different states should satisfy the orthonormal conditions:

$$\sum_{\alpha=1}^{n_x^M} \sum_{\beta=1}^{n_x^N} M_1 \left(\times \begin{array}{c} y_{\beta}^N \\ \times \\ x_{\alpha}^M \end{array} \right) M_2 = d_{\mathcal{A}} n_x^M \delta_{xy} \delta_{M_1 M_2 M_3} \delta_{MN} \sqrt{d_{M_1} d_{M_2} d_x} .$$
(C.7)

C.2 Duality

The duality map (C.3) can be simplified. We can represent the unitary duality map vertex by vertex:

$$\mathcal{D} := \frac{1}{d_{\mathcal{A}}^{N_P - 1 + \frac{g}{2}}} \prod_{\text{Edge } e} E_e \prod_{\text{Vertex } v} \mathcal{D}_v, \qquad (C.8)$$

where N_P is the number of plaquettes in the lattice, and \mathcal{D}_v acts on vertex v:

$$\mathcal{D}_{v} \qquad M_{2} \qquad M_{3} \qquad := \qquad M_{2} \qquad M_{3} \qquad = \qquad \sum_{x_{\chi} \in L_{M_{1}}} \sum_{y_{\upsilon} \in L_{M_{2}}} \sum_{z_{\zeta} \in L_{M_{3}}} \mathcal{V}_{M_{1}M_{2}M_{3}}^{x_{\chi}y_{\upsilon}z_{\zeta}} \qquad x_{\chi} \qquad (C.9)$$

Note that each edge connects two vertices that are acted upon by two \mathcal{D}_v operators independently. Nevertheless, an edge e can only carry one label. We use E_e to ensure this uniqueness:

$$E_{e} \quad v_{1} \xrightarrow{x_{\alpha}^{M_{1}} \quad y_{\beta}^{M_{2}}} v_{2} \qquad := \qquad \sum_{i} (A_{\alpha,i}^{x,M_{1}})^{*} A_{\beta,i}^{x,M_{2}} \xrightarrow{x} \quad . \tag{C.10}$$

The E_e moves erase the multiplicity indices of labels on edges. But the multiplicity labels on tails are retained.

The Hilbert space is not preserved under the duality map:

$$\mathcal{H}_{\mathscr{F}} \neq \mathcal{D}\mathcal{H}_{\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})},$$

where $\mathcal{H}_{\mathscr{F}}$ and $\mathcal{H}_{\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})}$ are the Hilbert spaces of the string-net model with input fusion category \mathscr{F} and $\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$, respectively, considered as subspaces of the enlarged Hilbert space. Nevertheless, since the two models describe the same topological order, the groundstate subspace \mathcal{H}_0 is preserved under the duality map:

$$\mathcal{H}_{0,\mathscr{F}} = \mathcal{H}_{0,\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})},\tag{C.11}$$

where $\mathcal{H}_{0,\mathscr{F}}$ and $\mathcal{H}_{0,\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})}$ are the ground-state subspaces of the string-net model with input fusion category \mathscr{F} and $\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$, respectively.

C.3 Symmetry Transformation

In particular cases, \mathscr{F} and $\operatorname{Bimod}_{\mathscr{F}}(\mathcal{A})$ are isomorphic fusion category. That is, there exists an isomorphic functor $\mathcal{F}_{\mathcal{A}}$ that maps simple objects of \mathscr{F} to simple objects in $\operatorname{Bimod}_{\mathscr{F}}(\mathcal{A})$:

$$\varphi_{\mathcal{A}}(a) = M_a \in \mathsf{Bimod}_{\mathscr{F}}(\mathcal{A}). \tag{C.12}$$

Such isomorphic functor induces a linear isomorphism φ_A between the Hilbert space of these two models that maps basic degrees of freedom on edges and tails to basic degrees of freedom:

$$\varphi_{\mathcal{A}} : \mathcal{H}_{\mathscr{F}} \to \mathcal{H}_{\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})}, \qquad \qquad \varphi_{\mathcal{A}} \quad a := \qquad M_a \quad . \tag{C.13}$$

The composition

$$\mathcal{G} := \mathcal{D} \circ i : \mathcal{H}_{\mathscr{F}} \to \mathcal{H}_{\mathscr{F}} \tag{C.14}$$

is just a unitary transformation of the same model with \mathscr{F} as the input fusion category, and the symmetry transformation is the composition of the unitary transformation and projection back into the original degrees of freedom. The set of all symmetry transformations in the Hilbert space of the string-net model with \mathscr{F} as the input fusion category forms the symmetry of the topological order.

In particular, consider the trivial Frobenius algebra \mathcal{A}_0 :

$$L_{\mathcal{A}_0} = \{0\}, \qquad f_{000} = 1, \tag{C.15}$$

whose simple objects are classified by simple objects in $L_{\mathscr{F}}$:

$$L_{M_a} = \{a\}, \qquad P_{aaa}^{00} = 1. \tag{C.16}$$

The gauge transformation induced by Frobenius algebra \mathcal{A}_0 is the identity transformation of the string-net model.

C.4 Braiding of Bimodules

The input fusion category \mathscr{F} is a fusion category that lacks a braiding structure for exchanging two simple objects a and b. Nevertheless, the braiding of the trivial object 0 with any other simple object a always exists as the trivial braiding, which can be represented graphically as:



The last equality holds because the fusion of 0 with any simple object a is also trivial.



Figure 3: Contrast a plaquette with a tail labeled by the trivial bimodule \mathcal{A} in the original model.

Similarly, in the bimodule category $\mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$, the trivial bimodule \mathcal{A} braids trivially with any other bimodule $M \in \mathsf{Bimod}_{\mathscr{F}}(\mathcal{A})$, based on the definition of bimodules:



As a practical example, consider the situation where we contrast a plaquette with a tail labeled by the trivial bimodule \mathcal{A} in the original model. Since the Pachner moves (A.3) can only contrast plaquettes without nontrivial tails inside them, one must first "pull" the trivial tail out of the plaquette and then annihilate the plaquettes, as shown in Fig. 3.

D The \mathbb{Z}_2 Global Symmetry transformation of the \mathbb{Z}_2 Toric Code String-Net Model

To provide another example of our symmetry transformation construction, and to contrast with the Fibonacci categorical gauge symmetry discussed in the main text, we briefly apply our method in this section to construct a well-known symmetry transformation: the \mathbb{Z}_2 global symmetry transformation of the \mathbb{Z}_2 toric code string-net model[6]. In the toric code case, we do not need to enlarge the Hilbert space. This transformation exchanges the charge and flux excitations.

D.1 The toric code string-net model

The input fusion category of the toric code string-net model is the \mathbb{Z}_2 fusion category, which uses the two group elements $\pm 1 \in \mathbb{Z}_2$ as its simple objects. The fusion rules capture the group multiplicity rules of \mathbb{Z}_2 :

$$\delta_{ijk} = \frac{ijk+1}{2},$$

and the quantum dimensions are $d_{\pm} = 1$. The nonzero 6*j* symbols are given by

$$G_{kln}^{ijm} = \delta_{ijm} \delta_{klm} \delta_{iln} \delta_{jkn}.$$

In the toric code string-net model, each edge or tail on the lattice carries a group element $\pm 1 \in \mathbb{Z}_2$, and the degrees of freedom i, j, k on any three edges or tails meeting at a vertex must satisfy the fusion rule $\delta_{ijk} = 1$. The Hamiltonian of the toric code string-net model is a sum of commutative projectors A_P and B_P ,

$$H_{\rm TC} := -\sum_{\rm Plaquettes \ P} (A_P + B_P), \tag{D.1}$$

where A_P acts on tails in plaquettes P, and B_P acts on edges surrounding plaquettes P:

$$A_{P} \xrightarrow{i_{0}}_{i_{6}} \xrightarrow{i_{1}}_{i_{5}} \xrightarrow{i_{4}}_{i_{6}} e_{4} := \frac{p+1}{2} \xrightarrow{i_{1}}_{i_{6}} \xrightarrow{i_{1}}_{i_{6}} \xrightarrow{i_{1}}_{i_{6}} e_{4} , \qquad (D.2)$$

$$B_{P} \xrightarrow{i_{0}}_{i_{6}} \underbrace{i_{1}}_{e_{6}} \underbrace{i_{2}}_{e_{5}} \underbrace{e_{3}}_{i_{4}} \underbrace{i_{2}}_{e_{4}} e_{4} := \frac{1}{2} \xrightarrow{i_{1}}_{i_{0}} \underbrace{e_{1}}_{i_{6}} \underbrace{i_{2}}_{e_{5}} \underbrace{e_{3}}_{i_{4}} \underbrace{i_{2}}_{e_{4}} \underbrace{e_{1}}_{e_{5}} \underbrace{i_{4}}_{e_{6}} \underbrace{e_{1}}_{e_{6}} \underbrace{i_{2}}_{e_{5}} \underbrace{e_{2}}_{e_{5}} \underbrace{e_{1}}_{e_{6}} \underbrace{e_{2}}_{e_{6}} \underbrace{e_{1}}_{e_{1}} \underbrace{e_{2}}_{e_{1}} \underbrace{e_{1}}_{e_{1}} \underbrace{e_{2}}_{e_{1}} \underbrace{e_{1}}_{e_{2}} \underbrace{e_{2}}_{e_{6}} \underbrace{e_{1}}_{e_{6}} \underbrace{e_{2}}_{e_{6}} \underbrace{e_{2}}_{e_{6}} \underbrace{e_{1}}_{e_{6}} \underbrace{e_{2}}_{e_{6}} \underbrace{e_{2}}_{e_{6}} \underbrace{e_{1}}_{e_{6}} \underbrace{e_{2}}_{e_{6}} \underbrace{e_{2}}_{e_{6}$$

The ground states are common eigenstates of all A_P and B_P operators with +1 eigenvalues, while an excited state $|\psi\rangle$ is another common eigenstate satisfying $A_P|\psi\rangle = 0$ (or $B_P|\psi\rangle =$ 0) for one or more plaquettes P, in each of which there resides a chargeon e (or a fluxon m). Unlike the original version of the string-net model, where chargeons reside on vertices, both chargeons and fluxons are situated in the plaquettes of our enlarged model. If $A_P|\psi\rangle =$ $B_P|\psi\rangle = 0$ in plaquette P, there is a dyon ϵ in plaquette P. We also refer to the ground state as the trivial excited state, in which there are trivial anyons 1 in all plaquettes.

D.2 The global symmetry transformation of the toric code topological order

The \mathbb{Z}_2 fusion category has a unique nontrivial Frobenius algebra \mathcal{A} , such that

$$L_{\mathcal{A}} = \{1, -1\}, \qquad f_{ijk} = \delta_{ijk},$$

There are two simple bimodules over \mathcal{A} , denoted as M_{\pm} , such that

$$L_{M_{+}} = L_{M_{-}} = \{0, 1\}, \qquad [P_{+}]_{xyz}^{ab} = \delta_{axy}\delta_{byz},$$
$$[P_{-}]_{+++}^{++} = [P_{-}]_{---}^{+-} = 1, \qquad [P_{-}]_{+-+}^{--} = [P_{-}]_{-+-}^{--} = -1,$$
$$[P_{-}]_{++-}^{+-} = [P_{-}]_{-++}^{-+} = i, \qquad [P_{-}]_{--+}^{+-} = [P_{-}]_{+--}^{-+} = -i.$$

The nonzero vertex coefficients are

$$\mathcal{V}_{M_{+}M_{+}M_{+}}^{ijk} = \delta_{ijk}, \quad \mathcal{V}_{M_{+}M_{-}M_{-}}^{+++} = \mathcal{V}_{M_{+}M_{-}M_{-}}^{-++} = 1, \quad \mathcal{V}_{M_{+}M_{-}M_{-}}^{+-+} = i, \quad \mathcal{V}_{M_{+}M_{-}M_{-}}^{++-} = -i.$$

The fusion category $\mathsf{Bimod}_{\mathbb{Z}_2}(\mathcal{A})$ is isomorphic to the \mathbb{Z}_2 fusion category:

$$\delta_{M_i M_j M_k} = \delta_{ijk} = 1, \qquad d_{M_i} = d_i, \qquad G_{M_k M_l M_n}^{M_i M_j M_m} = G_{kln}^{ijm}$$

By applying the above categorical data, we can formulate a symmetry transformation \mathcal{G} is

$$\mathcal{G} := \frac{1}{2^{N_{\mathrm{P}}-1+\frac{g}{2}}} \prod_{\text{Vertices } v} \mathcal{G}_{v},\tag{D.4}$$

where N_P is the number of plaquettes in the lattice, and g is the genus number of the surface in which the lattice is embedded. Each local transformation \mathcal{G}_v acts on vertex v of the lattice as

where + (-) refers to 1 (-1), and the three lines meeting at vertex v can be either edges or tails.

The symmetry transformation \mathcal{G} is a \mathbb{Z}_2 global symmetry transformation of the toric code string-net model because:

1. G is a unitary \mathbb{Z}_2 transformation:

$$\mathfrak{G}^{\dagger} = \mathfrak{G}^{-1} = \mathfrak{G}. \tag{D.6}$$

2. \mathcal{G} preserves the model's Hamiltonian H_{TC} :

$$\mathcal{G}^{\dagger}H_{\mathrm{TC}}\mathcal{G} = H_{\mathrm{TC}}.\tag{D.7}$$

3. Transformation \mathcal{G} preserves the ground-state Hilber space of the model but exchanges A_P and B_P operators:

$$\mathcal{G}^{\dagger}A_P\mathcal{G} = B_P, \qquad \qquad \mathcal{G}^{\dagger}B_P\mathcal{G} = A_P, \qquad (D.8)$$

and hence exchanges chargeons and fluxons because an $A_P(B_p)$ measures the chargeon (fluxon) in plaquette P^8 .

Note that the global symmetry transformation \mathcal{G} maps the degrees of freedom 0 and 1 the simple objects in the \mathbb{Z}_2 fusion category—to the degrees of freedom M_0 and M_1 , which are the simple objects in the bimodule category $\mathsf{Bimod}_{\mathbb{Z}_2}(\mathcal{A})$. Analogous to our discussion about the Fibonacci categorical gauge symmetry, this EM-exchange global symmetry of the toric code topological order is fundamentally a *categorical symmetry* described by a fusion 2-category—the \mathbb{Z}_2 fusion 2-category defined based on the \mathbb{Z}_2 fusion category. It reduces to a \mathbb{Z}_2 group symmetry because the transformation is invertible.

⁸In the original string-net model, chargeons reside on vertices and fluxons are located in plaquettes, necessitating lattice dualization after the symmetry transformation. In contrast, our model places both chargeons and fluxons in plaquettes, eliminating the need to alter the lattice shape.

D.3 Criterion for Distinguishing Gauge Symmetries from Global Symmetries

The symmetry transformation (D.4) of the \mathbb{Z}_2 toric code topological order is a global symmetry transformation that exchanges the anyon species of chargeons and fluxons. In contrast, the symmetry transformation of the doubled Fibonacci topological order in the main text is a gauge symmetry that preserves all anyon species but only transforms excited states within the internal Hilbert space of each anyon. To determine whether a symmetry transformation defined by a Frobenius algebra \mathcal{A} in a fusion category \mathscr{F} is a global symmetry or a gauge symmetry, we use the criterion based on Morita equivalence between Frobenius algebras[81]. This concept of Morita equivalence between two Frobenius algebras differs from the concept of Morita equivalence between fusion categories introduced in Sec. B.

Algebraic Morita equivalence implies that the two Frobenius algebras have the same modules. According to the boundary-bulk correspondence, two string-net models with bimodule categories over two Morita-equivalent Frobenius algebras not only have the same anyon species but also exhibit the same relationships between each anyon species and the fundamental degrees of freedom of the models. As previously noted, the fusion category \mathscr{F} itself is the bimodule category over the trivial one-dimensional Frobenius algebra $\mathcal{A}_0 = \{1\}$.

Therefore, a symmetry transformation is a gauge symmetry if and only if its defining Frobenius algebra \mathcal{A} is algebraically Morita equivalent to the trivial Frobenius algebra \mathcal{A}_0 . This is the case for the Frobenius algebra $\mathcal{A} = \{1, \tau\}$ in the Fibonacci fusion category. Otherwise, it is a global symmetry, as in the case of \mathbb{Z}_2 toric code.

E Frobenius Algebra of Fibonacci Fusion Categories and Its Simple Bimodules

In this section, we list the categorical data of the Fibonacci fusion category. The Fibonacci fusion category has two simple objects, denoted as 1 and τ . The nonzero fusion rules are $\delta_{111} = \delta_{1\tau\tau} = \delta_{\tau\tau\tau} = 1$, and the quantum dimensions are

$$d_1 = 1, \qquad d_\tau = \phi = \frac{\sqrt{5}+1}{2}.$$

The nonzero 6j symbols are

$$G_{111}^{111} = 1, \qquad G_{\tau\tau\tau}^{111} = \frac{1}{\sqrt{\phi}}, \qquad G_{1\tau\tau}^{1\tau\tau} = G_{\tau\tau\tau}^{1\tau\tau} = \frac{1}{\phi}, \qquad G_{\tau\tau\tau}^{\tau\tau\tau} = -\frac{1}{\phi^2}.$$

The Fibonacci fusion category has a nontrivial Frobenius algebra \mathcal{A} , such that

$$L_{\mathcal{A}} = \{1, \tau\}, \qquad f_{111} = f_{1\tau\tau} = f_{\tau 1\tau} = f_{\tau\tau 1} = 1, \qquad f_{\tau\tau\tau} = -\frac{1}{\phi^{\frac{3}{4}}}.$$

There are two simple bimodules over \mathcal{A} , denoted as M_0 and M_1 , such that

$$L_{M_0} = \{1, \tau\},$$
$$[P_{M_0}]_{111}^{11} = [P_{M_0}]_{\tau\tau\tau}^{11} = 1, \qquad [P_{M_0}]_{11\tau}^{1\tau} = [P_{M_0}]_{\tau\tau1}^{1\tau} = 1, \qquad [P_{M_0}]_{\tau\tau\tau}^{1\tau} = \frac{1}{\phi^{\frac{3}{4}}},$$

$$\begin{split} [P_{M_0}]_{\tau 11}^{\tau 1} &= [P_{M_0}]_{1\tau \tau}^{\tau 1} = 1, \qquad [P_{M_0}]_{\tau \tau \tau}^{\tau 1} = \frac{1}{\phi^{\frac{3}{4}}}, \\ [P_{M_0}]_{\tau \tau 1}^{\tau \tau} &= [P_{M_0}]_{\tau 1\tau}^{\tau \tau} = 1, \qquad [P_{M_0}]_{1\tau \tau}^{\tau \tau} = [P_{M_0}]_{\tau \tau \tau}^{\tau \tau} = \frac{1}{\phi^{\frac{3}{4}}}, \qquad [P_{M_0}]_{\tau \tau \tau}^{\tau \tau} = \frac{1}{\phi^{\frac{3}{2}}} \,. \\ L_{M_1} &= \{1, \tau_1, \tau_2\}, \\ [P_{M_1}]_{11\tau}^{11} &= [P_{M_1}]_{\tau_0 \tau \tau_0}^{11} = [P_{M_1}]_{\tau_1 \tau \tau_1}^{11} = 1 \,, \\ [P_{M_1}]_{1\tau \tau_0}^{1\tau} &= [P_{M_1}]_{\tau_1 \tau \tau}^{1\tau} = [P_{M_1}]_{\tau_0 11}^{\tau 1} = [P_{M_1}]_{1\tau \tau_1}^{\tau 1} = \frac{1}{2\phi} + \frac{\sqrt{\phi}}{2}i \,, \\ [P_{M_1}]_{1\tau \tau_1}^{1\tau} &= [P_{M_1}]_{\tau_0 \tau \tau_1}^{1\tau} = [P_{M_1}]_{\tau_1 \tau \tau_1}^{\tau 1} = [P_{M_1}]_{1\tau \tau_0}^{\tau 1} = \frac{1}{2\phi} - \frac{\sqrt{\phi}}{2}i \,, \\ [P_{M_1}]_{\tau_0 \tau \tau_0}^{1\tau} &= [P_{M_1}]_{\tau_1 \tau \tau_1}^{1\tau} = [P_{M_1}]_{\tau_0 \tau \tau_0}^{\tau 1} = [P_{M_1}]_{\tau_1 \tau \tau_1}^{\tau 1} = -\frac{\sqrt{\phi}}{2\phi^2} \,, \end{split}$$

$$[P_{M_1}]_{\tau_0\tau\tau_1}^{1\tau} = [P_{M_1}]_{\tau_1\tau\tau_0}^{\tau_1} = -\frac{\sqrt[4]{\phi}}{2} - \frac{\phi^{\frac{3}{4}}}{2}i , \qquad [P_{M_1}]_{\tau_1\tau\tau_0}^{1\tau} = [P_{M_1}]_{\tau_0\tau\tau_1}^{\tau_1} = -\frac{\sqrt[4]{\phi}}{2} + \frac{\phi^{\frac{3}{4}}}{2}i ,$$

$$[P_{M_1}]_{1\tau 1}^{\tau\tau} = -\frac{1}{\phi} , \qquad [P_{M_1}]_{1\tau\tau_0}^{\tau\tau} = [P_{M_1}]_{1\tau\tau_1}^{\tau\tau} = [P_{M_1}]_{\tau_0\tau 1}^{\tau\tau} = [P_{M_1}]_{\tau_1\tau 1}^{\tau\tau} = -\frac{\sqrt[4]{\phi}}{\phi} ,$$

$$[P_{M_1}]_{\tau_0 1 \tau_0}^{\tau\tau} = -\frac{1}{2\phi} + \frac{i}{2\sqrt{\phi}} , \qquad [P_{M_1}]_{\tau_1 1 \tau_1}^{\tau\tau} = -\frac{1}{2\phi} - \frac{i}{2\sqrt{\phi}} , \qquad [P_{M_1}]_{\tau_0 1 \tau_1}^{\tau\tau} = [P_{M_1}]_{\tau_1 1 \tau_0}^{\tau\tau} = \frac{1}{2} ,$$

$$[P_{M_1}]_{\tau_0\tau\tau_1}^{\tau\tau} = [P_{M_1}]_{\tau_1\tau\tau_0}^{\tau\tau} = \frac{\sqrt{\phi}}{2\phi^2} , \qquad [P_{M_1}]_{\tau_0\tau\tau_0}^{\tau\tau} = -\frac{\sqrt{\phi}}{2\phi^3} - \frac{\phi}{2}i , \qquad [P_{M_1}]_{\tau_1\tau\tau_1}^{\tau\tau} = -\frac{\sqrt{\phi}}{2\phi^3} + \frac{\phi}{2}i .$$

The nonzero vertex coefficients are

$$\mathcal{V}_{M_0M_0M_0}^{111} = \mathcal{V}_{M_0M_0M_0}^{1\tau\tau} = \mathcal{V}_{M_0M_0M_0}^{\tau1\tau} = \mathcal{V}_{M_0M_0M_0}^{\tau\tau1} = 1, \qquad \mathcal{V}_{M_0M_0M_0}^{\tau\tau\tau} = \frac{1}{\phi^{\frac{3}{4}}},$$

$$\begin{split} \mathcal{V}_{M_{0}M_{1}M_{1}}^{111} &= \mathcal{V}_{M_{0}M_{1}M_{1}}^{1\tau_{1}\tau_{1}} = \mathcal{V}_{M_{0}M_{1}M_{1}}^{1\tau_{2}\tau_{2}} = 1, \qquad \mathcal{V}_{M_{0}M_{1}M_{1}}^{\tau\tau_{1}\tau_{1}} = \mathcal{V}_{M_{0}M_{1}M_{1}}^{\tau\tau_{2}\tau_{2}} = -\frac{1}{2\phi^{\frac{7}{4}}}, \\ \mathcal{V}_{M_{0}M_{1}M_{1}}^{\tau_{1}\tau_{1}} &= \mathcal{V}_{M_{0}M_{1}M_{1}}^{\tau\tau_{2}1} = \frac{1}{2\phi} - \frac{\sqrt{\phi}}{2}i, \qquad \mathcal{V}_{M_{0}M_{1}M_{1}}^{\tau_{1}\tau_{2}} = \mathcal{V}_{M_{0}M_{1}M_{1}}^{\tau\tau_{1}1} = \frac{1}{2\phi} + \frac{\sqrt{\phi}}{2}i, \\ \mathcal{V}_{M_{0}M_{1}M_{1}}^{\tau\tau_{1}\tau_{2}} &= -\frac{\sqrt[4]{\phi}}{2} + \frac{\phi^{\frac{3}{4}}}{2}i, \qquad \mathcal{V}_{M_{0}M_{1}M_{1}}^{\tau\tau_{2}\tau_{1}} = -\frac{\sqrt[4]{\phi}}{2} - \frac{\phi^{\frac{3}{4}}}{2}i, \end{split}$$

$$\begin{split} \mathcal{V}_{M_{1}M_{1}M_{1}}^{111} &= \frac{1}{\phi^{\frac{3}{4}}}, \qquad \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{1}\tau_{1}\tau_{1}} = \mathcal{V}_{M_{1}M_{1}M_{1}M_{1}}^{\tau_{2}\tau_{2}\tau_{2}\tau_{2}} = -\frac{1}{2}\sqrt{\frac{5}{\phi}}, \\ \mathcal{V}_{M_{1}M_{1}M_{1}}^{1\tau_{1}\tau_{1}} &= \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{1}\tau_{1}} = \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{2}\tau_{2}} = \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{2}\tau_{2}} = \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{2}\tau_{2}\tau_{2}} = -\frac{1}{2\phi^{\frac{7}{4}}}, \\ \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{1}\tau_{1}} &= \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{2}\tau_{1}\tau_{1}} = \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{2}\tau_{2}\tau_{2}} = \mathcal{V}_{M_{1}M_{1}M_{1}M_{1}}^{\tau_{2}\tau_{2}\tau_{1}} = \mathcal{V}_{M_{1}M_{1}M_{1}M_{1}}^{\tau_{2}\tau_{2}\tau_{1}} = \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{2}\tau_{2}\tau_{1}} = \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{2}\tau_{2}\tau_{1}} = \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{2}\tau_{2}\tau_{1}} = \frac{1}{2sqrt\phi}, \\ \mathcal{V}_{M_{1}M_{1}M_{1}}^{1\tau_{1}\tau_{2}} &= \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{1}\tau_{2}} = \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{1}\tau_{2}} = -\frac{\sqrt[4]{\phi}}{2} - \frac{\phi^{\frac{3}{4}}}{2}i, \\ \mathcal{V}_{M_{1}M_{1}M_{1}}^{1\tau_{2}\tau_{1}} &= \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{2}\tau_{1}} = \mathcal{V}_{M_{1}M_{1}M_{1}}^{\tau_{1}\tau_{2}} = -\frac{\sqrt[4]{\phi}}{2} + \frac{\phi^{\frac{3}{4}}}{2}i. \end{split}$$

The fusion category $\operatorname{Bimod}_{\operatorname{Fibo}}(\mathcal{A})$ is isomorphic to the Fibonacci fusion category by replacing 1 as M_0 , and τ as M_1 :

$$\begin{split} \delta_{M_0M_0M_0} &= \delta_{M_0M_1M_1} = \delta_{M_1M_1M_1} = 1, \qquad d_{M_0} = 1, \qquad d_{M_1} = \phi, \\ G_{M_0M_0M_0}^{M_0M_0M_0} &= 1, \quad G_{M_1M_1M_1}^{M_0M_0M_0} = \frac{1}{\sqrt{\phi}}, \quad G_{M_0M_1M_1}^{M_0M_1M_1} = G_{M_1M_1M_1}^{M_0M_1M_1} = \frac{1}{\phi}, \quad G_{M_1M_1M_1}^{M_1M_1M_1} = -\frac{1}{\phi^2} \end{split}$$

Bimodule M_1 has multiplicity $n_{\tau}^{M_1} = 2$, so we have to enlarge the Hilbert spaces on tails from 2-dimensional local Hilbert spaces spanned by 1 and τ to 3-dimensional local Hilbert spaces spanned by 1, τ_1 and τ_2 . To ensure the orthonormality, the basic state $|\tau\rangle$ labeled by the simple object τ in the trivial bimodule M_0 is a superposition state of two basic state $|\tau_1\rangle, |\tau_2\rangle$ labeled by the simple object τ in the nontrivial bimodule M_1 :

This τ charge on a tail is unique up to exchanging τ_1 and τ_2 labels.

References

- E. Dijkgraaf, Robbert and Witten, Communications in Mathematical Physics 429, 393 (1990).
- J. Fuchs, C. Schweigert, and A. Valentino, Communications in Mathematical Physics 1015, 981 (2014), arXiv:1307.3632.
- [3] A. Kapustin and N. Seiberg, Journal of High Energy Physics 2014, 1 (2014).
- [4] I. Cong, M. Cheng, and Z. Wang, Physical Review B 96 (2017), 10.1103/PhysRevB.96.195129, arXiv:1703.03564.
- [5] S. B. Bravyi and A. Y. Kitaev, arXiv preprint quant-ph/9811052 (1998).
- [6] A. Kitaev, Annals of Physics **303**, 2 (2003).
- [7] S. Beigi, P. W. Shor, and D. Whalen, Communications in Mathematical Physics **306**, 663 (2011).

- [8] Y. Hu, Y. Wan, and Y.-S. Wu, Physical Review B 87, 125114 (2013), arXiv:1211.3695.
- [9] A. Bullivant, Y. Hu, and Y. Wan, Physical Review B 96, 165138 (2017), arXiv:1706.03611.
- [10] I. Cong, M. Cheng, and Z. Wang, Communications in Mathematical Physics 2017 355:2 355, 645 (2017).
- [11] H. Wang, Y. Li, Y. Hu, and Y. Wan, Journal of High Energy Physics 2020, 30 (2020).
- [12] H. Wang, Y. Li, Y. Hu, and Y. Wan, Journal of High Energy Physics 2020, 30 (2020), arXiv:1910.13441.
- [13] Y. Hu and Y. Wan, Journal of High Energy Physics **2020**, 170 (2020).
- [14] V. G. Turaev and O. Y. Viro, Topology **31**, 865 (1992).
- [15] V. G. Turaev, Quantum Invariants of Knots and 3-manifolds (Walter de Gruyter, Berlin, 1994).
- [16] L. Bhardwaj, D. Gaiotto, and A. Kapustin, Journal of High Energy Physics 2017, 1 (2017).
- [17] X. Cui, Higher categories and topological quantum field theories (University of California, Santa Barbara, 2016).
- [18] M. Levin and X.-g. Wen, Physical Review B 71, 21 (2005), arXiv:0404617 [cond-mat].
- [19] L.-Y. Hung and Y. Wan, Physical Review B 86, 235132 (2012), arXiv:1207.6169.
- [20] Y. Hu, S. D. Stirling, and Y.-s. Wu, Physical Review B 85, 075107 (2011), arXiv:1105.5771.
- [21] A. Kitaev and L. Kong, Communications in Mathematical Physics 313, 351 (2012).
- [22] M. D. Schulz, S. Dusuel, K. P. Schmidt, and J. Vidal, Physical review letters 110, 147203 (2013).
- [23] T. Lan and X. G. Wen, Physical Review B - Condensed Matter and Materials Physics 90, 115119 (2014), arXiv:1311.1784.
- [24] C.-H. Lin and M. Levin, Physical Review B 89, 195130 (2014).
- [25] Y. Hu, Y. Wan, and Y.-s. Wu, Chinese Physics Letters **34**, 077103 (2017), arXiv:1706.00650
- [26] Y. Hu, N. Geer, and Y.-S. Wu, Physical Review B 97, 195154 (2018).
- [27] J. C. Bridgeman and D. Barter, Quantum 4, 277 (2020).
- [28] H. Wang, Y. Hu, and Y. Wan, Journal of High Energy Physics 2022, 1 (2022).
- [29] Y. Zhao, S. Huang, H. Wang, Y. Hu, and Y. Wan, SciPost Physics Core 6, 076 (2023).
- [30] A. Polishchuk and E. Zaslow, arXiv preprint math/9801119 (1998).
- [31] L.-Y. Hung and Y. Wan, International Journal of Modern Physics B 28, 1450172 (2014).
- [32] L.-Y. Hung and Y. Wan, Physical Review B 87, 195103 (2013).
- [33] A. Baratin and L. Freidel, Journal of Mathematical Physics 56 (2015).
- [34] D. Gaiotto, A. Kapustin, N. Seiberg, and B. Willett, Journal of High Energy Physics 2015, 172 (2015), arXiv:1412.5148v2.
- [35] L. Bhardwaj and Y. Tachikawa, Journal of High Energy Physics 2018, 1 (2018).
- [36] W. Ji and X.-G. Wen, Physical Review Research 2, 033417 (2020).

- [37] W. Ji and X.-G. Wen, Physical Review Research 2, 033417 (2020).
- [38] L. Kong, T. Lan, X.-G. Wen, Z.-H. Zhang, and H. Zheng, Physical Review Research 2, 043086 (2020).
- [39] T. Bartsch, M. Bullimore, and A. Grigoletto, arXiv preprint arXiv:2305.17165 (2023).
- [40] L. Bhardwaj, L. E. Bottini, S. Schäfer-Nameki, and A. Tiwari, SciPost Physics 14, 007 (2023).
- [41] L. Bhardwaj, S. Schäfer-Nameki, and J. Wu, Fortschritte der Physik 70, 2200143 (2022).
- [42] Y. Choi, H. T. Lam, and S.-H. Shao, Physical Review Letters **129**, 161601 (2022).
- [43] Y. Choi, H. T. Lam, and S.-H. Shao, Physical Review Letters 130, 131602 (2023).
- [44] Y. Choi, M. Forslund, H. T. Lam, and S.-H. Shao, Physical Review Letters 132, 121601 (2024).
- [45] T. Bartsch, M. Bullimore, A. E. Ferrari, and J. Pearson, SciPost Physics 17, 015 (2024).
- [46] W. Ji and X.-G. Wen, arXiv preprint arXiv:2106.02069 (2021).
- [47] J. L. Jacobsen and H. Saleur, Journal of High Energy Physics 2023, 1 (2023).
- [48] C. Fechisin, N. Tantivasadakarn, and V. V. Albert, arXiv preprint arXiv:2312.09272 (2023).
- [49] W. Xi, T. Lan, L. Wang, C. Wang, and W.-Q. Chen, arXiv preprint arXiv:2312.15947 (2023).
- [50] Y. Li and M. Litvinov, arXiv preprint arXiv:2405.15951 (2024).
- [51] Y. Choi, Y. Sanghavi, S.-H. Shao, and Y. Zheng, arXiv preprint arXiv:2405.13105 (2024).
- [52] A. P. Mana, Y. Li, H. Sukeno, and T.-C. Wei, arXiv preprint arXiv:2402.09520 (2024).
- [53] K. Inamura and K. Ohmori, SciPost Physics 16, 143 (2024).
- [54] J. Kaidi, G. Zafrir, and Y. Zheng, Journal of High Energy Physics 2022, 1 (2022).
- [55] F. Apruzzi, I. Bah, F. Bonetti, and S. Schäfer-Nameki, Physical review letters 130, 121601 (2023).
- [56] M. van Beest, D. S. Gould, S. Schäfer-Nameki, and Y.-N. Wang, Journal of High Energy Physics 2023, 1 (2023).
- [57] J. Chen, W. Cui, B. Haghighat, and Y.-N. Wang, Journal of High Energy Physics 2023, 1 (2023).
- [58] O. Sela, Physical Review Letters 132, 201601 (2024).
- [59] F. A. Bais and J. Slingerland, Physical Review B 79, 045316 (2009).
- [60] I. S. Eliëns, J. C. Romers, and F. A. Bais, Physical Review B 90, 195130 (2014), arXiv:1310.6001.
- [61] Y. Wan and C. Wang, Journal of High Energy Physics 2017, 172 (2017).
- [62] F. Burnell, Annual Review of Condensed Matter Physics 9, 307 (2018).
- [63] Y. Hu, Z. Huang, L.-y. Hung, and Y. Wan, Journal of High Energy Physics 2022, 26 (2022), arXiv:2109.06145v1.
- [64] C.-H. Lin and F. J. Burnell, arXiv preprint arXiv:2303.07291 (2023).
- [65] R. B. Laughlin, Physical Review Letters **50**, 1395 (1983).

- [66] R. Tao and Y.-S. Wu, Physical Review B **30**, 1097 (1984).
- [67] J. Jain, Physical Review B 40, 8079 (1989).
- [68] X.-G. Wen, Physical Review B 41, 12838 (1990).
- [69] X.-g. Wen and Q. Niu, Physical Review B 41, 9377 (1990).
- [70] E. Tang, J.-W. Mei, and X.-G. Wen, Physical Review Letters 106, 236802 (2011).
- [71] D. J. Clarke, J. Alicea, and K. Shtengel, Nature Communications 2013 4:1 4, 1 (2013), arXiv:1204.5479.
- [72] B. Yang, N. Jiang, X. Wan, J. Wang, and Z.-X. Hu, Physical Review B 99, 161108 (2019).
- [73] K. Iguchi, Physical Review Letters 78, 3233 (1997).
- [74] S. Guruswamy and K. Schoutens, Nuclear Physics B 556, 530 (1999), arXiv:9903045 [cond-mat].
- [75] M. Barkeshli, C.-M. Jian, and X.-L. Qi, Physical Review B 87, 045130 (2013).
- [76] Y. Hu, S. D. Stirling, and Y.-s. Wu, Physical Review B 89, 115133 (2014), arXiv:1303.1586.
- [77] Y. Li, H. Wang, Y. Hu, and Y. Wan, Journal of High Energy Physics 2019, 1 (2019).
- [78] Y. Li, H. Wang, Y. Hu, and Y. Wan, Journal of High Energy Physics 2019, 78 (2019).
- [79] Y.-a. Fan, Y. Li, Y. Hu, Y. Li, X. Long, H. Liu, X. Yang, X. Nie, J. Li, T. Xin, et al., The Innovation 4 (2023), 10.1016/j.xinn.2023.100480.
- [80] Y. Zhao, H. Wang, Y. Hu, and Y. Wan, Journal of High Energy Physics 2024, 1 (2024).
- [81] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik, *Tensor categories*, Vol. 205 (American Mathematical Soc., 2016).
- [82] O. Buerschaper, M. Christandl, L. Kong, and M. Aguado, Nuclear Physics B 876, 619 (2013), arXiv:1006.5823.
- [83] C. L. Douglas and D. J. Reutter, arXiv preprint arXiv:1812.11933 (2018).